

Ins and outs of Ca^{2+} transport by acidic organelles and cell migration

Sandip Patel

Department of Cell and Developmental Biology, University College London, London, UK

ABSTRACT

Much contemporary evidence underscores the pathophysiological importance of Ca^{2+} handling by acidic organelles such as lysosomes. Whereas our knowledge of how Ca^{2+} is released from these acidic Ca^{2+} stores (the 'outs') is advancing, we know relatively little about how Ca^{2+} uptake is effected (the 'ins'). Here I highlight new work identifying animal $\text{Ca}^{2+}/\text{H}^{+}$ (CAX) exchangers that localize to acidic organelles, mediate Ca^{2+} uptake and regulate cell migration *in vivo*. Continued molecular definition of the acidic Ca^{2+} store toolkit provides new insight into Ca^{2+} -dependent function.

ARTICLE HISTORY

Received 8 May 2017
Accepted 15 May 2017

KEYWORDS

two-pore channels; Acidic Ca^{2+} stores; CAX; cell migration; lysosomes; NAADP; neural crest

The so called 'acidic Ca^{2+} stores' are a morphologically eclectic cohort of Ca^{2+} - and H^{+} -replete organelles that, in addition to fulfilling their canonical functions, also serve as mobilizable stores of Ca^{2+} .^{1,2} A well-studied example is the lysosome, acting as both a macronutrient recycling center and as a target Ca^{2+} store for the second messenger NAADP.³ The list of Ca^{2+} -permeable channels expressed on lysosomes—the 'outs'—is steadily increasing. It includes the archetypal TRP mucolipins,⁴ NAADP-regulated two-pore channels (TPCs)⁵ and other ion channels thought generally to reside on the plasma membrane (TRPM2,⁶ P2X4,⁷ Cav2.1⁸ and TRPA1⁹). These channels mediate a range of cellular processes including triggering of global Ca^{2+} changes and various trafficking events centered around endolysosomal fusion/fission.^{1,2,10,11} Given emerging links to disease,^{12,13} molecular and functional interest in Ca^{2+} release from acidic organelles is growing. But lagging behind (at least for the animal kingdom) is our understanding of the 'ins' i.e. how Ca^{2+} is taken up by acidic Ca^{2+} stores and its physiologic relevance.

Vacuoles are key acidic Ca^{2+} stores in plants, yeast and protists that are often likened to lysosomes of animal cells.¹ Vacuolar Ca^{2+} uptake is mediated by molecularly defined $\text{Ca}^{2+}/\text{H}^{+}$ exchangers (CAXs) and P-type Ca^{2+} ATPases.¹⁴ CAXs use the substantial proton gradient to drive the antiport of Ca^{2+} at high capacity in to the

lumen. Knockout of CAX genes, for example in *Arabidopsis*, leads to a significant reduction in vacuolar Ca^{2+} loading, an associated increase in apoplastic (cell wall) Ca^{2+} concentration and reduced gas exchange, growth and fitness.^{15,16} The existence of analogous CAXs have been proposed in animals based on early biochemical work in sea urchin eggs.¹⁷ This study showed blockade of Ca^{2+} uptake into target NAADP-sensitive stores by agents that collapsed proton gradients such as bafilomycin A1 (a V-type H^{+} ATPase inhibitor), but not by vanadate (a P-type Ca^{2+} ATPase inhibitor)¹⁷ consistent with secondary active transport of Ca^{2+} . Indeed, such a proposal has received widespread support where Ca^{2+} release by NAADP or NAADP-forming agonists is consistently blocked by bafilomycin A1 across a multitude of cells.^{18–20} However, the molecular identity of animal CAXs has been somewhat of a mystery.

Recent work identified several putative animal CAXs through searches of the ever-increasing genomic sequences now at our disposal.²¹ These genes were found in protostomes such as gastropods and polychaetes, and deuterostomes from basal animal species (reassuringly including the sea urchin, an echinoderm) through to amphibians and non-placental mammals. CAXs were cloned from the purple sea urchin and the African clawed frog. The latter was heterologously expressed in yeast where it was shown to possess Ca^{2+} - H^{+} activity

CONTACT Sandip Patel  patel.s@ucl.ac.uk  Department of Cell and Developmental Biology, University College London, Gower Street, London WC1E 6BT, UK.

Addendum to: Melchionda M, Pittman JK, Mayor R, Patel S. $\text{Ca}^{2+}/\text{H}^{+}$ exchange by acidic organelles regulates cell migration *in vivo*. *J Cell Biol* 2016; 212:803-13; <https://doi.org/10.1083/jcb.201510019>

© 2018 The Author. Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.




Figure 1. Ca²⁺/H⁺ exchange by acidic organelles regulates cell migration *in vivo*. (A) Structural model of *Xenopus laevis* CAX based on crystal structure of VCX1. (B) Transport of H⁺ into acidic Ca²⁺ stores by a V-type ATPase (bottom) and its exchange for Ca²⁺ by CAX (top). (C) Location of acidic Ca²⁺ stores in a migrating neural crest cell. (D) Migration of neural crest streams in a developing frog embryo. Direction of migration in C-D is depicted by the arrows.

dependent upon on a conserved acidic residue within the transport core and the N-terminal 316 amino acids. Frog CAX localized to lysosomes when expressed in mammalian cells and tempered receptor-mediated Ca²⁺ signals²¹ consistent with an analogous role for CAX in clearing Ca²⁺ elevations in response to osmotic stress in yeast.²² In both frog²¹ and zebrafish²³ embryos, transcripts for endogenous CAX were enriched in the neural crest. Within these highly migratory embryonic cells, CAX localized to yolk granules and small acidic vesicles (likely lysosomes).²¹ The former finding is interesting as it was these very same lysosome-like organelles that were originally identified as NAADP-sensitive Ca²⁺ stores in sea urchin eggs.¹⁷ Live cell imaging indicated that the smaller CAX-positive vesicles were highly mobile, often populating the leading edge of migrating neural crest cells.²¹ Indeed, knockdown of CAX disrupted multiple cell migration processes in neural crest explants including focal adhesion formation, cell spreading and chemotaxis. Importantly, migration of the neural crest was also disrupted *in vivo*. Thus, this study²¹ significantly furthers our understanding of how Ca²⁺ uptake by acidic Ca²⁺ stores impacts cellular function in animals and offers new opportunities for probing organellar Ca²⁺/H⁺ exchange by molecular means (Fig. 1).

But what about Ca²⁺ uptake into acidic Ca²⁺ stores in human cells given the apparent lack of CAX homologues in placental mammals? It is formally possible (although in my opinion unlikely) that CAXs within our lineage have substantially diverged in sequence to make them invisible to current homology searches. Alternatively, Ca²⁺ uptake could be mediated through a novel protein with Ca²⁺-H⁺ exchange activity,²⁴ or even through combinations of proteins (eg coupled Na⁺/H⁺ and Na⁺/Ca²⁺ exchange).²⁵ A recent study has suggested that lysosomal Ca²⁺ filling is achieved by neighboring IP₃-sensitive Ca²⁺ release channels on

the ER but via a route that does not require a proton gradient.²⁶ Certainly, identified membrane contact sites between the ER and lysosomes would facilitate this^{27,28} similar to their proposed role in amplifying lysosomal Ca²⁺ signals by the ER.²⁹ But the reported insensitivity of lysosomal Ca²⁺ uptake to bafilomycin A1 (assessed by monitoring Ca²⁺ signals evoked by the TRP mucolipin activator MLSA1),²⁶ is difficult to reconcile with previous studies; the findings are at odds not only with the aforementioned bafilomycin A1-sensitivity of NAADP action but also work by the same authors³⁰ and others³¹ showing inhibition of MLSA1-evoked Ca²⁺ signals upon V-type H⁺ ATPase blockade. Clearly further work is required to clarify the mechanisms mediating lysosomal Ca²⁺ uptake in mammalian cells, the importance of which is underscored by functional evidence linking signaling through acidic Ca²⁺ stores to cell migration.^{32,33}

Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

Acknowledgments

I thank Len Dahl, Bethan S. Kilpatrick and Christopher J. Penny for useful discussion.

Funding

Supported by BBSRC grants BB/K000942/1 and BB/N01524X/1.

References

- [1] Patel S, Docampo R. Acidic calcium stores open for business: expanding the potential for intracellular Ca²⁺ signaling. *Trends Cell Biol* 2010; 20:277-86;

- PMID:20303271; <https://doi.org/10.1016/j.tcb.2010.08.009>
- [2] Patel S, Cai X. Evolution of acid Ca^{2+} stores and their resident Ca^{2+} -permeable channels. *Cell Calcium* 2015; 57:222-30; PMID:25591931; <https://doi.org/10.1016/j.ceca.2014.12.005>
- [3] Galione A. A primer of NAADP-mediated Ca signalling: From sea urchin eggs to mammalian cells. *Cell Calcium* 2015; 58:27-47; PMID:25449298; <https://doi.org/10.1016/j.ceca.2014.09.010>
- [4] Venkatachalam K, Wong CO, Zhu MX. The role of TRPMLs in endolysosomal trafficking and function. *Cell Calcium* 2015; 58:48-56; PMID:25465891; <https://doi.org/10.1016/j.ceca.2014.10.008>
- [5] Patel S. Function and dysfunction of two-pore channels. *Sci Signal* 2015; 8:re7; PMID:26152696; <https://doi.org/10.1126/scisignal.aab3314>
- [6] Lange I, Yamamoto S, Partida-Sanchez S, Mori Y, Fleig A, Penner R. TRPM2 functions as a lysosomal Ca^{2+} -release channel in beta cells. *Sci Signal* 2009; 2:ra23. PMID:19454650; <https://doi.org/10.1126/scisignal.2000278>
- [7] Qureshi OS, Paramasivam A, Yu JC, Murrell-Lagnado RD. Regulation of $\text{P}2 \times 4$ receptors by lysosomal targeting, glycan protection and exocytosis. *J Cell Sci* 2007; 120:3838-49; PMID:17940064; <https://doi.org/10.1242/jcs.010348>
- [8] Tian X, Gala U, Zhang Y, Shang W, Nagarkar Jaiswal S, di Ronza A, Jaiswal M, Yamamoto S, Sandoval H, Duraine L, et al. A voltage-gated calcium channel regulates lysosomal fusion with endosomes and autophagosomes and is required for neuronal homeostasis. *PLoS Biol* 2015; 13:e1002103; PMID:25811491; <https://doi.org/10.1371/journal.pbio.1002103>
- [9] Shang S, Zhu F, Liu B, Chai Z, Wu Q, Hu M, Wang Y, Huang R, Zhang X, Wu X, et al. Intracellular TRPA1 mediates Ca^{2+} release from lysosomes in dorsal root ganglion neurons. *J Cell Biol* 2016; 215:369-81; PMID:27799370; <https://doi.org/10.1083/jcb.201603081>
- [10] Morgan AJ, Platt FM, Lloyd-Evans E, Galione A. Molecular mechanisms of endolysosomal Ca^{2+} signalling in health and disease. *Biochem J* 2011; 439:349-74; PMID:21992097; <https://doi.org/10.1042/BJ20110949>
- [11] Xu H, Ren D. Lysosomal physiology. *Annu Rev Physiol* 2015; 77:57-80; PMID:25668017; <https://doi.org/10.1146/annurev-physiol-021014-071649>
- [12] Hockey LN, Kilpatrick BS, Eden ER, Lin-Moshier Y, Brailoiu GC, Brailoiu E, Futter CE, Schapira AH, Marchant JS, Patel S. Dysregulation of lysosomal morphology by pathogenic LRRK2 is corrected by TPC2 inhibition. *J Cell Sci* 2015; 128:232-8; PMID:25416817; <https://doi.org/10.1242/jcs.164152>
- [13] Patel S. Deviant lysosomal Ca^{2+} signalling in neurodegeneration. An introduction. *Messenger* 2016; PMID:28529827; <https://doi.org/10.1166/msr.2016.1053>
- [14] Pittman JK. Vacuolar Ca^{2+} uptake. *Cell Calcium* 2011; 50:139-46; PMID:21310481; <https://doi.org/10.1016/j.ceca.2011.01.004>
- [15] Cheng NH, Pittman JK, Shigaki T, Lachmansingh J, LeClere S, Lahner B, Salt DE, Hirschi KD. Functional association of Arabidopsis CAX1 and CAX3 is required for normal growth and ion homeostasis. *Plant Physiol* 2005; 138:2048-60; PMID:16055687; <https://doi.org/10.1104/pp.105.061218>
- [16] Conn SJ, Gilliham M, Athman A, Schreiber AW, Baumann U, Moller I, Cheng NH, Stancombe MA, Hirschi KD, Webb AA, et al. Cell-specific vacuolar calcium storage mediated by CAX1 regulates apoplastic calcium concentration, gas exchange, and plant productivity in Arabidopsis. *Plant Cell* 2011; 23:240-57; PMID:21258004; <https://doi.org/10.1105/tpc.109.07.2769>
- [17] Churchill GC, Okada Y, Thomas JM, Genazzani AA, Patel S, Galione A. NAADP mobilizes Ca^{2+} from reserve granules, lysosome-related organelles, in sea urchin eggs. *Cell* 2002; 111:703-8; PMID:12464181; [https://doi.org/10.1016/S0092-8674\(02\)01082-6](https://doi.org/10.1016/S0092-8674(02)01082-6)
- [18] Yamasaki M, Masgrau R, Morgan AJ, Churchill GC, Patel S, Ashcroft SJ, Galione A. Organelle selection determines agonist-specific Ca^{2+} signals in pancreatic acinar and beta cells. *J Biol Chem* 2004; 279:7234-40; PMID:14660554; <https://doi.org/10.1074/jbc.M311088200>
- [19] Pandey V, Chuang CC, Lewis AM, Aley PK, Brailoiu E, Dun NJ, Churchill GC, Patel S. Recruitment of NAADP-sensitive acidic Ca^{2+} stores by glutamate. *Biochem J* 2009; 422:503-12; PMID:19548879; <https://doi.org/10.1042/BJ20090194>
- [20] Galione A, Morgan AJ, Arredouani A, Davis LC, Rietdorf K, Ruas M, Parrington J. NAADP as an intracellular messenger regulating lysosomal calcium-release channels. *Biochem Soc Trans* 2010; 38:1424-31; PMID:21118101; <https://doi.org/10.1042/BST0381424>
- [21] Melchionda M, Pittman JK, Mayor R, Patel S. $\text{Ca}^{2+}/\text{H}^{+}$ exchange by acidic organelles regulates cell migration *in vivo*. *J Cell Biol* 2016; 212:803-13; PMID:27002171; <https://doi.org/10.1083/jcb.201510019>
- [22] Denis V, Cyert MS. Internal Ca^{2+} release in yeast is triggered by hypertonic shock and mediated by a TRP channel homologue. *J Cell Biol* 2002; 156:29-34; PMID:11781332; <https://doi.org/10.1083/jcb.200111004>
- [23] Manohar M, Mei H, Franklin AJ, Sweet EM, Shigaki T, Riley BB, Macdiarmid CW, Hirschi K. Zebrafish (*Danio rerio*) endomembrane antiporter similar to a yeast cation/ H^{+} transporter is required for neural crest development. *Biochemistry* 2010; 49:6557-66; PMID:20578725; <https://doi.org/10.1021/bi100362k>
- [24] Demaegd D, Foulquier F, Colinet AS, Gremillon L, Legrand D, Mariot P, Peiter E, Van Schaftingen E, Matthijs G, Morsomme P. Newly characterized Golgi-localized family of proteins is involved in calcium and pH homeostasis in yeast and human cells. *Proc Natl Acad Sci USA* 2013; 110:6859-64; PMID:23569283; <https://doi.org/10.1073/pnas.1219871110>
- [25] Lamason RL, Mohideen MA, Mest JR, Wong AC, Norton HL, Aros MC, Jurynek MJ, Mao X, Humphreville VR, Humbert JE, et al. SLC24A5, a putative cation exchanger, affects pigmentation in zebrafish and humans. *Science* 2005; 310:1782-86; PMID:16357253; <https://doi.org/10.1126/science.1116238>
- [26] Garrity AG, et al. The endoplasmic reticulum, not the pH gradient, drives calcium refilling of lysosomes. *eLife* 2016; 5; PMID:27213518; <https://doi.org/10.7554/eLife.15887>

- [27] Kilpatrick BS, Eden ER, Schapira AH, Futter CE, Patel S. Direct mobilisation of lysosomal Ca^{2+} triggers complex Ca^{2+} signals. *J Cell Sci* 2013; 126:60-6; PMID:23108667; <https://doi.org/10.1242/jcs.118836>
- [28] Kilpatrick BS, Eden ER, Hockey LN, Yates E, Futter CE, Patel S. An endosomal NAADP-sensitive two-pore Ca^{2+} channel regulates ER-endosome membrane contact sites to control growth factor signaling. *Cell Rep* 2017; 18:1636-45; PMID:28199837; <https://doi.org/10.1016/j.celrep.2017.01.052>
- [29] Penny CJ, Kilpatrick BS, Han JM, Sneyd J, Patel S. A computational model of lysosome-ER Ca^{2+} microdomains. *J Cell Sci* 2014; 127:2934-43; PMID:24706947; <https://doi.org/10.1242/jcs.149047>
- [30] Shen D, Wang X, Li X, Zhang X, Yao Z, Dibble S, Dong XP, Yu T, Lieberman AP, Showalter HD, et al. Lipid storage disorders block lysosomal trafficking by inhibiting a TRP channel and lysosomal calcium release. *Nat Commun* 2012; 3:731; PMID:22415822; <https://doi.org/10.1038/ncomms1735>
- [31] Cao Q, Zhong XZ, Zou Y, Zhang Z, Toro L, Dong XP. BK channels alleviate lysosomal storage diseases by providing positive feedback regulation of lysosomal Ca^{2+} release. *Dev Cell* 2015; 33:427-41; PMID:25982675; <https://doi.org/10.1016/j.devcel.2015.04.010>
- [32] Sumoza-Toledo A, Lange I, Cortado H, Bhagat H, Mori Y, Fleig A, Penner R, Partida-Sánchez S. Dendritic cell maturation and chemotaxis is regulated by TRPM2-mediated lysosomal Ca^{2+} release. *FASEB J* 2011; 25:3529-42; PMID:21753080; <https://doi.org/10.1096/fj.10-178483>
- [33] Nguyen ON, Grimm C, Schneider LS, Chao YK, Atzberger C, Bartel K, Watermann A, Ulrich M, Mayr D, Wahl-Schott C. Two-pore channel function is crucial for the migration of invasive cancer cells. *Cancer Res* 2017; 77:1427-38.