

POSTER PRESENTATION

Open Access

Twist measurement of the left ventricle through radial tagging

Razieh Kaveh¹, Abbas N Moghaddam^{1,2*}, Sarah N Khan², J Paul Finn²

From 16th Annual SCMR Scientific Sessions
San Francisco, CA, USA. 31 January - 3 February 2013

Background

Assessment of left ventricular twist is of fundamental importance in evaluating cardiac pump function. The recently developed radial tagging method provides new insights into cardiac rotational motion and has facilitated quantifying LV twist. Herein, we suggest a method to simply obtain twist of radially tagged LV, directly in the image domain.


Methods

Short axis radially tagged images were acquired at apical and basal levels in 6 healthy subjects.

Contouring LV epicardial and endocardial borders is the only user interaction in the twist extraction process. Applying a proper threshold algorithm over the masked images results in binary images containing easily detectable tags. We tracked the mean position of tags over the cardiac cycle and utilized their spatial coordinates to calculate rotation by the deformation gradient tensor method. By calculating the rotation angle of apical and basal LV sections, we extract twist, as the difference of apical and basal rotation.

Results

The mean twist extracted from 6 healthy subjects is plotted in Figure 1. The values of peak twist and peak twist time are shown in Table 1. Compared to global circumferential strain, obtained with the method reported in



[1], peak twist and peak systolic strain occur at the same time for all 6 datasets. The results are consistent with those reported in the literature.

Conclusions

The proposed method takes advantages of novel radial tagging and is fast and reliable to quantify LV twist, providing additional insight into cardiac contractile function. As derived quantities, other related indices such as torsion, twisting rate and untwisting rate can be calculated.

Table 1 The values of peak twist and peak twist time

	Volunteer1	Volunteer2	Volunteer3	Volunteer4	Volunteer5	Volunteer6
Peak twist time (percentage of cardiac cycle)	%44	%43.8	%46.7	%40	%45	%50
Peak twist value (degree)	8.8	11.9	13.2	17.3	14.9	16.2

¹Biomedical Engineering, Tehran Polytechnic, Tehran, Islamic Republic of Iran
Full list of author information is available at the end of the article

Funding

Images were obtained through DCVI section at UCLA.

Author details

¹Biomedical Engineering, Tehran Polytechnic, Tehran, Islamic Republic of Iran.

²Radiological Science, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA.

Published: 30 January 2013

doi:10.1186/1532-429X-15-S1-E48

Cite this article as: Kaveh *et al.*: Twist measurement of the left ventricle through radial tagging. *Journal of Cardiovascular Magnetic Resonance* 2013 15(Suppl 1):E48.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

