

Poster presentation

Open Access

PI6-21. Altered T cell homeostasis and activation in HIV-1 elite suppressors with low CD4⁺ T-cell counts

KC Groves^{*1}, D Bibby², A Isaksen², J Deayton³, J Anderson⁴, C Orkin², AJ Stagg³ and Á McKnight³

Address: ¹Institute of Cell and Molecular Science, Barts and The London School of Medicine and Dentistry, London, UK, ²Barts and the London NHS Trust, London, UK, ³ICMS, Barts and The London School of Medicine and Dentistry, London, UK and ⁴Homerton University Hospital NHS Foundation Trust, London, UK

* Corresponding author

from AIDS Vaccine 2009
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, 6(Suppl 3):P250 doi:10.1186/1742-4690-6-S3-P250

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P250>

© 2009 Groves et al; licensee BioMed Central Ltd.

Background

HIV-1 elite-suppressors (ES) maintain low viral-load (VL) without therapy and classically maintain normal CD4⁺ counts. We have identified a rare subset with falling CD4⁺ counts (discord-ES). Disease progression in these patients may result from a viral influence on T-cell homeostasis and activation, rather than a direct consequence of virus-replication. Therefore, we compared T-cell homeostasis and activation in discord-ES and typical-ES.

Methods

Fresh whole blood samples were obtained from 8 discord-ES (VL < 2000 copies/ml > 1 year, CD4⁺ count < 450 × 10⁶/L), 12 typical-ES (VL < 2000 copies/ml > 1 year, CD4⁺ count > 450 × 10⁶/L), 5 progressors (VL > 5000 copies/ml, CD4⁺ count < 450 × 10⁶/L) and 5 uninfected individuals. Using multiparameter flow-cytometry we quantified naïve and memory T-cell subsets based on expression of CD45RA/RO and CD62L, and assessed activation based on co-expression of CD38 and HLA-DR.

Results

Blood from discord-ES contained fewer naïve (CD45RA⁺CD62L⁺) CD4⁺ T-cells than blood from typical-ES (median 80 [IQR 52–121] versus 277 [218–401]; *p* = 0.001). This lower number of naïve T-cells represented a lower proportion of total CD4⁺ T-cells. Numbers of naïve CD4⁺ T-cells in discord-ES did not differ significantly from those in progressors and were reduced (*p* = 0.002) com-

pared with uninfected controls. A similar pattern was observed with regard to numbers of CD4⁺ central-memory (CD45RO⁺CD62L⁺) and effector-memory (CD45RO⁺CD62L⁻) populations. Both activated and resting naïve CD4⁺ T-cells were reduced in number in discord-ES, but activated cells constituted an increased proportion of the naïve CD4⁺ pool compared with typical-ES (*p* = 0.003).

Conclusion

CD4⁺ T-cell subsets and their activation in discord-ES are distinct from those in typical-ES and indistinguishable from those in progressors, suggesting that CD4⁺ T-cell decline is more closely associated with perturbations in T-cell activation than with VL. The increased proportion of activated naïve CD4⁺ T-cells against a backdrop of declining numbers of these cells in discord-ES may suggest that activation contributes to progressive cell-loss despite only low-level viral-replication. However, impaired T-cell production or sequestration to non-blood compartments may also contribute.