

POSTER PRESENTATION

Open Access

Radiological characteristics of screen-detected lung cancers: predictive for histological subtype?

MA Heuvelmans^{1*}, E Salters¹, HJM Groen², P De Jong³, WPTM Mali³, M Oudkerk¹, R Vliegenthart¹

From International Cancer Imaging Society (ICIS) 14th Annual Teaching Course
Heidelberg, Germany. 9-11 October 2014

Aim

To evaluate CT-morphological features of lung cancers detected in a CT lung cancer screening trial, and to determine the correlation between CT-morphological features and the histopathological diagnosis of screen-detected cancers.

Methods

197 solid lung cancers (192 participants) detected in all four screening rounds of the Dutch-Belgian randomized lung cancer screening trial (NELSON) were included. CT-morphological features included nodule shape, margin, location, volume, and volume-doubling time (VDT). Based on histopathology, cancers were divided into four groups: adenocarcinoma (N=114), squamous cell carcinoma (N=37), large cell carcinoma (N=28) and neuro-endocrine cancers (N=18). Data were analyzed using ANOVA, Chi-square and Fischer's exact test.

Results

Mean participant age was 61.3 years (95%-confidence interval [CI]: 60.5-62.2), and 160/192 (83.3%) were male. In all four histopathologic groups, the majority of cancers had a spherical nodule shape (70.6-95.8%). Margins of malignant nodules were most often lobulated (39.3-45.9%) and spiculated (22.2-35.7%), without statistically significant difference between histopathological groups. Most cancers (63.5%) were located in the upper lobes, adenocarcinomas significantly more often (71.1%) than other types of cancers ($p=0.004$). Adenocarcinomas had a higher mean VDT than large cell carcinomas (214.8 days, 95%-CI: 186.2-243.4 days vs 96.8 days, 95%-CI: 15.8-177.9 days [$p<0.05$]).

VDTs of other histopathological subgroups did not differ significantly.

Conclusion

Only VDT and location in the upper versus lower lobes are associated with histopathological diagnosis of screen-detected lung cancers. No discrimination can be made between different histopathologic cancer types based on CT-morphological features alone.

Authors' details

¹Center for Medical Imaging – North East Netherlands, Department of Radiology, University of Groningen, University Medical Center Groningen, Groningen, Netherlands. ²Department of Pulmonology, University of Groningen, University Medical Center Groningen, Groningen, Netherlands.

³Department of Radiology, University Medical Center Utrecht, Utrecht, Netherlands.

Published: 9 October 2014

doi:10.1186/1470-7330-14-S1-P20

Cite this article as: Heuvelmans et al.: Radiological characteristics of screen-detected lung cancers: predictive for histological subtype? *Cancer Imaging* 2014 **14**(Suppl 1):P20.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Center for Medical Imaging – North East Netherlands, Department of Radiology, University of Groningen, University Medical Center Groningen, Groningen, Netherlands

Full list of author information is available at the end of the article