



Cite this: *RSC Adv.*, 2018, 8, 30957

Screening of potential IL-tolerant cellulases and their efficient saccharification of IL-pretreated lignocelluloses†

Yi-Xin Sun,^a Bing-Bing Shen,^a Hui-Ying Han,^a Yuan Lu,^a Bi-Xian Zhang,^b Yun-Fei Gao,^b Bao-Zhong Hu^{*c} and Xiao-Mei Hu^{*a}

Lignocellulosic biomass as one of the most abundant and renewable resources has great potential for biofuel production. The complete conversion of biomass to biofuel is achieved through the effective pretreatment process and the following enzyme saccharification. Ionic liquids (ILs) are considered as a green solvent for lignocellulose pretreatment. However, ILs exhibit an inhibitory effect on cellulase activity, leading to a subsequent decrease in the efficiency of saccharification. The screening of new potential IL-tolerant cellulases is important. In the current study, a fungal strain with a relatively high cellulase production was isolated and identified as *Penicillium oxalicum* HC6. The culture conditions were optimized using corn stover and peptone as the carbon source and nitrogen source at pH 4.0 and 30 °C with an inoculation size of 2% (v/v) for 8 days. It was found that *P. oxalicum* HC6 exhibited potential salt tolerance with the increase of the enzyme production at a salt concentration of 5.0% (w/v). In addition, high enzyme activities were obtained at pH 4.0–6.0 and 50–65 °C. The crude enzyme from *P. oxalicum* HC6 with good thermal stability was also stable in the presence of salt and ILs. Good yields of reducing sugar were obtained by the crude enzyme from *P. oxalicum* HC6 after the saccharification of corn stover that was pretreated by ILs. *P. oxalicum* HC6 with potentially salt-tolerant and IL-tolerant enzymes has great potential application in the enzymatic saccharification of lignocellulose.

Received 5th July 2018
Accepted 22nd August 2018

DOI: 10.1039/c8ra05729j

rsc.li/rsc-advances

1. Introduction

Lignocellulosic biomass can be considered as the most abundant polymer found on Earth. A substantial amount of lignocellulosic materials such as agricultural residues, agricultural by-products, and woody biomass are produced worldwide and their annual production rates are around 200 billion.¹ Among a variety of lignocellulosic biomasses, agricultural stover is essentially potential raw material for the production of high value products such as chemical reagents, biosurfactants and biofuel.^{2–4} Lignocellulosic biomass is mainly composed of three main biopolymers including cellulose, hemicellulose and lignin, which form a complex and recalcitrant structure.⁵

Generally, cellulose and hemicellulose, which comprise nearly two-thirds of lignocellulosic biomass, are preferred as the potential feed-stock for biofuel production.⁶ A complete cellulase system is required for the synergic action to further convert cellulose into monomeric sugars for the effective production of

biofuels. These cellulases comprise three major groups of enzymes: (1) endoglucanases (EC 3.2.1.4), which attack regions of low crystallinity in cellulose fibers, creating free chain-ends, (2) cellobiohydrolases (EC 3.2.1.91), which further degrade the molecule by cleaving cellobiose from the free-chain ends, and (3) β -glucosidases (EC 3.2.1.21) which hydrolyze cellobiose to produce glucose.⁷

Numbers of microorganisms are known as efficient cellulase producers and fungi are considered as potential sources for cellulase production, because of the large embedded accumulation of mycelium which is easy to separate and hence it helps to reduce the cost of separation process from the fermentation medium.⁸ *Aspergillus* sp. and *Trichoderma* sp. are known as the model fungi among a variety of fungal organisms.⁹ *Trichoderma reesei* and its mutants, including certain hyper-producing strains for example, Rut C-30, possess low β -glucosidase activity relative to the total cellulase activity, which impairs performance during the hydrolysis of lignocellulosic materials.¹⁰ Considering the lack of a complete cellulase system offered by these fungi, researchers are focusing on the isolation and screening of novel fungi with improved cellulase systems which are capable of acting synergically on the cellulosic biomass.⁸

Nevertheless, the recalcitrant nature of lignocelluloses is the obstacle in the production of biofuels. The amount of lignin

^aCollege of Life Science, Northeast Agricultural University, Harbin, 150030, China. E-mail: huxiaomei1982@163.com

^bHeilongjiang Academy of Agricultural Sciences, Harbin, 150086, China

^cHarbin University, Harbin, 150086, China. E-mail: bzhu@neau.edu.cn

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c8ra05729j



present in the biomass and the de-polymerization (DP) of cellulose limit the transformation efficiency.⁸ The pretreatment process such as steam explosion, acid or alkaline hydrolysis and ionic liquids-pretreatment are always employed.

Ionic liquids (ILs) have been recognized as an alternative green solvent for lignocellulose pretreatment, which possess several advantages including high thermal stability, negligible vapor pressure, no flammability, and widely accessible temperature range.^{11,12} For instance, 1-ethyl-3-methylimidazolium acetate ([EMIM]CH₃COOH) was reported widely to perform the effective decomposition of lignocelluloses and to provide the high cellulose digestibility.^{13–16}

Although ILs are effective for breaking down lignocelluloses, they can also inhibit the enzymes used in the subsequent saccharification, resulting the inefficient saccharification of cellulose. Therefore, a washing process is required to remove any residual ILs after ILs-pretreatment. Large scale washing requires extra energy, increasing more processing cost.^{17,18} However, there are very limited reports concerning on the ILs-tolerant cellulase.

In this study, a cellulose-degradable strain was isolated and identified. Factors governing the enzyme production were investigated. Studies on evaluation of the capability of salt-tolerant microbes were carried out. The capability of the potential microbes for producing salt-tolerant and ILs-tolerant cellulase was investigated. We also compared the enzymatic hydrolysis of lignocellulosic substrates by the crude enzyme and commercial cellulase.

2. Materials and methods

2.1 Samples collection

The soil was collected from wheat and corn rotation farm of Shanxian county in Shandong province in China. The surface soil (1–5 cm) was collected and stored at 4 °C. Corn stover and rice straw were provided by Xiangfang farm in Harbin city in China and they were dried, ground and passed through 40 mesh sieves. Wheat bran was provided by Xiangfang Flour Factories of Harbin in China. All chemicals were purchased from Aladdin company in China and were used as received.

2.2 Screening procedure

The colonies of various fungi were isolated from the soil using PDA medium. The isolated single colony was incubated on the CMC medium (K₂HPO₄ 1.0 g, NaCl 0.5 g, (NH₄)₂SO₄ 2.0 g, MgSO₄ 0.5 g, CMC-Na 10 g, agar 15 g) at 30 °C for 5 days. Following that, all plates were stained with Congo-red solution (0.1%, w/v) for 15 min and were discolored with 1 mol L⁻¹ NaCl.¹⁹ The strain with a large hydrolysis zone was selected and cultured on PDA medium for 4 days. After that, 1 cm of agar piece was taken and transferred into 30 mL of PD medium. The solutions were then cultured at 30 °C for 3 days with shaking at 200 rpm. 1 mL of the suspension (4–6 mg dry mycelia wt per mL) was added into 100 mL of Hutchinson inorganic salt medium (KH₂PO₄ 1.0 g, MgSO₄ 0.3 g, NaCl 0.1 g, CaCl₂ 0.1 g, FeCl₃ 0.01 g) with cover stover of 1% (w/v) and

peptone of 0.2% (w/v), which was incubated at 30 °C for 5 days with shaking at 150 rpm. The resulting solution was then centrifuged at 8000 rpm for 5 min at 4 °C to give the crude enzyme.

2.3 Enzyme assays

Enzyme assays, including filter paper cellulase (FPase), endo-glucanase (CMCase) and β-glucosidase, were determined as International Union of Pure and Applied Chemistry (IUPAC) standard procedure.²⁰ FPase activity was assayed by incubating the 0.5 mL of suitably diluted enzyme with Whatman no. 1 filter paper (1.0 × 6.0 cm) containing 1.0 mL of sodium citrate buffer (pH 4.8) for 60 min at 50 °C. The CMCase activity was determined using (1%, w/v) sodium carboxymethyl cellulose (CMC-Na) for 30 min at 50 °C. β-glucosidase activity was measured using salicin solution (0.5%, w/v) for 30 min at 50 °C. The reducing sugar was determined by 3,5-dinitrosalicylic acid (DNS) method.²¹ One unit (U) of enzyme activity was defined as the amount of the enzyme which released 1 μmol of reducing sugar/min under the conditions indicated.

2.4 Molecular identification

The DNA of the enriched fungus was extracted using a DNA kit according to the manufacturer's instructions (OMEGA, USA). Universal primers of ITS1: (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4: (5'-TCCTCGCCTTATTGATATGC-3') were used for fungal analysis. The ITS sequence of the gene was amplified by polymerase chain reaction (PCR). The reaction mixture of 50 μL contained 2 μL template DNA, 2 μL forward primer, 2 μL reverse primer, 25 μL 2 × Taq PCR MasterMix (TIANGEN, China) and 19 μL ddH₂O. The samples were amplified with the following conditions: initial denaturation at 95 °C for 5 min, followed by 35 cycles of denaturation at 95 °C for 30 s, annealing at 55 °C for 35 s, and elongation at 72 °C for 2 min, and final elongation at 72 °C for 10 min. The PCR products were sequenced in Huada gene company (Beijing, China). The sequence was compared to the sequences in the GenBank database. A homology search was performed using bioinformatics tools available online: BLAST (www.ncbi.nlm.nih.gov/BLAST). MEGA program (version 5.0) with the neighbor-joining (NJ) algorithm was used to build phylogenetic trees based on the ITS sequence.²² The ITS sequence data for fungal sample were deposited into the GenBank database.

2.5 Optimization of the conditions for cellulase production

Various factors were investigated in order to optimize the environmental conditions of cellulase production. The effects of carbon sources (corn stover, rice straw, wheat bran, CMC-Na, avicel) at a concentration of 1% (w/w) and nitrogen sources (peptone, beef extract, yeast extract, NH₄NO₃, urea) at a concentration of 0.2% (w/w) on the enzyme activity were determined on the basis of Hutchinson inorganic salt medium. Other factors including pH value, inoculum size, incubation temperature and time were evaluated.

2.6 Determination of salt-tolerance of the fungi

The capability of the fungi on cellulase production under different salt concentrations were determined. According to Wang *et al.* and Taechapoempol *et al.*,^{23,24} a modified method was used in our work. Based on the optimal conditions for cellulase production, different NaCl concentrations including 0%, 2.5%, 5.0%, 7.5% and 10.0% (w/v) was added and investigated. FPase, CMCase and β -glucosidase activity were measured as the method described in 2.3.

2.7 Optimization of the conditions for enzyme activities for enzymatic hydrolysis

The crude enzyme solution was obtained under the optimal conditions for cellulase production. The effect of pH value in the range of pH 3.0–11.0 was estimated, which was prepared by citrate buffer (pH 3.0–6.0), sodium phosphate buffer (pH 7.0–8.0) and glycine–NaOH buffer (pH 9.0–11.0). The incubation temperature (30–80 °C) with 5 °C intervals was studied. The mixture was cultured for 1 h. FPase, CMCase and β -glucosidase activities were measured as the method described in 2.3.

2.8 Determination of enzyme activities in the presence of salt and ionic liquid

The 1-ethyl-3-methylimidazolium acetate ([EMIM]CH₃COOH) was prepared using the published methods in the literature.²⁵ According to Gunny *et al.*,²⁶ a modified method was used in our work. To evaluate the stability of cellulase at different salt and ILs concentrations, the crude enzyme solution was mixed with citrate buffer at the pH 5.0 with the addition of NaCl or [EMIM]CH₃COOH in the range of 0–15.0% (w/v) and was cultured for 1 h at 60 °C. FPase, CMCase and β -glucosidase activities were measured as the method described in 2.3.

2.9 Pretreatment of corn stover

The pretreatment of corn stover (1 g) was performed in [EMIM]CH₃COOH (20 g) at 120 °C for 3 h in an 100 mL autoclave. After that, the mixture was cooled to room temperature quickly and 80 mL of acetone/deionized water (1/1, v/v) was added into the mixture. The resulting solid was filtered and washed with 80 mL of acetone/deionized water (1/1, v/v), followed by 300 mL of deionized water in order to remove any residual ionic liquid. The solid was then dried in a vacuum at 60 °C to give the cellulose-rich materials. The yield of cellulose-rich materials was determined as the ratio of the mass of cellulose-rich materials in the mass of the corn stover that was subjected to the pretreatment process according to eqn (1), and the yield was 60.23%.

$$\text{Cellulose - rich materials(\%)} = \frac{\text{cellulose - rich materials(mg)}}{\text{corn stover subjected to pretreatment(mg)}} \quad (1)$$

2.10 Analysis of corn stover after ILs-pretreatment

According to National Renewable Energy Lab (NREL) method, 300 mg of corn stover that was pretreated by [EMIM]CH₃COOH or corn stover was mixed with 3 mL 72% H₂SO₄ in a water bath at 30 °C for 60 min. The mixture was stirred strongly every 10 min. Following that, 84 mL deionized water was added into the mixture, which is diluted into 4% H₂SO₄ aqueous solution. The mixture was put in an autoclave at 121 °C for 60 min. After the mixture was cooled to room temperature, the mixture was centrifuged. The filtrate was then determined for glucose and xylose using HPLC analysis. Measure the absorbance of the sample at 320 nm with $\epsilon=(30 \text{ L g}^{-1} \text{ cm}^{-1})$ on a UV-visible spectrophotometer. The filtered solids were put into a muffle furnace at 575 °C to give the acid-insoluble lignin and ash. The content of cellulose and hemicellulose was calculated from glucose and xylose by conversion factors of 0.90 and 0.88, respectively.^{11,27} There are cellulose (30.55%), hemicellulose (14.40%), lignin (22.47%) and ash (1.3%) in original corn stover. There are cellulose (33.20%), hemicellulose (20.54%), lignin (21.98%) and ash (1.0%) in regenerated cellulose-rich materials.

2.11 Enzymatic hydrolysis

50 mg of corn stover that was pretreated by [EMIM]CH₃COOH was mixed with an appropriate amount of the crude enzyme (0.02 U FPase/mg of substrate) that was filtered through 0.22 μm filter membrane in a 50 mL Erlenmeyer flask. The total volume of the reaction mixture was replenished to 20 mL using citrate buffer (0.05 mol L⁻¹) at pH 5.0. The reaction mixture was incubated under a rotary shaker at 60 °C for 72 h with 150 rpm. The commercial cellulase (from *Trichoderma reesei* ATCC26921) was used as a positive control. A commercial cellulase reaction system (enzyme dosage equal to crude enzyme) in 20 mL of citrate buffer (0.05 mol L⁻¹, pH 4.8) served as a substitute of the culture supernatants from *T. reesei* ATCC26921. The enzymatic hydrolysis was carried out for 72 h at 50 °C with 150 rpm. The control experiments were carried out under the same conditions using the denatured commercial enzymes or the denatured crude enzyme that was boiled for 30 min. The reducing sugars were measured by 3,5-dinitrosalicylic acid (DNS) method as described by Miller.²¹ The yields of glucose, xylose and cellobiose were measured by HPLC. The sugar saccharification rate of 1U FPase was calculated according to eqn (2).

$$\% \text{ Saccharification} = \frac{\text{reducing sugar(mg)}}{\text{cellulose - rich materials(mg)}} \quad (2)$$

3. Results and discussion

The detailed schematic representation has been provided as Fig. 1.

3.1 Isolation and identification of the microbe

Five microbial strains with discoloration zones of Congo-red were isolated as the positive microbes, indicating their ability to secrete cellulase. The microbe with a large discoloration zone




Fig. 1 The schematic representation in this study.

was selected and determined for its enzyme activities, which were 0.11 U mL^{-1} (FPase), 0.21 U mL^{-1} (CMCase) and 0.43 U mL^{-1} (β -glucosidase), respectively.

This fungal strain was determined based on its ITS sequence of DNA. The obtained nucleotide sequence has been submitted to NCBI GenBank with under accession no. MH558553. Phylogenetic tree was shown in Fig. 1. The ITS sequence of HC6 showed 100% sequence similarity with that of *Penicillium oxalicum* strain 68 (accession no. GU078430). Thus, this fungal strain was identified as *Penicillium oxalicum* HC6.

Many strains of *Penicillium* sp. such as *Penicillium janthinellum* EMS-UV-8 and *Penicillium decumbens* JU-A10 were increasingly being reported to produce cellulase at a commercial level.^{28,29} 13 species of *Penicillium* sp. exhibiting avicelase activity among 31 fungal strain were isolated from subtropical and tropical forests of China.³⁰ The newly isolated *P. oxalicum* HC6 has a potential to possess the cellulase production (Fig. 2).

3.2 Optimization of the conditions for the cellulase production

3.2.1 Effects of carbon and nitrogen sources. Cellulase production was found to be dependent on the nature of the




Fig. 2 Phylogenetic relationship of fungal strain HC6 based on ITS sequence.




Fig. 3 Effects of carbon sources and nitrogen sources on the cellulase production.

carbon source. Among the natural lignocellulosic materials (corn stover, rice straw, wheat bran) and the commercial polymeric materials (CMC-Na, avicel), corn stover was found to be the most effective carbon source. High enzyme activities of CMCase and β -glucosidase were obtained at 30°C and pH 7.2 with peptone and corn stover under 150 rpm with inoculation size of 1% (v/v) for 5 days. Good FPase activity was observed, when rice straw was used as the carbon source. Unfortunately, the enzyme activity was notably lowest when CMC-Na and avicel were used as the carbon source.

Recently, lignocellulosic biomasses have been studied as potential substrates for the enzyme production using microorganisms, especially filamentous fungi. Similar results were reported that high amounts were obtained from *Penicillium oxalicum* GZ-2 and *Penicillium* sp. GD01 from untreated lignocellulosic materials compared with commercial polymeric materials.^{31,32} The chemical composition of lignocellulosic biomass and the accessibility of various components of corn stover and rice straw, especially hemicelluloses, improved the enzyme hydrolysis by *P. oxalicum* HC6. The use of economical agricultural wastes instead of expensive cellulose is of the great interest for industrial application.

Nitrogen source is also one of the crucial factors that affect microbial metabolism.³³ In this work, three nitrogen organic sources (peptone, beef extract, yeast extract) and two inorganic nitrogen sources (NH_4NO_3 , urea) were selected for the optimization of nitrogen sources at 30°C and pH 7.2 with corn stover and incubation size of 1% (v/v) under 150 rpm for 5 days. As shown in Fig. 3b, urea was the most effective nitrogen source for CMCase production of *P. oxalicum* HC6. High production of FPase and β -glucosidase were obtained, when peptone was employed as the nitrogen source.

Variation in the nitrogen source can affect the metabolic processes of the microbes significantly to produce different cellulase production. Previous studies have shown that organic nitrogen sources can induce cellulase secretion better than inorganic nitrogen sources.^{34,35} Peptone was the suitable organic nitrogen source for most cellulases for *P. oxalicum* HC6.

3.2.2 Effects of pH, inoculum size, temperature and incubation time. Initial pH analysis of *P. oxalicum* HC6 was carried out at 30°C with inoculation size of 1% (v/v) for 5 days. As shown in Fig. 4a, both FPase and CMCase showed good stability in the range of pH 3.0–11.0 and the optimal pH of *P. oxalicum* HC6 appeared at pH 5.0 and pH 4.0, respectively. β -glucosidase activity was increased from pH 3.0 to pH 4.0 and the optimal pH showed at pH 4.0.




Fig. 4 Effects of pH (a), inoculation size (b), temperature (c) and incubation time (d) on the cellulase production.

Many enzyme systems and the transport of several enzyme species across the cell membrane are influenced by the initial pH of medium.³⁶ Moreover, the synthesis and expression of certain genes, and the microbial metabolic activities, are influenced by the internal pH, in response to that of the external environment.³⁷ The preference for acidic pH for cellulase producing microbes has been reported for ascomycetes fungi *e.g.* *T. reesei*,³⁸ *Talaromyces verruculosus*,³⁹ and *Chaetomium globosum*.⁴⁰ It was reported that good CMCase activity was observed with *Penicillium funiculosum* (pH 4.82), *Penicillium citrinum* (pH 5.5), *Penicillium chrysogenum* (pH 5.0), and *Aspergillus niger* (pH 5.0).^{41–44} As reported, filamentous fungi have reasonably good growth over a broad range of pH 2–9, with an optimal range of 3.8 to 6.0.⁴⁵ Each microorganism possesses a pH range for its growth and activity with an optimum value within the range. The acidic pH 4 was favorable for the *P. oxalicum* HC6.

The investigation of inoculation size was performed at 30 °C and pH 4.0 for 5 days and the result was shown in Fig. 4b. The enzyme production was increased with the inoculation size from 1% to 2% and reduced from 2% to 3%, which was then stable from 3% to 5%. Good CMCase, FPase and β -glucosidase activities were obtained with the inoculation size of 2%. Lower inoculation size requires longer time for fungal multiplication and substrate utilization, whereas higher inoculation size increases the spore density as well as the water content in the medium causing hindrance in oxygen penetration resulting in the inhibited fungal growth and enzyme production.⁴⁶ A balance between nutrients and growing microbes is necessary for optimum enzyme production.

As presented in Fig. 3c, the maximum CMCase and β -glucosidase activities were found at 30 °C with inoculation size of 2% at pH 4.0 for 5 days. Good FPase activity was obtained when the fermentation temperature was 25 °C. When the temperature was increased more than 30 °C, the enzyme activities were reduced significantly. At high temperature, a lower amount of enzyme will be formed due to the thermal denaturation of the metabolic pathway of enzymes.⁴⁷ However, at low temperature, metabolic activities of microorganisms slow down resulting in low enzyme yield.⁴⁸

The effect of fermentation time on the cellulase production of *P. oxalicum* HC6 was illustrated in Fig. 4d. As the




Fig. 5 The analysis of salt tolerance of *P. oxalicum* HC6 for cellulase production.

fermentation time was increased from 2 days to 8 days, nutrients were efficiently used by microbes that resulted in high enzyme production. After 8 days, the enzyme yield was reduced due to the depletion of nutrients. In addition, higher β -glucosidase/FPase ratio has been previously reported to reduce enzyme inhibition due to lesser cellobiose accumulation during hydrolysis.^{49,50} Therefore, the crude enzyme from *P. oxalicum* HC6 with high β -glucosidase/FPase ratio possibly help to decrease enzyme inhibition and to improve glucose production instead of cellobiose.

Based on the optimum conditions for cellulase production of *P. oxalicum* HC6, the activities of FPase, CMCase and β -glucosidase were 0.20 U mL⁻¹, 0.63 U mL⁻¹, 1.55 U mL⁻¹, when cron stover and peptone were used as the carbon source and nitrogen source, at pH 4.0 and 30 °C with inoculation size of 2% for 8 days.

3.3 Analysis of salt tolerant capability of *P. oxalicum* HC6

The salt tolerance of *P. oxalicum* HC6 was determined by selecting different NaCl concentrations of 0–10.0% (w/v). The results were shown in Fig. 5. It was obvious that the activities of CMCase and β -glucosidase of *P. oxalicum* HC6 were increased with an increase of NaCl concentration in the range of 0–5.0%. High activities of CMCase and β -glucosidase were observed at 0.73 U mL⁻¹ and 1.83 U mL⁻¹ with 5.0% NaCl, respectively, which were increased to 115.87% and 118.06% compared with the enzyme activities in the absence of NaCl. The results showed that CMCase and β -glucosidase could be promoted at an appropriate NaCl concentration. However, FPase activity was inhibited at the presence of NaCl. Smolyanyuk *et al.* described that salt tolerant fungi could grow at the salt concentration of more than 0.5 mol L⁻¹.⁵¹ *P. oxalicum* HC6 with the salt tolerance of more than 0.7 mol L⁻¹ in this study has an effective salt tolerance capability with high cellulase activity.

3.4 Optimization of the conditions for enzyme activities for enzymatic hydrolysis

The study for enzyme activities for enzymatic hydrolysis under different pH was carried out at 50 °C (Fig. 6a). CMCase and β -glucosidase showed good activities over a broad acidic pH range of 3.0–6.0. The optimal pH was obtained at pH 4.0 for CMCase (0.65 U mL⁻¹) and pH 5.0 for β -glucosidase (1.67 U mL⁻¹). It was




Fig. 6 The enzyme activities of *P. oxalicum* HC6 at different pH (a) and temperature (b).

reported that the highest activity for CMCase of *Penicillium* sp. was 0.409 U mL^{-1} at the optimal pH 4.2 and highest activity of β -glucosidase was 0.038 U mL^{-1} at pH 6.0.⁵² The degradation of lignocellulose requires the cooperation of several cellulases and their synergistic effect. Considering this, pH 5.0 was selected for the following studies.

In addition, the thermostable cellulase is an ideal candidate for bioprocessing industry, as a relatively high temperature is favorable for the solubility of the substrate and the efficiency of the enzymatic hydrolysis.⁵³ High temperature could also reduce the risk of contamination by undesirable microorganisms.⁵⁴ Castro *et al.* reported an optimum temperature of $60 \text{ }^\circ\text{C}$ for both endoglucanase and β -glucosidase activities with *Penicillium funiculosum* ATCC 11797.⁵⁵

Generally, thermophilic microorganisms having growth temperature ranges from 50 to $80 \text{ }^\circ\text{C}$ are potential candidates to produce highly thermostable cellulase. Interestingly, several mesophilic microorganisms were reported to produce a considerable thermostable cellulases.⁵⁶ In this study, although the high enzyme production of *P. oxalicum* HC6 were obtained at $30 \text{ }^\circ\text{C}$, CMCase and β -glucosidase reached their maximum activities at $60 \text{ }^\circ\text{C}$, which were 0.71 U mL^{-1} and 7.44 U mL^{-1} , respectively. It is worth to note that when the temperature was increased to $60 \text{ }^\circ\text{C}$, the activities of β -glucosidase and CMCase were increased to 445.51% and 129.09%, respectively. The cellulase of *P. oxalicum* HC6 with high pH and thermal stability has great potential for the industrial application.

3.5 Stability of enzyme activity of *P. oxalicum* HC6 under NaCl and ILS

As reported, the endoglucanase from *Bacillus agaradhaerens* JAM-KU023 was a salt-activated endoglucanase by adding 0.2 – 2.0 mol L^{-1} NaCl.⁵⁷ The cellulase from *Aspergillus terreus* UniMAP AA-6, was reported to be stable under NaCl and [EMIM] CH_3COOH solution. The highest FPase activity of *A. terreus* UniMAP AA-6 was 0.0625 U mL^{-1} in the presence of 7.70% NaCl (w/v).⁵⁸

The crude enzyme of *P. oxalicum* HC6 in this work was determined under different NaCl solution (0–15.0%, w/v) at $60 \text{ }^\circ\text{C}$ with initial pH 5.0 for 1 h. Every enzyme activity (FPase, CMCase and β -glucosidase) was considered as 100% at NaCl concentration of 0%. As shown in Fig. 7, β -glucosidase activity was increased as NaCl concentration was increased from 0% to 2.5%. The optimal β -glucosidase activity reached 8.00 U mL^{-1} , corresponding to 107.53% of the enzyme activity. At a higher NaCl solution of 15.0%, β -glucosidase activity still remained 4.47 U mL^{-1} , corresponding to 60% of the enzyme activity. The




Fig. 7 The enzyme activity of *P. oxalicum* HC6 under different salt concentrations.

optimal CMCase activity was achieved at 0.79 U mL^{-1} , corresponding to 111.27% of the enzyme activity. At a higher NaCl solution of 15.0%, CMCase activity still remained 0.50 U mL^{-1} , corresponding to 70.42% of the enzyme activity.

It was described that the excess acidic amino acids on the cellulases could provide a negatively charge on its surface. The negative charges keep the protein soluble in a high salt solution either by forming a hydrated ion network with cations or by inhibiting the formation of protein aggregation through electrostatic repulsive charges at the protein surface.⁵⁹

The crude enzyme from *P. oxalicum* HC6 was further incubated in [EMIM] CH_3COOH with different concentrations. The result was shown in Fig. 8. Every enzyme activity (CMCase, FPase and β -glucosidase) was considered as 100% at ILS concentration of 0%. It was found that CMCase and β -glucosidase showed good stability under the ILS concentration of 0–2.5% (w/v), which maintained almost 100% of the enzyme activity. However, β -glucosidase activity was reduced from 100.02% to 25.67%, when the concentration of ILS was increased from 2.5% to 10.0%. CMCase activity was reduced from 100% to 26.76%, when the concentration of ILS was increased from 2.5% to 10.0%. A relatively lower stability was observed for FPase in the presence of [EMIM] CH_3COOH .

ILs as effective pretreated method for lignocelluloses was reported widely. However, numbers of extensive washing with water after ILS-pretreatment were required, because the commercial fungal cellulases were inhibited by some ILS,^{60,61} which increased the production cost. In order to decrease the number of washing steps and increase the yields of sugars, it is necessary to identify ILS-tolerant enzymes for enzymatic hydrolysis of the regenerated cellulose in the presence of ILS.




Fig. 8 The enzyme activity of *P. oxalicum* HC6 under different [EMIM] CH_3COOH concentrations.




Fig. 9 SEM analysis of untreated (a) corn stover, (b) corn stover pretreated by [EMIM]CH₃COOH, (c) cellulose-rich materials treated by the crude of *P. oxalicum* HC6, (d) cellulose-rich materials treated by the commercial cellulose.

The cellulase from *P. oxalicum* HC6 in this work was potentially ILs-tolerant enzymes for further application.

It has been suggested that ILs inhibit enzymatic activity by disrupting hydrogen bonding and hydrophobic interactions and depriving the water hydration shell of the protein.^{62,63} This is similar to the denaturing effect caused by salt on mesophilic proteins. High ionic strength could be the possible reason for both salt and ILs that caused the denaturing effect of enzymes.

3.6 Analysis of scanning electron microscopy

The structural changes in morphology of the corn stover that was pretreated by [EMIM]CH₃COOH and the following crude enzyme of *P. oxalicum* HC6 were investigated. SEM images were taken on a Hitachi S-3400N microscope. The original corn stover showed the compact ordered and rigid structures (Fig. 9a). The significant morphological modification was observed after [EMIM]CH₃COOH-pretreatment, which was loose and less compact (Fig. 9b). After the treatment by the crude enzyme (Fig. 9c) and the enzyme from *T. reesei* ATCC26921 (Fig. 9d), the surface of the residue become more loose and porous. The alteration of morphology indicated the disruption of linkages in lignocellulose, providing a more accessible surface to cellulase, leading to enhanced enzymatic digestibility.

3.7 Enzymatic saccharification of the crude enzyme of *P. oxalicum* HC6

FPase activity was added as 1 U for both the commercial enzyme and the crude enzyme of *P. oxalicum* HC6. After enzymatic hydrolysis, a high yield of reducing sugar (12.29 mg) was obtained after the saccharification by the crude enzyme of *P. oxalicum* HC6. The total saccharification degree of cellulose-rich materials was up to 24.58%. Sugars were determined by HPLC (Agilent 1200 Series, USA) analysis. There are 8.93 mg of glucose and 1.64 mg of xylose. In addition, a good yield of reducing

sugar (10.70 mg) was obtained after the saccharification by the commercial cellulase and the total saccharification degree was up to 21.40%. 3.46 mg of glucose, 1.73 mg of xylose and 5.05 mg of cellobiose were achieved. No significant difference was observed between *P. oxalicum* HC6 and commercial cellulase with the same FPase activity.

Specifically, higher yield of glucose was obtained, but no cellobiose was observed for the treatment by the crude enzyme of *P. oxalicum* HC6 compared with the commercial cellulase *T. reesei* ATCC26921, because of the capability of *P. oxalicum* HC6 to produce a high β -glucosidase activity. An optimal ratio of β -glucosidase/FPase is critical. Although *Trichoderma* sp. has been reported to produce good quantities of cellulolytic enzymes, it possess low β -glucosidase activity, which reduce the hydrolysis rate of lignocellulose to glucose.¹⁰ Moreover, the lack of β -glucosidase activity of *T. reesei* cellulase could result in inhibition of the cellulase by accumulated cellobiose in the reaction.⁶⁴ As a result, the addition of external β -glucosidase enzymes from different microbial origins was always required. *Penicillium* sp. have the ability to produce complete enzyme mixture with a better β -glucosidase/FPase ratio than *Trichoderma* sp.⁶⁵

It was reported that the crude enzyme from *Penicillium citrinum* CBMAI 1186, *Aspergillus sydowii* CBMAI 934, *Aspergillus sydowii* CBMAI 935 and *Mucor racemosus* CBMAI 847d produced 19%, 19%, 78% and 24% of the saccharification with the addition of FPase (1 U mL⁻¹) for sugarcane bagasse after the alkaline treatment.⁶⁶ The crude enzyme from *Geobacillus stearothermophilus* resulted in 71.03% of the saccharification with the addition of FPase (30 U mL⁻¹) for the date palm wastes after the alkaline treatment.⁶⁷ *P. oxalicum* HC6 in this study has a potential for its application in enzymatic saccharification.

4. Conclusions

In this study, a fungal strain was newly isolated and identified as *Penicillium oxalicum* HC6. Under the optimal conditions using corn stover and peptone as the carbon source and nitrogen source at pH 4.0 and 30 °C with inoculation size of 2% (v/v) for 8 days, a relatively high production of FPase, CMCase and β -glucosidase were obtained as 0.20 U mL⁻¹, 0.63 U mL⁻¹ and 1.55 U mL⁻¹, respectively. It was found that the enzyme production of CMCase and β -glucosidase was increased to 0.73 U mL⁻¹ and 1.83 U mL⁻¹ at the presence of NaCl (5.0%, w/v). Good activities of FPase, CMCase and β -glucosidase were observed at 0.20 U mL⁻¹, 0.71 U mL⁻¹ and 7.44 U mL⁻¹ at the optimal pH 5.0 and the optimal temperature of 60 °C. The enzyme activity was stable in the presence of salt and ILs. Hydrolytic efficiency of *Penicillium oxalicum* HC6 was comparable to the commercial cellulase preparation, which indicated its great potential for its application in the conversion of lignocelluloses to biofuels.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was financially supported by University Nursing Program for Young Scholars with Creative Talents in Heilongjiang Province (UNPYSCT-2017021).

References

- 1 Z. Anwar, M. Gulfranz and M. Irshad, *J. Radiat. Res. Appl. Sci.*, 2014, **7**, 163–173.
- 2 A. Pandey, C. R. Soccol, P. Nigam and V. T. Soccol, *Bioresour. Technol.*, 2000, **74**, 69–80.
- 3 S. S. Behera and R. C. Ray, *Int. J. Biol. Macromol.*, 2016, **86**, 656–669.
- 4 O. Kirk, T. V. Borchert and C. C. Fuglsang, *Curr. Opin. Biotechnol.*, 2002, **13**, 345–351.
- 5 M. Knauf and M. Moniruzzaman, *Int. Sugar J.*, 2004, **106**, 147–150.
- 6 M. Chandra, A. Kalra, P. K. Sharma, H. Kumar and R. S. Sangwan, *Biomass Bioenergy*, 2010, **34**, 805–811.
- 7 R. L. Howard, E. Abotsi, E. L. J. van Rensburg and S. Y. Howard, *Afr. J. Biotechnol.*, 2003, **2**, 602–619.
- 8 N. Srivastava, M. Srivastava, P. K. Mishra, V. K. Gupta, G. Molina, S. Rodriguez-Couto, A. Manikanta and P. W. Ramteke, *Renewable Sustainable Energy Rev.*, 2018, **82**, 2379–2386.
- 9 R. Rawat, N. Srivastava, B. S. Chadha and H. S. Oberoi, *Energy Fuels*, 2014, **28**, 5067–5075.
- 10 H. Jorgensen, A. Morkeberg, K. B. R. Krogh and L. Olsson, *Enzyme Microb. Technol.*, 2005, **36**, 42–48.
- 11 P. Zhang, S. J. Dong, H. H. Ma, B. X. Zhang, Y. F. Wang and X. M. Hu, *Ind. Crops Prod.*, 2015, **76**, 688–696.
- 12 P. Mäki-Arvela, I. Anugwom, P. Virtanen, R. Sjöholm and J. P. Mikkola, *Ind. Crops Prod.*, 2010, **32**, 175–201.
- 13 C. G. Liu, J. C. Qin, L. Y. Liu, B. W. Jin and F. W. Bai, *ACS Sustainable Chem. Eng.*, 2016, **S1**, 577–582.
- 14 S. D. Wang, J. C. Chen, G. H. Yang, W. H. Gao and K. F. Chen, *Bioresour. Technol.*, 2017, **243**, 335–338.
- 15 S. M. Raeisi, M. Tabatabaei, A. Bitá, A. Ghafari and S. H. Mood, *Waste Biomass Valorization*, 2016, **7**, 97–107.
- 16 F. Xu, Y. C. Shi and D. Wang, *Bioresour. Technol.*, 2012, **114**, 720–724.
- 17 P. Engel, R. Mladenov, H. Wulffhorst, G. Jager and A. C. Spiess, *Green Chem.*, 2010, **12**, 1959–1966.
- 18 T. Zhang, S. Datta, J. Eichler, N. Ivanova, S. D. Axen, C. A. Kerfeld, F. Chen, N. Kyrpides, P. Hugenholtz and J. F. Cheng, *Green Chem.*, 2011, **13**, 2083–2090.
- 19 R. M. Teather and P. J. Wood, *Appl. Environ. Microbiol.*, 1982, **43**, 777–780.
- 20 V. K. Ghose, *Pure Appl. Chem.*, 1987, **59**, 257–268.
- 21 G. L. Miller, *Anal. Chem.*, 1959, **31**, 426–428.
- 22 J. Felsenstein, *Evolution*, 1985, **39**, 739–791.
- 23 C. Wang, Y. Hsieh, C. Ng, H. Chan, H. Lin, W. Tzeng and Y. Shyu, *Enzyme Microb. Technol.*, 2009, **44**, 373–379.
- 24 K. Taechapoempol, T. Sreethawong, P. Rangsunvigit, W. Namprohm, B. Thamprajamchit, S. Rengpipat and S. Chavadej, *Appl. Biochem. Biotechnol.*, 2011, **164**, 204–219.
- 25 P. Wasserscheid and T. Welton, *Ionic liquids in synthesis*, WILEY-VCH, Federal republic of Germany, 2003, vol. 2, ISBN:3-527-30515-7.
- 26 A. A. N. Gunny, D. Arbain, R. E. Gumba, B. C. Jong and P. Jamal, *Bioresour. Technol.*, 2014, **155**, 177–181.
- 27 A. Sluiter, B. Hames, R. Ruiz, C. Scarlata, J. Sluiter, D. Templeton and D. Crocker, *National Renewable Energy Laboratory*, 2011, 1–15.
- 28 M. G. Adsul, K. B. Bastawde, A. J. Varma and D. V. Gokhale, *Bioresour. Technol.*, 2007, **98**, 1467–1473.
- 29 Y. Cheng, X. Song, Y. Qin and Y. Qu, *J. Appl. Microbiol.*, 2009, **107**, 1837–1846.
- 30 J. Zhang, J. L. Liu, J. Y. Lan, C. J. Duan, Q. S. Ma and J. X. Feng, *Biotechnol. Biofuels.*, 2014, **7**, 107.
- 31 H. P. Liao, X. T. Fan, X. L. Mei, Z. Wei, W. Raza, Q. Shen and Y. C. Xu, *Biomass Bioenerg.*, 2015, **74**, 122–134.
- 32 D. M. Kim, E. J. Cho, J. W. Kim, Y. W. Lee and H. J. Chung, *Afr. J. Biotechnol.*, 2014, **13**, 145–155.
- 33 N. Kulkarni, A. Shendye and M. Rao, *FEMS Microbiol. Rev.*, 1999, **23**, 411–456.
- 34 M. Sakthivel, N. Karthikeyan, R. Jayaveny and P. Palani, *J. Ecobiotechnol.*, 2010, **2**, 6–13.
- 35 D. J. M. Kumar, P. D. Poovai, C. L. P. Kumar, Y. S. Saroja, A. Manimaran and P. T. Kalaichelvan, *Der Pharmacia Lett.*, 2012, **4**, 881–888.
- 36 U. Gupta and R. Kar, *Jordan J. Biosci.*, 2008, **1**, 129–134.
- 37 E. R. Olson, *Mol. Microbiol.*, 1993, **8**, 5–14.
- 38 C. Li, Z. Yang, R. Z. C. He, D. Zhang, S. Chen and L. Ma, *J. Biotechnol.*, 2013, **168**, 470–477.
- 39 S. Goyari, S. H. Devi, L. Bengyella, M. Khan, C. K. Sharma, M. C. Kalita and N. C. Talukdar, *J. Appl. Microbiol.*, 2015, **119**, 88–98.
- 40 A. H. M. El-Said and A. Saleem, *Mycobiology*, 2008, **36**, 1–9.
- 41 T. Dutta, R. Sahoo, R. Sengupta, S. S. Ray, A. Bhattacharjee and S. Ghosh, *J. Ind. Microbiol. Biotechnol.*, 2008, **35**, 275–282.
- 42 F. El Bergadi, F. Laachari, M. Sadiki, S. Elabed, M. H. Iraqui and S. K. Ibnsouda, *Microbiology*, 2016, **85**, 47–55.
- 43 S. Ahmed, M. I. Ghorí, M. A. Malana and A. Jamil, *Afr. J. Biotechnol.*, 2012, **11**, 7227–7231.
- 44 A. M. Castro, M. L. Carvalho, S. G. F. Leite and N. Pereira, *J. Ind. Microbiol. Biotechnol.*, 2010, **37**, 151–158.
- 45 M. K. Gowthaman, C. Krishna and M. Moo-Young, *Appl. Mycol. Biotechnol.*, 2001, **1**, 305–352.
- 46 V. H. Vu, T. A. Pham and K. Kim, *Mycobiology*, 2011, **39**, 20–25.
- 47 S. P. Shah, K. S. Kalia and J. S. Patel, *J. Gen. Appl. Microbiol.*, 2015, **61**, 35–43.
- 48 P. F. Omojasola and O. P. Jilani, *Res. J. Microbiol.*, 2009, **4**, 67–74.
- 49 M. Galbe and G. Zacchi, *Appl. Microbiol. Biotechnol.*, 2002, **59**, 618–628.
- 50 M. Camassola, L. R. de Bittencourt, N. T. Shenem, J. Andreaus and A. J. P. Dillon, *Biocatal. Biotransform.*, 2004, **22**, 391–396.
- 51 E. V. Smolyanyuk and E. N. Bilanenko, *Microbiology*, 2011, **80**, 877–883.

- 52 P. S. Santa-Rosa, A. L. Souza, R. A. Roque, E. V. Andrade, S. Astolfi-Filho, A. J. Mota and C. G. Nunes-Silva, *Electron. J. Biotechnol.*, 2018, **31**, 84–92.
- 53 O. Khelila and B. Cheba, *Procedia. Technol.*, 2014, **12**, 519–528.
- 54 M. J. Liszka, M. E. Clark, E. Schneider and D. S. Clark, *Annu. Rev. Chem. Biomol. Eng.*, 2012, **3**, 77–102.
- 55 V. P. Zambare, A. Bhalla, K. Muthukumarappan, R. K. Sani and L. P. Christophe, *Extremophiles*, 2011, **15**, 611–618.
- 56 A. M. de Castro, M. L. D. de Carvalho, S. G. F. Leite and N. Pereira, *J. Ind. Microbiol. Biotechnol.*, 2010, **37**, 151–158.
- 57 K. Hirasawa, K. Uchimura, M. Kashiwa, W. D. Grant, S. Ito, T. Kobayashi and K. Horikoshi, *Antonie van Leeuwenhoek*, 2006, **89**, 211–219.
- 58 A. A. N. Gunny, D. Arbain, P. Jamal and R. E. Gumba, *Saudi J. Biol. Sci.*, 2015, **22**, 476–483.
- 59 A. Ventosa, A. Oren and Y. Ma, *Halophiles and Hypersaline Environments Current Research and Future Trends*, Springer-Verlag, Berlin, Heidelberg, 2011, vol. 18, ISBN: 978–3–642–20197–4.
- 60 N. Kamiya, Y. Matsushita, M. Hanaki, K. Nakashima, M. Narita, M. Goto and H. Takahashi, *Biotechnol. Lett.*, 2008, **30**, 1037–1040.
- 61 M. B. Turner, S. K. Spear, J. G. Huddleston, J. D. Holbrey and R. D. Rogers, *Green Chem.*, 2003, **5**, 443–447.
- 62 S. Bose, D. W. Armstrong and J. W. Petrich, *J. Phys. Chem. B*, 2010, **114**, 8221–8227.
- 63 M. Moniruzzaman, N. Kamiya and M. Goto, *Org. Biomol. Chem.*, 2010, **8**, 2887–2899.
- 64 M. Dashtban and W. Qin, *Microb. Cell Fact.*, 2012, **11**, 63.
- 65 H. Jorgensen and L. Olsson, *Enzyme Microb. Technol.*, 2006, **38**, 381–390.
- 66 D. A. Santos, M. M. Oliveira, A. A. S. Curvelo, L. P. Fonseca and A. L. M. Porto, *Int. Biodeterior. Biodegrad.*, 2017, **121**, 66–78.
- 67 S. A. Alrumman, *Braz. J. Microbiol.*, 2016, **47**, 110–119.