ANTHROPOLOGY

Comment on "Evidence of prehistoric human activity in the Falkland Islands"

T. J. Clark¹*, Jason Newton², Ewan D. Wakefield³

Stable isotopes from archaic Falkland Islands wolves (*Dusicyon australis*) indicate a high trophic, marine diet. Hamley *et al.* argue that this is consistent with mutualism with Yaghan people. However, most *D. australis* had similar isotopic signatures in the European era, despite human persecution. These data therefore neither support nor refute human-mediated introduction of *D. australis* to the Falklands.

Before their extirpation by European settlers ~1876, *Dusicyon australis* was the Falkland's only endemic land mammal. It remains unclear how it populated the Falklands, the two main theories being that it crossed from continental South America during the Last Glacial Maximum, when the sea level was lower and seasonal ice was more extensive (1), or that it arrived from the continent with humans, with whom it had a mutualistic relationship (2). (We note that these two hypotheses are not necessarily mutually exclusive.) Hamley *et al.* (3) provide valuable new evidence of pre-European bone deposits, the discovery of a stone projectile of apparently local manufacture, and sedimentary charcoal accumulation in the Falklands. Together with previously reported lithics (4) and charcoal accumulation (5), these data are consistent with the hypothesis that people indigenous to continental South America were present in the archipelago before European colonization.

Hamley et al. (3) additionally report stable isotope ratios in subfossil D. australis collagen, citing these in support of the pre-European occupation hypothesis. Their reasoning is as follows: Stable isotope ratios in the tissues of consumers predictably reflect those in their diet (6). High *D. australis* δ^{13} C and δ^{15} N values indicate a relatively marine, high trophic level diet consistent with mutualism with humans such as the Yaghan (Yámana) people, who hunted and gathered marine organisms, including seals, penguins, and shellfish. Although they acknowledge that the observed *D. australis* isotopic signatures could also arise from scavenging such prey, the weight of their argument is in favor of the former hypothesis. For example, they state that "The δ^{15} N values reported here reflect a marine diet at higher trophic levels, including seals, which the small canid was unlikely to have hunted" (3). However, D. australis was a mediumsized canid, with body mass of 12 to 14 kg, body length without tail of ~77 cm, height at shoulder of 48 cm (7), and smaller or similarly sized canids routinely predating pinniped pups [e.g., (8, 9)]. They then say that "Future studies of warrah [D. australis] remains collected during the European period, when abundant terrestrial food sources (e.g., sheep, cattle, horses, rabbits, pigs, rats, and mice) had been introduced to the islands, may provide further insight into the warrah's dietary preferences and potential interaction with humans" (3).

*Corresponding author. Email: tjclark1873@gmail.com

Here, Hamley et al. (3) omit an important piece of evidence that detracts from their argument. Previously, we analyzed stable isotope ratios in 19th century D. australis hair tissues from eight museum specimens (7). This showed that despite being persecuted by humans, eventually to extinction, the majority of D. australis in the European period also had a high δ^{13} C, high δ^{15} N diet. Furthermore, with their consent, we included collagen isotope data from Hamley et al. (3) before the submission of their paper and found that their stable isotope data were within or near the probable isotopic niche of 19th century D. australis (7). By implication, the most parsimonious explanation of the currently available stable isotope data is that it does not imply mutualism with humans and therefore has little bearing on the question of how D. australis first arrived in the Falklands. Other evidence supports the supposition that D. australis were able to obtain higher trophic marine prey for themselves. This includes not only morphological analysis, which indicates adaptation to efficient consumption of large prey (10), but also direct contemporary accounts of D. australis consuming penguins and sea lions (11, 12).

In conclusion, while we applaud continued investigations of the human and ecological history of the Falkland Islands, we caution that it is unparsimonious to conclude from current stable isotope data that *D. australis* had a mutualistic relationship with humans and therefore arrived in the islands by human agency. Equally, we acknowledge that both hypotheses remain open and deserving of testing further.

REFERENCES AND NOTES

- J. J. Austin, J. Soubrier, F. J. Prevosti, L. Prates, V. Trejo, F. Mena, A. Cooper, The origins of the enigmatic Falkland Islands wolf. *Nat. Commun.* 4, 1552 (2013).
- J. Clutton-Brock, Man-made dogs: The origins of the domestic dog, South American canids, and the enigma of the Falkland Island wolf are discussed. *Science* **197**, 1340–1342 (1977).
- K. M. Hamley, J. L. Gill, K. E. Krasinski, D. V. Groff, B. L. Hall, D. H. Sandweiss, J. R. Southon, P. Brickle, T. V. Lowell, Evidence of prehistoric human activity in the Falkland Islands. *Sci. Adv.* 7, eabh3803 (2021).
- R. A. Philpott, R. E. Eshelman, A Fuegian projectile point found on Steeple Jason, Falkland Islands. *Falkl. Islands J.* 8, 46–58 (2004).
- P. C. Buckland, K. J. Edwards, Palaeoecological evidence for possible pre-European settlement in the Falkland Islands. J. Archaeological Sci. 25, 599–602 (1998).
- D. L. Phillips, R. Inger, S. Bearhop, A. L. Jackson, J. W. Moore, A. C. Parnell, B. X. Semmens, E. J. Ward, Best practices for use of stable isotope mixing models in food-web studies. *Can. J. Zool.* 92, 823–835 (2014).
- T. J. Clark, B. Vick, J. Newton, I. Marengo, E. D. Wakefield, A wolf in fox's clothing? Using stable isotopes to quantify ecological replacement. *Conserv. Lett.* 14, e12791 (2021).
- W. H. Oosthuizen, M. A. Meyer, J. H. M. David, N. M. Summers, P. G. H. Kotze,
 S. W. Swanson, P. O. Shaughnessy, Variation in jackal numbers at the Van Reenen Bay seal

The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution NonCommercial License 4.0 (CC BY-NC).

Copyright © 2022

¹Center for Ecosystem Sentinels, Department of Biology, University of Washington, Seattle, WA 98195, USA. ²National Environmental Isotope Facility, Scottish Universities Environmental Research Centre, Rankine Avenue, East Kilbride G75 0QF, UK. ³Department of Geography, Durham University, Lower Mountjoy, South Road, Durham DH1 3LE, UK.

colony with comment on likely importance of jackals as predators. S. Afr. J. Wildl. Res. 27, 26–29 (1997).

- 9. T. Smith, Predation of ringed seal pups (Phoca hispida) by the Arctic fox (Alopex lagopus). *Can. J. Zool.* **54**, 1610–1616 (1976).
- C. Meloro, J. Hunter, L. Tomsett, R. Portela Miguez, F. J. Prevosti, R. P. Brown, Evolutionary ecomorphology of the Falkland Islands wolf *Dusicyon australis*. *Mamm. Rev.* 47, 159–163 (2017).
- 11. R. Philpott, Investigation of the Site of Discovery of Remains of the Falklands Wolf at Spring Point, West Falkland, February–March 2011 (Falkland Islands Museum and National Trust, 2011).
- 12. R. Gallagher, Byron's Journal of His Circumnavigation (Hakluyt Society, 1964).

Acknowledgments

Funding: The authors acknowledge that they received no funding in support of this work. Author contributions: All authors contributed equally to this paper. Competing interests: The authors declare that they have no competing interests. Data and materials availability: No data/materials are needed to evaluate the conclusions presented in this paper.

Submitted 12 January 2022 Accepted 14 March 2022 Published 29 April 2022 10.1126/sciadv.abo0928