

Clinical Study

Comparative Approach to Define Increased Regulatory T Cells in Different Cancer Subtypes by Combined Assessment of CD127 and FOXP3

Marc Beyer,¹ Sabine Classen,¹ Elmar Endl,² Matthias Kochanek,³ Martin R. Weihrauch,³ Svenja Debey-Pascher,¹ Percy A. Knolle,² and Joachim L. Schultze³

¹Laboratory for Genomics and Immunoregulation, LIMES-Institute, University of Bonn, Carl-Troll-Street 31, 53115 Bonn, Germany

²Institute for Molecular Medicine and Experimental Immunology, University of Bonn, 53105 Bonn, Germany

³Clinic I for Internal Medicine, University of Cologne, Kerpenerstr. 62, 50924 Cologne, Germany

Correspondence should be addressed to Marc Beyer, marc.beyer@uni-bonn.de

Received 26 May 2011; Accepted 29 June 2011

Academic Editor: D. Craig Hooper

Copyright © 2011 Marc Beyer et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In recent years an increase of functional CD4⁺CD25⁺ regulatory T cells (T_{reg} cells) has been established for patients with solid tumors, acute leukemias, and lymphomas. We have reported an expanded pool of CD4⁺CD25^{high} T_{reg} cells in patients with chronic lymphatic leukemia (CLL), multiple myeloma (MM) as well as its premalignant precursor monoclonal gammopathy of undetermined significance (MGUS). In healthy individuals, low-level expression of CD127 on T cells in addition to the expression of FOXP3 has been associated with T_{reg} cells. Here, we demonstrate that the expanded FOXP3⁺ T-cell population in patients with colorectal cancer, CLL, MGUS, MM, follicular lymphoma, and Hodgkin's disease are exclusively CD127^{low} T_{reg} cells and were strongly suppressive. A significant portion of CD127^{low}FOXP3⁺ T_{reg} cells expressed only low levels of CD25 suggesting that the previously reported expansion of CD25⁺ T_{reg} cells underestimates the true expansion. The assessment of CCR7 and CD45RA expression on the expanded CD4⁺CD127^{low}FOXP3⁺ T_{reg} cells revealed an increase of both naïve as well as central and effector memory T_{reg} cells in peripheral blood. Our data strongly support superiority of combined CD127 and FOXP3 analysis in comparison to CD25 and FOXP3 assessment for further quantification of T_{reg} cells in malignant diseases.

1. Introduction

CD4⁺CD25⁺ regulatory T cells (T_{reg} cells) are expanded in murine tumor models, and their deletion can lead to complete tumor regression [1]. In humans, T_{reg} cells are mostly enriched in the CD4⁺CD25^{high} T-cell population [2]. We and others have reported increased frequencies of CD4⁺CD25^{high}FOXP3⁺ T_{reg} cells in cancer patients [1, 3]. However, the expansion of T_{reg} cells based on the assessment of CD25 is likely to underestimate the true expansion since FOXP3⁺ T cells are also present in the CD25^{-/low} fraction [4, 5]. Furthermore, molecular and functional characterization of this population is hampered by the inability to separate CD25⁺ T_{reg} cells from activated effector T cells. Two recent studies, however, have shown that reciprocal expression of the IL7 receptor (CD127) on FOXP3⁺ T_{reg} cells is most likely

a more specific way to quantify FOXP3⁺ T_{reg} cells [5, 6]. This has been adopted lately for the quantification of T_{reg} cells in solid tumors [7–10] and hematologic malignancies [11–13], with one of the reports establishing CD127 as an even superior marker for the identification of T_{reg} cells in cancer patients [9]. However, no systematic analysis has been undertaken to establish CD127 as a superior marker for T_{reg}-cell enumeration in cancer patients, and only one initial report of malignant melanoma patients has addressed reciprocal expression of CD127 and FOXP3 on T_{reg} cells in cancer patients independently of CD25 [9]. It is, therefore, necessary to determine whether CD127 is also a better marker for enumerating FOXP3⁺ T_{reg} cells in cancer patients in general by comparing T_{reg} cell numbers in a larger number of different tumor subtypes. Besides the integration of CD25^{low/-} FOXP3-expressing T_{reg} cells, analysis of CD127

might, furthermore, clarify contradictory results concerning frequencies as well as prognostic value of T_{reg} cells in cancer patients [14–16].

Similarly, there is still debate whether human $CD4^+CD25^{high}FOXP3^+ T_{reg}$ solely belong to the memory T-cell compartment [17]. Valmori et al. were the first to identify a T_{reg} -cell population with a naïve phenotype ($CCR7^+CD45RA^+$), which they termed natural naïve T_{reg} cells [18]. As expected, the frequency of these naïve T_{reg} cells was relatively low in healthy individuals [19]. More recently, Seddiki et al. have described the persistence of a population of naïve $CD45RA^+ T_{reg}$ cells in adult life [20], which was further characterized by resistance to CD95L-induced cell death [21]. Recent data further supports that a population of naïve T_{reg} cells exist in healthy individuals that exerts suppressive function [22]. So far, our own observations suggested an increased frequency of naïve $CD4^+CD25^{high}FOXP3^+ T_{reg}$ cells in MM and MGUS [23]. However, previous findings were restricted to the $CD4^+CD25^{high}$ subpopulation excluding a significant fraction of T_{reg} cells from analysis. With the emergence of CD127 as a new marker separating T_{reg} cells from conventional T cells, the question whether the expanded T_{reg} cells in cancer patients are mainly antigen-experienced memory cells or also naïve T_{reg} cells needs reevaluation.

Here, we present clear evidence that $FOXP3^+$ T cells derived from patients with CLL, MGUS, MM, follicular lymphoma (FL), Hodgkin's disease (HD), and colorectal cancer (CRC) are lacking CD127. This newly defined fully functional $CD4^+CD127^{low}FOXP3^+ T_{reg}$ -cell population is expanded in all tumor entities as well as the premalignant MGUS supporting the hypothesis of increased T_{reg} cells as a rather early event during tumor development. Moreover, we demonstrate a significant increase of naïve $CD4^+CD127^{low}FOXP3^+ T_{reg}$ cells in peripheral blood of cancer patients while we could not detect an increase in lymph node biopsies of lymphoma patients. Finally, these data strongly support the assessment of CD127 expression—instead of CD25—in combination with FOXP3 for a more precise enumeration of T_{reg} cells in malignant diseases while functional characterization still relies on the combination of CD127 and CD25.

2. Material and Methods

2.1. Patients and Clinical Parameters. Following approval by the institutional review board of the University of Cologne, peripheral blood from 10 healthy individuals, 7 MGUS, 10 MM, 10 CLL and 6 patients with CRC (2 time points at least 1 month apart) was obtained after informed consent. For the assessment of T_{reg} cell numbers in lymph node biopsies, lymph nodes from 7 healthy donors, 6 patients with HD, and 7 patients with FL were analyzed following approval by the institutional review board of the University of Cologne. Patients were either untreated or had not received cytoreductive treatment for a period of at least 1 month prior to investigation. Characteristics of the patients studied are summarized in Tables 1 and 2.

2.2. Antibodies and FACS Analysis. Phenotype of T cells was defined by flow cytometry using the following antibodies: CD45RA-PE-Cy5 (HI100), CD127-PE (hIL-7R-M21), CD4-APC-Cy7 (RPA-T4), CD25-PE-Cy7 (M-A251, all from Becton Dickinson), CCR7-FITC (150503, R&D) as well as the corresponding isotype control antibodies. Intracellular staining for FOXP3 was performed with FOXP3-APC (PCH101, eBioscience) according to the manufacturer's recommendations [23]. Samples were acquired on a FACS LSR II and analyzed with FlowJo software (TreeStar Inc). Frequencies of $CD4^+CD127^{low}FOXP3^+$ T cells are shown as percent values of $CD4^+$ T cells.

2.3. Isolation of $CD4^+CD127^{low}CD25^{+/-}$ and $CD4^+CD127^+CD25^-$ T Cells and Assessment of Inhibitory Function. To assess the suppressive activity of $CD4^+CD127^{low}CD25^{+/-}$ T cells, a modified MLR was performed as previously described [23]. Briefly, $CD4^+CD127^{low}CD25^{+/-}$ and $CD4^+CD127^+CD25^-$ T cells were stained with CD4, CD25, and CD127 mAb and sorted on a FACSDiVa or FACS Aria III (both BD Biosciences) and incubated for 20 hours with 10 U/mL IL-2 (Proleukin) and 0.5 μ g/mL anti-CD3 (OKT3) in X-VIVO 15 (BioWhittaker) [24]. Subsequently, $CD4^+CD127^+CD25^-$ T cells were stained with 5,6-Carboxyfluorescein-Diacetat-Succinimidyl-Ester (CFSE, Sigma-Aldrich) and stimulated in X-VIVO 15 supplemented with 10% fetal calf serum, 100 U/mL penicillin/streptomycin and 2 mM glutamine (Invitrogen) with magnetic beads (DynaL Biotech,) coated with 5% anti-CD3, 14% anti-CD28 (9.3), and 81% anti-MHC class I (W6/32) at a ratio of 3:1 (cells:beads). To assess inhibitory capacity of T_{reg} cells from cancer patients, autologous $CD4^+CD127^{low}CD25^{+/-}$ T_{reg} cells were added at a 1:1 ratio to the culture, and the proliferation of $CD4^+CD127^+CD25^-$ T cells was determined by assessing CFSE dilution after four days of culture as described previously [23].

2.4. RNA Preparation and Quantitative Real-Time PCR. For analysis of CD127 mRNA expression, $CD4^+CD25^-$ and $CD4^+CD25^{high}$ T cells from five healthy donors and five CLL patients were purified as previously described [24]. The described technique is optimized for the isolation of human $CD4^+CD25^{high}$ T cells with high purity [23, 24]. Cells were reanalyzed after sorting and routinely showed >95% purity. Subsequently, the cells were lysed in TRIzol reagent (Invitrogen). 50–100 ng RNA were reverse transcribed using the Transcriptor First Strand cDNA Synthesis Kit (Roche, Penzberg, Germany). Rt-PCR was performed with the LightCyclerTaqman master kit and Universal ProbeLibrary Assay on a Light Cycler 480 II. Analysis was performed using Light-Cycler3 and RelQuant software using a calibrator normalized relative quantification based on β -2 microglobulin (B2M) expression. Primers used: CD127 forward, 5'-AAAGTTTAAATGCACGATGTAGCTT-3'; CD127 reverse, 5'-TGTGCTGGATAAATTCACATGC-3'; Probe 72; B2M forward, 5'-TTCTGGCCTGGAGGCTAT-3'; B2M reverse, 5'-TCAGGAAATTTGACTTTCCATTC-3'; Probe 42.

TABLE 1: Patient characteristics for T_{reg}-cell assessment in peripheral blood.

(a) MGUS (peripheral blood)						
ID	Gender	Age (yr)	Stage	Therapy	Paraprotein	T _{reg} (%)
1	M	37	MGUS	no tx	IgG/λ	3.6
2	F	84	MGUS	no tx	IgA/κ	7.0
3	F	86	MGUS	no tx	IgG/κ	7.5
4	F	62	MGUS	no tx	IgA	6.0
5	M	52	MGUS	no tx	n.a.	6.8
6	F	79	MGUS	no tx	n.a.	2.4
7	M	65	MGUS	no tx	IgM/κ	8.0
(b) Multiple myeloma (peripheral blood)						
ID	Gender	Age (yr)	Stage	Therapy	Paraprotein	T _{reg} (%)
8	M	62	MM I A	M	IgG/κ	16.5
9	F	39	MM I A	VAD, CAD, M, BMT	IgG/κ	7.4
10	M	59	MM II A	no tx	IgG/κ	10.2
11	F	74	MM I A	no tx	IgG/λ	4.5
12	M	86	MM I A	no tx	IgG/κ	6.7
13	M	60	MM II A	VAD, CAD, M, BMT	IgG/κ	16.9
14	F	52	MM II A	VID, C	IgG/κ	4.4
15	M	59	MM II A	TAD, CAD, M, BMT	IgG/κ	8.6
16	F	67	MM II A	VAD, CAD, M, BMT	IgG/κ	22.2
17	M	53	MM II A	no tx	IgG/λ	8.0
(c) Chronic lymphocytic leukemia (peripheral blood)						
ID	Gender	Age (yr)	Stage	Therapy	T _{reg} (%)	
18	m	72	CLL A	no tx	6.1	
19	m	62	CLL A	no tx	5.7	
20	f	73	CLL A	no tx	5.6	
21	m	60	CLL B	no tx	9.8	
22	f	55	CLL B	no tx	10.6	
23	m	73	CLL B	no tx	16.4	
24	m	64	CLL B	no tx	7.4	
25	m	39	CLL C	no tx	15.3	
26	m	60	CLL C	no tx	6.2	
27	m	54	CLL C	no tx	6.4	
(d) Colorectal cancer (peripheral blood)						
ID	Gender	Age (yr)	Stage	Primary tumor	Sites of metastases	T _{reg} (%)
28	f	43	D	Rectum	Liver, bone, pararectal, para-aortal lymph nodes	7.5/10.4
29	f	32	D	Colon	Liver, spleen, ovaries, pelvis, peritoneum	5.7/9.3
30	m	57	D	Rectum	Lung	5.7/9.9
31	m	66	D	Colon	Liver	5.0/6.2
32	m	75	D	Colon	Liver	9.0/4.8
33	f	44	D	Colon	Liver	9.9/3.3
(e) Healthy donors (peripheral blood)						
ID	Gender	Age (yr)	T _{reg} (%)			
34	m	66	5.4			
35	m	67	6.6			
36	m	55	3.9			
37	m	50	6.5			
38	m	47	4.5			
39	m	46	3.5			

(e) Continued.

ID	Gender	Age (yr)	T _{reg} (%)
40	m	46	4.8
41	m	62	4.1
42	m	45	2.6
43	f	44	4.0

Patient characteristics including gender, age at analysis, Durie and Salmon, Binet or Dukes stage, first diagnosis, primary tumor, sites of metastases, therapy, paraprotein, and frequency of T_{reg} cells. (f: female, m: male; therapy: A: Doxorubicin, BMT: autologous bone-marrow transplantation, C: Cyclophosphamide, I: Idarubicin, M: Melphalan, V: Vincristine, D: Prednisone, T: Thalidomide, no tx: no therapy, n.a.: not accessible.)

TABLE 2: Patient characteristics for T_{reg}-cell assessment in lymph node biopsies.

(a) Follicular lymphoma (lymph node)

ID	Gender	Age (yr)	Stage	Therapy	T _{reg} (%)
44	m	59	FL I	no tx	23.9
45	m	46	FL I	no tx	13.4
46	f	58	FL I	no tx	26.8
47	m	73	FL II	no tx	19.6
48	f	66	FL II	no tx	20.0
49	m	59	FL II	no tx	13.2
50	m	57	FL II	no tx	22.5
51	m	65	FL II	no tx	37.5

(b) Hodgkin's disease (lymph node)

ID	Gender	Age (yr)	Entity	Therapy	T _{reg} (%)
52	f	53	HD (ns)	no tx	37.8
53	m	44	HD (ns)	no tx	16.8
54	m	51	HD (ns)	no tx	43.9
55	m	19	HD (ns)	no tx	18.5
56	m	34	HD (ns)	no tx	16.9
57	f	25	HD (mc)	no tx	12.4

(c) Healthy donors (reactive lymph nodes)

ID	Gender	Age (yr)	T _{reg} (%)
58	m	35	4.9
59	f	18	11.7
60	f	17	4.9
61	m	22	9.7
62	f	45	16.7
63	m	39	8.9
64	m	24	14.0

Patient characteristics including gender, age at analysis, first diagnosis, therapy, and frequency of T_{reg} cells. (f: female, m: male; no tx: no therapy; mc: mixed cellularity; ns: nodular sclerosing).

2.5. Statistical Analysis. Comparison between paired or unpaired groups was performed using the appropriate Student's *t*-test. A *P*-value < 0.05 was defined as statistically significant. Due to the explorative nature of this study, no multiplicity adjustment procedures were performed. All statistical analyses were performed using the SPSS statistical software package (SPSS 19.0, SPSS Inc.).

3. Results

3.1. Downregulation of CD127 mRNA Expression in CD4⁺ CD25^{high} T_{reg} Cells from CLL Patients. As CD25 is not solely

expressed on T_{reg} cells but also on activated conventional CD4⁺ T cells, and the downregulation of CD127 expression in CD4⁺CD25^{high} FOXP3⁺ T_{reg} cells from healthy donors has been reported [5, 6], we first assessed if CD127 downregulation is also apparent in CD4⁺CD25^{high} T_{reg} cells from cancer patients. We detected a significant downregulation of CD127 mRNA expression in CD4⁺CD25^{high} T cells from healthy donors (*n* = 5) as well as CLL patients (*n* = 5, *P* < 0.05, Figure 1(a)) by quantitative PCR indicating that CD127 expression might also be used to specifically identify CD4⁺FOXP3⁺ T_{reg} cells in cancer patients.




FIGURE 1: CD127 mRNA expression in CD4⁺CD25^{high} T_{reg} cells and integration of CD127 in the analysis of human T_{reg} cells. (a) Expression of CD127 mRNA in CD4⁺CD25^{high} T_{reg} cells and conventional CD4⁺CD25⁻ T cells in healthy donors ($n = 5$, control) and CLL patients ($n = 5$, CLL) as determined by qPCR (*, $P < 0.05$, Student's t -test). (b) Gating strategies for analysis of expression of CD127 in CD4⁺CD25^{high}FOXP3⁺ T_{reg} cells or (c) CD25 expression in CD4⁺CD127^{low}FOXP3⁺ T_{reg} cells.

3.2. Coassessment of CD127 and FOXP3 for the Enumeration of Human T_{reg} Cells. Next the expression of CD127 in relation to FOXP3 and CD25 was evaluated by flow cytometry on CD4⁺ T cells. Gating on CD4 and CD25 with subsequent analysis of the CD4⁺CD25^{high} T_{reg}-cell population for expression of FOXP3 and CD127 confirmed the downregulation of CD127 in CD4⁺CD25^{high}FOXP3⁺ T_{reg} cells on protein level in healthy individuals (Figure 1(b)). However, assessing coexpression of CD127 and FOXP3 by CD4⁺ T cells without gating beforehand on the CD4⁺CD25^{high} T-cell population clearly revealed a significantly higher percentage of cells expressing FOXP3 but lacking CD127 (Figure 1(c)). Subsequent analysis of the CD127^{low}FOXP3⁺ T_{reg}-cell population for expression of CD25 demonstrated that gating on CD127 and FOXP3 identifies not only CD4⁺CD25^{high} T_{reg} cells but also T_{reg} cells expressing only low levels of CD25 (Figure 1(c)). The identification of this subpopulation of T_{reg} cells is of specific interest as up to now only T_{reg} cells expressing high amounts of CD25 were accessible to functional analysis.

3.3. Increase of CD4⁺CD127^{low}FOXP3⁺ T_{reg} Cells in Cancer Patients. Inclusion of the CD25^{low} T_{reg}-cell subpopulation in the enumeration of T_{reg} cells by defining human T_{reg} cells as CD4⁺CD127^{low}FOXP3⁺ demands the reassessment of T_{reg}-cell frequencies in cancer patients as the actual frequencies were probably underestimated until now. Comparison of healthy individuals with cancer patients revealed elevated levels of CD4⁺CD127^{low}FOXP3⁺ T_{reg} cells in cancer and MGUS patients, as exemplified for individual patients in Figures 2(a) and 2(b). In total, frequencies of T_{reg} cells derived from peripheral blood of 12 patients with CRC, 10 CLL patients, 7 MGUS, and 10 MM patients as well as

10 healthy individuals were evaluated. In addition, lymph node biopsies from 7 patients with follicular lymphoma, 6 patients with Hodgkin's disease, and 7 reactive lymph nodes from healthy individuals were assessed for expanded T_{reg}-cell numbers. Gating on CD4 and CD25 with subsequent gating on FOXP3 confirmed the already described increase of T_{reg} cells in patients with CRC, CLL, MGUS, MM, FL, and HD (Figures 3(a) and 3(b) and Tables 3 and 4). More important, when gating on FOXP3 and CD127 without using CD25 as primary inclusion criteria, frequencies of CD4⁺CD127^{low}FOXP3⁺ T_{reg} cells in controls ($4.1\% \pm 0.7\%$) were similar to previously published results (Figure 3(c) and Table 3) [2, 5, 6, 24]. In contrast, individuals with CRC ($7.2\% \pm 2.4\%$, $P < 0.005$), CLL ($8.9\% \pm 4.0\%$, $P < 0.005$), as well as MM ($11.7\% \pm 5.4\%$, $P < 0.005$) showed significantly increased frequencies of CD127^{low}FOXP3⁺ T_{reg} cells compared to healthy individuals (Figure 3(c) and Table 3). Even in MGUS patients, a significantly higher frequency of T_{reg} cells ($6.0\% \pm 1.8\%$, $P < 0.05$) was observed (Figure 3(c) and Table 3), which is indicative of T_{reg}-cell expansion as an early event in tumorigenesis. Similarly, we observed significantly increased frequencies of CD127^{low}FOXP3⁺ T_{reg} cells in patients with FL ($21.8\% \pm 8.0\%$, $P < 0.01$) and HD ($24.4\% \pm 13.1\%$, $P < 0.05$) in comparison to reactive lymph node specimens from healthy individuals ($10.1\% \pm 4.4$, Figure 3(d) and Table 4). Moreover, the percentage of FOXP3⁺ cells within the CD4⁺CD127^{low} T-cell population was always higher than within the CD4⁺CD25^{high} population, suggesting that previous data only assessing a CD4⁺CD25^{high} phenotype have underestimated the absolute increase of FOXP3⁺ T_{reg} cells in cancer patients (Tables 3 and 4).




FIGURE 2: Frequency of $CD4^+CD127^{low}FOXP3^+$ T_{reg} cells. Flow cytometric analysis of CD127 and FOXP3 expression in $CD4^+$ T cells from (a) peripheral blood of a representative healthy individual (control) and representative patients with colorectal cancer (CRC), CLL, MGUS, and multiple myeloma (MM) and (b) lymph node biopsies from a healthy individual (reactive) and patients with follicular lymphoma (FL) and Hodgkin's diseases (HD).

TABLE 3: Assessment of T_{reg} -cell frequencies in peripheral blood.

	Control	Colon		CLL		MGUS		MM	
	Mean (SD)	Mean (SD)	<i>P</i>	Mean (SD)	<i>P</i>	Mean (SD)	<i>P</i>	Mean (SD)	<i>P</i>
$CD4^+CD127^{low}$	4.6 (1.3)	7.6 (1.5)	<0.001	8.3 (2.5)	<0.005	5.4 (1.3)	n.s.	11.2 (5.9)	<0.01
$CD4^+CD127^{low}FOXP3^+$	4.1 (0.7)	7.2 (2.4)	<0.005	8.9 (4.0)	<0.005	6.0 (1.8)	<0.05	11.7 (5.4)	<0.005
$CD4^+CD127^{low}FOXP3^+CD25^{high}$	2.5 (0.6)	4.3 (1.6)	<0.005	4.7 (2.7)	<0.05	3.9 (1.3)	<0.05	7.1 (4.9)	<0.05
$CD4^+CD25^{high}$	2.8 (0.9)	7.6 (1.2)	<0.001	6.4 (1.8)	<0.001	4.5 (1.1)	<0.05	9.0 (5.3)	<0.01
$CD4^+CD25^{high}CD127^{low}$	2.9 (0.9)	4.5 (1.2)	<0.005	4.5 (2.0)	<0.05	3.5 (1.3)	n.s.	7.0 (5.1)	<0.05
$CD4^+CD25^{high}FOXP3^+$	2.1 (0.8)	4.2 (1.2)	<0.001	3.6 (1.7)	<0.05	2.5 (0.6)	n.s.	6.3 (4.5)	<0.05
$CD4^+FOXP3^+$	2.8 (0.9)	4.7 (2.1)	<0.05	4.7 (2.4)	<0.05	3.6 (1.1)	n.s.	7.7 (5.1)	<0.05

Definition of subpopulations based on expression of CD25, CD127, and FOXP3 (SD: standard deviation, n.s.: not significant).

3.4. $CD4^+CD127^{low}CD25^{+/-} T_{reg}$ Cells are Fully Functional in Cancer Patients. As intracellular FOXP3 staining is not applicable for functional analysis of T_{reg} cells, classification of $FOXP3^+$ T_{reg} cells based solely on cell surface markers is necessary. The characterization of $FOXP3^+$ T_{reg} cells was best achieved when combining CD127 and CD25 (Figures 4(a) and 4(b)). We, therefore, used this combination of cell surface markers to sort T_{reg} cells for functional analysis. Staining for FOXP3 expression after sorting routinely showed purities of $CD4^+CD127^{low}FOXP3^+CD25^{+/-}$ T_{reg} cells >95 percent (Figure 4(b)). To determine whether the $CD4^+CD127^{low}CD25^{+/-}$ T_{reg} cells from cancer patients are functional, we used an *in vitro* suppression assay. When activated with CD3/CD28 beads conventional $CD4^+CD127^+CD25^-$ T cells, but not $CD4^+CD127^{low}CD25^{+/-}$ T_{reg} cells, proliferate strongly. In the presence of $CD4^+CD127^{low}CD25^{+/-}$ T_{reg} cells, this proliferation is suppressed (Figure 4(c)). These data clearly demonstrate that

$CD4^+CD127^{low}CD25^{+/-}$ T cells are $FOXP3^+$ and that these cells are fully functional in CRC patients.

3.5. Naïve $CD4^+CD127^{low}FOXP3^+$ T_{reg} Cells are Increased in Peripheral Blood of Cancer Patients. In healthy individuals, T_{reg} cells have been shown to exist at all differentiation states, namely, naïve, central, and effector memory T_{reg} cells [18, 20, 25]. To determine which T_{reg} -cell subpopulation is responsible for the increase of $CD4^+CD127^{low}FOXP3^+$ T_{reg} cells in cancer patients, we determined the frequency of naïve, central, and effector memory cells within the T_{reg} -cell compartment from healthy individuals, CRC, CLL, MGUS, and MM patients (Figure 5(a)) and compared these data with those previously described for $CD4^+CD25^{high}$ T_{reg} cells in healthy individuals as well as MGUS and MM patients [18, 23]. In healthy individuals, naïve $CCR7^+CD45RA^+CD4^+CD127^{low}FOXP3^+$ T_{reg} cells were hardly detectable (Figures 5(b) and 5(c)). T_{reg} cells were




FIGURE 3: Assessment of T_{reg}-cell frequencies. Frequency of CD4⁺CD25^{high}FOXP3⁺ T_{reg} cells in (a) peripheral blood of 10 healthy donors (control), 12 colorectal cancer (CRC), 10 CLL, 7 MGUS, and 10 multiple myeloma (MM) patients and (b) 7 reactive lymph node biopsies from healthy individuals (reactive), 7 patients with follicular lymphoma (FL), and 6 patients with Hodgkin's disease (HD). (c) and (d) Frequencies of CD4⁺CD127^{low}FOXP3⁺ T_{reg} cells in the respective groups. Error bars represent standard deviation (*, $P < 0.05$, Student's t test).

almost exclusively of memory phenotype (Figures 5(b) and 5(c)). In contrast, in peripheral blood of CRC, CLL, and MM patients, a significant expansion of CD4⁺CD127^{low}FOXP3⁺ T_{reg} cells with a naïve phenotype was observed (Figures 5(b) and 5(c)). The expansion of naïve T_{reg} cells was apparent as part of the T_{reg}-cell pool as well as in relation to the total number of CD4⁺ T cells in cancer patients. This increase in naïve T_{reg} cells was further accompanied by an expansion of T_{reg} cells with a central as well as

effector memory phenotype in all patient groups (Figures 5(b) and 5(c)). Interestingly, the observed expansion of naïve CD4⁺CD127^{low}FOXP3⁺ T_{reg} cells was also detectable in MGUS patients (Figures 5(b) and 5(c)) further underlining that frequencies of naïve T_{reg} cells increase rather early during tumor development and progression. When assessing subpopulations of T_{reg} cells in lymph node specimens, we observed a predominance of CD4⁺CD127^{low}FOXP3⁺ T_{reg} cells with a central-memory phenotype, with a significantly

TABLE 4: Assessment of T_{reg} -cell frequencies in lymph node biopsies.

	Control	FL		HD	
	Mean (SD)	Mean (SD)	<i>P</i>	Mean (SD)	<i>P</i>
$CD4^+CD127^{low}$	54.7 (23.1)	68.6 (15.8)	>0.05	68.0 (12.7)	>0.05
$CD4^+CD127^{low}FOXP3^+$	10.1 (4.4)	21.8 (8.0)	<0.01	24.4 (13.1)	<0.05
$CD4^+CD127^{low}FOXP3^+CD25^{high}$	3.1 (1.9)	11.7 (5.5)	<0.005	6.4 (2.3)	<0.05
$CD4^+CD25^{high}$	5.1 (2.9)	13.8 (7.1)	<0.05	11.2 (3.3)	<0.005
$CD4^+CD25^{high}CD127^{low}$	4.1 (3.0)	16.3 (7.1)	<0.005	11.3 (5.2)	<0.05
$CD4^+CD25^{high}FOXP3^+$	2.9 (2.6)	10.0 (5.4)	<0.01	5.6 (1.2)	<0.05
$CD4^+FOXP3^+$	10.3 (5.5)	19.4 (8.5)	<0.05	23.4 (12.0)	<0.05

Definition of subpopulations based on expression of CD25, CD127, and FOXP3 (SD: standard deviation, n.s.: not significant).




FIGURE 4: Functional analysis of $CD4^+CD127^{low}CD25^+$ T_{reg} cells in cancer patients. (a) Sorting strategy for isolating $CD4^+CD127^{low}CD25^+$ T_{reg} cells (T_{reg}) as well as conventional $CD4^+CD127^+CD25^-$ T cells (T_{conv}). (b) Expression of FOXP3 in the corresponding T-cell populations. (c) Percentage of proliferation of $CD4^+CD25^-CD127^+$ T_{conv} cells (black bar) alone or cultivated with $CD4^+CD127^{low}CD25^+$ T_{reg} cells derived from CRC patients ($n = 4$) at a 1 : 1 ratio (white bar) both in the presence of CD3/CD28 mAb coated beads. Error bars represent standard deviation (*, $P < 0.05$, Student's *t* test).

expanded population of central-memory T_{reg} cells apparent in patients with FL and HD (Figure 5(d)). In addition, we could also detect an increase in effector-memory T_{reg} cells (Figure 5(d)) while the pool of naïve T_{reg} cells was basically absent independent if reactive or diseased lymph nodes were analyzed (Figure 5(d)).

4. Discussion

Expansion of $CD4^+CD25^{high}$ T_{reg} cells within the tumor microenvironment and peripheral blood has so far been accepted as a hallmark of cancer [1, 26, 27]. Moreover, augmented T_{reg} -cell frequencies have been linked to tumor stage, prognosis, and survival [1, 26, 27]. We present new evidence that the increase of T_{reg} cells in cancer was even underestimated previously due to suboptimal classification of T_{reg} cells. Integrating analysis of FOXP3 with the cell-surface molecule CD127 clearly demonstrates that significantly higher numbers of $CD127^{low}FOXP3^+$ T_{reg} cells are expanded in cancer patients in general. The assessment of

CD127 instead of CD25 is clearly superior in enumerating T_{reg} cells in the diseased state.

Natural T_{reg} cells have been described as $CD4^+CD25^+$ T cells in mice [28], and initial reports in cancer patients relied solely on the assessment of CD4 and CD25 expression for the identification of T_{reg} cells [3, 29]. Only since the identification of the transcription factor FOXP3 lineage-specific marker of T_{reg} cells a more specific characterization of T_{reg} cells is possible [28]. In murine models, FOXP3 expression is strongly associated with the $CD25^+$ T_{reg} -cell population. However, even the inclusion of FOXP3 assessment has been interpreted differentially when assessing frequencies of T_{reg} cells in healthy individuals and cancer patients [23, 30]. The analysis of T_{reg} cells in humans has been further complicated as several studies reported FOXP3⁺ cells within the $CD4^+CD25^{low}$ or even $CD4^+CD25^-$ population [5], and even the reprogramming of T_{reg} cells into effector T cells has been reported [31]. Therefore, a more specific definition of T_{reg} cells based on unique or additional T_{reg} -cell marker molecules is urgently needed. The introduction of CD127 as a new marker to distinguish T_{reg} cells from




FIGURE 5: Assessment of naive CCR7⁺CD45RA⁺CD4⁺CD127^{low}FOXP3⁺ T_{reg} cells. (a) Frequencies of CCR7⁺CD45RA⁺ naive CD4⁺CD127^{low}FOXP3⁺ T_{reg} cells (T_{naive}), CCR7⁺CD45RA⁻CD4⁺CD127^{low}FOXP3⁺ central memory T_{reg} cells (T_{CM}), and CCR7⁻CD45RA⁻CD4⁺CD127^{low}FOXP3⁺ effector memory T_{reg} cells (T_{EM}) were assessed in peripheral blood using gating on CD127 and FOXP3 with successive gating on CCR7 and CD45RA. (b) Flow cytometric analysis of naive, central memory, and effector memory CD4⁺CD127^{low}FOXP3⁺ T_{reg} cells in peripheral blood from a representative healthy individual (control) as well as representative patients with colorectal cancer (CRC), CLL, MGUS, and multiple myeloma (MM). Frequencies of regulatory T_{naive}, T_{CM}, and T_{EM} cells were assessed in (c) peripheral blood of CRC (CRC, *n* = 12), CLL (CLL, *n* = 10), MGUS (MGUS, *n* = 7), MM (MM, *n* = 10), and healthy individuals (control, *n* = 10) and (d) 7 reactive lymph node biopsies from healthy individuals (reactive), 7 patients with follicular lymphoma (FL), and 6 patients with Hodgkin's disease (HD). Error bars represent standard deviation (*, *P* < 0.05, Student's *t*-test).

conventional T cells is an important improvement and will help to clarify several previous conflicting results in human T_{reg}-cell biology, particularly in cancer patients.

Several recent studies have adopted the approach to use CD127, CD25, and FOXP3 for the quantification of T_{reg} cells in tumor-bearing individuals and could demonstrate increased numbers of CD4⁺CD25^{high}CD127^{low} T_{reg} cells in patients with solid tumors [7–10] and hematologic malignancies [11–13]. However, the majority of these reports

focused solely on the enumeration of the T_{reg}-cell compartment while at the same time focusing on only one tumor subtype. Only one study assessed T_{reg}-cell numbers in more than one tumor subtype showing similar numbers of T_{reg} cells for all gastrointestinal tumor subtypes analyzed [8]. Furthermore, these studies did not systematically compare possible marker combinations to establish the most suitable approach to identify T_{reg} cells. This was analyzed in more detail in only one of the reports with the combination of CD127 and

FOXP3 being the most appropriate combination to identify T_{reg} cells in patients with malignant melanoma [9].

The integration of CD127 permits to redefine the importance of CD25 expression on human T_{reg} cells. Up to now, high expression of CD25 allowed for an enrichment of $CD4^+$ T cells with regulatory properties [2]. However, it is undisputed that neither all human T_{reg} cells are included by this approach nor that activated T cells expressing CD25 are excluded. Zelenay et al. could demonstrate a population of $CD4^+CD25^-FOXP3^+$ T cells which can upregulate CD25 upon the depletion of all CD25 expressing cells and are able to replace the original T_{reg} -cell population [4]. These data were a first hint that the expression of CD25 on T_{reg} cells is similarly regulated like its expression on conventional T cells [4]. Human T_{reg} cells need IL-2 for their survival and proliferation, and expression of the IL2R- α chain is certainly a prerequisite for IL-2 to exert its biological function [32]. However, the expression of CD25 is not homogenous and might also be dependent on the activation status and other exogenous factors [33].

Using CD127 and FOXP3 to define human T_{reg} cells demonstrates varying expression of CD25 in the $CD4^+CD127^{low}FOXP3^+$ T_{reg} -cell population. Additionally, the newly defined T_{reg} -cell population comprises of significantly more T_{reg} cells compared to the traditionally defined $CD4^+CD25^{high}$ T_{reg} cells as demonstrated recently for malignant melanoma [9]. Coassessment of CD127 and FOXP3 to determine T_{reg} cells also resolves the uncertainty to differentiate between activated conventional T cells and T_{reg} cells in patients with active disease. This is of particular importance when only using CD4 and CD25 for the identification of T_{reg} cells in cancer patients, as contamination with effector T cells most frequently occurs when solely these two markers are used for analysis. As functional assessment of the $CD4^+CD127^{low}FOXP3^+$ T_{reg} -cell population is not possible as FOXP3 cannot be used for live studies of human T_{reg} cells, using expression of CD4, CD25, and CD127 is the best possible approximation. T cells isolated by this approach almost exclusively express FOXP3. Moreover, when isolated from cancer patients, this T_{reg} -cell population exerts strong inhibition.

Using a comparative approach analyzing different tumor subtypes from hematologic as well as epithelial origin, we demonstrate that all independent cancer patient groups studied uniformly show an expanded pool of $CD4^+CD127^{low}FOXP3^+$ T_{reg} cells. We therefore postulate that expansion of T_{reg} cells is a general phenomenon in cancer patients. Moreover, since MGUS patients already have increased frequencies of T_{reg} cells, it is very likely that expansion of T_{reg} cells is an early event in the development of human tumors. Elevated T_{reg} -cell levels might be associated with the progression from premalignant lesions that are still under control of the immune system to the uninhibited growth of malignant tumors.

The findings that naïve T_{reg} cells are increased both in the premalignant state as well as in cancer patients might further support this hypothesis. T_{reg} cells were first identified as antigen-experienced memory cells expressing CD45RO [2]. Only recently the existence of naïve T_{reg} cells in human adults

has been reported [18, 20, 22, 23, 34], and the naïve T_{reg} -cell population can be expanded *in vitro* while retaining its suppressive function [35, 36]. However, the physiological function of the naïve T_{reg} -cell population remains unclear. Definition of T_{reg} cells as $CD4^+CD127^{low}FOXP3^+$ has enabled us to verify the increase of naïve T_{reg} cells in MGUS and MM patients [23] and to extend these findings to CLL and CRC.

The identification of an expanded pool of naïve T_{reg} cells in cancer patients opens new avenues to better understand the role of T_{reg} cells in malignant disease. Memory T_{reg} cells apparently cannot undergo self-renewal [37]. Therefore, the replenishment of an increased memory T_{reg} -cell pool by differentiation of naïve T_{reg} cells into memory T_{reg} cells might be an alternative to the recently proposed conversion of conventional memory T cells to T_{reg} cells [37]. In fact, the increased pool of naïve T_{reg} cells with an unaltered frequency of memory T_{reg} cells in premalignant MGUS suggests that expansion of naïve T_{reg} cells is indeed preceding the expansion of memory T_{reg} cells following differentiation during tumor development. Besides the expansion of naïve T_{reg} cells through enhanced self-renewal and differentiation, other mechanisms have been proposed amongst them the interaction of CCR4 on T_{reg} cells with CCL22 released in the tumor microenvironment [38] as well as the conversion of conventional $CD4^+CD25^-$ T cells to T_{reg} cells through TGF- β [39] or prostaglandin E_2 [40]. How these factors influence the expansion of naïve T_{reg} cells needs further clarification and might in the end result in better strategies to target expanded T_{reg} cells in tumor patients.

In conclusion this study demonstrates that $CD4^+CD127^{low}FOXP3^+$ T_{reg} cells are increased in cancer patients. Definition of T_{reg} cells by combining CD127 and FOXP3 has the advantage of including not only T_{reg} cells expressing high levels of CD25 but also T_{reg} cells with low CD25 expression and excluding at the same time activated conventional T cells. Furthermore, the naïve T_{reg} -cell population is expanded in all tumor bearing individuals.

Abbreviations

CLL:	Chronic lymphatic leukemia;
MM:	Multiple myeloma
MGUS:	Monoclonal gammopathy of undetermined significance
CRC:	Colorectal cancer
FL:	Follicular lymphoma
HD:	Hodgkin's disease
FOXP3:	Forkhead box protein 3
T_{reg} cells:	Regulatory T cells.

Acknowledgments

The authors are indebted to our patients for their commitment to this study. They thank I. Büchmann for excellent technical assistance, A. Dolf for cell sorting, K.-H. Grips and D. Gerecke for referral of patients, C. June and J. Riley for providing them with the 9.3 antibody, and B. Gathof and the Division of Transfusion Medicine of the University of Cologne for providing them with blood samples

from healthy individuals. This work was supported by a Sofja Kovalevskaja Award of the Alexander von Humboldt-Foundation (JLS), the Wilhelm-Sander Stiftung (JLS), the German Jose-Carreras Foundation (MB & JLS), and the German Research Foundation (SFB704: JLS, EE; PAK, SFB832: MB & JLS, INST 217/576-1: JLS, INST 217/577-1: JLS).

References

- [1] M. Beyer and J. L. Schultze, "Regulatory T cells in cancer," *Blood*, vol. 108, no. 3, pp. 804–811, 2006.
- [2] C. Baecher-Allan, J. A. Brown, G. J. Freeman, and D. A. Hafler, "CD4⁺ CD25^{high} regulatory cells in human peripheral blood," *Journal of Immunology*, vol. 167, no. 3, pp. 1245–1253, 2001.
- [3] E. Y. Woo, C. S. Chu, T. J. Goletz et al., "Regulatory CD4⁺ CD25⁺ T cells in tumors from patients with early-stage non-small cell lung cancer and late-stage ovarian cancer," *Cancer Research*, vol. 61, no. 12, pp. 4766–4772, 2001.
- [4] S. Zelenay, T. Lopes-Carvalho, I. Caramalho, M. F. Moraes-Fontes, M. Rebelo, and J. Demengeot, "Foxp3⁺ CD25⁻ CD4 T cells constitute a reservoir of committed regulatory cells that regain CD25 expression upon homeostatic expansion," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 102, no. 11, pp. 4091–4096, 2005.
- [5] W. Liu, A. L. Putnam, Z. Xu-yu et al., "CD127 expression inversely correlates with FoxP3 and suppressive function of human CD4⁺ T reg cells," *Journal of Experimental Medicine*, vol. 203, no. 7, pp. 1701–1711, 2006.
- [6] N. Seddiki, B. Santner-Nanan, J. Martinson et al., "Expression of interleukin (IL)-2 and IL-7 receptors discriminates between human regulatory and activated T cells," *Journal of Experimental Medicine*, vol. 203, no. 7, pp. 1693–1700, 2006.
- [7] L. S. Shen, J. Wang, D. F. Shen et al., "CD4⁺ CD25⁺ CD127^{low/-} regulatory T cells express Foxp3 and suppress effector T cell proliferation and contribute to gastric cancers progression," *Clinical Immunology*, vol. 131, no. 1, pp. 109–118, 2009.
- [8] R. F. Gabitass, N. E. Annels, D. D. Stocken, H. A. Pandha, and G. W. Middleton, "Elevated myeloid-derived suppressor cells in pancreatic, esophageal and gastric cancer are an independent prognostic factor and are associated with significant elevation of the Th2 cytokine interleukin-13," *Cancer Immunology, Immunotherapy*. In press.
- [9] A. Correll, A. Tüettenberg, C. Becker, and H. Jonuleit, "Increased regulatory T-cell frequencies in patients with advanced melanoma correlate with a generally impaired T-cell responsiveness and are restored after dendritic cell-based vaccination," *Experimental Dermatology*, vol. 19, no. 8, pp. e213–e221, 2010.
- [10] M. Vergati, V. Cereda, R. A. Madan et al., "Analysis of circulating regulatory T cells in patients with metastatic prostate cancer pre- versus post-vaccination," *Cancer Immunology, Immunotherapy*, vol. 62, no. 2, pp. 197–206, 2010.
- [11] S. Mittal, N. A. Marshall, L. Duncan, D. J. Culligan, R. N. Barker, and M. A. Vickers, "Local and systemic induction of CD4⁺ CD25⁺ regulatory T-cell population by non-Hodgkin lymphoma," *Blood*, vol. 111, no. 11, pp. 5359–5370, 2008.
- [12] L. Weiss, T. Melchardt, A. Egle, C. Grabmer, R. Greil, and I. Tinhofer, "Regulatory T cells predict the time to initial treatment in early stage chronic lymphocytic leukemia," *Cancer*, vol. 117, no. 10, pp. 2163–2169, 2011.
- [13] J. M. Rojas, L. Wang, S. Owen, K. Knight, S. J. Watmough, and R. E. Clark, "Naturally occurring CD4⁺ CD25⁺ FOXP3⁺ T-regulatory cells are increased in chronic myeloid leukemia patients not in complete cytogenetic remission and can be immunosuppressive," *Experimental Hematology*, vol. 38, no. 12, pp. 1209–1218, 2010.
- [14] D. Wolf, A. M. Wolf, H. Rumpold et al., "The expression of the regulatory T cell-specific forkhead box transcription factor FoxP3 is associated with poor prognosis in ovarian cancer," *Clinical Cancer Research*, vol. 11, no. 23, pp. 8326–8331, 2005.
- [15] J. Carreras, A. Lopez-Guillermo, B. C. Fox et al., "High numbers of tumor-infiltrating FOXP3-positive regulatory T cells are associated with improved overall survival in follicular lymphoma," *Blood*, vol. 108, no. 9, pp. 2957–2964, 2006.
- [16] A. M. Lee, A. J. Clear, M. Calaminici et al., "Number of CD4⁺ cells and location of forkhead box protein P3-positive cells in diagnostic follicular lymphoma tissue microarrays correlates with outcome," *Journal of Clinical Oncology*, vol. 24, no. 31, pp. 5052–5059, 2006.
- [17] L. S. Taams, J. Smith, M. H. Rustin, M. Salmon, L. W. Poulter, and A. N. Akbar, "Human anergic/suppressive CD4⁺ CD25⁺ T cells: a highly differentiated and apoptosis-prone population," *European Journal of Immunology*, vol. 31, no. 4, pp. 1122–1131, 2001.
- [18] D. Valmori, A. Merlo, N. E. Souleimanian, C. S. Hesdorffer, and M. Ayyoub, "A peripheral circulating compartment of natural naive CD4⁺ Tregs," *Journal of Clinical Investigation*, vol. 115, no. 7, pp. 1953–1962, 2005.
- [19] M. Beyer and J. L. Schultze, "CD4⁺ CD25^{high} FOXP3⁺ regulatory T cells in peripheral blood are primarily of effector memory phenotype," *Journal of Clinical Oncology*, vol. 25, no. 18, pp. 2628–2630, 2007.
- [20] N. Seddiki, B. Santner-Nanan, S. G. Tangye et al., "Persistence of naive CD45RA⁺ regulatory T cells in adult life," *Blood*, vol. 107, no. 7, pp. 2830–2838, 2006.
- [21] B. Fritzsching, N. Oberle, E. Pauly et al., "Naive regulatory T cells: a novel subpopulation defined by resistance toward CD95L-mediated cell death," *Blood*, vol. 108, no. 10, pp. 3371–3378, 2006.
- [22] M. Miyara, Y. Yoshioka, A. Kitoh et al., "Functional delineation and differentiation dynamics of human CD4⁺ T cells expressing the FoxP3 transcription factor," *Immunity*, vol. 30, no. 6, pp. 899–911, 2009.
- [23] M. Beyer, M. Kochanek, T. Giese et al., "In vivo peripheral expansion of naive CD4⁺ CD25^{high} FoxP3⁺ regulatory T cells in patients with multiple myeloma," *Blood*, vol. 107, no. 10, pp. 3940–3949, 2006.
- [24] M. Beyer, M. Kochanek, K. Darabi et al., "Reduced frequencies and suppressive function of CD4⁺ CD25^{hi} regulatory T cells in patients with chronic lymphocytic leukemia after therapy with fludarabine," *Blood*, vol. 106, no. 6, pp. 2018–2025, 2005.
- [25] K. A. Kasow, X. Chen, J. Knowles, D. Wichlan, R. Handgretinger, and J. M. Riberdy, "Human CD4⁺ CD25⁺ regulatory T cells share equally complex and comparable repertoires with CD4⁺ CD25⁻ counterparts," *Journal of Immunology*, vol. 172, no. 10, pp. 6123–6128, 2004.
- [26] W. Zou, "Regulatory T cells, tumour immunity and immunotherapy," *Nature Reviews Immunology*, vol. 6, no. 4, pp. 295–307, 2006.
- [27] M. Beyer and J. L. Schultze, "Regulatory T cells: major players in the tumor microenvironment," *Current Pharmaceutical Design*, vol. 15, no. 16, pp. 1879–1892, 2009.
- [28] S. Sakaguchi, T. Yamaguchi, T. Nomura, and M. Ono, "Regulatory T cells and immune tolerance," *Cell*, vol. 133, no. 5, pp. 775–787, 2008.
- [29] E. Y. Woo, H. Yeh, C. S. Chu et al., "Cutting edge: regulatory T cells from lung cancer patients directly inhibit autologous

- T cell proliferation,” *Journal of Immunology*, vol. 168, no. 9, pp. 4272–4276, 2002.
- [30] R. H. Prabhala, P. Neri, J. E. Bae et al., “Dysfunctional T regulatory cells in multiple myeloma,” *Blood*, vol. 107, no. 1, pp. 301–304, 2006.
- [31] M. Beyrer and J. L. Schultze, “Plasticity of T_{reg} cells: is reprogramming of T_{reg} cells possible in the presence of FOXP3?” *International Immunopharmacology*, vol. 11, no. 5, pp. 555–560, 2010.
- [32] E. Zorn, E. A. Nelson, M. Mohseni et al., “IL-2 regulates FOXP3 expression in human CD4⁺ CD25⁺ regulatory T cells through a STAT-dependent mechanism and induces the expansion of these cells in vivo,” *Blood*, vol. 108, no. 5, pp. 1571–1579, 2006.
- [33] Y. Kuniyasu, T. Takahashi, M. Itoh, J. Shimizu, G. Toda, and S. Sakaguchi, “Naturally anergic and suppressive CD25⁺ CD4⁺ T cells as a functionally and phenotypically distinct immunoregulatory T cell subpopulation,” *International Immunology*, vol. 12, no. 8, pp. 1145–1155, 2000.
- [34] H. W. Lim, H. E. Broxmeyer, and C. H. Kim, “Regulation of trafficking receptor expression in human forkhead box P3+ regulatory T cells,” *Journal of Immunology*, vol. 177, no. 2, pp. 840–851, 2006.
- [35] P. Hoffmann, R. Eder, T. J. Boeld et al., “Only the CD45RA⁺ subpopulation of CD4⁺ CD25^{high} T cells gives rise to homogeneous regulatory T-cell lines upon in vitro expansion,” *Blood*, vol. 108, no. 13, pp. 4260–4267, 2006.
- [36] P. Hoffmann, T. J. Boeld, R. Eder et al., “Loss of FOXP3 expression in natural human CD4⁺ CD25⁺ regulatory T cells upon repetitive in vitro stimulation,” *European Journal of Immunology*, vol. 39, no. 4, pp. 1088–1097, 2009.
- [37] M. Vukmanovic-Stejic, Y. Zhang, J. E. Cook et al., “Human CD4⁺ CD25^{hi} Foxp3⁺ regulatory T cells are derived by rapid turnover of memory populations in vivo,” *Journal of Clinical Investigation*, vol. 116, no. 9, pp. 2423–2433, 2006.
- [38] T. J. Curiel, G. Coukos, L. Zou et al., “Specific recruitment of regulatory T cells in ovarian carcinoma fosters immune privilege and predicts reduced survival,” *Nature Medicine*, vol. 10, no. 9, pp. 942–949, 2004.
- [39] W. Chen, W. Jin, N. Hardegen et al., “Conversion of peripheral CD4⁺ CD25⁻ naive T cells to CD4⁺ CD25⁺ regulatory T cells by TGF- β induction of transcription factor Foxp3,” *Journal of Experimental Medicine*, vol. 198, no. 12, pp. 1875–1886, 2003.
- [40] S. Sharma, S. C. Yang, L. Zhu et al., “Tumor cyclooxygenase-2/prostaglandin E2-dependent promotion of FOXP3 expression and CD4⁺ CD25⁺ T regulatory cell activities in lung cancer,” *Cancer Research*, vol. 65, no. 12, pp. 5211–5220, 2005.