


BMJ Open Knowledge, attitude and practice concerning healthcare-associated infections among healthcare workers in Wuhan, China: cross-sectional study

Wenwen Wu,¹ Wenru Wang,² Yufeng Yuan,³ Likai Lin,⁴ Yibin Tan,³ Jinru Yang,⁵ Li Dai,⁶ Ying Wang ³

To cite: Wu W, Wang W, Yuan Y, *et al.* Knowledge, attitude and practice concerning healthcare-associated infections among healthcare workers in Wuhan, China: cross-sectional study. *BMJ Open* 2021;**11**:e042333. doi:10.1136/bmjopen-2020-042333

► Prepublication history and supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-042333>).

WWu and WWa contributed equally.

WWu and WWa are joint first authors.

Received 01 July 2020

Revised 15 November 2020

Accepted 20 November 2020



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Ying Wang;
wangying19900621@126.com

ABSTRACT

Objectives To assess the knowledge, attitude and practice (KAP) concerning healthcare-associated infections (HAIs) among healthcare givers and to identify the factors influencing KAP.

Design This was a hospital-based, cross-sectional study.

Setting Two public hospitals in Wuhan, central China.

Participants Participants were recruited among healthcare workers (HCWs) of one general hospital and one children's hospital in Wuhan city between 1 June and 30 September 2019.

Primary and secondary outcome measures The outcomes were KAPs concerning HAIs.

Results Data from 455 HCWs were included in the final data analysis. The mean scores of KAP and total KAP were 15.67 ± 3.32 , 25.00 ± 2.75 , 43.44 ± 5.15 and 84.76 ± 6.72 , respectively. The following factors were significantly associated with the total KAP score concerning HAIs, explaining 61% of the variance ($p < 0.001$): gender ($\beta = 2.36$, 95% CI 1.11 to 4.40), age ($\beta = 6.65$, 95% CI 5.07 to 7.74), position ($\beta = 7.02$, 95% CI 3.88 to 8.45), type of employment ($\beta = -1.08$, 95% CI -2.08 to -0.07), with HAI education within last year ($\beta = -2.98$, 95% CI -4.23 to -1.72), with invasive operation authority ($\beta = -4.22$, 95% CI -5.46 to -2.99), antibacterial drug training ($\beta = -4.38$, 95% CI -5.45 to -3.31) and with antibacterial drug training and clinical consultation ($\beta = -4.35$, 95% CI -5.38 to -3.32).

Conclusion The controllable factors identified in this study can be used by hospital managers to implement measures that improve KAP among HCWs. Moreover, these measures should be customised, based on uncontrollable factors to suit the specific characteristics of medical staff and to improve KAP. Training programmes should be designed for medical workers to increase their awareness of HAIs and to foster positive attitudes and practices.

INTRODUCTION

Healthcare-associated infections (HAIs) refer to the infections acquired in hospitals but are neither present nor incubating at the time of a patient's admission.¹ HAIs represent significant challenges to the effective delivery of healthcare services and can result in prolonged hospital stays, microbial

Strengths and limitations of this study

- A large sample was used to investigate knowledge, attitude and practice (KAP) concerning healthcare-associated infections and to identify the significant influencing factors of KAP among healthcare workers in central China.
- The use of self-reporting data can cause response bias, which potentially affected the accuracy of the findings.
- This study was cross-sectional, so a causal relationship could not be confirmed.

resistance, exacerbation of existing conditions, worsening of patients' economic burdens, stretching available healthcare resources and even deaths.²⁻⁴ According to the WHO, at any moment, 1.4 million patients bear the consequences of HAIs globally.⁵ It has been estimated that nearly 10% of inpatients suffer the consequences of HAIs.⁶ Healthcare challenges emerging from HAIs are currently among the most significant public global health issues.⁷ The risk of acquiring an HAI in low-income countries is 2–20 times higher than that in high-income countries.^{8,9} Wang and colleagues reported that the weighted prevalence of HAIs varies between 1.73% and 5.45% in Chinese municipalities and provinces.¹⁰ The direct economic burden of hospital infections in China ranges from \$1.5 billion to \$2.3 billion annually.¹¹ Therefore, prevention and management of HAIs in China in the presence of competing interests remain an important clinical and public health topic.^{12,13}

One of the primary causes of HAIs is the contact and transmission of contaminated hand and medical equipment by healthcare workers (HCWs) who do not properly comply with hospital hygiene practices.¹⁴ For example, after evaluating or caring for

one patient, HCWs occasionally touch another patient without properly washing their hands. A previous study reported that adherence to hand hygiene recommendations among HCWs remains suboptimal, yet the compliance rate is approximately 30%.¹⁵ In fact, nearly 42% of COVID-19 infections in HCWs are related to the inappropriate use of personal protective equipment (PPE), masks and gloves.¹⁶

Effective prevention and control measures should always be observed, specifically by HCWs, to minimise the risk of HAIs.¹⁷ According to Kelman's theory of knowledge, attitude and practice (KAP), knowledge is the basis for changing practice, and attitude is the driving force of change.¹⁸ Therefore, understanding KAP of HCWs in relation to HAIs is essential in establishing these measures. Identifying the factors that significantly affect KAP is important and can provide a basis for implementing intervention measures by HAI managers. Few studies have investigated the relationship between KAP and HAIs among HCWs, or investigated the relationship between KAP and HAIs among HCWs.^{19–21} However, these studies have some limitations. First, they described only the current KAP status, but the factors influencing KAP remain poorly understood. Second, the majority of published KAP reports have focused only on hand hygiene. To the best of our knowledge, no studies have assessed KAP and identified its influencing factors among Chinese HCWs concerning HAIs in various healthcare settings.

Hence, this study aimed to assess KAP associated with HAIs and to identify the factors that significantly influence KAP among HCWs at two university-affiliated hospitals in China. Based on Kelman's theory of KAP,¹⁸ the stated hypothesis was that the factors significantly affecting the knowledge and attitudes of HCWs would be partially coincident with the factors influencing their practices concerning HAIs. Specifically, sociodemographic and job-related factors would significantly influence the knowledge and practice of HCWs toward HAIs, whereas the factors significantly affecting the attitudes of HCWs concerning HAIs would be primarily job-related.

METHODS

Study design and participants

A cross-sectional questionnaire survey was conducted in Wuhan, from 1 June 2019 to 30 September 2019. A total of 49 tertiary public hospitals were located in Wuhan, with 8.41 hospital beds per 1000 patients.²² The following multistage stratified sampling approach was employed: (1) 2 out of the 13 administrative regions of Wuhan were randomly selected for the study; (2) for each of the 2 selected regions, one hospital out of all the grade III level A hospitals in the region was randomly chosen; (3) with the support of the department of human resources, each study hospital's potential participants were randomly selected from the list of job numbers of HCWs and were provided the online link for questionnaires; and

(4) the HCWs who received questionnaires voluntarily completed and returned them online. The term 'HCWs' referred to doctors and nurses only and excluded interns, nurse assistants and medical students because some of the information requested could only be provided by them. To be included in the study, HCWs were required to meet the following criteria: (1) formal doctors and nurses registered at two hospitals, (2) professional qualification certificates and (3) voluntary participation in the study. HCWs who were on leave at the time of the survey and non-clinical staff were excluded from this study.

The sample size for the study was calculated using statistical power analysis. According to Cohen's guidelines,²³ in multiple linear regression analyses with an estimate of 10 independent variables,²⁴ a minimum of 120 subjects would be required to achieve a median effect size (0.15) at 80% statistical power and a significance level of 0.05.^{25 26} A total of 468 HCWs completed the online questionnaire, and incomplete questionnaires were excluded. The 455 completed questionnaires were used for downstream analyses. The larger samples increased the statistical power of the study.

Measures

The questionnaire was based on standard precaution knowledge questions and the core content of HAI prevention and control systems in China, due to a lack of prior research on the KAP of HAIs among HCWs.^{27–30}

The questionnaire consisted of two sections: the first section covered general information, and the second one included KAP concerning HAIs. The general information section comprised 16 questions to collect the participants' sociodemographic data, including age, gender, clinical work experience, marital status, educational level, occupation, department, position, professional title, employment, hospital type, HAI education within the last year, occupational exposure within the past 6 months, invasive operation authority, antibacterial drug training and attended clinical consultation.

The HAI knowledge domain consisted of six questions to assess the participants' knowledge of hand hygiene, HAIs, multidrug resistance, standard precautions and surgery site infection. The HAI attitude domain included eight questions to assess the participants' attitudes about personal and social motivation, which addressed the aspects of responsibility, attention, necessity and initiative among HAIs. The HAI practice domain consisted of 12 questions to assess the participants' practice of aseptic operation, standard precautions and antibiotic use. The responses were scored on a 5-point Likert scale ranging from 1 (consistent with my cognition) to 5 (very inconsistent with my cognition).

Pilot study

Thirty participants were recruited for the pilot study from 1 to 15 May 2019 to test the trial version of the quick response code for this study. The pilot participant responses were then analysed for clarity, understandability

and applicability of the questionnaire. The time to complete the questionnaire and any technical difficulties while scanning the quick response code were recorded.

Cronbach's alpha values were 0.662 (domain A, knowledge), 0.784 (domain B, attitudes) and 0.806 (domain C, practice). In addition, six experts in the field of nosocomial infection were invited to review each item using a 4-point rating scale (1=not relevant, 2=somewhat relevant, 3=quite relevant and 4=very relevant) and to test the content validity of the KAP. The overall content validity index was 0.95, which indicated that the content validity of the KAP questionnaire was reliable.

Data collection procedure

With the support of the hospital's human resources department, potential participants were contacted. After the questionnaire's reliability and validity assessment, web links to the questionnaire and informed consent forms were emailed to the qualifying potential participants by the researchers. The estimated time needed to complete the survey was 15 min. After the questionnaires were completed by the participants, responses were submitted online, and their electronic informed consent was returned via email. The questionnaires were then carefully reviewed for data analysis, and incomplete or incorrectly completed questionnaires were excluded.

Data analysis

For continuous variables, the means and SD were calculated, whereas frequencies and percentages were calculated for categorical variables. The scores of KAP for general characteristics were analysed via t-test or analysis of variance for continuous data. Multiple linear regression analysis was performed to determine the significant factors influencing KAP and KAP total scores. Variables with $p < 0.05$ determined from univariate analysis were included as independent variables in the regression model. Unstandardised coefficients and R^2 were used to interpret the effects and variability of the significant dependent variables, respectively. Statistical analyses were performed using STATA V.14.0. Statistical tests were two-sided, and statistical significance was set at a p value of < 0.05 .

Patient and public involvement

No patients or members of the public were involved in the design or planning of the study.

RESULTS

Descriptive statistics of participant characteristics

A total of 500 HCWs were invited to participate in the study. A total of 468 HCWs completed the online questionnaire (response rate=93.6%). After the incomplete questionnaires were excluded, the data from 455 HCWs (395 nurses and 60 doctors) were included in the final analysis. The age of the study participants ranged from 22 to 59 years (mean age=31.35 years). Most of the

participants were female (91%), and the mean duration of working experience was 9.45 years. Most participants were married (75.6%) and had attained a bachelor's degree (68.1%). More than a quarter of the participants worked in the operating room (28.1%). The majority (96%) of the participants were general staff, and 66.4% had a junior professional title. More than half of the participants (52.3%) were contractual employees, and 70.1% worked in the general hospital. Among the participants, 60.2% had received HAI education within the previous year; 62% experienced occupational exposures; and most participants received invasive operation authority (71.6%). In addition, 56.5% received antibacterial drug training, and 52.3% attended clinical consultation. The participants' scores of KAP and total KAP were 15.67 ± 3.32 , 25.00 ± 2.75 , 43.44 ± 5.15 and 84.76 ± 6.72 , respectively. The demographics and general characteristics of the participating groups are presented in [table 1](#).

Univariate analysis

Univariate analyses were performed to identify the factors influencing KAP, and the results are presented in online supplemental tables A1–A4.

The mean score of knowledge was significantly higher among the following groups of participants: received HAI education within the previous year, received antibacterial drug training, worked in the operating room or infectious diseases department, and had more than 10 years of work experience (all factors $p < 0.05$). There were significant differences in knowledge scores between the following groups: gender, age group, type of employment, received invasive operation authority and participated in clinical consultations with infectious disease doctors ($p < 0.001$) (online supplemental table A1).

Online supplemental table A2 presents the factors associated with the mean attitude score. The participants who were contractual employees, were married, had higher education levels, had antibacterial drug training and had a higher education level reported a significantly higher score on attitude (all factors $p < 0.05$). In addition, the attitude score was significantly associated with age, HAI education within the previous year, skin or mucous membrane exposure to patient bodily fluids within the previous 6 months, invasive operation authority, work department, clinical work experience and job position (all factors $p < 0.001$).

Univariate analysis also revealed that being female, having worked in general hospitals, being the department head, having more than 10 years of working experience and holding a senior technical job position, were associated with higher mean scores of practice (all factors $p < 0.05$). In addition, the mean scores of practice were significantly higher among the following groups of participants: those aged 40–59 years, contractual employees, individuals with higher education levels, those who received HAI education within the previous year, those who had skin or mucous membrane exposure to patient bodily fluids within the previous 6 months, individuals with invasive

Table 1 General characteristics of the participants	
Variables	n (%)
Age (years), mean±SD	31.35±7.12
Gender	
Male	41 (9)
Female	414 (91)
Clinical work experience (years), mean±SD	9.45±8.35
Marital status	
Unmarried	99 (21.8)
Married	344 (75.6)
Widowed/divorced	12 (2.6)
Educational level	
Junior college	37 (8.2)
Bachelor's degree	310 (68.1)
Master's degree or higher	108 (23.7)
Occupation	
Doctor	60 (13.2)
Nurse	395 (86.8)
Department	
Internal medicine	16 (3.5)
Surgery	83 (18.2)
Obstetrics	20 (4.4)
Intensive care unit	87 (19.1)
Emergency	21 (4.6)
Outpatient	11 (2.4)
Operating room	128 (28.1)
Infectious diseases	68 (14.9)
Other	21 (4.6)
Position	
Staff	437 (96)
Head	18 (4)
Professional title	
Senior	23 (5.1)
Middle	130 (28.6)
Junior	302 (66.4)
Type of employment	
Contract	238 (52.3)
Permanent	217 (47.7)
Type of hospital	
Children's hospital	136 (29.9)
General hospital	319 (70.1)
Received HAI education within the previous year	
Yes	274 (60.2)
No	181 (39.8)
Occupational exposures (impaired skin or mucosa to blood, body fluid, secretion and excretion of patients within 6 months)	

Continued

Table 1 Continued	
Variables	n (%)
Yes	282 (62)
No	173 (38)
Received invasive operation authority	
Yes	326 (71.6)
No	129 (28.4)
Received antibacterial drug training	
Yes	257 (56.5)
No	198 (43.5)
Attended consultation of nosocomial infection disease	
Yes	238 (52.3)
No	217 (47.7)
Knowledge score, mean±SD	15.67±3.32
Attitude score, mean±SD	25.00±2.75
Practice score, mean±SD	43.44±5.15
KAP, mean±SD	84.76±6.72

HAI, healthcare-associated infection; KAP, knowledge, attitude and practice.

operation authority, those who received antibacterial drug training; individuals working in an operating room, surgery department, intensive care unit or the department of infectious diseases; and those who participated in clinical consultations with infectious disease doctors (all factors $p < 0.001$) (online supplemental table A3).

Being female, working as a nurse and having worked in general hospitals were significantly associated with higher total KAP scores (all factors $p < 0.05$). Furthermore, the participants with the following characteristics reported significantly higher scores of the total KAP: 40–59 years of age, department head, contractual employees, received HAI education within the previous year, had skin or mucous membrane exposure to patient bodily fluids within the previous 6 months, had invasive operation authority, received antibacterial drug training and participated in clinical consultations with infectious disease doctors (all factors $p < 0.001$) (online supplemental table A4).

Multiple linear regression analysis

The results of the assessed regression models are reported in tables 2–5. Gender, age group, type of employment and clinical work experience were identified as significant predictors of knowledge in the multivariate regression analysis model, assuming knowledge as the outcome variable, and accounted for 21.4% of variance (adjusted $R^2 = 0.214$, $p < 0.001$). Being female, older age and 16–20 years of clinical work experience were significantly and positively associated with knowledge scores, whereas permanent staff was significantly and negatively associated with knowledge scores (table 2).

Table 2 Multiple linear regression analysis of the influencing factors for knowledge scores

Independent variables	B (95% CI)	SD	β	T	P value
Intercept	13.20 (11.03 to 15.36)	1.10		11.99	<0.001
Gender					
Female (vs male)	2.36 (1.24 to 3.47)	0.57	0.19	4.15	<0.001
Age group (years)					
40–59 (vs 18–39)	3.04 (1.84 to 4.24)	0.61	0.27	4.98	<0.001
Type of employment					
Permanent staff (vs contractual)	–1.27 (–1.82 to –0.56)	0.32	–0.18	–3.93	<0.001
Clinical work experience (years)					
6–10 (vs 1–5)	–0.17 (–0.93 to 0.59)	0.39	–0.02	–0.44	0.660
11–15 (vs 1–5)	0.65 (–0.47 to 1.77)	0.57	0.05	1.14	0.253
16–20 (vs 1–5)	1.54 (0.40 to 2.68)	0.58	0.13	2.66	0.008
≥21 (vs 1–5)	0.87 (–0.34 to 2.08)	0.61	0.08	1.41	0.158

The independent variables included in the regression model were gender, age group, occupation, type of employment, having received healthcare-associated infection education within the previous year, occupational exposure within 6 months, having received invasive operation authority, having received antibacterial drug training, having department and clinical work experience, and having attended consultation of nosocomial infection disease.

Adjusted $R^2=0.214$ (p value of <0.001).

A significant model was set up through multiple linear regression analysis ($p<0.001$), explaining 14.3% of the variance in attitude score (adjusted $R^2=0.143$). The

following aspects were positively associated with attitude scores [table 3](#): received HAI education within the last year, experienced occupational exposure within 6 months,

Table 3 Multiple linear regression analysis of the influencing factors for attitude scores

Independent variables	B (95% CI)	SD	β	T	P value
Intercept	25.20 (22.89 to 27.51)	1.18		21.44	<0.001
Received HAI education within the previous year					
No (vs yes)	–0.97 (–1.64 to –0.29)	0.34	–0.17	–2.82	0.005
Occupational exposure within 6 months					
Yes (vs no)	0.90 (0.15 to 1.66)	0.38	0.16	2.36	0.019
Received invasive operation authority					
No (vs yes)	–1.04 (–2.05 to –0.65)	0.33	–0.17	–3.12	0.002
Attended consultation of nosocomial infection disease					
No (vs yes)	–0.73 (–1.27 to –0.19)	0.28	–0.13	–2.65	0.008
Department					
Surgery (vs internal medicine)	0.20 (–1.21 to 1.62)	0.72	0.03	0.28	0.778
Obstetrics (vs internal medicine)	–0.87 (–2.57 to 0.84)	0.87	–0.06	–1.00	0.319
Intensive care unit (vs internal medicine)	0.47 (–0.96 to 1.91)	0.73	0.07	0.65	0.517
Emergency (vs internal medicine)	–0.99 (–2.67 to 0.68)	0.85	–0.08	–1.16	0.245
Outpatient (vs internal medicine)	–2.11 (–4.13 to –0.09)	1.03	–0.12	–2.05	0.041
Operating room (vs internal medicine)	0.38 (–1.02 to 1.78)	0.71	0.06	0.54	0.591
Infectious diseases (vs internal medicine)	0.46 (–1.02 to 1.94)	0.75	0.06	0.61	0.543
Other (vs internal medicine)	–0.94 (–2.64 to 0.76)	0.87	–0.07	–1.08	0.280

The independent variables included in the regression model were age group, type of employment, clinical work experience (years), educational level, marital status, professional title, having received healthcare-associated infection education within the previous year, occupational exposure within 6 months, having received invasive operation authority, having attended consultation of nosocomial infection disease and work department. Adjusted $R^2=0.1434$ (p value of <0.001).

HAI, healthcare-associated infection.

Table 4 Multiple linear regression analysis of the influencing factors for practice scores

Independent variables	B (95% CI)	SD	β	T	P value
Intercept	40.71 (37.31 to 44.10)	1.73		23.58	<0.001
Gender					
Female (vs male)	1.55 (0.19 to 2.90)	0.69	0.09	2.24	0.025
Occupational exposure within 6 months					
No (vs yes)	-1.49 (-2.60 to -0.38)	0.56	-0.14	-2.64	0.009
Received invasive operation authority					
No (vs yes)	-1.70 (-2.67 to -0.74)	0.49	-0.15	-3.47	0.001
Received antibacterial drug training					
No (vs yes)	-3.01 (-3.85 to -2.17)	0.43	-0.29	-7.06	<0.001
Educational level					
Bachelor's degree (vs college degree)	3.40 (2.02 to 4.78)	0.70	0.31	4.85	<0.001
Master's degree or above (vs college degree)	3.74 (2.15 to 5.33)	0.81	0.31	4.62	<0.001
Attended consultation of nosocomial infection disease					
No (vs yes)	-2.60 (-3.40 to -1.80)	0.41	-0.25	-6.40	<0.001
Department					
Surgery (vs internal medicine)	2.78 (0.70 to 4.86)	1.06	0.21	2.62	0.009
Obstetrics (vs internal medicine)	-1.06 (-3.59 to 1.47)	1.29	-0.04	-0.82	0.412
Intensive care unit (vs internal medicine)	1.70 (-0.41 to 3.82)	1.08	0.13	1.58	0.114
Emergency (vs internal medicine)	0.91 (-1.56 to 3.38)	1.26	0.04	0.73	0.468
Outpatient (vs internal medicine)	2.18 (-0.78 to 5.14)	1.51	0.07	1.45	0.148
Operating room (vs internal medicine)	2.76 (0.71 to 4.81)	1.04	0.24	2.65	0.008
Infectious diseases (vs internal medicine)	2.70 (0.52 to 4.87)	1.11	0.19	2.43	0.015
Other (vs internal medicine)	0.08 (-2.44 to 2.60)	1.28	0	0.06	0.951

The independent variables included in the regression model included: gender, age group, type of hospital, position, type of employment, having received healthcare-associated infection education within the previous year, occupational exposure within 6 months, having received invasive operation authority, having received antibacterial drug training, educational level, having attended consultation of nosocomial infection disease, department, clinical work experience and professional title. Adjusted $R^2=0.4705$ (p value of <0.001).

received invasive operation authority and attended clinical consultation.

The results of the multiple linear regression analysis on practices are shown in table 4. Gender, education level, work department, occupational exposure within 6 months, invasive operation authority, antibacterial drug training and attending clinical consultation were identified as significant predictors of practice and explained 47.05% (adjusted $R^2=0.471$) of variance. Being female, experiencing occupational exposure within 6 months, having invasive operation authority, having antibacterial drug training, achieving higher education level, attending clinical consultation, and working in surgery, operating room or infectious disease department were significantly and positively associated with the practice of HCWs.

Another significant model was established through multiple linear regression analysis ($p<0.001$), explaining 61% of the variance in the total KAP scores (adjusted $R^2=0.61$). Being male, younger age, general staff and permanent staff had a significantly negative influence on KAP total scores. In contrast, the following aspects were

positively associated with the total KAP scores: received HAI education within the previous year, received invasive operation authority, received antibacterial drug training and attended clinical consultation (table 5).

DISCUSSION

This study appears to be the first to describe the KAPs in relation to HAIs and their influencing factors among HCWs in central China. Although increased awareness and stricter regulations on the control of hospital infections have been observed, the study survey found that limitations still exist in HCWs' knowledge and practices, in terms of HAIs. With the current COVID-19 pandemic, understanding HCWs' KAP concerning HAIs and the significant factors influencing their KAP is essential. These findings may provide a basis for designing and implementing targeted intervention programmes to promote the KAP of HCWs and to establish the basis for conducting future studies.

Table 5 Multiple linear regression analysis of the influencing factors for knowledge, attitude and practice total scores

Independent variables	B (95% CI)	SD	β	T	P value
Intercept	87.06 (85.12 to 88.99)	0.99		88.31	<0.001
Gender					
Female (vs male)	2.36 (1.11 to 4.40)	0.84	0.09	3.29	0.008
Age group (years)					
40–59 (vs 18–39)	6.65 (5.07 to 7.74)	0.68	0.30	9.44	<0.001
Position					
Head (vs Staff)	7.02 (3.88 to 8.45)	1.16	0.18	5.30	<0.001
Type of employment					
Permanent staff (vs contractual)	–1.08 (–2.08 to –0.07)	0.51	–0.07	–2.11	0.035
Received HAI education within the previous year					
No (vs yes)	–2.98 (–4.23 to –1.72)	0.64	–0.20	–4.65	<0.001
Received invasive operation authority					
No (vs yes)	–4.22 (–5.46 to –2.99)	0.63	–0.26	–6.71	<0.001
Received antibacterial drug training					
No (vs yes)	–4.38 (–5.45 to –3.31)	0.55	–0.29	–8.03	<0.001
Attended consultation of nosocomial infection disease					
No (vs yes)	–4.35 (–5.38 to –3.32)	0.52	–0.29	–8.31	<0.001

The independent variables included in the regression model were gender, age group, type of hospital, occupation, position, type of employment, having received HAI education within the previous year, occupational exposure within 6 months, having received invasive operation authority, having received antibacterial drug training and having attended consultation of nosocomial infection disease. Adjusted $R^2=0.61$ (p value of <0.001). HAI, healthcare-associated infection.

Results indicated that the HCWs' sociodemographic factors, such as gender, age, employment and clinical work experience, significantly affected their knowledge of HAIs. Although some of these factors are unchangeable (eg, age and gender), continuous education on HAIs remains essential to improve knowledge of HAIs. Previous studies similarly demonstrated that participants who underwent training within the previous 5 years obtained higher knowledge scores.³¹ Another previous study on the KAP, associated with central vascular catheters, proved this hypothesis and reported that knowledge scores were significantly higher among respondents who underwent active formal training than those who did not.³² However, career seniority and gender were not identified as significant factors influencing the knowledge level among UK HCWs, and this observation was partly inconsistent with the finding of this present study.³³

Possessing HAI education, experiencing occupational exposure within 6 months, having the authority to perform invasive operations and participating in clinical consultations promote positive HAI attitudes; however, working in outpatient clinics is not conducive to developing positive HAI attitudes. Respondents' attitudes toward prevention-related HAIs are significantly high among HCWs who are assigned in intensive care units and have appropriate knowledge and training.³² In a multi-centre study conducted in Shanghai, China, independent associations between older age or higher education and

categorical knowledge are observed among physicians.²⁷ A longer working experience is inversely and independently associated with the knowledge and attitudes of HCWs.²⁷ However, age, education level and working experience were not identified as significant influencing factors of attitudes concerning HAIs in this study. In contrast, receiving HAI education was the most significantly influential factor of attitudes. The inconsistent findings between this study and the study conducted in Shanghai may be due to the difference in the selection of the study hospitals. The study conducted in Shanghai recruited HCWs from community hospitals, where the HCWs typically possess a lower education level, compared with those from acute hospitals. During the COVID-19 pandemic, a high proportion of HCWs admitted fear of working.³⁴ As such, periodic educational interventions and training programmes regarding infection control practices for COVID-19 must be implemented among all HCWs, especially those who encounter new emerging infectious diseases.³⁵

In the practice domain, education level had the highest influence on the ability of HCWs to implement the prevention and control of HAIs. Other positive factors included gender, occupational exposure within the previous 6 months, authority to perform invasive operations, antibacterial drug training and attendance of clinical consultations. Previous studies largely focused on hand hygiene practices, and most of them reported poor

compliance with hand hygiene recommendations.^{15 36} Other studies have shown that factors such as perceived severity, subjective norms and job demands also influence practices significantly.³⁷ However, to some extent, influencing factors, such as occupational exposure and training, also relate to self-perception in this study.

Many medical professionals have become infected during the COVID-19 pandemic due to the lack of PPE. Statistical data have shown that more than 100 000 HCWs have been infected worldwide.³⁸ The adequate and correct use of PPE is the best measure to prevent HCWs from acquiring COVID-19 infection.³⁹ However, at the early stage of the outbreak, a global shortage of PPE occurred, and HCWs lacked practice on the proper donning and doffing of PPE.⁴⁰ Insufficient knowledge and skills related to the isolation of respiratory diseases pose a high risk of infection with HCWs. Although this study did not specifically focus on COVID-19, this pandemic demands awareness and attention to prepare HCWs with adequate knowledge, positive attitudes and practice in preventing and controlling transmitted infections and diseases.

Biases, especially those associated with participants' behaviour and practices, may exist in self-reported surveys. Consequently, participants may overstate their good practices. This study was cross-sectional, so inferences drawn from self-reported practices may vary from direct observation evidence. Moreover, no causal relationship was found.

CONCLUSION

In this study, KAP is closely associated with uncontrollable factors (such as gender, age, job position, employment type, educational level and clinical work experience) and controllable ones (such as HAI education within the previous year, occupational exposure within the previous 6 months, antibacterial drug training and participation in clinical consultations). Controllable factors indicate that hospital managers can respond appropriately for all HCWs to promote the improvement of KAP. Furthermore, uncontrollable factors imply that when taking measures to improve KAP, hospital managers should consider the backgrounds of individual HCWs. In addition, some sociodemographic and job-related factors significantly influence the knowledge and practices of HAIs among Chinese HCWs, whereas job-related factors significantly affect the attitudes of HCWs concerning HAIs. This result supports the hypotheses of this study. However, further studies should be performed to establish the benchmark of KPA for HAIs among HCWs.

Author affiliations

¹School of Public Health and Management, Hubei University of Medicine, Shiyan, Hubei, China

²Yong Loo Lin School of Medicine, National University of Singapore, Singapore

³Department of Nosocomial Infection Management, Wuhan University Zhongnan Hospital, Wuhan, Hubei, China

⁴Institute of Hospital Management, Wuhan University, Wuhan, Hubei, China

⁵School of Medicine, Wuhan University of Science and Technology, Wuhan, Hubei, China

⁶Department of Cardiovascular Medicine, Affiliated Dongfeng Hospital, Hubei University of Medicine, Shiyan, Hubei, China

Acknowledgements We thank all study participants for their valuable time and their support of our study. Furthermore, we would like to thank in particular the two hospitals which allowed us to perform survey with their staff.

Contributors WWu conceived the study. WWu, WWa, YY, LL, YT, JY and YW contributed to the survey design and data collection. LD contributed to the data analysis. All authors contributed to the interpretation of data and intellectual revision of multiple drafts. WWu and WWa drafted the manuscript. All authors approved the final version of the manuscript.

Funding This research was funded by Health commission of Hubei Province scientific research project (number WJ2019F161) and Faculty Development Grants From Hubei University of Medicine (number 2020QDJRW003).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval Ethical approval was obtained from the institutional ethics board of Wuhan University Zhongnan Hospital (number 2019191).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Some or all data, models or code generated or used during the study are available from the corresponding author (YW) by request. Reuse of the data is permitted for non-commercial purposes. Contact details: email: wangying19900621@126.com.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Ying Wang <http://orcid.org/0000-0003-2094-9347>

REFERENCES

- Breathnach AS. Nosocomial infections and infection control. *Medicine* 2013;41:649–53.
- Daxboeck F, Budic T, Assadian O, *et al*. Economic burden associated with multi-resistant gram-negative organisms compared with that for methicillin-resistant *Staphylococcus aureus* in a university teaching hospital. *J Hosp Infect* 2006;62:214–8.
- Rattanaumpawan P, Thamlikitkul V. Epidemiology and economic impact of health care-associated infections and cost-effectiveness of infection control measures at a Thai university hospital. *Am J Infect Control* 2017;45:145–50.
- Hollmeyer HG, Hayden F, Poland G, *et al*. Influenza vaccination of health care workers in hospitals—a review of studies on attitudes and predictors. *Vaccine* 2009;27:3935–44.
- Global patient safety challenge: 2005–2006/World alliance for patient safety. Available: <https://apps.who.int/iris/handle/10665/43358> [Accessed 16 Jun 2006].
- Humphreys H, Newcombe RG, Enstone J, *et al*. Four country healthcare associated infection prevalence survey 2006: risk factor analysis. *J Hosp Infect* 2008;69:249–57.
- Rosenthal VD. Health-care-associated infections in developing countries. *Lancet* 2011;377:186–8.
- Xie D-shuang, Fu X-yun, Wang H-fang, *et al*. Annual point-prevalence of healthcare-associated infection surveys in a university hospital in China, 2007–2011. *J Infect Public Health* 2013;6:416–22.

- 9 Liu JY, Wu YH, Cai M, *et al.* Point-Prevalence survey of healthcare-associated infections in Beijing, China: a survey and analysis in 2014. *J Hosp Infect* 2016;93:271–9.
- 10 Wang J, Liu F, Tartari E, *et al.* The prevalence of healthcare-associated infections in mainland China: a systematic review and meta-analysis. *Infect Control Hosp Epidemiol* 2018;39:701–9.
- 11 Sun B. Nosocomial infection in China: management status and solutions. *Am J Infect Control* 2016;44:851–2.
- 12 Byarugaba DK. A view on antimicrobial resistance in developing countries and responsible risk factors. *Int J Antimicrob Agents* 2004;24:105–10 <https://pubmed.ncbi.nlm.nih.gov/15288307>
- 13 Mir F, Zaidi AKM. Hospital infections by antimicrobial-resistant organisms in developing countries. *Springer* 2010;317:199–232.
- 14 Adebimpe WO, Asekun-Olarinmoye EO, Bamidele JO, *et al.* A comparative study of awareness and attitude to nosocomial infections among levels of health care workers in southwestern Nigeria. *Continent J Trop Med* 2011;5:5.
- 15 Hien H, Drabo M, Ouédraogo L, *et al.* [Knowledge and practices of health care workers in the area of healthcare-associated infection risks. A case study in a district hospital in Burkina Faso]. *Sante Publique* 2013;25:219–26.
- 16 Jin Y-H, Huang Q, Wang Y-Y, *et al.* Perceived infection transmission routes, infection control practices, psychosocial changes, and management of COVID-19 infected healthcare workers in a tertiary acute care hospital in Wuhan: a cross-sectional survey. *Mil Med Res* 2020;7:24.
- 17 World Health Organization. *Guidelines for the prevention and control of carbapenem-resistant Enterobacteriaceae, Acinetobacter baumannii and Pseudomonas aeruginosa in health care facilities*. Geneva, 2017.
- 18 Jiang H, Zhang S, Ding Y, *et al.* Development and validation of college students' tuberculosis knowledge, attitudes and practices questionnaire (CS-TBKAPQ). *BMC Public Health* 2017;17:949.
- 19 Angelillo IF, Mazziotta A, Nicotera G. Nurses and hospital infection control: knowledge, attitudes and behaviour of Italian operating theatre staff. *J Hosp Infect* 1999;42:105–12.
- 20 Balarabe SA, Joshua IA, Danjuma A, *et al.* Knowledge of healthcare workers on nosocomial infection in selected secondary health institutions in Zaria, Nigeria. *J Prevent Med* 2015;3:1–6.
- 21 Adegboye MB, Zakari S, Ahmed BA, *et al.* Knowledge, awareness and practice of infection control by health care workers in the intensive care units of a tertiary hospital in Nigeria. *Afr Health Sci* 2018;18:72–8.
- 22 Wuhan WSB. Wuhan health statistical Yearbook, 2019. Available: <http://tjj.wuhan.gov.cn/tjfw/tjnj/202004/P020200426461240969401.pdf> [Accessed 7 Sep 2020].
- 23 Cohen J. A power primer. *Psychol Bull* 1992;112:155–9.
- 24 Brown MT, Bussell JK. Medication adherence: who cares? *Mayo Clin Proc* 2011;86:304–14.
- 25 Glader E-L, Sjölander M, Eriksson M, *et al.* Persistent use of secondary preventive drugs declines rapidly during the first 2 years after stroke. *Stroke* 2010;41:397–401.
- 26 Maxwell SE. Sample size and multiple regression analysis. *Psychol Methods* 2000;5:434–58.
- 27 Zhou Y, Zhang D, Chen Y, *et al.* Healthcare-Associated infections and Shanghai clinicians: a multicenter cross-sectional study. *PLoS One* 2014;9:e105838.
- 28 Luo Y, He G-P, Zhou J-W, *et al.* Factors impacting compliance with standard precautions in nursing, China. *Int J Infect Dis* 2010;14:e1106–14.
- 29 Tavolacci M-P, Ladner J, Bailly L, *et al.* Prevention of nosocomial infection and standard precautions: knowledge and source of information among healthcare students. *Infect Control Hosp Epidemiol* 2008;29:642–7.
- 30 Circular of the general office of the National health Commission on further strengthening the prevention and control of infection in medical institution. Available: <http://www.cha.org.cn/plus/view.php?aid=15223> [Accessed 15 Mar 2017].
- 31 Taffurelli C, Sollami A, Camera C, *et al.* Healthcare associated infection: good practices, knowledge and the locus of control in healthcare professionals. *Acta Biomed* 2017;88:31–6.
- 32 Bianco A, Coscarelli P, Nobile CGA, *et al.* The reduction of risk in central line-associated bloodstream infections: knowledge, attitudes, and evidence-based practices in health care workers. *Am J Infect Control* 2013;41:107–12.
- 33 Brady RRW, McDermott C, Cameron F, *et al.* UK healthcare workers' knowledge of meticillin-resistant Staphylococcus aureus practice guidelines: a questionnaire study. *J Hosp Infect* 2009;73:264–70.
- 34 Rymarowicz J, Stefura T, Major P, *et al.* General surgeons' attitudes towards COVID-19: A national survey during the SARS-CoV-2 virus outbreak. *Eur Surg* 2020;21:1–6.
- 35 Neupane HC, Shrestha N, Adhikari S, *et al.* Knowledge of health care professionals and medical students regarding Covid-19 in a tertiary care hospital in Nepal. *JNMA J Nepal Med Assoc* 2020;58:480–6.
- 36 Ellingson K, Haas JP, Aiello AE, *et al.* Strategies to prevent healthcare-associated infections through hand hygiene. *Infect Control Hosp Epidemiol* 2014;35 Suppl 2:S155–78 <https://pubmed.ncbi.nlm.nih.gov/25026608>
- 37 Brazzell BD. Improving high Hand-hygiene compliance and reducing healthcare-associated infection in eight nursing units. *Am J Infect Control* 2014;42:S25–6.
- 38 Stubblefield WB, Talbot HK, Feldstein L, *et al.* Seroprevalence of SARS-CoV-2 Among Frontline Healthcare Personnel During the First Month of Caring for COVID-19 Patients - Nashville, Tennessee. *Clin Infect Dis* 2020:ciaa936.
- 39 Eyre DW, Lumley SF, O'Donnell D, *et al.* Differential occupational risks to healthcare workers from SARS-CoV-2 observed during a prospective observational study [published online ahead of print, 2020 Aug 21]. *Elife* 2020;9:e60675.
- 40 Wang Y, Wu W, Cheng Z, *et al.* Super-factors associated with transmission of occupational COVID-19 infection among healthcare staff in Wuhan, China. *J Hosp Infect* 2020;106:25–34.