

Diabetes mellitus, diabetes insipidus, optic atrophy, and deafness: A case of Wolfram (DIDMOAD) syndrome

Nasrollah Maleki ^{a,*}, Bahman Bashardoust ^a, Anahita Zakeri ^a, Azita Salehifar ^b, Zahra Tavosi ^b

^a Department of Internal Medicine, Imam Khomeini Hospital, Ardabil University of Medical Sciences, Ardabil, Iran

^b Department of Internal Medicine, Shohadaye Khalije Fars Hospital, Bushehr University of Medical Sciences Bushehr, Iran

Received 14 October 2015; revised 14 November 2015; accepted 21 November 2015

Available online 2 January 2016

Abstract

Purpose: To report a case of Wolfram syndrome (WS) characterized by diabetes mellitus, diabetes insipidus, progressive optic atrophy, and deafness.

Case report: A 19-year-old female patient, a known case of diabetes mellitus type I from six years before, presented with progressive vision loss since four years earlier. On fundoscopic examination, she had bilateral optic atrophy without diabetic retinopathy. The patient also had diabetes insipidus, neurosensory deafness, and neurogenic bladder.

Conclusion: WS should be considered a differential diagnosis in patients with diabetes mellitus who present with optic atrophy, and it is necessary to perform a hearing test as well as collecting 24-h urine output.

Copyright © 2015, Iranian Society of Ophthalmology. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Diabetes mellitus; Diabetes insipidus; Optic atrophy; Deafness

Introduction

Wolfram syndrome (WS) is an autosomal recessive neurodegenerative disorder characterized by Diabetes Insipidus, Diabetes Mellitus (non-autoimmune), Optic Atrophy, and Deafness (a set of conditions referred to as DIDMOAD).^{1–3} The gene responsible for the WS, named WSF1, encodes an endoplasmic reticulum membrane-embedded protein called wolframin that is expressed in pancreatic beta cells and neurons.^{4–6} The estimated prevalence of WS is 1 in 770,000, and it is believed to occur in 1 of 150 patients with type 1 diabetes.⁶ Affected patients usually develop insulin-requiring diabetes and optic atrophy in early childhood, and diabetes insipidus as they become teenagers or young adults. In WS patients with diabetes insipidus, not only

does vasopressin neuron loss occur in the supraoptic nucleus, but there is also a defect in vasopressin precursor processing.⁷ Anterior pituitary dysfunction has also been reported.⁶ Other manifestations of WS include progressive sensorineural deafness, hydronephrosis (due in part to the high urine flow in diabetes insipidus), and neurologic dysfunction.⁸ Why severe insulin-requiring diabetes develops is not known; immunologic factors do not appear to be important.³ Affected individuals have a median life expectancy of 30 years, with early mortality credited to neurological disorders, urological abnormalities and infection.⁹ There is currently no cure for WS. Differential diagnosis and treatment are difficult due to the variable presentation of symptoms.¹⁰ In the present study, we report one patient with WS from Ardabil, Iran.

Case report

A 19-year-old female patient, a known case of type I diabetes mellitus from six years before, presented to a private ophthalmology office. She complained of progressive

* Corresponding author. Department of Internal Medicine, Imam Khomeini Hospital, Ardabil University of Medical Sciences, Ardabil, Iran. Tel.: +98 9163057715.

E-mail address: malekinasrollah@yahoo.com (N. Maleki).

Peer review under responsibility of the Iranian Society of Ophthalmology.

vision loss from four years earlier. Her visual acuity reduced to 1/10 on both eyes. She had a history of nocturia and high urine output for five years, and after that, she had incontinency. In the previous two years, she had been avoiding difficulty and using self-intermittent catheterization four times a day.

On ophthalmic examination, intraocular pressure was 15 mmHg in both eyes, and biomicroscopic examinations were completely normal. On fundoscopic examination, she had bilateral optic atrophy without diabetic retinopathy (Figs. 1 and 2). Accordingly, WS was suspected by the ophthalmologist, who referred her to an endocrinologist.

Investigations revealed polyuria (14 L per day), hypernatremia (156 mEq/L), and hyperglycemia (351 mg/dL). Other laboratory values were: creatinine, 1.1 mg/dL; urea nitrogen, 28 mg/dL; K, 4.5 mEq/L; TSH, 2.5 mU/L. A random urine analysis showed a urine specific gravity of 1.006, and the results of 24-h urine collection obtained a urine specific gravity of 1.005 [normal values: 1.010–1.030]. It was suspected that there might be an underlying cause for diabetes insipidus, due to the presence of a low urine specific gravity ... Subsequently, a urine and plasma osmolality was performed, which were 150 and 312 mmol/kg, respectively. These findings suggested a disturbance in the regulation of vasopressin. Therefore, a standardized water deprivation test was done. There was no change either in urine specific gravity or urine osmolality before and after the water deprivation test. After nasal desmopressin administration, urine osmolality increased from 125 mOsm/kg to 600 mOsm/kg and serum osmolality increased from 281 mOsm/kg to 328 mOsm/kg. Given to an increase in urine osmolality of more than 100% measured by the desmopressin stimulation test, the patient was diagnosed with complete central diabetes insipidus. Treatment with oral desmopressin was prescribed 100 mcg three times daily. A few days later, the volume of urine collected in 24 h was within acceptable range. A Magnetic Resonance Imaging (MRI) of the brain revealed the loss of the physiological



Fig. 1. Optic atrophy without diabetic retinopathy in the right eye.



Fig. 2. Optic atrophy without diabetic retinopathy in the left eye.

hyperintense signal of the posterior pituitary gland (bright spot) on T1-weighted images, suggesting the diagnosis of central diabetes insipidus (Fig. 3).

An audiologic examination revealed neurosensory deafness, and audiometry showed a bilateral high-frequency sensorineural hearing loss. Renal sonography showed pelvicalyceal dilatation in both kidneys, equal to hydronephrosis grade II. The bladder wall had a trabecular pattern, and the urinary tract was dilated on both sides. The patient had 150 ml post-voiding bladder residue. Urologic examination and sonographic findings were consistent with a neurogenic bladder.




Fig. 3. Magnetic Resonance Imaging (MRI) of the brain revealed the loss of posterior pituitary bright spot, suggesting the diagnosis of central diabetes insipidus.

The patient's sister had similar symptoms, but she did not cooperate for further evaluation.

Discussion

WS was first described in 1938 by Wolfram and Wagener¹¹ as a hereditary syndrome characterized by diabetes mellitus and optic atrophy acquired early in life. Subsequent reports added diabetes insipidus and deafness to the syndrome, which developed in approximately 73 and 62% of sufferers, respectively.^{3,12}

The minimal criteria for diagnosis are juvenile-onset diabetes and optic atrophy; however patients also present with additional complications, hence the acronym DIDMOAD (Diabetes Insipidus, Diabetes Mellitus, Optic Atrophy, and Deafness) is used. The diagnostic criteria for WS are shown in Table 1. The diagnosis is established in individuals of all ages in whom two pathological WFS1 or CISD2 mutations are identified.¹³

Diabetes mellitus is a key feature of WS, which is usually the first sign of the disease (median age at diagnosis, 6–15 years), and optic atrophy (median age at diagnosis 11 years). The mean age of presentation of diabetes mellitus has been reported to be 5 ± 4 years.¹⁴ There are multiple differences between the presentation and course of autoimmune Type 1 diabetes and diabetes of WS. Namely, patients affected with WS have a low incidence of diabetic ketoacidosis at the time of diagnosis (only 3% compared to 30% in Type 1 diabetes), a much lower insulin requirement in the first several years after diagnosis, rare microvascular complications, and rare presence of diabetes antibodies.¹⁵ Patients with WS show smaller glycemic variability than individuals with type 1 diabetes mellitus, and this may be associated with persistent residual insulin secretion.¹⁶

Diabetes insipidus is another component of WS. A loss of vasopressin-producing neurons in the hypothalamus has been well documented in this disease and is thought to be the cause of WS-associated diabetes insipidus.^{17,18} Neuroradiologists have reported a loss of signal intensity in the posterior pituitary, indicating an absence or degeneration of the neurohypophysis^{18,19}; some patients have been found to have “empty sella”.²⁰

Patients with WS demonstrate progressive ophthalmologic symptoms that usually occur after diabetes mellitus. Besides optic atrophy, constriction of visual fields and declined visual acuity and color vision are the other ophthalmological findings

in WS²¹; however, diabetic retinopathy is rarely observed.²² Only five cases with WS have been reported without optic atrophy.²² In 2002, Al-Till et al²³ performed a study to investigate ophthalmologic abnormalities in fifteen patients with WS. Of 15 patients with WS reviewed in this study, 93.3% had optic atrophy, 92.9% had color deficit, 66.6% had cataract, 30% had pigmentary retinopathy, and 20% had diabetic retinopathy. Abnormal pupillary light reflexes and nystagmus were also reported.²³ In another study, eye complications were described in four patients with WS. The ophthalmic signs were a progressive decrease in visual acuity, constriction of the peripheral visual field with or without central scotoma, color vision disturbances, and bilateral optic disc atrophy. Diabetic retinopathy was a rare complication.²⁴ Recently, the retinal thinning has been proposed as a marker of disease progression in patients with WS.²⁵

Deafness in WS is commonly a high-frequency, symmetric, sensorineural hearing loss, usually manifesting in the second or third decade with a relatively slow rate of deterioration.^{26,27} Some studies have reported urinary tract abnormalities (hydro-ureter, urinary incontinence, recurrent infections) in patients with WS, that these abnormalities are expected in about 66% of patients. Urinary tract abnormalities include obstruction of the ureters, an atonic bladder with a large capacity, high pressure bladder with sphincteric dyssynergia, and incontinence.²⁸ Urinary tract dilatation is presented in 45% of patients with WS, which may be secondary to chronic high urine flow rate or neurodegeneration in different levels of the urinary tract.²⁹

Differential diagnosis of WS includes¹³: (1) Mitochondrial disorders: Maternally Inherited Diabetes mellitus and Deafness, Leber Hereditary Optic Neuropathy; (2) Thiamine-responsive megaloblastic anemia, diabetes, and deafness; (3) Autosomal Dominant Optic Atrophy; (4) X-linked Charcot-Marie-Tooth disease type 5; (5) Deafness, Dystonia, Optic Neuropathy syndrome; (6) Friedreich ataxia; (7) Bardet-Biedl syndrome; (8) Alstrom syndrome; and (9) Congenital rubella syndrome. The association of diabetes mellitus with optic atrophy also occurs in Friedreich's ataxia, Refsum disease, Alstrom syndrome, Lawrence-Moon syndrome, Kearn-Sayre syndrome, and deafness and diabetes in the “3243” mitochondrial DNA mutation.³⁰

The pathogenesis of the disease remains unknown, but genetic linkage studies have shown that the mutations of the WFS1 gene are responsible for the symptoms in 90% of cases

Table 1
The diagnostic criteria for Wolfram syndrome.

Major criteria	Minor criteria	Minimum required	Other variable suggestive evidence:
1. Diabetes mellitus <16 yrs, 2. Optic atrophy <16 yrs	1. Diabetes insipidus, 2. Diabetes mellitus >16 yrs, 3. Optic atrophy >16 yrs, 4. sensorineural deafness, 5. Neurological signs (ataxia, epilepsy, neuropathy, cognitive impairment), 6. Renal tract abnormalities, 7. 1 loss of function mutation in _WFS1/CISD2_ AND/OR family history of Wolfram syndrome	2 major OR 1 major plus 2 minor criteria	- Hypogonadism (males), - Absence of type 1 diabetes auto-antibodies, - Bilateral cataracts, - Psychiatric disorder, - Gastrointestinal disorders

meeting specific WFS diagnostic criteria. This gene is located on the short arm of chromosome 4 (4p16). The WFS1 gene encodes an 890-amino-acid protein called wolframin, a putative multispinning membrane glycoprotein of the endoplasmic reticulum.³¹ Although no function has yet been ascribed to wolframin; however, its localization to the endoplasmic reticulum suggests it can play a role in calcium homeostasis, membrane trafficking and protein processing.³²

There is currently no cure for WS, but there are treatments for some of the features. Diabetes mellitus is treated with insulin, and diabetes insipidus with vasopressin. While hearing aids can help with hearing loss, there is unfortunately no treatment for vision loss. Renal problems may be treated by catheterization (passing a thin, flexible tube into the bladder to drain away urine), and some of the neurological symptoms can be treated with medication.²⁹

Based on the findings of the present case, it is highly recommended that, in patients with diabetes mellitus who have optimal blood sugar presenting with polyuria and polydipsia, we need to do urine specific gravity to prove the presence of diabetes insipidus. In such patients, performing an ophthalmologic evaluation can probably reveal this rare disorder of Wolfram syndrome. In addition, ophthalmologists should be aware that in patients presenting with diabetes mellitus and optic atrophy without any signs of diabetic retinopathy, it is necessary to perform a hearing test as well as collect and analyze 24-h urine output. Thus, awareness of this condition and timely diagnosis is important for improving patient prognosis, anticipating associated complications, and enabling timely genetic counseling for family members.

References

- Hardy C, Khanim F, Torres R, et al. Clinical and molecular genetic analysis of 19 Wolfram syndrome kindreds demonstrating a wide spectrum of mutations in WFS1. *Am J Hum Genet.* 1999;65:1279–1290.
- Rando TA, Horton JC, Layzer RC. Wolfram syndrome: evidence of a diffuse neurodegenerative disease by magnetic resonance imaging. *Neurology.* 1992;42:1220–1224.
- Barrett TG, Bunday SE, Macleod AF. Neurodegeneration and diabetes: UK nationwide study of Wolfram (DIDMOAD) syndrome. *Lancet.* 1995;346:1458–1463.
- Inoue H, Tanizawa Y, Wasson J, et al. A gene encoding a transmembrane protein is mutated in patients with diabetes mellitus and optic atrophy (Wolfram syndrome). *Nat Genet.* 1998;20:143.
- Strom TM, Hörtnagel K, Hofmann S, et al. Diabetes insipidus, diabetes mellitus, optic atrophy and deafness (DIDMOAD) caused by mutations in a novel gene (wolframin) coding for a predicted transmembrane protein. *Hum Mol Genet.* 1998;7:2021.
- Boutzios G, Livadas S, Marinakis E, et al. Endocrine and metabolic aspects of the Wolfram syndrome. *Endocrine.* 2011;40:10.
- Gabreëls BA, Swaab DF, de Kleijn DP, et al. The vasopressin precursor is not processed in the hypothalamus of Wolfram syndrome patients with diabetes insipidus: evidence for the involvement of PC2 and 7B2. *J Clin Endocrinol Metab.* 1998;83:4026.
- Chausseot A, Bannwarth S, Rouzier C, et al. Neurologic features and genotype-phenotype correlation in Wolfram syndrome. *Ann Neurol.* 2011;69:501.
- Kinsley BT, Swift M, Dumont RH, Swift RG. Morbidity and mortality in the wolfram syndrome. *Diabetes Care.* 1995;18:1566–1570.
- Ganie MA, Bhat D. Current developments in wolfram syndrome. *J Pediatr Endocrinol Metab.* 2009;22:3–10.
- Wolfram DJ, Wagener HP. Diabetes mellitus and simple optic atrophy among siblings: report of four cases. *Mayo Clin Proc.* 1938;13:715–718.
- Hannon P, Becker L, Deck J. Evidence of widespread axonal pathology in Wolfram syndrome. *Acta Neuropathol.* 1999;98:304–308.
- Barrett T, Gharanei S. *Wolfram Syndrome* [internet]. Diapedia 4104640117 rev. no. 16. Available from: 2014 Aug 13 <http://dx.doi.org/10.14496/dia.4104640117.16>.
- Rohayem J, Ehlers C, Wiedeman B, et al. Diabetes and neurodegeneration in Wolfram syndrome: a multicenter study of phenotype and genotype. *Diabetes Care.* 2011;11503–11510.
- Garcia-Luna PP, Villechenous E, Leal-Cerro A, et al. Contrasting features of insulin dependent diabetes mellitus associated with neuroectodermal defects and classical insulin dependent diabetes mellitus. *Acta Paediatr Scand.* 1988;77:413–418.
- Zmyslowska A, Fendler W, Szadkowska A, et al. Glycemic variability in patients with Wolfram syndrome is lower than in type 1 diabetes. *Acta Diabetol.* 2015;52:1057–1062.
- Medlej R, Wasson J, Baz P, et al. Diabetes mellitus and optic atrophy: a study of Wolfram syndrome in the Lebanese population. *J Clin Endocrinol Metab.* 2004;89:1656–1661.
- Scolding NJ, Kellar-Wood HF, Shaw C, Shneerson JM, Anton Nagui. Wolfram syndrome: hereditary diabetes mellitus with brainstem and optic atrophy. *Ann Neurol.* 1996;29:352–360.
- Galluzzi P, Filisomi G, Vallone IM, Bardelli AM, Venturi C. MRI of Wolfram syndrome (DIDMOAD). *Neuroradiology.* 1999;41:729–731.
- Soliman AT, Bappal B, Darwish A, Rajab A, Asfour M. Growth hormone deficiency and empty sella in DIDMOAD syndrome: an endocrine study. *Arch Dis Child.* 1995;73:251–253.
- Fishman L, Ehrlich RM. Wolfram syndrome: report of four new cases and a review of literature. *Diabetes Care.* 1986;9:405.
- Seyhmus A, Ugur K, Ihsan C, Kaan U, Hasan K. Wolfram syndrome: case report and review of the literature. *Compr Ther.* 2007;33:18–20.
- Al-Till M, Jarrah NS, Ajlouni KM. Ophthalmologic findings in fifteen patients with Wolfram syndrome. *Eur J Ophthalmol.* 2002;12:84–88.
- Seynaeve H, Vermeiren A, Leys A, Dralands L. Four cases of Wolfram syndrome: ophthalmologic findings and complications. *Bull Soc Belge Ophthalmol.* 1994;252:75–80.
- Zmyslowska A, Fendler W, Niwald A, et al. Retinal thinning as a marker of disease progression in patients with Wolfram syndrome. *Diabetes Care.* 2015;38:e36–e37.
- Pennings RJ, Huygen PL, van den Ouweland JM, et al. Sex-related hearing impairment in Wolfram syndrome patients identified by inactivating WFS1 mutations. *Audiol Neurootol.* 2004;9:51–62.
- Plantinga RF, Pennings RJ, Huygen PL, et al. Hearing impairment in genotyped Wolfram syndrome patients. *Ann Otol Rhinol Laryngol.* 2008;117:494–500.
- Viswanathan V, Medempudi S, Kadiri M. Wolfram syndrome. *J Assoc Physicians India.* 2008;56:197–199.
- Tekgöl S, Oge O, Simsek E, Yordam N, Kendi S. Urological manifestations of the Wolfram syndrome: observations in 14 patients. *J Urol.* 1999;161:616–617.
- Barrett TG, Bunday SE. Wolfram (DIDMOAD) syndrome. *J Med Genet.* 1997;34:838–841.
- Van ven Ouweland JM, Cryns K, Pennings RJ, et al. Molecular characterization of WFS1 in patients with Wolfram syndrome. *J Mol Diagn.* 2003;5:88–95.
- Ribeiro MR, Crispim F, Vendramini MF, Moisés RS. Wolfram syndrome: from definition to molecular bases. *Arq Bras Endocrinol Metabol.* 2006;50:839–844.