

assessed with a 10-question quiz on PrEP facts. No incentives to complete the survey were offered. Data were summarized by frequency (%) for categorical variables and with means (SD) for continuous variables. Chi-squared tests were used to examine differences between knowledge of PrEP and other categorical variables.

Results. The study population ($N = 1588$) included women (53%), non-Caucasians (27%), and non-heterosexuals (15%). Median age was 25 (range 21–53). Forty-eight home states, including D.C., and 21 home countries were represented. 18% of fourth-year students were never taught about PrEP in medical school, compared with 40% of first-year students ($P < 0.001$). Overall, 28% of students were unaware of PrEP. Those unaware believed patients without HIV will not adhere to PrEP ($P < 0.001$). Awareness was associated with knowing someone with HIV besides a patient and experience caring for HIV-infected, intravenous drug-using, and LGBTQ people ($P < 0.001$). Higher knowledge scores were associated with confidence in determining a patient's candidacy for PrEP and, for third- and fourth-year students, having recommended PrEP in the clinical setting ($P < 0.001$). Overall, 57% believed that behavioral intervention should be tried before prescribing PrEP, 45% believed that patients would not adhere to PrEP, and 22% worried that PrEP is not effective.

Conclusion. We show in an 18-site study that medical students have limited awareness, knowledge, positive beliefs, and experiences of PrEP. Given these findings and the underutilization of PrEP by current practitioners, we recommend increasing the inclusion of PrEP in medical student education.

Disclosures. All authors: No reported disclosures.

880. Characterization of New HIV Infections among Adults ≥ 45 Years—New York City, North Carolina, San Francisco, 2011–2013

Mary Evans, MD, MPH¹; Anne Patala, MPH^{2,3}; Ellsworth Campbell, MS³; Emily Westheimer, MS³; Cynthia L. Gay, MD, MPH²; Stephanie E. Cohen, MD, MPH⁶; William M. Switzer, MPH²; Philip J. Peters, MD³; ¹Division of HIV and AIDS Prevention, Centers for Disease Control and Prevention, Atlanta, Georgia; ²Division of HIV and AIDS Prevention, ICF International, Atlanta, Georgia; ³Centers for Disease Control and Prevention, Atlanta, Georgia; ⁴Bureau of HIV Prevention and Control, New York City Department of Health and Mental Hygiene, Queens, New York; ⁵North Carolina Department of Public Health, Raleigh, North Carolina; ⁶Population Health, San Francisco Department of Public Health, San Francisco, California

Session: 93. Preventing and Identifying New HIV Infections


Thursday, October 5, 2017: 2:00 PM

Background. While HIV prevention activities are often focused on younger people, older people can also be at risk for HIV infection. We aimed to characterize HIV transmission in older adults.

Methods. The STOP study was a multi-site prospective study of persons with acute HIV infection (AHI) from 2011 to 2013. Older adults were defined as ≥ 45 years and younger persons were 13–44 years. AHI was defined by a negative rapid test but a reactive antigen/antibody or HIV RNA test. We performed bivariate analysis using Pearson's chi-square and odds ratios to examine associations between older age and transmission characteristics. Among persons with HIV-1 polymerase (*pol*) sequences, transmission linkages were inferred when the genetic distance between sequences was $< 1.5\%$ and did not indicate directionality of transmission.

Results. Among 86,836 participants (median age, 29 years; 75.0% male; 51.8% MSM), HIV infection was diagnosed in 176 (1.46%) of 12,036 older adults compared with 1,150 (1.53%) of 74,800 younger people ($P = 0.56$). Among HIV-infected persons, AHI was diagnosed in similar proportions of older and younger people (13.1% vs. 12.6%; $P = 0.86$). Among HIV-infected persons who participated in partner notification ($n = 1,326$), older adults were less likely to report meeting a sex partner online (11.3% vs. 26.9%; OR 0.52, 95% CI = 0.35–0.78) and were less likely to name ≥ 2 sex partners (31.5% vs. 46.8%; OR = 0.28, 95% CI = 0.15–0.53) compared with younger people. Among HIV-infected persons with HIV-1 *pol* sequences ($n = 537$), similar proportions of older and younger people had viruses that genetically linked with another study participant (15.9% vs. 23.5%; OR 0.62, 95% CI 0.31–1.22) (Figure).

Conclusion. In this study, older adults had a similar frequency of newly diagnosed HIV infection, acute infection, and genetic linkage compared with younger people, suggesting that increased HIV prevention efforts may be needed in this population.



Disclosures. All authors: No reported disclosures.

881. HIV Transmission Rates and Factors Associated with Recent HIV Infection:

Results from the Ndihiwa HIV Impact Assessment, South Nyanza, Kenya, 2012
 Elfriede Aggemang, MD, MPH¹; Clement Zeh, PhD²; Irene Mukui, MBChB³; David Maman, PhD⁴; Andrea Kim, PhD, MPH⁵; ¹Division of Global HIV/AIDS and Tuberculosis, Centers for Disease Control and Prevention, Atlanta, Georgia; ²Centers for Disease Control and Prevention, Atlanta, Georgia; ³National AIDS and STI Control Programme (NASCOP), Nairobi, Kenya; ⁴Médecins Sans Frontières, Paris, France; ⁵Division of Global HIV/AIDS and Tuberculosis, CDC, Atlanta, Georgia

Session: 93. Preventing and Identifying New HIV Infections

Thursday, October 5, 2017: 2:00 PM

Background. Identifying populations with high HIV transmission rates is important for prevention and treatment strategies. Persons with recently acquired HIV infection are drivers of HIV transmission due to high levels of HIV viral load (VL). We assessed annual HIV transmission rates and factors associated with recent infection to inform targeted interventions in a hyperendemic region in Kenya.

Methods. The Ndihiwa HIV impact assessment was a population-based survey among persons aged 15–59 years living in South Nyanza, Kenya in 2012. Respondents were tested for HIV using rapid tests per national guidelines and provided blood for centralized testing. Specimens from HIV+ persons were tested for VL and recent infection. Recent infection was defined as normalized optical density value < 1.5 on the Limiting Antigen Enzyme Immunoassay, VL $> 1,000$ copies/mL, and no report of HIV treatment. The annual HIV transmission rate per 100 persons living with HIV (PLHIV) was calculated as HIV incidence divided by HIV prevalence, multiplied by 100. Annualized HIV incidence was estimated, assuming a mean duration of recent infection of 141 days (confidence interval [CI] 123–160). Multivariate analysis identified independent factors associated with recent infection. Estimates were adjusted for survey design.

Results. Of 6,076 persons tested, 1,457 were HIV+, and 28 were recently infected. HIV incidence and prevalence were 1.7% (CI 1.5–2.0) and 24.1% (CI 22.6–25.5), respectively. Per 100 PLHIV, the annual HIV transmission rate was 7.0 and varied by sex (4.6 male vs. 8.3 female), age (5.2 aged 30+ vs. 10.4 aged < 30), and residence (1.4 Kobama vs. 12.0 Riana vs. 12.1 Pala divisions). After controlling for age, sex, and residence, recently infected persons were significantly more likely to reside in Pala division (AOR 8.3, CI 1.1–62.9) than HIV-uninfected persons.

Conclusion. Approximately 7 in 100 PLHIV transmitted to HIV-uninfected persons in South Nyanza in 2012, similar to national rates observed in the 2012 Kenya AIDS Indicator Survey. HIV transmission rates were higher in females than males, younger than older, and Riana and Pala than other divisions. Residence in Pala was a risk factor for recent infection. These findings could guide prioritization of interventions to interrupt HIV transmission in this hyperendemic setting.

Disclosures. All authors: No reported disclosures.

882. Recent HIV Risk Behavior and Partnership Type Predict PrEP Adherence in Men Who Have Sex with Men

Jill Blumenthal, MD¹; Sonia Jain, PhD²; Michael Dube, MD³; Xiaoying Sun, MS²; Eric Ellorin, MAS¹; Martin Hoenig, MD⁴; Katya Corado, MD⁵; David Moore, PhD⁶; Sheldon Morris, MD, MPH¹ and California Collaborative Treatment Group; ¹Medicine, University of California, San Diego, San Diego, California; ²Family and Preventive Medicine, University of California, San Diego, La Jolla, California; ³USC, Los Angeles, California; ⁴Section of Infectious Diseases and Tropical Medicine, Medical University of Graz, Graz, Austria; ⁵Los Angeles Biomedical Research Institute at Harbor-University of California at Los Angeles, Torrance, California; ⁶Psychiatry, University of California at San Diego, San Diego, California

Session: 93. Preventing and Identifying New HIV Infections

Thursday, October 5, 2017: 2:00 PM

Background. Individuals engaging in higher risk behavior are often more adherent to PrEP but it is unclear if partnership type itself affects PrEP adherence. We examined the effect of recent HIV risk behaviors and partnership type on PrEP adherence in men who have sex with men (MSM) taking PrEP.

Methods. CCTG 595 is a 48-week PrEP demonstration study of 398 HIV- at-risk MSM. At baseline and week 48, HIV risk score was estimated as the probability of seroconversion over the next year based on number of condomless anal sex acts with HIV+/unknown partners in the last month and any STI diagnosed at study visit. HIV risk score was categorized as low (< 0.12), moderate (0.12–0.59) and high (> 0.59) risk based on population seroconversion probabilities. Partnership type was assigned as no/single HIV- partner, single HIV+ partner, or multiple partners of any serostatus in the past 3 months. PrEP adherence was estimated by intracellular tenofovir-diphosphate (TFV-DP) levels as a continuous variable at week 48. Statistical methods included McNemar's test, Wilcoxon rank-sum test, and linear regression model where appropriate.

Results. Of 313 MSM who completed week 48, there was no significant change in HIV risk category from baseline to week 48 (low: 44 to 42%; moderate: 27 to 24%; high: 28 to 34%; $P = 0.25$). There was a significant change in partnership type, with the proportion of those with no or single HIV- partnerships increasing (1 to 9%, $P < 0.001$). In univariate analysis, moderate and high-risk groups had higher TFV-DP levels than the low-risk group at week 48 ($P = 0.018$). Participants with no/single HIV- partner had significantly lower TFV-DP levels than those with one HIV+ partner or multiple partners ($P = 0.007$). In a multivariable linear regression model, only low-risk partnerships remained significant where no/single HIV- partnerships were associated with lower TFV-DP levels (mean difference = -344 fmol/punch [$-617, -71$], $P = 0.014$).