


CORRECTION

Correction: Expanding syphilis test uptake using rapid dual self-testing for syphilis and HIV among men who have sex with men in China: A multiarm randomized controlled trial

Cheng Wang, Jason J. Ong, Peizhen Zhao, Ann Marie Weideman, Weiming Tang, M. Kumi Smith, Michael Marks, Hongyun Fu, Weibin Cheng, Fern Terris-Prestholt, Heping Zheng, Joseph D. Tucker, Bin Yang

In [Table 4](#), the first two heading rows are not properly aligned. Please see the correct [Table 4](#) below.



OPEN ACCESS

Citation: Wang C, Ong JJ, Zhao P, Weideman AM, Tang W, Smith MK, et al. (2022) Correction: Expanding syphilis test uptake using rapid dual self-testing for syphilis and HIV among men who have sex with men in China: A multiarm randomized controlled trial. *PLoS Med* 19(8): e1004071. <https://doi.org/10.1371/journal.pmed.1004071>

Published: August 4, 2022

Copyright: © 2022 Wang et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 4. HIV/STI testing and sexual behaviors self-reported by men who had syphilis testing during the trial and initiated at least 1 follow-up survey.

	SOC group	Standard SST group	Lottery incentivized SST group	Standard SST versus SOC		Lottery incentivized SST versus SOC		Lottery incentivized SST versus standard SST	
	Mean (SD)	Mean (SD)	Mean (SD)	MD (95% CI) ^a	<i>p</i> -Value	MD (95% CI)	<i>p</i> -Value	MD (95% CI)	<i>p</i> -Value
Number of male sex partners in the past 3 months									
Month 0 to 3	2.0 (1.0)	2.2 (1.7)	2.3 (1.9)	0.2 (-0.8, 1.1)	0.67	0.3 (-0.6, 1.3)	0.45	0.2 (-0.4, 0.8)	0.60
Month 4 to 6	1.6 (1.0)	2.1 (1.7)	2.0 (1.6)	0.4 (-0.3, 1.1)	0.26	0.3 (-0.4, 1.0)	0.33	-0.1 (-0.7, 0.5)	0.78
	<i>n/N</i> (%)	<i>n/N</i> (%)	<i>n/N</i> (%)	RD (95% CI)	<i>p</i> -Value	RD (95% CI)	<i>p</i> -Value	RD (95% CI)	<i>p</i> -Value
Male sex partners without condoms in the past 3 months									
Month 0 to 3	2/7 (28.6)	22/66 (33.3)	26/68 (38.2)	4.8 (-37.1, 32.4)	0.98	9.7 (-32.8, 37.4)	0.81	4.9 (-11.7, 21.2)	0.60
Month 4 to 6	5/13 (38.5)	20/45 (44.4)	24/62 (38.7)	6.0 (-26.2, 34.1)	0.82	0.3 (-31.5, 26.8)	1.00	-5.7 (-24.8, 13.5)	0.61
Anal group sex in the past 3 months									
Month 0 to 3	2/7 (28.6)	5/74 (6.8)	6/72 (8.3)	-21.8 (-63.1, 3.2)	0.08	-20.2 (-61.3, 5.1)	0.10	1.6 (-7.9, 11.4)	0.77
Month 4 to 6	1/14 (7.1)	5/51 (9.8)	9/69 (13.0)	2.7 (-24.9, 17.1)	0.84	5.9 (-22.1, 19.1)	0.70	3.2 (-9.8, 15.2)	0.61
Ever used substances before or during sex in the past 3 months									
Month 0 to 3	3/7 (42.9)	31/74 (41.9)	40/72 (55.6)	-1.0 (-39.6, 32.7)	0.99	12.7 (-26.3, 46.2)	0.66	13.7 (-3.0, 29.7)	0.11
Month 4 to 6	5/14 (35.7)	18/51 (35.3)	33/69 (47.8)	-0.4 (-31.0, 25.9)	1.00	12.1 (-19.0, 37.4)	0.47	12.5 (-5.8, 29.9)	0.17
Tested for HIV									
Month 0 to 3	7/7 (100.0)	60/74 (81.8)	55/72 (76.4)	-18.9 (-31.3, 25.5)	0.26	-23.6 (-36.6, 20.4)	0.17	-4.7 (-18.3, 8.8)	0.53
Month 4 to 6	14/14 (100.0)	48/51 (94.1)	62/69 (89.9)	-5.9 (-17.1, 18.6)	0.46	-10.1 (-20.6, 15.8)	0.24	-4.3 (-14.9, 7.5)	0.56
Overall	19/20 (95.0)	83/90 (92.2)	80/90 (88.9)	-2.8 (-12.4, 18.3)	0.82	-6.1 (-16.5, 14.7)	0.50	-3.3 (-12.9, 5.7)	0.53
Tested for other STIs^b									
Month 0 to 3	1/7 (14.3)	6/74 (8.1)	6/72 (8.3)	-6.2 (-49.1, 11.2)	0.79	-6.0 (-49.2, 11.5)	0.86	0.2 (-9.7, 10.1)	1.00
Month 4 to 6	4/14 (28.6)	3/51 (5.9)	9/69 (13.0)	-22.7 (-51.7, -0.6)	0.02	-15.5 (-44.8, 6.6)	0.15	7.2 (-4.7, 18.4)	0.22
Overall	8/20 (40.0)	14/90 (15.6)	20/90 (22.2)	-24.4 (-48.5, -0.9)	0.02	-17.8 (-42.5, 5.5)	0.10	6.7 (-5.2, 18.4)	0.28

MD = mean difference. RD = risk (probability) difference expressed as a percentage.

^aDue to rounding error, mean differences may differ by one decimal place from differences obtained by subtracting means listed in table.

^bOther STIs included gonorrhoea, chlamydia, human papillomavirus, and herpes simplex virus.

CI, confidence interval; SOC, standard of care; SST, syphilis self-testing; STI, sexually transmitted infection.

<https://doi.org/10.1371/journal.pmed.1004071.t001>

Reference

1. Wang C, Ong JJ, Zhao P, Weideman AM, Tang W, Smith MK, et al. (2022) Expanding syphilis test uptake using rapid dual self-testing for syphilis and HIV among men who have sex with men in China: A multiarm randomized controlled trial. *PLoS Med* 19(3): e1003930. <https://doi.org/10.1371/journal.pmed.1003930> PMID: 35235573