

The First Report of *Escherichia coli* and *Klebsiella pneumoniae* Strains That Produce Both NDM-5 and OXA-181 in Jiangsu Province, China

Guixiang Tao^{1,*}, Hua Tan^{2,*}, Qian Chen¹

¹Institute of Pediatrics, Children's Hospital of Nanjing Medical University, Nanjing, People's Republic of China; ²Department of Clinical Laboratory, Children's Hospital of Nanjing Medical University, Nanjing, People's Republic of China

*These authors contributed equally to this work

Correspondence: Qian Chen, Institute of Pediatrics, Children's Hospital of Nanjing Medical University, Nanjing, People's Republic of China, Tel +8618951768213, Email js_chenqian@sina.com

Objective: The aim of this study was to analyze the genetic characteristics of three Enterobacteriaceae strains (one strain of *Escherichia coli* and two strains of *Klebsiella pneumoniae*) that produce both the NDM-5 and OXA-181 carbapenemases in pediatric patients.

Methods: Carbapenem-resistant Enterobacteriaceae (CRE) strains were collected from the Children's Hospital Affiliated to Nanjing Medical University in 2022. Resistance genes were detected by PCR. CRE strains that produced both the *bla*_{NDM-5} and *bla*_{OXA-181} genes were further characterized by antimicrobial susceptibility testing, multilocus sequence typing (MLST), plasmid conjugation assay, S1 nuclease-PFGE, Southern blotting and whole-genome sequencing.

Results: Three Enterobacteriaceae strains carrying both the *bla*_{NDM-5} and *bla*_{OXA-181} resistance genes were screened. MLST results showed that the strain of *Escherichia coli* carrying both *bla*_{NDM-5} and *bla*_{OXA-181} was ST410; the two strains of *Klebsiella pneumoniae* with both *bla*_{NDM-5} and *bla*_{OXA-181} were ST2601 and ST759. Conjugation assays showed that the plasmids harboring the *bla*_{NDM-5} and *bla*_{OXA-181} genes were self-transmissible. S1-PFGE and Southern blotting showed that the *bla*_{NDM-5} and *bla*_{OXA-181} genes were located on the plasmid with the size of about 60kb~. The genotyping results showed that the plasmid types were ColKP3 and IncX3.

Conclusion: This is the first report of Enterobacteriaceae strains that produce both NDM-5 and OXA-181 isolated from pediatric patients in China. Active infection control measures are urgently needed to prevent the spread of bacteria in children.

Keywords: *Escherichia coli*, *Klebsiella pneumoniae*, NDM-5, OXA-181, plasmid

Introduction

In recent years, the continuous emergence of carbapenem-resistant Enterobacteriaceae (CRE) strains threatens public health worldwide. The main mechanism of carbapenem resistance in Enterobacteriaceae is the production of KPC (Ambler class A), NDM, VIM, IMP (class B) and OXA-48-like (class D) β -lactamases. New Delhi gold β -lactamase (NDM-1) was first discovered in 2008,¹ and since then, Enterobacteriaceae carrying the *bla*_{NDM} gene have been found around the world. To date, 31 gene subtypes have been found globally.² In 2011, Hornsey et al first identified *bla*_{NDM-5} in a strain of *Escherichia coli* ST648 with a multidrug resistance phenotype in the UK.³ Since then, strains carrying *bla*_{NDM-5} have been found in many countries around the world, including Egypt,⁴ South Korea,⁵ Italy⁶ and China.⁷ Compared with NDM-1, the NDM-5 enzyme has two amino acid substitutions (Val88Leu and Met154Leu), and it has been shown that the NDM-5 enzyme causes a higher level of resistance to carbapenems and broad-spectrum cephalosporins.³

Since 2004, when *Klebsiella pneumoniae* carrying *bla*_{OXA-48} was isolated from a patient living in Turkey,⁸ Enterobacteriaceae strains carrying *bla*_{OXA-48-like} have been found around the world. Several OXA-48-like

carbapenemases have been reported, including OXA-48, OXA-162, OXA-181, OXA-204, OXA-232, OXA-244, and OXA-245.⁹ OXA-181, a variant of OXA-48 with four amino acid substitutions, was first reported in India in 2007.¹⁰ Since then, it has been found mainly in *Escherichia coli* and *Klebsiella pneumoniae* strains in several countries (UK, USA, etc.). The gene encoding OXA-181 is usually located on the IncX3-type plasmid.¹¹ This plasmid carries several carbapenemase genes, including *bla*_{KPC}¹² and *bla*_{NDM}.¹³ To date, OXA-181-producing *Escherichia coli* has been reported in Sichuan¹⁴ and Henan,¹⁵ and OXA-181-producing *Klebsiella pneumoniae* has been reported in Zhejiang.¹⁶ The first strain containing *bla*_{OXA-48-like} and *bla*_{NDM} was reported in Singapore in 2013, followed by Egypt,¹⁷ the United States,¹⁸ Bangladesh,¹⁹ Italy²⁰ and other locations. Here, we report the first *Escherichia coli* and *Klebsiella pneumoniae* strains in China that produce both the NDM-5 and OXA-181 enzymes to analyze the genetic characteristics and environment in pediatric patients.

Materials and Methods

Clinical Data

On January 6, 2022, a 24-day-old newborn with “poor response, abdominal distension” was admitted to the children’s hospital of Nanjing Medical University. After admission, abdominal distension did not improve after symptomatic treatments, including fasting, anti-infection and defecation, so an emergency exploratory laparotomy was performed that night. Necrotizing enterocolitis was confirmed during the operation, and most of the small intestine was necrotic. “Intestinal resection, jejunostomy and intestinal adhesiolysis” were performed; the operation was successful, and the treatment was continued in the neonatal Medical Center after the operation. The patient was diagnosed with neonatal necrotizing enterocolitis, peritonitis and neonatal sepsis and was given meropenem combined with vancomycin to prevent infection. Sputum cultures and drug sensitivity testing on January 8 revealed that the identified *Escherichia coli* (EC73) strain was multidrug resistant and only sensitive to amikacin, polymyxin and tigecycline. The infant was placed in the isolation room for MDRO isolation. On January 10, vancomycin was discontinued, and immunoglobulin was given as a supportive therapy. On March 28, the infant had fever with a peak of 40.5°C and some fatigue. On March 30, a blood-bacterial culture and drug sensitivity testing revealed that there were gram-negative bacilli present. On April 1, blood-bacterial culture and drug sensitivity assays revealed that the identified *Klebsiella pneumoniae* (KP92) strain was sensitive to aztreonam. Meropenem was discontinued, and the anti-bacterial treatment aztreonam was given. After 8 days, the blood cultures were negative for CRE.

On April 24, the third strain (KP100) was extracted from the abdominal effusion of an 8-month-old child three days after enterostomy. As the drug sensitivity results showed sensitivity to aztreonam, the child was given this treatment as an anti-infective. Neither of the patients had traveled in the 30 days before their admission.

Antimicrobial Susceptibility Testing and Identification of Antibiotic Resistance Genes

This study was conducted at the Children’s Hospital of Nanjing Medical University in Jiangsu Province, China. Enterobacteriaceae was identified by a VITEK-2 Compact system (bioMérieux, Marcy-L ‘Etoile, France). Antimicrobial susceptibility testing was performed by broth microdilution. Susceptibility breakpoints were set according to criteria from the Institute of Clinical and Laboratory Standards (CLSI),²¹ except for polymyxin E and tigecycline, whose susceptibility breakpoints were set based on criteria from the European Committee for Antimicrobial Susceptibility Testing (EUCAST) 10.0.²² The quality control (QC) strain *Escherichia coli* ATCC 25922 was used in all tests.

PCR was used to identify *bla*_{NDM-5} and *bla*_{OXA-181} using the following primers previously described in the literature: NDM-5-F, 5’-GAAGCTGAGCACCGCATTAG-3’, NDM-5-R, 5’-GGGCCGTATGAGTGATTGC-3’;²³ OXA-48-like-F, 5’-GCGTGGTTAAGGATGAACAC-3’ and OXA-48-like-R, 5’-CATCAAGTTCAACCCAACCG-3’.²⁴

The amplification system consisted of 50 µL of ddH₂O (20 µL), Multiplex Buffer (25 µL), Primer1 (10 µM) (2 µL), Primer2 (10 µM) (2 µL), and template (1 µL). The PCR amplification conditions were 95 °C for 3 min; 35 cycles of 95 °C for 15s, 55°C for 15s, and 72°C for 15s; and 72 °C for 5 min. The PCR products were identified using a 1.0% agarose gel for electrophoresis, and the positive products were sequenced by Sanger sequencing. The complete sequences were

compared with those reported in GenBank using the BLAST algorithm (<http://www.ncbi.nlm.nih.gov/blast/>). Enterobacteriaceae strains carrying both *bla*_{NDM-5} and *bla*_{OXA-181} were collected for further study.

Multilocus Sequence Typing (MLST)

Referring to the Pasteur MLST website (<http://bigsdB.Pasteur.fr/>), seven housekeeping genes from *Klebsiella pneumoniae*, *rpoB*, *infB*, *phoE*, *mdh*, *pgi*, *gapA*, and *tonB*, and seven housekeeping genes from *Escherichia coli*, *adk*, *fumC*, *gyrB*, *icd*, *mdh*, *purA*, and *recA*, were amplified. DNA sequencing was performed on the positive products. The allelic profiles of the seven housekeeping genes for the respective strains were obtained from the MLST database and then submitted to the MLST website to determine the sequence typing of clinical isolates.

Conjugation Assay

Conjugation assays were performed to test the ability of plasmids carrying *bla*_{NDM-5} and *bla*_{OXA-181} to transfer between strains, and the rifampicin-resistant *E. coli* strain EC600 was used as the receptor. Overnight cultures of donor and recipient strains were mixed and dropped on sterile filter paper at a 1:3 ratio and incubated overnight at 37°C on MH agar plates. *bla*_{NDM-5} and *bla*_{OXA-181} conjugates were selected by screening on dual antibody plates containing imipenem (4mg/L) and rifampicin (600 mg/L).

S1-PFGE and Southern Blotting

Drug-resistant plasmids carrying the *bla*_{NDM-5} and *bla*_{OXA-181} genes were isolated and mapped by S1-PFGE and Southern blotting. Briefly, isolates with *bla*_{NDM-5} and *bla*_{OXA-181} were embedded in gold agarose and digested with S1 nuclease (TaKaRa Biotechnology Co., Dalian, China), and plasmid DNA (Bio-Rad, USA) was analyzed by PFGE electrophoresis. Then, DNA fragments were transferred to positively charged nylon membranes and hybridized specifically with *bla*_{NDM-5} and *bla*_{OXA-181} probes. Probes were synthesized using the DIG High Prime DNA Labeling and Detection Starter Kit I (Roche Applied Sciences, Penzberg, Germany) according to the manufacturer's instructions.

Whole-Genome Sequencing and Analysis

To get a comprehensive picture of the strains that produced both *bla*_{NDM-5} and *bla*_{OXA-181}, the three strains that were identified in this study (EC73, KP92 and KP100) were selected for whole-genome sequencing. Genome sequencing was performed by the Personal Biotechnology Company (Shanghai, China) using the Pacific Biosciences platform and the Illumina MiSeq/HiSeq/Novaseq platform.

The whole-genome shotgun (WGS) strategy and next-generation sequencing (NGS) technology were used to construct libraries of different insertions. The libraries were sequenced on the PacBio Sequel sequencing platform which is based on the Illumina NovaSeq sequencing platform and third-generation single-molecule sequencing technology. FastQC was used for quality control. Using CARD (The Comprehensive Antibiotic Resistance Database), antibiotic resistance genes in the genome sequence can be identified through analysis of antibiotic resistance profiles. PlasmidFinder 2.1 was used to analyze plasmid replicon types. Reference sequences were searched through the NCBI blast database, comparative gene cluster analysis was performed using Easyfig, and comparative genomic circle maps were constructed using CGView.

Results

Antimicrobial Susceptibility and Resistance Gene Testing

The three CRE strains were all resistant to imipenem, meropenem, cephalosporin and enzyme inhibitor compounds, and the *Escherichia coli* strain was only sensitive to amikacin, polymyxin and tigecycline. Both *Klebsiella pneumoniae* strains were sensitive to aztreonam, amikacin, ciprofloxacin, levofloxacin, gentamicin, tobramycin, polymyxin and tigecycline. The minimum inhibitory concentration (MIC) assay results for the three CRE strains are shown in Table 1.

All three CRE strains were found to carry the carbapenemase resistance genes *bla*_{NDM-5} and *bla*_{OXA-181}.

Table 1 The Minimum Inhibitory Concentration of Donor Bacteria, Recipient Bacteria and Conjugant (MIC, $\mu\text{g}/\text{MI}$)

Antibiotics	EC73	J73	KP92	J92	KP100	J100	EC600
Ampicillin	≥ 32	≥ 32	≥ 32	≥ 32	≥ 32	≥ 32	4
Ampicillin/Sulbactam	≥ 32	≥ 32	≥ 32	≥ 32	≥ 32	≥ 32	≤ 2
Piperacillin	≥ 128	≥ 128	≥ 128	≥ 128	≥ 128	≥ 128	≤ 4
Piperacillin/tazobactam	≥ 128	≥ 128	≥ 128	64	≥ 128	64	≤ 4
Cefuroxime	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	4
Cefuroxime Axetil	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	4
Cefotetan	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	≤ 4
Ceftazidime	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	≤ 1
Ceftriaxone	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	≤ 1
Cefepime	≥ 64	16	≥ 32	32	≥ 32	32	≤ 1
Aztreonam	16	≤ 1	≤ 1	≤ 1	≤ 1	≤ 1	≤ 1
Imipenem	8	4	≥ 16	8	≥ 16	≥ 16	≤ 1
Meropenem	8	8	8	4	8	8	≤ 0.25
Amikacin	8	≤ 2	≤ 2	≤ 2	≤ 2	≤ 2	≤ 2
Gentamicin	≥ 16	≤ 1	≤ 1	≤ 1	≤ 1	≤ 1	≤ 1
Tobramycin	≥ 16	≤ 1	≤ 1	≤ 1	≤ 1	≤ 1	≤ 1
Ciprofloxacin	≥ 4	0.5	1	1	1	1	≤ 0.25
Levofloxacin	≥ 8	1	1	1	1	1	≤ 0.25
Trimethoprim/Sulfamethoxazole	≥ 320	≤ 20	≥ 320	≤ 20	≥ 320	≤ 20	≤ 20
Cefazolin	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	≥ 64	≤ 4
Nitrofurantoin	64	≤ 16	128	≤ 16	64	≤ 16	≤ 16
Colistin	≤ 0.5	≤ 0.5	≤ 0.5	≤ 0.5	≤ 0.5	≤ 0.5	≤ 0.5
Tigecycline	≤ 0.5	≤ 0.5	≤ 0.5	≤ 0.5	≤ 0.5	≤ 0.5	≤ 0.5

Notes: 1) In order to avoid duplication, only the first isolate was taken from the same specimen of the same child. In this study, a total of 3 strains were collected after removing the duplicate specimen: *Escherichia coli* (from sputum culture specimens) is denoted by EC73, *Klebsiella pneumoniae* (from blood culture specimens) is denoted by KP92, and *Klebsiella pneumoniae* (from ascites culture specimens) is denoted by KP100. Their conjugation products were denoted by J73, J92 and J100 respectively, and the donor bacteria was EC600. 2) The results for drug resistance have been bolded.

MLST

MLST analysis showed that the EC73 strain is ST410, the KP92 strain ST2601 and the KP100 strain ST759.

Plasmid Conjugation Assay and Gene Location

The plasmids carrying the *bla*_{NDM-5} and *bla*_{OXA-181} genes could be transferred from the donor strain to the recipient strain EC600. The corresponding conjugants are named J73, J92, and J100. The results of PCR amplification showed that all the conjugants carried the *bla*_{NDM-5} and *bla*_{OXA-181} genes. The drug sensitivity results showed that the three conjugants were resistant to carbapenems and enzyme inhibitors, as shown in Table 1.

The results of S1-PFGE and Southern blotting showed that *bla*_{NDM-5} and *bla*_{OXA-181} were located on plasmids of approximately 60 kb in size in the three donor strains and in the three conjugants, as shown in Figure 1.




Figure 1 S1-PFGE profiles and Southern blotting results. From left to right: S1-PFGE profiles; Southern blotting results: The red arrow indicates the hybridization and localization of the plasmid carrying the *bla*_{OXA-181} gene. Southern blotting results: The blue arrow indicates the hybridization and localization of the plasmid carrying the *bla*_{NDM-5} gene.

Note: M: Salmonella H9812 Marker.

Plasmid Sequence Analysis Based on Whole-Genome Sequencing

The CARD database was used to analyze the whole-genome sequencing results, and the predicted resistance genes in the EC73, KP92 and KP100 strains were identical as follows: 1) the β -lactam resistance genes: *bla*_{NDM-5} and *bla*_{OXA-181}; 2) the quinolone resistance gene: *qnrS1*; 3) the aminoglycoside resistance gene: *aadA2*; 4) the sulfa resistance gene: *sul*; 5) the methylene benacil resistance gene: *dfxA12*; and 6) the quaternary ammonium resistance gene: *qacEA1*. The carbapenemase resistance gene *bla*_{OXA-181} was located on the 61,973 bp plasmid 2. The length of EC73 plasmid 2 was 61,973 bp, the average GC content was 48.06%, and there were 81 open reading frames. The length of KP92 plasmid 2 was 61,973 bp, the average GC content was 48.06%, and there were 81 open reading frames. The length of KP100 plasmid 2 was 61,973 bp, the average GC content was 48.06%, and there were 83 open reading frames. According to the PlasmidFinder database from the Center for Genomic Epidemiology, the bacterial drug resistance genes *bla*_{NDM-5} and *bla*_{OXA-181} were on the IncX3 and ColKP3 plasmids in this study. The skeleton structure of these plasmids includes the plasmid conjugation T4SS secretion system, which consists of the Vir gene family (*virB2*, *virB3*, *virB9*, *virB10*, *virB11*), the protein-related genes *parA* and *topB* that help maintain plasmid stability, and the *traG* and *trbM* elements that play a role in conjugation and transfer.

The *bla*_{NDM-5} and *bla*_{OXA-181} carrier genes are located on the Tn3 gene transposable unit. The complete sequences of the three strains (EC73, KP92 and KP100) have been deposited in GenBank, and the accession numbers are SRP399849, SRP398461 and SRP398463. Further BLAST plasmid sequence alignment analysis showed that the nucleotide similarity between EC73 plasmid 2 and NZ_CP024825.1 (*Escherichia coli* isolated from Korea in 2015, *bla*_{NDM-5} located on the IncX3 plasmid) was more than 99%, and KP92 plasmid 2 and NZ_CP026727.1 (*Escherichia coli* isolated from the United Kingdom in 2018, *bla*_{OXA-181} is located on the IncX3 plasmid) had more than 99% nucleotide similarity. The genome comparisons are shown in Figures 2 and 3.

Discussion

Carbapenems have long been considered the last line of defense for multidrug-resistant gram-negative bacterial infections. The emergence of carbapenem-resistant *Klebsiella pneumoniae* is a serious challenge as there are few clinical anti-




Figure 2 Comparison of NZ_CP024825.1, NZ_CP026727.1, KPI100, KP92 and EC73 genomes from the outside in.

infective treatments, especially in pediatric patients. The purpose of this study was to investigate the resistance-related characteristics of CRE strains isolated in the Children's Hospital Affiliated to Nanjing Medical University that produce both the NDM-5 and OXA-181 enzymes to help prevent large-scale hospitalization of patients with CRE in the Children's Hospital. To the best of our knowledge, this is the first time in China that *Escherichia coli* and *Klebsiella pneumoniae* strains that produce both the NDM-5 and OXA-181 enzymes have been found in children.

The two children whose samples were used in this study were both preterm infants who had very low birth weights. Since preterm birth and low birth weight are both risk factors for CRE infection in children,²⁵ children with high risk factors should receive more attention when implementing prevention and control measures in the future. In this study, all clinical strains were sensitive to polymyxin and tigecycline in vitro. Tigecycline is not recommended for children because of the risk of tooth coloring, and polymyxin is not used for the clinical treatment of children in China.²⁶ In addition, aminoglycoside antibiotics and quinolones are rarely used in children because of their ototoxicity and nephrotoxicity.^{27,28} Clinical data showed that both children had good clinical effects after treatment with aztreonam,




Figure 3 Comparative gene cluster analysis. Arrows indicate open reading frames arranged in the direction of transcription. The gene functions are represented by different colors. The red region represents drug resistance genes, the green region represents genes related to plasmid conjugation, the Orange regions represent insertion elements, the blue regions represent transposons, and the black regions encode other genes. Gray shaded areas indicate sequences that are highly similar sequences between plasmids.

which was consistent with the drug resistant profiles of the clinical isolates. Therefore, it is possible that the strains isolated from sputum specimens may be a part of the colonizing flora in the respiratory tract rather than true pathogens. Because this study was retrospective, we were unable to conduct an in-depth analysis of the environment surrounding the patients and the source of the strains.

According to the MLST results in this study, the EC73 strain is ST410, the KP92 strain is ST2601, and the KP100 strain is ST759. The *Escherichia coli* strain identified in this study is ST410, which belongs to the ST23 clonal complex and has been considered the origin of the ST23 complex. *Escherichia coli* strains that are ST410 are widely distributed in Europe (Poland,²⁹ Denmark,³⁰ Italy,³¹ etc.), Asia (South Korea,³² Singapore,³³ India,³⁴ China,³⁵ etc.), Africa,³⁶ North America (United States,³⁷ Mexico,³⁸ etc.) and South America (Brazil,³⁹ Chile⁴⁰ et al), etc. However, *Klebsiella pneumoniae* strains that are ST2601 and ST759 and that produce both NDM-5 and OXA-181, such as the strains used in this study, have never been reported globally.

The results of conjugation experiments showed that the resistance genes bla_{NDM-5} and $bla_{OXA-181}$ in the three CRE strains could be successfully transferred into the recipient strain EC600 by plasmid, which made the conjugants resistant to carbapenems. Plasmid analysis showed that bla_{NDM-5} and $bla_{OXA-181}$ are located on the IncX3 and ColKP3 plasmids, respectively, both of which are the same size (approximately 61 kb). The ColKP3 plasmid is not self-transmitted but can move with the help of other plasmids.⁹ The IncX3 plasmid plays an important role in the transmission of the bla_{NDM-5} gene among Enterobacteriaceae.⁷ IncX3-type plasmids have also been frequently reported to mediate the spread of other NDM

subtypes, including *bla*_{NDM-1}, *bla*_{NDM-4}, *bla*_{NDM-5}, *bla*_{NDM-6}, *bla*_{NDM-7}, *bla*_{NDM-13}, *bla*_{NDM-17}, *bla*_{NDM-19}, *bla*_{NDM-20}, and *bla*_{NDM-21}.⁴¹ Therefore, we need to take effective measures to control the spread of this resistance plasmid among strains.

Whole-genome sequencing of the CRE strains revealed *bla*_{NDM-5} and *bla*_{OXA-181} to be present on the IncX3 and ColKP3 plasmids that were 61,973 bp in size. BLAST analysis of plasmid sequences showed that plasmid 2 of EC73 is highly similar to that in NZ_CP024825.1 (*Escherichia coli* isolated from Korea in 2015) and that plasmid 2 of KP92 is highly similar to that in NZ_CP026727.1 (*Escherichia coli* isolated from the UK in 2018). These findings suggest that the CRE plasmids that carry both of the carbapenemase genes may be derived from *Escherichia coli*. In plasmids, drug resistance genes are often associated with mobile genetic elements, such as transposons (Tn) and insertion sequences (IS). IS26 belongs to the IS6 family, which plays a pivotal role in the transmission of antibiotic resistance genes among gram-negative bacilli.⁴² IS26 encloses multiple resistance genes and lays the groundwork for expression of resistance genes.⁴³ In the three CRE strains in this study, *bla*_{NDM-5} is surrounded by two IS26 on both sides to form the complex transposon in the core region of *dfrA12-aadA2-qacEΔ1-sul*-ISCR1-*dsbD-trpF-bleMBL-bla*_{NDM-5} (not seen in NZ_CP024825.1), and *bla*_{OXA-181} is flanked by two IS26 envelopes to form the *qnrS1-bla*_{OXA-181} resistance gene cassette (also present in NZ_CP026727.1), with the *qnrS1* gene being associated with low levels of fluoroquinolone resistance. The downstream elements of the *bla*_{NDM-5} gene (*bleMBL*, *trpF* and *dsbD*) often appeared in various plasmids, and a large number of drug resistance genes were found to coexist upstream and downstream of NDM-5.^{44,45} The drug resistance genes included *dfrA12*, *aadA2*, *qacEΔ1* and *sul*. The ISCR1 element was first reported by the Australian scholar Stokes.⁴⁶ The ISCR1 element can recognize and carry downstream drug resistance genes via transposition by rolling loop replication. Because ISCR1 can easily carry drug resistance genes, it leads to the spread of drug resistance.⁴⁷ In addition, the three CRE strains in this study were found to carry the disinfectant resistance gene *qacEΔ1*, indicating that Enterobacteriaceae strains have increased resistance to disinfectants that are commonly used in hospitals. The disinfectants commonly used in clinical practice include iodophor, Pasteurian disinfectant, 533 (chlorine-containing disinfectant), Aijiajia (chlorhexidine gluconate), chlorhexidine, and benzalkonium chloride. However, due to the widespread and unreasonable use of various disinfectants, selective pressure on bacteria causes them to become insensitive to disinfectants or makes conventionally used disinfectant concentrations ineffective. The *qacEΔ1* gene encodes a transmembrane protein that is resistant to quaternary ammonium disinfectants (such as benzylammonium bromide and neogeramine). Due to the linkage of the *qacEΔ1* and *sulI* genes, these bacteria are resistant to sulfonamides.⁴⁸ Therefore, we should switch the disinfectants that are regularly used in the clinic to minimize bacterial tolerance to disinfectants.

Although international travel has been reported to accelerate the spread of NDM and OXA-48-like enzymes, neither the children nor the mothers in this study had a history of international travel. At present, more epidemiological studies are urgently needed to better understand the emergence and transmission mechanism of the *bla*_{NDM-5} and *bla*_{OXA-181} genes in China.

Conclusion

In conclusion, this study is the first report of *Escherichia coli* and *Klebsiella pneumoniae* strains that produce carbapenemases (NDM-5 and OXA-181) in pediatric patients in Jiangsu Province, China. These strains are a major threat because they make treatment and recovery more difficult in young children. Plasmid analysis showed the *bla*_{NDM-5} and *bla*_{OXA-181} genes to be located on the IncX3 and ColKP3 plasmids with a size of 61 kb. The IncX3 plasmid, which is a carrier of drug resistance genes and is associated with a risk of spread in pediatric wards, showed a high degree of autonomous transfer between strains. Therefore, we need to actively monitor the genetics of CRE strains, especially in immunocompromised children, and take more stringent control measures to prevent the further spread of these multi-drug-resistant strains in pediatric wards. In addition, clinicians need to fully understand the characteristics of disinfectant resistance genes and their current status to establish sound norms for the use of disinfectants and to reduce the spread of these resistance genes caused by irregular use or abuse of disinfectants.

Ethical Approval

The guardian of the pediatric patients signed informed consent to participate in the study before the study began and this study was conducted in accordance with the Declaration of Helsinki. The Clinical Research Ethics Committee of the Children's Hospital of Nanjing Medical University approved the study (202205054-1), as all samples collected in this work were initially used to diagnose patient care without increasing the patient's medical costs and suffering.

Consent for Publication

Written informed consent was provided by the patient guardian for the publication of the case details.

Funding

This research was supported by the Second Level of the Nanjing Health Youth Talents Training Project (QRX17078).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Yong D, Toleman MA, Giske CG, et al. Characterization of a new metallo-beta-lactamase gene, bla(NDM-1), and a novel erythromycin esterase gene carried on a unique genetic structure in Klebsiella pneumoniae sequence type 14 from India. *Antimicrob Agents Chemother.* 2009;53(12):5046–5054. doi:10.1128/AAC.00774-09
2. Li X, Zhao D, Li W, Sun J, Zhang X. Enzyme inhibitors: the best strategy to tackle superbug NDM-1 and its variants. *Int J Mol Sci.* 2021;23(1):197. doi:10.3390/ijms23010197
3. Hornsey M, Phee L, Wareham DW. A novel variant, NDM-5, of the New Delhi metallo-β-lactamase in a multidrug-resistant Escherichia coli ST648 isolate recovered from a patient in the United Kingdom. *Antimicrob Agents Chemother.* 2011;55(12):5952–5954. doi:10.1128/AAC.05108-11
4. Ramadan H, Gupta SK, Sharma P, et al. Circulation of emerging NDM-5-producing Escherichia coli among humans and dogs in Egypt. *Zoonoses Public Health.* 2020;67(3):324–329. doi:10.1111/zph.12676
5. Hong JS, Song W, Jeong SH. Molecular characteristics of NDM-5-producing Escherichia coli from a cat and a dog in South Korea. *Microb Drug Resist.* 2020;26(8):1005–1008. doi:10.1089/mdr.2019.0382
6. Alba P, Taddei R, Cordaro G, et al. Carbapenemase IncF-borne bla_{NDM-5} gene in the E. coli ST167 high-risk clone from canine clinical infection, Italy. *Vet Microbiol.* 2021;256:109045. doi:10.1016/j.vetmic.2021.109045
7. Tian D, Wang B, Zhang H, et al. Dissemination of the bla_{NDM-5} gene via IncX3-type plasmid among Enterobacteriaceae in children. *mSphere.* 2020;5(1):e00699–19. doi:10.1128/mSphere.00699-19
8. Poirel L, Heritier C, Tolun V, Nordmann P. Emergence of oxacillinase-mediated resistance to imipenem in Klebsiella pneumoniae. *Antimicrob Agents Chemother.* 2004;48(1):15–22. doi:10.1128/AAC.48.1.15-22.2004
9. Pitout J, Peirano G, Kock MM, Strydom K-A, Matsumura Y. The global ascendancy of OXA-48-type carbapenemases. *Clin Microbiol Rev.* 2019;33(1):e00102–19. doi:10.1128/CMR.00102-19
10. Castanheira M, Deshpande LM, Mathai D, Bell JM, Jones RN, Mendes RE. Early dissemination of NDM-1- and OXA-181-producing Enterobacteriaceae in Indian hospitals: report from the SENTRY Antimicrobial Surveillance Program, 2006–2007. *Antimicrob Agents Chemother.* 2011;55(3):1274–1278. doi:10.1128/AAC.01497-10
11. Izdebski R, Baraniak A, Zabicka D, et al. Enterobacteriaceae producing OXA-48-like carbapenemases in Poland, 2013–January 2017. *J Antimicrob Chemother.* 2018;73(3):620–625. doi:10.1093/jac/dkx457
12. Fuga B, Ferreira ML, Cerdeira LT, et al. Novel small IncX3 plasmid carrying the bla_{KPC-2} gene in high-risk Klebsiella pneumoniae ST11/CG258. *Diagn Microbiol Infect Dis.* 2020;96(2):114900. doi:10.1016/j.diagmicrobio.2019.114900
13. Zhu W, Wang X, Qin J, Liang W, Shen Z. Dissemination and stability of the bla_{NDM-5}-carrying IncX3-type plasmid among multiclonal Klebsiella pneumoniae isolates. *mSphere.* 2020;5(6):e00917–20. doi:10.1128/mSphere.00917-20
14. Liu Y, Feng Y, Wu W, et al. First report of OXA-181-producing Escherichia coli in china and characterization of the isolate using whole-genome sequencing. *Antimicrob Agents Chemother.* 2015;59(8):5022–5025. doi:10.1128/AAC.00442-15
15. Qin S, Cheng J, Wang P, Feng X, Liu HM. Early emergence of OXA-181-producing Escherichia coli ST410 in China. *J Glob Antimicrob Resist.* 2018;15:215–218. doi:10.1016/j.jgar.2018.06.017
16. Liu C, Fang Y, Zeng Y, et al. First report of OXA-181-producing Klebsiella pneumoniae in China. *Infect Drug Resist.* 2020;13:995–998. doi:10.2147/IDR.S237793
17. Gamal D, Fernández-Martínez M, El-Defrawy I, Ocampo-Sosa AA, Martínez-Martínez L. First identification of NDM-5 associated with OXA-181 in Escherichia coli from Egypt. *Emerg Microbes Infect.* 2016;5(4):e30. doi:10.1038/emi.2016.24
18. Rojas LJ, Hujer AM, Rudin SD, et al. NDM-5 and OXA-181 beta-lactamases, a significant threat continues to spread in the Americas. *Antimicrob Agents Chemother.* 2017;61(7):e00454–17. doi:10.1128/AAC.00454-17
19. Okanda T, Haque A, Koshikawa T, et al. Characteristics of carbapenemase-producing Klebsiella pneumoniae isolated in the intensive care unit of the largest tertiary hospital in Bangladesh. *Front Microbiol.* 2021;11:612020. doi:10.3389/fmicb.2020.612020
20. Marchetti VM, Bitar I, Mercato A, et al. Complete nucleotide sequence of plasmids of two Escherichia coli strains carrying bla_{NDM-5} and bla_{NDM-5} and bla_{OXA-181} from the same patient. *Front Microbiol.* 2020;10:3095. doi:10.3389/fmicb.2019.03095

21. Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing*. 32nd ed. Wayne, PA: Clinical and Laboratory Standards Institute; 2022.
22. European Committee on Antimicrobial Susceptibility Testing. *Breakpoint Tables for Interpretation of MICs and Zone Diameters. Version 12.0*. Växjö, Sweden: European Committee on Antimicrobial Susceptibility Testing; 2022.
23. Rogers BA, Sidjabat HE, Silvey A, et al. Treatment options for New Delhi metallo-beta-lactamase-harboring Enterobacteriaceae. *Microb Drug Resist*. 2013;19(2):100–103. doi:10.1089/mdr.2012.0063
24. Wang X, Wang Y, Zhou Y, et al. Emergence of a novel mobile colistin resistance gene, mcr-8, in NDM-producing *Klebsiella pneumoniae*. *Emerg Microbes Infect*. 2018;7(1):122. doi:10.1038/s41426-018-0124-z
25. Nour I, Eldeglia HE, Nasef N, Shouman B, Abdel-Hady H, Shabaan AE. Risk factors and clinical outcomes for carbapenem-resistant Gram-negative late-onset sepsis in a neonatal intensive care unit. *J Hosp Infect*. 2017;97(1):52–58. doi:10.1016/j.jhin.2017.05.025
26. Chiotos K, Hayes M, Gerber JS, Tamma PD. Treatment of carbapenem-resistant Enterobacteriaceae infections in Children. *J Pediatric Infect Dis Soc*. 2020;9(1):56–66. doi:10.1093/jpids/piz085
27. Montagnani C, Tersigni C, D'Arienzo S, et al. Resistance patterns from urine cultures in children aged 0 to 6 years: implications for empirical antibiotic choice. *Infect Drug Resist*. 2021;14:2341–2348. doi:10.2147/IDR.S293279
28. Li S, Chen Z, Huang L, et al. Safety of quinolones in children: a systematic review and meta-analysis. *Paediatr Drugs*. 2022;24(5):447–464. doi:10.1007/s40272-022-00513-2
29. Rzeczowska M, Wolkowicz T, Zacharczuk K, et al. Draft genome sequence of an *Escherichia coli* ST410 isolate co-harboring bla_{CTX-M-15}, bla_{CMY-42}, bla_{OXA-1}, aac(3)-IIa and aac(6')-Ib-cr genes with gyrA and parC mutations isolated from a paediatric patient in Poland. *J Glob Antimicrob Resist*. 2019;16:120–122. doi:10.1016/j.jgar.2018.11.024
30. Roer L, Overballe-Petersen S, Hansen F, et al. *Escherichia coli* sequence type 410 is causing new international high-risk clones. *mSphere*. 2018;3(4):e00337–18. doi:10.1128/mSphere.00337-18
31. Piazza A, Comandatore F, Romeri F, et al. First report of an ST410 OXA-181 and CTX-M-15 coproducing *Escherichia coli* clone in Italy: a whole-genome sequence characterization. *Microb Drug Resist*. 2018;24(8):1207–1209. doi:10.1089/mdr.2017.0366
32. Kim JS, Yu JK, Jeon SJ, et al. Dissemination of an international high-risk clone of *Escherichia coli* ST410 co-producing NDM-5 and OXA-181 carbapenemases in Seoul, Republic of Korea. *Int J Antimicrob Agents*. 2021;58(6):106448. doi:10.1016/j.ijantimicag.2021.106448
33. Zhong Y, Guo S, Schlundt J, Kwa AL. Identification and genomic characterization of a bla_{NDM-5}-harbouring MDR plasmid in a carbapenem-resistant *Escherichia coli* ST410 strain isolated from a natural water environmental source. *JAC Antimicrob Resist*. 2022;4(4):dlac071.
34. Devanga Ragupathi NK, Vasudevan K, Venkatesan M, Veeraraghavan B. First Indian report on B4/H24RxC ST410 multidrug-resistant *Escherichia coli* from bloodstream infection harbouring bla_{OXA-181} and bla_{CTX-M-15}. *J Glob Antimicrob Resist*. 2020;22:568–570. doi:10.1016/j.jgar.2020.06.013
35. Gu J-N, Chen L, Weng X-B, Yang X-Y, Pan D-M. Clinical and microbiological characteristics of a community-acquired carbapenem-resistant *Escherichia coli* ST410 isolate harbouring bla_{NDM-5}-encoding IncX3-type plasmid from blood. *Front Med*. 2021;8:658058. doi:10.3389/fmed.2021.658058
36. Negeri AA, Mamo H, Gurung JM, et al. Antimicrobial resistance profiling and molecular epidemiological analysis of extended spectrum beta-lactamases produced by extraintestinal invasive *Escherichia coli* isolates from Ethiopia: the presence of international high-risk clones ST131 and ST410 revealed. *Front Microbiol*. 2021;12:706846. doi:10.3389/fmicb.2021.706846
37. Chen L, Peirano G, Kreiswirth BN, Devinney R, Pitout JDD. Acquisition of genomic elements were pivotal for the success of *Escherichia coli* ST410. *J Antimicrob Chemother*. 2022;77(12):3399–3407. doi:10.1093/jac/dkac329
38. Magaña-Lizárraga JA, Gómez-Gil B, Rendón-Maldonado JG, Delgado-Vargas F, Vega-López IF, Báez-Flores ME. Genomic profiling of antibiotic-resistant *Escherichia coli* isolates from surface water of agricultural drainage in North-Western Mexico: detection of the international high-risk lineages ST410 and ST617. *Microorganisms*. 2022;10(3):662.
39. Furlan J, Gonzalez I, Ramos PL, Stehling EG. International high-risk clone of multidrug-resistant CTX-M-8-producing *Escherichia coli* C-ST410 infecting an elephant (*Loxodonta africana*) in a zoo. *J Glob Antimicrob Resist*. 2020;22:643–645. doi:10.1016/j.jgar.2020.06.018
40. Ortega-Paredes D, Barba P, Mena-López S, Espinel N, Zurita J. *Escherichia coli* hyperepidemic clone ST410-A harboring bla_{CTX-M-15} isolated from fresh vegetables in a municipal market in Quito-Ecuador. *Int J Food Microbiol*. 2018;280:41–45. doi:10.1016/j.ijfoodmicro.2018.04.037
41. Wu W, Feng Y, Tang G, Qiao F, McNally A, Zong Z. NDM metallo-beta-lactamases and their bacterial producers in health care settings. *Clin Microbiol Rev*. 2019;32(2):e00115–18. doi:10.1128/CMR.00115-18
42. Varani A, He S, Siguiet P, Ross K, Chandler M. The IS6 family, a clinically important group of insertion sequences including IS26. *Mob DNA*. 2021;12(1):11. doi:10.1186/s13100-021-00239-x
43. Pong CH, Harmer CJ, Ataide SF, Hall RM. An IS 26 variant with enhanced activity. *FEMS Microbiol Lett*. 2019;366(3):fnz031. doi:10.1093/femsle/fnz031
44. Chowdhury G, Ramamurthy T, Das B, et al. Characterization of NDM-5 carbapenemase-encoding gene (bla_{NDM-5}) – positive multidrug resistant commensal *Escherichia coli* from diarrheal patients. *Infect Drug Resist*. 2022;15:3631–3642. doi:10.2147/IDR.S364526
45. Zou H, Berglund B, Wang S, et al. Emergence of bla_{NDM-1}, bla_{NDM-5}, bla_{KPC-2} and bla_{IMP-4} carrying plasmids in *Raoultella* spp. in the environment. *Environ Pollut*. 2022;306:119437. doi:10.1016/j.envpol.2022.119437
46. Stokes HW, Tomaras C, Parsons Y, Hall RM. The partial 3'-conserved segment duplications in the integrons In6 from pSa and In7 from pDGO100 have a common origin. *Plasmid*. 1993;30(1):39–50. doi:10.1006/plas.1993.1032
47. Lallement C, Pasternak C, Ploy M-C, Jové T. The role of ISCR1-borne POUT promoters in the expression of antibiotic resistance genes. *Front Microbiol*. 2018;9:2579. doi:10.3389/fmicb.2018.02579
48. Nowrotek M, Kotlarska E, Luczkiewicz A, Felis E, Sochacki A, Miksch K. The treatment of wastewater containing pharmaceuticals in microcosm constructed wetlands: the occurrence of integrons (int1–2) and associated resistance genes (sul1–3, qacEA1). *Environ Sci Pollut Res Int*. 2017;24(17):15055–15066. doi:10.1007/s11356-017-9079-1

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>