

Role of Flippases in Protein Glycosylation in the Endoplasmic Reticulum

Supplementary Issue: Cellular Anatomy of Lipid Traffic

Jeffrey S. Rush

Department of Molecular and Cellular Biochemistry, University of Kentucky College of Medicine, Lexington, KY, USA.

ABSTRACT: Glycosylation is essential to the synthesis, folding, and function of glycoproteins in eukaryotes. Proteins are co- and posttranslationally modified by a variety of glycans in the endoplasmic reticulum (ER); modifications include C- and O-mannosylation, N-glycosylation, and the addition of glycosylphosphatidylinositol membrane anchors. Protein glycosylation in the ER of eukaryotes involves enzymatic steps on both the cytosolic and luminal surfaces of the ER membrane. The glycans are first assembled as precursor glycolipids, on the cytosolic surface of the ER, which are tethered to the membrane by attachment to a long-chain polyisoprenyl phosphate (dolichol) containing a reduced α -isoprene. The lipid-anchored *building blocks* then migrate transversely (flip) across the ER membrane to the luminal surface, where final assembly of the glycan is completed. This strategy allows the cell to export high-energy biosynthetic intermediates as lipid-bound glycans, while constraining the glycosyl donors to the site of assembly on the membrane surface. This review focuses on the flippases that participate in protein glycosylation in the ER.

KEYWORDS: flippase, glycosylation, dolichol, congenital disorder of glycosylation

SUPPLEMENT: Cellular Anatomy of Lipid Traffic

CITATION: Rush. Role of Flippases in Protein Glycosylation in the Endoplasmic Reticulum. *Lipid Insights* 2015;8(S1) 45–53 doi:10.4137/LPI.S31784.

TYPE: Review

RECEIVED: October 20, 2015. **RESUBMITTED:** January 12, 2016. **ACCEPTED FOR PUBLICATION:** January 15, 2016.

ACADEMIC EDITOR: Tim Levine, Editor in Chief

PEER REVIEW: Three peer reviewers contributed to the peer review report. Reviewers' reports totaled 1233 words, excluding any confidential comments to the academic editor.

FUNDING: Parts of this work were supported by NIH grants R01 GM36065 and GM102129 (C. Waechter). The author confirms that the funder had no influence over the study design, content of the article, or selection of this journal.

COMPETING INTERESTS: Author discloses no potential conflicts of interest.

COPYRIGHT: © the authors, publisher and licensee Libertas Academica Limited. This is an open-access article distributed under the terms of the Creative Commons CC-BY-NC 3.0 License.

CORRESPONDENCE: jrush@uky.edu

Paper subject to independent expert single-blind peer review. All editorial decisions made by independent academic editor. Upon submission manuscript was subject to anti-plagiarism scanning. Prior to publication all authors have given signed confirmation of agreement to article publication and compliance with all applicable ethical and legal requirements, including the accuracy of author and contributor information, disclosure of competing interests and funding sources, compliance with ethical requirements relating to human and animal study participants, and compliance with any copyright requirements of third parties. This journal is a member of the Committee on Publication Ethics (COPE).

Published by Libertas Academica. Learn more about this journal.

Introduction

Eukaryotic cells are subdivided into organelles by lipid bilayers, with a variety of unique functions and lipid compositions.¹ In addition, the lipid components in the plasma membrane are distributed asymmetrically between the two monolayers with the aminophospholipids, phosphatidylserine and phosphatidylethanolamine, located on the cytoplasmic leaflet, and the choline-containing lipids, sphingomyelin and phosphatidylcholine, in the exoplasmic leaflet. As the cytoplasmic surface of the endoplasmic reticulum (ER) is the site of synthesis of the majority of eukaryotic phospholipids, at least 50% of the newly synthesized phospholipids must flip to the luminal monolayer to support the uniform expansion of the membrane bilayer. The observed final asymmetric distribution of phospholipids in the plasma membrane is the net result of competing bidirectional, energy-independent flip/floppases and ATP-requiring, phospholipid pumps catalyzing the unidirectional transport of phospholipids across the membrane bilayers. Current models suggest that energy-independent flip/floppases mediate the transmembrane movement of phospholipids in biogenic (self-assembling) membranes, such as the ER, and the P-type and ATP-binding cassette transporters function to maintain the transbilayer phospholipid asymmetry in membranes, such as the plasma membrane.^{2–5} The glycosyl donors employed in protein N-glycosylation,

C- and O-mannosylation, and glycosylphosphatidylinositol membrane (GPI) anchoring are assembled on both sides of the ER membrane and require the participation of the bidirectional, energy-independent class of flippase proteins to enter the lumen of the ER and participate as glycosyl donors.^{5–7} While more properly functioning as *scramblases*, these transporters will be referred to as flippases, but they should not be confused with the phospholipid flip- and floppases.

Protein N-glycosylation requires the synthesis of a precursor oligosaccharide, containing three glucose (Glc), nine mannose (Man), and two N-acetyl glucosamine (GlcNAc) residues, attached to a long-chain polyisoprenyl alcohol (dolichol, Dol) via a diphosphate bridge (Glc₃Man₉GlcNAc₂-P-P-Dol, G3-DLO [dolichol-linked oligosaccharide]), as illustrated in Figure 1.^{6,8} G3-DLO synthesis is initiated (Step 1, Fig. 1) on the cytoplasmic leaflet of the ER with the synthesis of a dolichol-diphosphate-linked oligosaccharide intermediate, Man₅GlcNAc₂-P-P-Dol (M5-DLO),⁹ formed by the sequential addition of sugars to dolichyl phosphate (Dol-P) using UDP-GlcNAc (catalyzed by a complex of DPAGT1 and ALG13/14 [asparagine-linked glycosylation])¹⁰ and GDP-Man (catalyzed by a second complex composed of ALG1, ALG2, and ALG11)¹¹ as glycosyl donors (Step 1, Fig. 1). The M5-DLO intermediate then migrates (*flips*) from the cytosolic leaflet of the ER to the luminal monolayer (Flippase I, Fig. 1), where




Figure 1. Topological model for DLO synthesis in the ER. M5-DLO (Step 1), Man-P-Dol (Step 2), and Glc-P-Dol (Step 3) are synthesized on the cytosolic surface of the ER membrane from appropriate sugar-nucleotide donors and Dol-P and diffuse transversely to the luminal monolayer where they participate in the synthesis of full-length G3-DLO. The polar head groups of the dolichol-linked intermediates do not spontaneously cross the ER membrane but are escorted across the highly hydrophobic lipid bilayer by a specialized class of membrane transporters referred to as flippases, numbered arbitrarily as indicated (Flippase 1: M5-DLO Flippase; Flippase 2: Man-P-Dol Flippase; Flippase 3: Glc-P-Dol Flippase; and Flippase 4: Dol-P Flippase). The G3 oligosaccharide is transferred to protein nascent chains releasing a molecule of Dol-P-P, which is dephosphorylated by DolPP1; Dol-P is then returned to the cytosolic monolayer by Flippase 4 (Step 4) for utilization in further rounds of DLO synthesis.

four more Man residues are added using mannosyl phosphoryldolichol (Man-P-Dol) as mannosyl donor and three Glc residues are added using glucosyl phosphoryldolichol (Glc-P-Dol) as glucosyl donor to form the full-length G3-DLO. Man-P-Dol and Glc-P-Dol are also synthesized on the cytosolic face of the ER membrane, catalyzed by DPM1/2/3 (Step 2, Fig. 1) and ALG5 (Step 3, Fig. 1), respectively. Following synthesis on the cytosolic monolayer, the dolichol-linked intermediates are *flipped* to the luminal monolayer by Flippase 2 and Flippase 3 (Fig. 1), where they function as glycosyl donors in this pathway. Following en bloc transfer of the precursor oligosaccharide to protein nascent chains, the carrier lipid, Dol-P-P, is discharged on the luminal leaflet. Dol-P-P is dephosphorylated by Dol-P-P phosphatase (DolPP1),^{12,13} to form Dol-P, which is then flipped back to the cytosolic monolayer by Flippase 4 (Step 4, Fig. 1), to be reutilized for further rounds of DLO synthesis.¹⁴

Man-P-Dol also provides the mannosyl units for protein C- and O-mannosylation and GPI anchor biosynthesis, as illustrated in Figure 2.¹⁵ Similar to DLO assembly, this Man-P-Dol is synthesized on the cytosolic surface of the ER membrane and must be transported to the luminal monolayer to function as mannosyl donor. Presumably, the Dol-P molecules released by the mannosyltransferase reactions are returned to the cytosolic surface for reutilization by the same mechanism as the Dol-P formed from Dol-P-P and the Man-P-Dol and Glc-P-Dol mediated reactions in G3-DLO synthesis on the luminal leaflet.⁶

There is substantial evidence supporting the hypothesis that the transbilayer diffusion of the dolichol-linked intermediates is facilitated by specialized membrane transporters, termed “flippases”. The highly hydrophobic nature of the inner core of the lipid bilayer presents a substantial biophysical barrier to the spontaneous passage of the hydrophilic head




Figure 2. Man-P-Dol acts as mannose donor for the synthesis of M9-DLO, GPI anchors, and C- and O-mannosylated proteins. Man-P-Dol is synthesized on the cytosolic monolayer of the ER membrane from GDP-Man and Dol-P and flips (Flippase 2) to the luminal monolayer where it functions as mannose donor for four mannose transferases in the DLO pathway and three mannose transferases in GPI anchor assembly. At least seven O-mannose transferases have been characterized in yeasts and mammals. The number of C-mannose transferases remains unknown. Following mannose transfer, Dol-P is released and returned to the cytosolic monolayer by Flippase 4.

group of the amphipathic dolichol-linked intermediates.^{5,16–18} Although all of the glycosyltransferases required to assemble the DLO intermediates have been described, the identities of the proteins that catalyze the transmembrane movements of the dolichol-linked intermediates and the precise roles of two proteins involved in DLO assembly (Lec35 and RFT1) have not yet been clearly defined.

The description of this enigmatic class of membrane transporters remains a high priority. It is likely that currently undiagnosed congenital disorders of glycosylation (CDG)^{19–24} will be traced to defects in flippases in the protein glycosylation pathways. CDG is a rapidly expanding family of genetic diseases attributed to the defective synthesis of glycoproteins and glycolipids, and there are presently over 100 genetic diseases resulting from mutations in glycosylation-related genes.²³ CDG patients present with a wide range of clinical symptoms varying from relatively mild, with primarily gastrointestinal effects, to very severe, involving multi-organ dysfunctions, skeletal abnormalities, and mental and psychomotor delay, which often result in embryonic lethality or infant mortality.²¹ As many of these disorders are extremely

rare, it is not presently known if the range of clinical symptoms can be considered diagnostic of a specific disorder or are due to the *leakiness* of the defect.²⁰ It is speculative to attempt to predict the clinical features anticipated from specific flippase defects; however, some general expectations can be proposed. CDGs arising from defects in Man-P-Dol synthesis and utilization (DPM1, 2, and 3-CDG and MPDU1-CDG) appear to be quite severe due to the additive effects on protein O-mannosylation (α -dystroglycanopathies), GPI anchoring, and DLO assembly, which would imply that a Man-P-Dol flippase deficiency would have severe consequences. On the other hand, defects in Glc-P-Dol utilization (ALG6-CDG and ALG8-CDG), are much less severe, suggesting that a defect in Glc-P-Dol flippase might be less severe than the Man-P-Dol flippase deficiency. M5-DLO flippase deficiency might be expected to be similar to CDGs involving the early stages in DLO synthesis (DPAGT1-CDG, ALG13/14-CDG, ALG1-CDG, ALG2-CDG, and ALG11-CDG), which are typically severe. The consequences of the prospective Dol-P recycling flippase are somewhat harder to predict. Dol-P flippase deficiency may be similar in scope to those



arising from inadequate Dol-P synthesis (DHDDS-CDG, SRD5A3-CDG, and DK-CDG), and it is possible that deficiencies in Dol-P recycling could be partially alleviated by the upregulation of Dol-P synthesis de novo.²⁵ It is interesting to note that a CDG attributed to the recycling DolPP1 has not yet been reported.

Despite continued keen interest in the putative flippases of the protein glycosylation pathways of the ER, they remain undescribed due, primarily, to technical difficulties inherent in assaying their biochemical activity. Current analytical approaches are based on localizing a specific DLO intermediate to one leaflet of the bilayer by using chemical modification with a membrane-impermeant reagent,^{5,26,27} or lectin capture,^{5,28} or transport of water-soluble analogs containing short isoprenyl chains.^{29–32} The existing evidence suggests that the DLO flippases are energy independent and bidirectional and catalyze an extremely rapid equilibration of DLO intermediates between the monolayers, with half-times in the range of a minute or less. These properties, along with the inherent fragility of transbilayer asymmetry, pose formidable experimental challenges to using classical biochemical methodologies. This article reviews the information that is currently available concerning the flippase activities of the protein glycosylation pathways of the eukaryotic ER. Results and observations from some previously unpublished studies are included. All studies were conducted in accordance with federal guidelines and approved by the Institutional Animal Care and Use Committee of the University of Kentucky.

Man-P-Dol Flippase

As noted earlier, Man-P-Dol is synthesized on the cytosolic surface of the ER from GDP-Man and dolichol phosphate by Dpm1 in *Saccharomyces cerevisiae* and a complex of three polypeptide subunits (Dpm1, Dpm2, and Dpm3) in other eukaryotes.^{15,33} Following synthesis, Man-P-Dol diffuses transversely through the ER membrane to the luminal monolayer, where it functions as a mannosyl donor for the C-mannosylation of tryptophan residues in thrombospondin repeats of cell surface receptors, the O-mannosylation of serine and threonine units of α -dystroglycan, the addition of four Man residues of the DLO intermediate of the N-glycosylation pathway, and three (or more) Man units of GPI anchor intermediates,^{15,34} as illustrated in Figure 2.

The molecular description of the proteins involved in the transmembrane diffusion of Man-P-Dol has been a subject of interest for a number of years.^{35,36} However, due to the absence of a convenient biochemical assay for Man-P-Dol flippase activity, progress has been slow. Genetic approaches have also failed to identify likely flippase candidate proteins, including two recent screens in Hap1 cells (haploid mammalian cell lines) for defects in GPI assembly. Both of these screens identified all of the proteins known to participate in GPI formation but failed to identify any proteins that could plausibly be involved in the transmembrane flipping of

either Man-P-Dol or glucosamine-PI.^{37,38} There are a number of potential explanations for this observation, including the possibility, as is true in yeast, that an essential protein is synthesized by a flippase requiring pathway, in this case an essential GPI-anchored protein,^{39,40} or that multiple genomic copies of flippase genes are present, making it difficult to eliminate all of the active transcripts. It is possible that flippases from other pathways possess overlapping substrate specificity, allowing complementation of the defect by other flippases, or the flippase activity may reside in a multitasking protein with another essential function as a second *moonlighting* activity.⁴¹ This final possibility is supported by the recent discovery that Rhodopsin, in addition to its role as a visual pigment and G protein-coupled receptor, also functions as an active phospholipid flippase in photoreceptor disc membranes.^{27,42}

Two strategies for the in vitro measurement of Man-P-Dol flippase activity have been developed. One approach, relies on the use of a water-soluble version of Man-P-Dol containing citronellyl phosphate (Cit-P), a short-chain analog of Dol-P with a 10 carbon isoprenyl chain and a saturated α -isoprene. This assay relies on the assumption that recognition of the polar head group and its movement across the bilayer are the most critical aspects of the flipping process and measures Man-P-Dol flipping as a simple transport process into sealed rat liver ER vesicles.^{29,32,43} This is a variation of the approach used by Bishop and Bell to assay flip-flopping of phosphatidylcholine in ER vesicles.⁴⁴

The selection of Man-P-Cit as an appropriate analog for Man-P-Dol is strongly supported by the observation that Man-P-Cit is a suitable substrate for all the Man-P-Dol-mediated mannosyltransferases in the DLO pathway⁴⁵ and in GPI assembly,⁴⁶ in O-mannosylation,⁴⁷ and in C-mannosylation (J. S. Rush, C. J. Waechter, and J. Hofsteenge, 2008, unpublished observation). Furthermore, Man-P-Cit was able to allosterically stimulate GlcNAc-P-P-Dol synthesis,⁴⁸ similar to native Man-P-Dol, and was able to cross the ER membrane and function as a mannosyl donor in streptolysin-O-permeabilized Lec35 cells under conditions in which the Man-P-Dol utilization defect was preserved.³⁴ Lec35 cells exhibit a defect in Man-P-Dol and Glc-P-Dol utilization that, apparently, does not involve dolichol-intermediate flipping.³⁴

Man-P-Cit is transported into the lumen of intact rat liver ER vesicles by a process that is time dependent, reversible, saturable, protease sensitive, stereoselective, enriched in the ER, sensitive to incubation with diethylpyrocarbonate (DEPC), and energy independent.²⁹ The activity was not observed in intact erythrocytes or synthetic liposomes²⁹ but was observed in proteoliposomes reconstituted from octyl glucoside and Triton X-100 soluble detergent extracts of rat liver ER membrane proteins.⁴³ Furthermore, the soluble detergent extracts could be fractionated by ion exchange and partially purified.⁴³ It is observed that when Man-P-Cit uptake in intact rat liver ER vesicles is measured over a range of temperatures, an Arrhenius plot of the rate of transport reveals an inflection point



at $\sim 16^\circ\text{C}$, correlating exactly with a corresponding inflection point in the rate of increase in the incorporation of $[\text{^3H}]$ mannose units into $[\text{^3H}]\text{Man}_{6-9}\text{-DLO}$ from $\text{GDP-}[\text{^3H}]\text{Man}$ (which requires the participation of $[\text{^3H}]\text{Man-P-Dol}$ as an intermediate), performed in a parallel incubation (J. S. Rush and C. J. Waechter, 2007, unpublished observation). In this experiment, the increase in the initial rate of incorporation of $[\text{^3H}]\text{mannose}$ into $[\text{^3H}]\text{Man-P-Dol}$ is linear with respect to temperature, as would be anticipated for a process occurring on only one side of the membrane bilayer. This observation is consistent with Man-P-Cit uptake and Man-P-Dol flipping occurring by a transport process, which is affected similarly by membrane fluidity and supports the conclusion that Man-P-Cit transport accurately reflects flippase activity.

An alternative approach utilized native $[\text{^3H}]\text{Man-P-Dol}$ symmetrically reconstituted into sealed synthetic proteoliposomes and a membrane-impermeant oxidant (carboxy-2,2,6,6-tetramethylpiperidine 1-oxyl NO^+) to selectively modify the $[\text{^3H}]\text{Man-P-Dol}$ molecules located in the outer monolayer of the synthetic vesicles.^{5,26} In this method, 50% of the $[\text{^3H}]\text{Man-P-Dol}$ in phospholipid vesicles lacking an active flippase will be oxidized. The remaining 50% is sequestered within the inner monolayer of the vesicle and is protected from reaction with the oxidant. However, in vesicles equipped with an active flippase, the $[\text{^3H}]\text{Man-P-Dol}$ facing the lumen of the proteoliposome is able to equilibrate into the outer monolayer and be oxidized. The result is that a larger fraction of the $[\text{^3H}]\text{Man-P-Dol}$ will be subject to oxidation. This study demonstrated rapid transbilayer movement of Man-P-Dol that is protein dependent, energy independent, DEPC sensitive, and highly specific for both the anomeric configuration of Man and a reduced α -isoprene unit.²⁶

Further studies showed that Man-P-Dol flippase activity could be resolved from flippase activities that mediate the transmembrane movements of M5-DLO (see $\text{Man}_5\text{GlcNAc}_2\text{-P-P-Dol}$ flippase section) and glycerophospholipid flippase by concanavalin A (ConA) affinity and ion-exchange chromatography.²⁶ Limiting dilution of these detergent-soluble extracts into phospholipid vesicles leads to the conclusion that the Man-P-Dol flippase constitutes $\sim 1\%$ of the proteins in the detergent extract.²⁶

Flippase assays based on water-soluble analog transport have the advantages of simplicity and rapidity and allow near-initial rate estimations in the absence of detergents.^{29,32} The alternative method employing full-length Man-P-Dol avoids potential concerns arising from the use of structural analogs,^{5,26} but relies on the reconstitution of detergent-solubilized membrane preparations, and is potentially limited by the time constraint imposed by reaction with the mannosyl lipid, which may significantly exceed the extremely rapid time course of transmembrane diffusion, even at 0°C ($\tau < 7$ minutes).^{5,26} In this regard, it is important to note that the properties of the Man-P-Cit transport activity in rat liver ER vesicles were identical to the activity reconstituted from detergent extracts, so the probability that some essential intrinsic property of the flippase

system is affected by detergent solubilization and reconstitution is very low. Although neither method has yet yielded a full molecular description of the Man-P-Dol flippase proteins, some preliminary properties have been identified and it is reassuring that the apparent properties of the flippase protein determined by the two methods are very similar. The complementary analytical methods identify an activity that is energy independent, bidirectional, stereoselective, sensitive to proteolysis and protein alkylation by DEPC and NEM, and amenable to fractionation by a variety of chromatographic approaches.

Glc-P-Dol Flippase

Glc-P-Dol is synthesized by Glc-P-Dol synthase (ALG5) using UDP-Glc as glucosyl donor and Dol-P as the lipid acceptor on the cytosolic surface of the ER but functions as a glucosyl donor in the synthesis of G3-DLO on the luminal monolayer.^{49,50} The topological reorientation of Glc-P-Dol has been studied in sealed, ER-enriched microsomal vesicles from pig brain and rat liver by following the transport of a water-soluble Glc-P-Dol analog, Glc-P-citronellol (Glc-P-Cit).³⁰ Similar to the properties of the Man-P-Dol flippase reported earlier, Glc-P-Cit uptake in rat liver ER vesicles is time dependent, saturable, bidirectional, trypsin sensitive, and completely dependent on an intact permeability barrier. Furthermore, Glc-P-Cit is not taken up by synthetic liposomes or erythrocytes and Glc-P-Cit transport activity fractionates with heavy microsomes (ER-enriched) in brain.³⁰ The molecular description of the Glc-P-Dol flippase protein remains a high priority, as severely affected human patients with CDG attributable to defects in the enzymes that catalyze attachment of Glc units to the DLO intermediate (ALG6 and ALG8) have been identified and genetic defects in Glc-P-Dol flipping would be expected to have similar pathologies.⁵¹

$\text{Man}_5\text{GlcNAc}_2\text{-P-P-Dol}$ Flippase

M5-DLO is synthesized from Dol-P , UDP-GlcNAc , and GDP-Man on the cytosolic surface of the ER (Fig. 1) by two multienzyme complexes, consisting of DPAGT1 , ALG13 , and ALG14 that form $\text{GlcNAc}_2\text{-P-P-Dol}$ ¹⁰ and ALG1 , ALG2 , and ALG11 that add one β -mannosyl and four α -mannosyl units to form M5-DLO .¹¹ The transmembrane diffusion of the M5-DLO intermediate is particularly interesting due to the presumed difficulty involved in the passage of this large, extremely hydrophilic head group through the hydrophobic core of the ER membrane.^{28,52,53} The identification of the yeast RFT1 protein as a candidate M5-DLO flippase, primarily based on genetic evidence by Helenius et al, was an exciting development.⁵⁴ RFT1 -depleted yeast cells accumulate large quantities of M5-DLO and are severely defective in protein N-glycosylation despite possessing normal levels of Man-P-Dol synthase and no apparent defect in the Man-P-Dol : M5-DLO mannosyltransferases (ALG3 , ALG9 , and ALG12).⁵⁴ However, direct biochemical assay of M5-DLO flippase activity by two independent



methods indicates that although RFT1 is clearly required for proper assembly of the full-length DLO intermediate, it is dispensable for M5-DLO flipping *in vitro*.^{28,55–57}

M5-DLO flipping has been measured directly in reconstituted proteoliposomes, employing a ConA binding assay.^{26,28,55,56} In this assay, proteoliposomes are reconstituted from Triton X-100 soluble cellular extracts and exogenous phospholipids with traces of [³H]M5-DLO equally distributed between the two monolayers. When incubated with soluble ConA, the radiolabeled DLO in the outer leaflet of the proteoliposome is bound by ConA and cannot be extracted into organic solvent, whereas since ConA cannot spontaneously cross the membrane bilayer, the DLO in the inner leaflet is sequestered from the ConA and, as a result, is fully extractable. Ideally, in proteoliposomes lacking an active flippase, only 50% of the [³H]M5-DLO is extractable. In proteoliposomes containing an active flippase, unbound M5-DLO is able to equilibrate with the outer monolayer and become bound by ConA. Consequently, a smaller fraction of the intermediate is extractable in liposomes containing a flippase activity. Using this assay, M5-DLO flipping was shown to be extremely rapid (a rate constant of $> 2 \times 10^{-3} \text{ s}^{-1}$), protein dependent, energy independent, bidirectional, and protease sensitive. Significantly, the M5-DLO flippase activity could be resolved from an unrelated phospholipid flippase activity by several chromatographic methods.⁵⁶ Furthermore, the M5-DLO flippase activity preferred the natural M5-DLO isomer over a catabolic M5-DLO intermediate, with an alternative branching pattern, and a biosynthetic M7-DLO intermediate demonstrating a high degree of substrate specificity.⁵⁶ A further report showed that M5-DLO flipping was inhibited by the inclusion of GlcNAc₂-P-P-Cit, a water-soluble M5-DLO analog, and that whereas M5-DLO was rapidly flipped with a time constant $\tau < 2$ minutes, the triantennary intermediate M3-DLO flipped much more slowly with $\tau < 200$ minutes. DLOs larger than M5-DLO also flipped slowly with τ values ranging from 10 to > 200 minutes.²⁸ Importantly, this activity was present in Triton X-100 soluble extracts derived from cells in which RFT1 had been depleted and was not immunoprecipitated with specific antibody to epitope-tagged RFT1.^{28,55} In addition, M5-DLO flippase activity could be resolved from the epitope-tagged RFT1 protein by velocity sedimentation, ion-exchange, and blue dye resin chromatography.^{28,55}

In another set of studies, sealed ER vesicles from RFT1-depleted yeast cells were fully capable of elongating [³H]GlcNAc₂-P-P-Dol, synthesized *in situ*, to form [³H]M9-DLO *in vitro* when incubated with nonradioactive GDP-Man.⁵⁷ As this elongation reaction cannot occur without translocation of the DLO intermediate to the luminal compartment, this is strong evidence that the RFT1-depleted microsomes are still capable of flipping M5-DLO. GlcNAc₂-P-P-Dol has been shown to be flipped, presumably by the M5-DLO flippase in microsomes from a variety of tissues and yeasts.^{58–63} To measure M5-DLO flipping directly in sealed

yeast ER vesicles, the transport of [³H]GlcNAc₂-P-P-Dol₁₅, a water-soluble analog of GlcNAc₂-P-P-Dol with a 15-carbon α -reduced isoprenol chain, was measured. RFT1-depleted, ER-enriched yeast microsomes transported water-soluble [³H]GlcNAc₂-P-P-Dol₁₅ into the lumen of the sealed vesicles at an even faster rate than intact vesicles from undepleted cells.⁵⁷ Furthermore, when the sealed RFT1-depleted microsomal vesicles were incubated with a membrane-permeant N-glycosylation acceptor peptide and [³H]GlcNAc₂-P-P-Dol₁₅, the peptide was glycosylated with [³H]GlcNAc₂, presumably by the luminal oligosaccharyltransferase (OST), demonstrating that the water-soluble analog has access to the luminal compartment.⁵⁷ Finally, in an indirect assay for the recycling of Dol-P,¹⁴ it is observed that when sealed RFT1-depleted yeast microsomal vesicles are incubated with a membrane-permeant N-glycosylation acceptor peptide, a very large amount of newly formed Dol-P is generated (J. S. Rush and C. J. Waechter, 2008, unpublished observation). This assay was previously used to successfully establish the biochemical mechanism, by which Dol-P derived from Dol-P-P, generated by the N-glycosylation reaction, is returned to the cytosolic monolayer for further rounds of DLO synthesis¹⁴ and is strong evidence that a large portion of the accumulated M5-DLO can enter the lumen of the ER in vesicles prepared from RFT1-depleted cells and become accessible to OST.

This biochemical evidence strongly suggests that RFT1 is not an essential subunit of the M5-DLO flippase in ER vesicles isolated from disrupted cells. Recently, this conclusion has been further supported by the observation that mutant *Trypanosoma brucei* cells lacking RFT1 are viable and N-glycosylate protein normally despite accumulating a large amount of M5-DLO.⁶⁴ The exact role of RFT1 in the synthesis of full-length DLOs *in vivo* and the identity of the M5-DLO flippase remain to be elucidated, but RFT1's role may be similar to that of the CHO Lec35 protein (MPDU1), which appears to be required for the utilization of both Man-P-Dol and Glc-P-Dol, *in vivo*, but is dispensable *in vitro*.³⁴

Dol-P Flippase

Dol-P is formed on the luminal surface of the ER membrane from several biosynthetic processes, including (1) cleavage of Dol-P-P generated during the OST reaction of protein N-glycosylation, (2) the three glucosyltransferase reactions utilizing Glc-P-Dol, and (3) by all of the Man-P-Dol-mediated mannosyltransferase reactions (the four DLO mannosyltransferases, the three GPI anchor mannosyltransferases and C- and O-mannosylation; see Fig. 2) utilizing Man-P-Dol as mannosyl donor.^{6,14} This lumenally oriented Dol-P must be returned to the cytosolic surface of the ER in order to participate in further rounds of glycan synthesis. The recycling of Dol-P substantially contributes to the maintenance of biologically relevant rates of protein N-glycosylation, as a significant reduction of glycosylation is observed in the yeast DolPP1 mutant, CWH8, even though this mutation presumably




Figure 3. Transport of ^{32}P -Cit into the lumen of intact rat liver ER vesicles. Panel A, open circle (\circ), ^{32}P -Cit was incubated with sealed rat liver ER vesicles at 22°C for the indicated periods of time. Following incubation, the transport reactions were diluted with 0.5 mL ice-cold buffer (10 mM Tris-Cl, pH 7.4, 0.25 M sucrose) and filtered through a Millipore $0.45\ \mu\text{m}$ HA filter. The filters were washed with an additional 10 mL ice-cold buffer and assayed for radioactivity, as described earlier.²⁹ Panel A, closed circle (\bullet), sealed rat liver ER vesicles were pre-incubated with ^{32}P -Cit for 10 minutes and diluted with 0.5 mL, 10 mM Tris-Cl, pH 7.4, and 0.25 M sucrose all at 22°C . At the indicated times, the diluted reactions were filtered through a $0.45\ \mu\text{m}$ HA filter, washed with an additional 10 mL ice-cold buffer, and assayed for radioactivity. Panel B, large unilamellar synthetic phospholipid vesicles were reconstituted from octyl glucoside soluble phospholipids in the presence of ^{32}P -Cit, as described elsewhere.^{43,67} The vesicles containing ^{32}P -Cit entrapped in the lumen were incubated with 10 mM Tris-Cl, pH 7.4, and 0.25 M sucrose at 22°C for the indicated periods of time and chromatographed on a 30 mL column of Sephadex G-50 equilibrated in 10 mM Tris-Cl, pH 7.4, and 0.25 M sucrose. Fractions of 1 mL were collected and monitored for radioactivity by scintillation spectrometry. The percentage of the radioactivity eluting in the void volume of the column is plotted versus time of incubation.

affects only the recycling of Dol-P from DLO and not the substantial number of Dol-P molecules arising from the other glycosyltransferase reactions.¹⁴ Despite the clear significance of the recycling of the lipid anchor, the molecular and biochemical details of this process remain virtually unknown. It will be worth exploring the possibility that CWH8 (DolPP1) plays a role in the translocation of Dol-P back to the cytosolic surface of the ER.

Presumably, Dol-P is returned to the cytoplasmic monolayer by a flippase-mediated process, as the amphipathic lipid anchor is unable to spontaneously cross the hydrophobic interior of the membrane bilayer. However, the identity of this flippase has not been established. Indeed, it is not known if multiple, unique flippases are associated with the various glycosyltransferases and the DolPP1 involved in the production of luminal Dol-P mentioned earlier. A membrane-associated activity that transports Cit- ^{32}P (a water-soluble analog of Dol-P) into the lumen of sealed vesicles has been detected (J. S. Rush and C. J. Waechter, 2006, unpublished observation), as shown in Figure 3. When sealed rat liver ER vesicles are incubated with Cit- ^{32}P , the analog is rapidly transported into the lumen of the vesicle with an apparent half-time of uptake of ~ 23 seconds (Fig. 3, panel A). When the ER vesicles are first loaded with Cit- ^{32}P by preincubation and then diluted into isotonic buffer at room temperature, the internalized Cit- ^{32}P effluxes from the vesicles with a half-time of exit of ~ 30 seconds, showing that the uptake process is fully reversible (Fig. 3, panel A). Cit- ^{32}P does not spontaneously leak out of protein-free phospholipid

vesicles, as shown in Figure 3, panel B (half-time of exit ~ 94 hours). Preliminary studies indicate that as with the other ER-associated flippase activities, this activity is membrane protein dependent, time dependent, protease sensitive, and energy independent (J. S. Rush and C. J. Waechter, 2006, unpublished observation). The identity of the protein mediating the return of Dol-P to the cytosolic monolayer has not yet been determined, but recycling by an antiport mechanism via one of the other flippase activities seems unlikely, as Cit-P does not have any appreciable effect on the uptake or efflux of any of the other water-soluble analogs employed in flippase assays.

Concluding Comments

The transbilayer diffusion of Man-P-Dol in the ER membrane was first investigated by Haselbeck and Tanner using an elegant enzyme-coupled assay over 40 years ago.^{35,36} Despite continued interest in the mechanism of Man-P-Dol flipping in the ER, the proteins that participate in this essential step involved in C- and O-mannosylation, DLO synthesis, and GPI anchor formation in the ER remain unidentified. The fragility of the topological asymmetry of the lipid bilayer and the apparent physical properties of the putative flippases have presented substantial experimental difficulties to the direct measurement of flippase activity *in vitro*. The biochemical assays reviewed here are important advances and have led to the identification of a number of physical properties of the prospective flippases. In common with the glycerophospholipid flippases present in biogenic membranes,



the ER dolichol-intermediate flippases are bidirectional and energy independent and catalyze an extremely rapid reequilibration of the intermediates between the two monolayers. Hypothetical models for flippase action have been proposed by Kol et al,⁶⁵ Sanyal and Menon,^{5,26} and Ernst and Menon.⁶⁶ It is possible that the translocation events are energy independent for lipid intermediates because the flipping is driven by mass action, as the intermediates are rapidly utilized at the luminal sites. It is also reassuring that although the experimental approaches employed in these assays differ significantly, the individual studies identify very similar physical properties, which are also consistent with observations from coupled in vitro enzymatic systems. The development of reliable methods for the solubilization, fractionation, and functional reconstitution of the flippase activities coupled to proteomic approaches paves the way for the purification, identification, and characterization of these final undescribed steps in protein glycosylation in the ER.

Acknowledgments

I am grateful to Skip Waechter (University of Kentucky College of Medicine) for his support and encouragement and for the opportunity to participate in a portion of the work reviewed in this article and to Dr. Dean C. Crick (Colorado State University), Dr. Hudson H. Freeze (Burnham Institute, La Jolla, CA), and Dr. Anant K. Menon (Weill Cornell Medical College) for critical comments.

Author Contributions

Conceived and designed the experiments: JSR. Analyzed the data: JSR. Wrote the first draft of the manuscript: JSR. Developed the structure and arguments for the paper: JSR. Made critical revisions: JSR. The author reviewed and approved of the final manuscript.

REFERENCES

- Holthuis JC, Menon AK. Lipid landscapes and pipelines in membrane homeostasis. *Nature*. 2014;510(7503):48–57.
- Pomorski T, Holthuis JC, Herrmann A, van Meer G. Tracking down lipid flippases and their biological functions. *J Cell Sci*. 2004;117(pt 6):805–813.
- Pomorski T, Menon AK. Lipid flippases and their biological functions. *Cell Mol Life Sci*. 2006;63(24):2908–2921.
- Panatala R, Hennrich H, Holthuis JC. Inner workings and biological impact of phospholipid flippases. *J Cell Sci*. 2015;128(11):2021–2032.
- Sanyal S, Menon AK. Flipping lipids: why an' what's the reason for? *ACS Chem Biol*. 2009;4(11):895–909.
- Schenk B, Fernandez F, Waechter CJ. The ins(ide) and out(side) of dolichyl phosphate biosynthesis and recycling in the endoplasmic reticulum. *Glycobiology*. 2001;11(5):61R–70R.
- Breitling J, Aebi M. N-linked protein glycosylation in the endoplasmic reticulum. *Cold Spring Harb Perspect Biol*. 2013;5(8):a013359.
- Helenius A, Aebi M. Roles of N-linked glycans in the endoplasmic reticulum. *Annu Rev Biochem*. 2004;73:1019–1049.
- Snider MD, Rogers OC. Transmembrane movement of oligosaccharide-lipids during glycoprotein synthesis. *Cell*. 1984;36(3):753–761.
- Noffz C, Keppler-Ross S, Dean N. Hetero-oligomeric interactions between early glycosyltransferases of the dolichol cycle. *Glycobiology*. 2009;19(5):472–478.
- Gao XD, Nishikawa A, Dean N. Physical interactions between the Alg1, Alg2, and Alg11 mannosyltransferases of the endoplasmic reticulum. *Glycobiology*. 2004;14(6):559–570.
- Fernandez F, Rush JS, Toke DA, et al. The CWH8 gene encodes a dolichyl pyrophosphate phosphatase with a luminally oriented active site in the endoplasmic reticulum of *Saccharomyces cerevisiae*. *J Biol Chem*. 2001;276(44):41455–41464.
- Rush JS, Cho SK, Jiang S, Hofmann SL, Waechter CJ. Identification and characterization of a cDNA encoding a dolichyl pyrophosphate phosphatase located in the endoplasmic reticulum of mammalian cells. *J Biol Chem*. 2002;277(47):45226–45234.
- Rush JS, Gao N, Lehrman MA, Waechter CJ. Recycling of dolichyl monophosphate to the cytoplasmic leaflet of the endoplasmic reticulum after the cleavage of dolichyl pyrophosphate on the luminal monolayer. *J Biol Chem*. 2008;283(7):4087–4093.
- Maeda Y, Kinoshita T. Dolichol-phosphate mannose synthase: structure, function and regulation. *Biochim Biophys Acta*. 2008;1780(6):861–868.
- Hanover JA, Lennarz WJ. The topological orientation of N,N'-diacetylchitobiosylpyrophosphoryldolichol in artificial and natural membranes. *J Biol Chem*. 1979;254(18):9237–9246.
- McCloskey MA, Troy FA. Paramagnetic isoprenoid carrier lipids. 1. Chemical synthesis and incorporation into model membranes. *Biochemistry*. 1980;19(10):2056–2060.
- Bretscher MS. Membrane structure: some general principles. *Science*. 1973;181(4100):622–629.
- Freeze HH. Genetic defects in the human glycome. *Nat Rev Genet*. 2006;7(7):537–551.
- Schachter H, Freeze HH. Glycosylation diseases: quo vadis? *Biochim Biophys Acta*. 2009;1792(9):925–930.
- Jaeken J. Congenital disorders of glycosylation. *Handb Clin Neurol*. 2013;113:1737–1743.
- Hennet T. Diseases of glycosylation beyond classical congenital disorders of glycosylation. *Biochim Biophys Acta*. 2012;1820(9):1306–1317.
- Freeze HH, Chong JX, Bamshad MJ, Ng BG. Solving glycosylation disorders: fundamental approaches reveal complicated pathways. *Am J Hum Genet*. 2014;94(2):161–175.
- Scott K, Gadomski T, Kozicz T, Morava E. Congenital disorders of glycosylation: new defects and still counting. *J Inherit Metab Dis*. 2014;37(4):609–617.
- Buczowska A, Swiezewska E, Lefeber DJ. Genetic defects in dolichol metabolism. *J Inherit Metab Dis*. 2015;38(1):157–169.
- Sanyal S, Menon AK. Stereoselective transbilayer translocation of mannosyl phosphoryl dolichol by an endoplasmic reticulum flippase. *Proc Natl Acad Sci U S A*. 2010;107(25):11289–11294.
- Menon I, Huber T, Sanyal S, et al. Opsin is a phospholipid flippase. *Curr Biol*. 2011;21(2):149–153.
- Sanyal S, Menon AK. Specific transbilayer translocation of dolichol-linked oligosaccharides by an endoplasmic reticulum flippase. *Proc Natl Acad Sci U S A*. 2009;106(3):767–772.
- Rush JS, Waechter CJ. Transmembrane movement of a water-soluble analogue of mannosylphosphoryldolichol is mediated by an endoplasmic reticulum protein. *J Cell Biol*. 1995;130(3):529–536.
- Rush JS, van Leyen K, Ouerfelli O, Wolucka B, Waechter CJ. Transbilayer movement of Glc-P-dolichol and its function as a glucosyl donor: protein-mediated transport of a water-soluble analog into sealed ER vesicles from pig brain. *Glycobiology*. 1998;8(12):1195–1205.
- Rick PD, Barr K, Sankaran K, Kajimura J, Rush JS, Waechter CJ. Evidence that the wzxE gene of *Escherichia coli* K-12 encodes a protein involved in the transbilayer movement of a trisaccharide-lipid intermediate in the assembly of enterobacterial common antigen. *J Biol Chem*. 2003;278(19):16534–16542.
- Rush JS, Waechter CJ. Assay for the transbilayer movement of polyisoprenoid-linked saccharides based on the transport of water-soluble analogues. *Methods*. 2005;35(4):316–322.
- Maeda Y, Tanaka S, Hino J, Kangawa K, Kinoshita T. Human dolichol-phosphate-mannose synthase consists of three subunits, DPM1, DPM2 and DPM3. *EMBO J*. 2000;19(11):2475–2482.
- Anand M, Rush JS, Ray S, et al. Requirement of the Lec35 gene for all known classes of monosaccharide-P-dolichol-dependent glycosyltransferase reactions in mammals. *Mol Biol Cell*. 2001;12(2):487–501.
- Haselbeck A, Tanner W. Dolichyl phosphate-mediated mannosyl transfer through liposomal membranes. *Proc Natl Acad Sci U S A*. 1982;79(5):1520–1524.
- Haselbeck A, Tanner W. Further evidence for dolichyl phosphate-mediated glycosyl translocation through membranes. *FEMS Microbiol Lett*. 1984;21(3):305–308.
- Rong Y, Nakamura S, Hirata T, et al. Genome-wide screening of genes required for glycosylphosphatidylinositol biosynthesis. *PLoS One*. 2015;10(9):e0138553.
- Davis EM, Kim J, Menasche BL, et al. Comparative haploid genetic screens reveal divergent pathways in the biogenesis and trafficking of glycosylphosphatidylinositol-anchored proteins. *Cell Rep*. 2015;11(11):1727–1736.
- Leidich SD, Drapp DA, Orlean P. A conditionally lethal yeast mutant blocked at the first step in glycosyl phosphatidylinositol anchor synthesis. *J Biol Chem*. 1994;269(14):10193–10196.



40. Orlean P, Menon AK. Thematic review series: lipid posttranslational modifications. GPI anchoring of protein in yeast and mammalian cells, or: how we learned to stop worrying and love glycospholipids. *J Lipid Res.* 2007;48(5):993–1011.
41. Jeffery CJ. An introduction to protein moonlighting. *Biochem Soc Trans.* 2014; 42(6):1679–1683.
42. Ernst OP, Menon AK. Phospholipid scrambling by rhodopsin. *Photochem Photobiol Sci.* 2015;14(11):1922–1931.
43. Rush JS, Waechter CJ. Functional reconstitution into proteoliposomes and partial purification of a rat liver ER transport system for a water-soluble analogue of mannosylphosphoryldolichol. *Biochemistry.* 2004;43(23):7643–7652.
44. Bishop WR, Bell RM. Assembly of the endoplasmic reticulum phospholipid bilayer: the phosphatidylcholine transporter. *Cell.* 1985;42(1):51–60.
45. Rush JS, Shelling JG, Zingg NS, Ray PH, Waechter CJ. Mannosylphosphoryldolichol-mediated reactions in oligosaccharide-P-P-dolichol biosynthesis. Recognition of the saturated alpha-isoprene unit of the mannosyl donor by pig brain mannosyltransferases. *J Biol Chem.* 1993;268(18):13110–13117.
46. DeLuca AW, Rush JS, Lehrman MA, Waechter CJ. Mannolipid donor specificity of glycosylphosphatidylinositol mannosyltransferase-I (GPIMT-I) determined with an assay system utilizing mutant CHO-K1 cells. *Glycobiology.* 1994; 4(6):909–915.
47. Dotson SB, Rush JS, Ricketts AD, Waechter CJ. Mannosylphosphoryldolichol-mediated O-mannosylation of yeast glycoproteins: stereospecificity and recognition of the alpha-isoprene unit by a purified mannosyltransferase. *Arch Biochem Biophys.* 1995;316(2):773–779.
48. Kean EL, Rush JS, Waechter CJ. Activation of GlcNAc-P-P-dolichol synthesis by mannosylphosphoryldolichol is stereospecific and requires a saturated alpha-isoprene unit. *Biochemistry.* 1994;33(34):10508–10512.
49. Hirschberg CB, Snider MD. Topography of glycosylation in the rough endoplasmic reticulum and Golgi apparatus. *Annu Rev Biochem.* 1987;56:63–87.
50. Rush JS, Waechter CJ. Topological studies on the enzymes catalyzing the biosynthesis of Glc-P-dolichol and the triglycosyl cap of Glc3Man9GlcNAc2-P-P-dolichol in microsomal vesicles from pig brain: use of the processing glucosidases I/II as latency markers. *Glycobiology.* 1998;8(12):1207–1213.
51. Freeze HH. Understanding human glycosylation disorders: biochemistry leads the charge. *J Biol Chem.* 2013;288(10):6936–6945.
52. Vijayraghavan U, Company M, Abelson J. Isolation and characterization of pre-mRNA splicing mutants of *Saccharomyces cerevisiae*. *Genes Dev.* 1989;3(8): 1206–1216.
53. Aebi M, Gassenhuber J, Domdey H, te Heesen S. Cloning and characterization of the ALG3 gene of *Saccharomyces cerevisiae*. *Glycobiology.* 1996;6(4):439–444.
54. Helenius J, Ng DT, Marolda CL, Walter P, Valvano MA, Aebi M. Translocation of lipid-linked oligosaccharides across the ER membrane requires Rft1 protein. *Nature.* 2002;415(6870):447–450.
55. Frank CG, Sanyal S, Rush JS, Waechter CJ, Menon AK. Does Rft1 flip an N-glycan lipid precursor? *Nature.* 2008;454(7204):E3–E4. discussion E4–5.
56. Sanyal S, Frank CG, Menon AK. Distinct flippases translocate glycerophospholipids and oligosaccharide diphosphate dolichols across the endoplasmic reticulum. *Biochemistry.* 2008;47(30):7937–7946.
57. Rush JS, Gao N, Lehrman MA, Matveev S, Waechter CJ. Suppression of Rft1 expression does not impair the transbilayer movement of Man5GlcNAc2-P-P-dolichol in sealed microsomes from yeast. *J Biol Chem.* 2009;284(30): 19835–19842.
58. Reuvers F, Habets-Willems C, Reinking A, Boer P. Glycolipid intermediates involved in the transfer of N-acetylglucosamine to endogenous proteins in a yeast membrane preparation. *Biochim Biophys Acta.* 1977;486(3):541–552.
59. Lehle L, Tanner W. Glycosyl transfer from dolichyl phosphate sugars to endogenous and exogenous glycoprotein acceptors in yeast. *Eur J Biochem.* 1978;83(2): 563–570.
60. Chen WW, Lennarz WJ. Metabolism of lipid-linked N-acetylglucosamine intermediates. *J Biol Chem.* 1977;252(10):3473–3479.
61. Struck DK, Lennarz WJ, Brew K. Primary structural requirements for the enzymatic formation of the N-glycosidic bond in glycoproteins. Studies with alpha-lactalbumin. *J Biol Chem.* 1978;253(16):5786–5794.
62. Harford JB, Waechter CJ. Transfer of N, N'-diacetylchitobiose from dolichyl diphosphate into a gray matter membrane glycoprotein. *Arch Biochem Biophys.* 1979;197(2):424–435.
63. Cueva R, Cotano C, Larriga G. N-glycosylation by transfer of GlcNAc2 from dolichol-PP-GlcNAc2 to the protein moiety of the major yeast exoglucanase. *Yeast.* 1998;14(8):773–781.
64. Jelk J, Gao N, Serricchio M, et al. Glycoprotein biosynthesis in a eukaryote lacking the membrane protein Rft1. *J Biol Chem.* 2013;288(28):20616–20623.
65. Kol MA, de Kroon AI, Killian JA, de Kruijff B. Transbilayer movement of phospholipids in biogenic membranes. *Biochemistry.* 2004;43(10):2673–2681.
66. Ernst OP, Menon AK. Phospholipid scrambling by rhodopsin. *Photochem Photobiol Sci.* 2015;14(11):1922–1931.
67. Mimms LT, Zampighi G, Nozaki Y, Tanford C, Reynolds JA. Phospholipid vesicle formation and transmembrane protein incorporation using octyl glucoside. *Biochemistry.* 1981;20(4):833–840.