



OPEN ACCESS

Better reporting standards are needed to enhance the quality of hop testing in the setting of ACL return to sport decisions: a narrative review

Paul Read ,^{1,2} Sean Mc Auliffe ,³ Mathew G Wilson ,^{1,4}
Gregory D Myer ,^{5,6,7}

¹Research Department, Aspetar Orthopaedic and Sports Medicine Hospital, Doha, Qatar

²School of Sport and Exercise, University of Gloucestershire, Gloucester, UK

³Department of Physical Therapy and Rehabilitation Sciences, Qatar University, Doha, Qatar

⁴Institute of Sport Exercise and Health (ISEH), University College London, London, UK

⁵Division of Sports Medicine, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA

⁶Department of Pediatrics and Orthopaedic Surgery, College of Medicine, College of Medicine, University of Cincinnati, Cincinnati, Ohio, USA

⁷The Micheli Center for Sports Injury Prevention, Boston, MA, USA

Correspondence to

Dr Paul Read, Research Department, Aspetar Orthopaedic and Sports Medicine Hospital, PO Box number: 29222, Doha, Ad Dawhah, Qatar; paulread10@hotmail.com

Accepted 19 May 2020

Published Online First

10 June 2020

ABSTRACT

Background/aim There is a lack of consistency in return to sport (RTS) assessments, in particular hop tests to predict who will sustain a reinjury following anterior cruciate ligament (ACL) reconstruction. Inconsistent test battery content and methodological heterogeneity might contribute to variable associations between hop test performance and subsequent injury. Our aim was to investigate whether commonly used hop tests are administered in a consistent manner and in accordance with reported guidelines.

Methods We conducted a narrative review of studies that examined whether hop testing could differentiate RTS pass rates, reinjury and rerupture in athletes after ACL reconstruction. Our specific focus was on the methodological procedures of hop testing as this component is widely used to evaluate patients' function and readiness to RTS.

Main findings Substantial variation exists in RTS hop test administration, scoring and interpretation. Authors often failed to report important details of methods such as warm up activities, randomisation, number of trials, rest periods and landing requirements.

Conclusion We recommend researchers provide clearer descriptions of how hop tests are performed to increase standardisation and promote accurate data collection. Absence of reporting to describe test methods and using different test procedures makes it difficult to compare study findings.

INTRODUCTION

Return to sport (RTS) decision making following anterior cruciate ligament (ACL) reconstruction is a complex process involving many factors. A criterion-based approach is now accepted,¹ where a range of tests are used in various combinations. Passing a test battery including a series of single leg hop and isokinetic tests was associated with lower rerupture rates following RTS,^{1,2} and an increased likelihood of returning to previous sporting levels.³

However, hop and isokinetic tests do not consistently predict successful outcomes following ACL rehabilitation. Losciale *et al*⁴ reported no associations between the use of RTS discharge tests and greater risk of reinjury, stating the low quality of evidence affects our ability to make definitive conclusions. Toole *et al*⁵ reported that many young athletes had been cleared to RTS by their surgeon and rehabilitation specialist but failed to pass the RTS cut-offs cited in the literature. A recent 2019 systematic review⁶ demonstrated that only 23% of

patients passed RTS test batteries. These authors⁶ also suggested an apparent paradox that 'passing' an RTS battery was associated with a greater risk of injury to the contralateral limb.

That conclusion led us, and others,⁷ to question whether the existing data relating to RTS and subsequent reinjury displays too much clinical variability among the patient groups (race, age, sex, level of performance, type of ACL surgery, other associated injuries) and inconsistent application of test batteries (different clinicians and studies use a widely ranging set of tests). On deeper reflection, and after scoping the literature, we wondered whether the tests themselves (eg, the hop tests) are described clearly and used in a reproducible manner. For RTS tests to be to be valid and generalised across clinical settings, standardised methods are required for administration, scoring and interpretation.

The primary aim of this narrative review was to provide an overview of the RTS testing batteries of the studies included^{8–20} in the 2019 synthesis of the literature⁶ (table 1), and the cited studies in the methods sections of these manuscripts^{21–35} (table 2). Our focus was applied to hop testing methods and administration procedures following ACL reconstruction. This current review⁶ provided the most comprehensive and up-to-date body of literature in the area of RTS testing and numerous discussions followed its publication. Our second aim was to demonstrate how even minor alterations to the test protocol might affect interpretation and subsequent RTS decisions.

RESULTS

Table 1 shows that a wide range of approaches in test administration, scoring and interpretation were used, and important methodological details were infrequently reported. To illustrate this point, we created a colour code system to indicate the methodological quality of each of the included studies (green=aligned with current evidence; yellow=reported but not aligned with current evidence; red=not stated) (tables 1 and 2). A discussion of why these factors affect test outcomes is provided below.

Test order

Substantial variation existed across the included studies in relation to testing order. No studies randomised the order of testing, seven did not state the sequence performed, and the remaining studies included additional hop/jump tests.



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Read P, Mc Auliffe S, Wilson MG, *et al*. *Br J Sports Med* 2021;**55**:23–29.

Table 1 Studies cited in the 2019 systematic review of RTS outcomes following ACL reconstruction⁶

Study	Tests included	Testing order	Warm up activities	Familiarisation session	# Practice trials	# Trials on each limb	Limb order	Best or mean score	Rest period	Hand placement	Landing requirements	Point of measurement	Checks for systematic bias	Studies cited for procedures
Beischer <i>et al</i> ⁸	Isometric quad strength, isokinetics, vertical hop, single hop, side hop	Controlled as stated	10 mins exercise bike	Not stated	3–5	3	Not stated	Best	3 mins between trials for side hops, other tests not stated	Not stated	Not stated	Not stated	Yes	Gustavsson <i>et al</i> ¹⁵
Di Stasi <i>et al</i> ⁹	Quadriceps strength index, single hop, triple-hop, cross-over hop, timed hop	Order not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Fitzgerald <i>et al</i> ²³
Ebert <i>et al</i> ¹⁰	Single hop, timed hop, triple-hop, cross-over hop	Controlled as stated	6 mins walk test, optional unstandardised stretching	Not stated	2–3	2–4, test dependent	Uninvolved then involved	Mean	Based on patient readiness	Not stated	Controlled landing	Not stated	Not stated	Reid <i>et al</i> ²⁴
Fältström <i>et al</i> ¹¹	Star excursion balance, single hop, 5 jump test, side hop, drop jump, tuck jump	Controlled as stated	5–10 mins run, 10 squats, toe raises, 1 min skipping	Not stated	A few	3	Uninvolved then involved	Best	3 mins between tests, between trials not stated	Not stated	Not stated	Not stated	Yes	Gustavsson <i>et al</i> ¹⁵ , Neeter <i>et al</i> ²⁵
Gokeler <i>et al</i> ¹²	Drop jump, single hop, triple hop, side hop, isokinetics	Controlled as stated	10 mins exercise bike	Not stated	3	Not stated	Not stated	Not stated	30 s between trials; 3–5 mins between tests	Not stated	Not stated	Not stated	Not stated	Gustavsson <i>et al</i> ¹⁵ , Noyes <i>et al</i> ²²
Graziano <i>et al</i> ¹³	Isokinetics, single hop and subjective movement test battery	Order not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	None
Grindem <i>et al</i> ¹⁴	Isokinetics, single hop, triple-hop, cross-over hop, timed hop	Controlled as stated	Not stated	Not stated	1	2	Uninvolved then involved	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Logerstedt <i>et al</i> ¹⁵ , Grindem <i>et al</i> ²⁴
Krych <i>et al</i> ¹⁴	Isokinetic, vertical jump, single hop, triple-hop	Order not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Janvela <i>et al</i> ¹⁷ , Petschnig <i>et al</i> ²³ , Reid <i>et al</i> ²⁴
Kyritsis <i>et al</i> ¹²	Isokinetics, single hop, triple-hop, cross-over hop, agility T-test	Controlled for hops, order not stated for other tests	Not stated	Not stated	Not stated	3	Not stated	Not stated	Not stated	Not stated	Land without losing balance	Not stated	Not stated	Hopper <i>et al</i> ²⁶ , Reid <i>et al</i> ²⁴ , Ross <i>et al</i> ²⁵
Logerstedt <i>et al</i> ¹⁵	Isokinetics, single hop, cross-over hop, triple-hop, timed Hop, IKDC	Controlled as stated	Not stated	Not stated	1–2	2	Uninvolved then involved	Mean	Not stated	Not stated	Landing was stable	Heel	Not stated	Noyes <i>et al</i> ²²
Luo <i>et al</i> ¹⁶	Isokinetics, vertical jump, single hop, triple-hop	Order not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Petschnig <i>et al</i> ²³
Nawastreh <i>et al</i> ¹⁷	Isometric quad strength, single hop, triple-hop, cross-over hop, timed Hop*	Order not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Nawastreh <i>et al</i> ²¹ , Noyes <i>et al</i> ²²
Sousa <i>et al</i> ¹⁷	Isokinetic, vertical jump, single hop, triple-hop	Order not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Janvela <i>et al</i> ¹⁷ , Petschnig <i>et al</i> ²³
Thomeé <i>et al</i> ¹⁸	Vertical jump, single hop, side hop	Controlled as stated	10 mins exercise bike	Not stated	3–5	3	Uninvolved then involved	Best	3 mins between side hops only	Not stated	Not stated	Not stated	Yes	Gustavsson <i>et al</i> ¹⁵
Toole <i>et al</i> ¹⁵	Isokinetics, single hop, triple hop, cross-over hop, timed Hop, IKDC	Controlled as stated	Not stated	Not stated	1	2	Randomised	Mean	Not stated	Not stated	Not stated	Not stated	Not stated	Noyes <i>et al</i> ²²

Continued

Table 1 Continued

Study	Tests included	Testing order	Warm up activities	Familiarisation session	# Practice trials	# Trials on each limb	Limb order	Best or mean score	Rest period	Hand placement	Landing requirements	Point of measurement	Checks for systematic bias	Studies cited for procedures
Welling <i>et al</i> ²⁹	Drop jump, single hop, triple-hop, side hop, isokinetics	Controlled as stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Not stated	Kockum <i>et al</i> ²⁶ , Mumro and Herrington ³⁰
Wellsandt <i>et al</i> ²⁰	Isometric quad strength, single hop, triple-hop, cross-over hop, timed Hop	Order not stated	Not stated	Not stated	2	2	Uninvolved then involved	Mean	Not stated	Not stated	Not stated	Not stated	Not stated	Noyes <i>et al</i> ³²

Heibst *et al*. Functional assessments for decision making regarding return to sports following ACL reconstruction. Part II: clinical application of a new test battery. *Knee Surg Sports Traumatol* 2015;23:1288–1291 was included in the systematic review⁶ but the hop tests described in our review were not used in their study. Consequently, the details were not provided here.

⁶Procedural review applied to hop tests only.

^{*}Hop tests performed in a knee brace.

Green, aligned with current evidence; red, not stated; yellow, reported but not aligned with current evidence.

Randomisation of test order will reduce the potential of an order effect and control fatigue. However, if a ‘battery’ of related tests is being used, it is prudent to standardise the order. A logical sequence is to perform non-fatiguing tests requiring high skill movements and coordination first. Therefore, power tests (hops) may be preferentially performed prior to other RTS tests such as those requiring maximal strength (isokinetic dynamometry).³⁶ In athletes following ACL reconstruction, considering the loading requirements for each hop test is important. We suggest moving in sequence from the least to most demanding task and propose the following test order: (1) timed hop; (2) single hop; (3) triple-hop and (4) cross-over hop.

Influence of preceding tests measuring different performance constructs

Test order should consider the physiological energy systems targeted, as well as the athletes training age/experience. Performing hop tests soon after or even before maximal strength tests can have a pronounced effect on the test outcomes. Table 1 indicates that four studies performed isokinetic testing first, while two studies employed isokinetic testing following hop tests. Other studies also included either isokinetic or isometric knee strength assessments, but the test order was not stated.

Following a bout of resistance training, involved musculature are in both a fatigued and potentiated state. The balance between these two factors determines the subsequent muscle performance.³⁷ An acute enhancement in muscle function following intense activity is defined as postactivation potentiation,³⁸ with significant improvements frequently shown in resistance exercise performance following plyometric activities.^{39–41} However, modulating factors can influence these positive benefits. A larger effect is shown for stronger individuals and those with greater resistance training experience. The length of the rest interval is also dependent on the individual’s level of strength, with weaker athletes requiring longer rest periods following the previous activity.⁴¹

Quadriceps muscle fatigue can affect hop performance in athletes following ACL reconstruction.⁴² Single leg hop performance was compared in both a non-fatigued state, and immediately following a pre-exhaustion set (as many reps as possible) of knee extensions at 50% of 1 repetition maximum (RM) strength. All participants displayed ‘pass’ limb symmetry scores (LSI>90%) in the non-fatigued condition, but only 68% ‘passed’ the test following the pre-exhaustion protocol. Thus, pre-exhaustion of the quadriceps may be deleterious to single leg hop performance after ACL reconstruction.

Assessing athletes under conditions of fatigue may provide valuable information to the tester; however, the inconsistencies in test order displayed in the studies reviewed limits the interpretation of results and our ability to generalise the findings. The number of warm up trials for isokinetic protocols was also frequently not stated, and a range of strength test modes and speeds were used (60°/s, 90°/s and 180°/s), further limiting extrapolation of findings across all studies.

Limb testing order

Most studies tested the uninvolved before the involved limb. Only one study randomised limb order, and the others did not state the order limbs were tested. If the task is always performed on the uninvolved limb first as was the case in six studies, performance on the involved limb may benefit from a learning effect from the preceding hops on the other side. Research studies should counterbalance the limb order where the participant

Table 2 Studies cited in the methods of manuscripts included in the 2019 systematic review⁶

Study	Tests included	Testing order	Warm up activities	Familiarisation session	# Practice trials	# trials on each limb	Limb order	Best or mean score	Rest period	Hand placement	Landing requirements	Point of measurement	Checks for systematic bias
Barber <i>et al</i> ²¹	Isokinetics, single hop, timed hop, vertical jump, shuttle run, shuttle run	Order not stated	Not stated	Not stated	Not stated	2	Not stated	Mean	Not stated	Not stated	Not stated	Not stated	Not stated
Bolgla and Kekula ²²	Single hop, triple-hop, cross-over hop, timed hop	Randomised order	3 mins cycling, static stretching & practice trials	Not stated	3	3	Dominant limb only	Mean	1 min between trials	Not stated	Not stated	Not stated	Not stated
Frizgerald <i>et al</i> ²³	Single hop, cross-over hop, triple-hop, timed hop, isometric quad strength	Controlled as stated	Not stated	Not stated	2	2	Not stated	Mean	Not stated	Not stated	Not stated	Not stated	Not stated
Grindem <i>et al</i> ²⁴	Single hop, cross-over hop, triple-hop, timed hop	Controlled as stated	Not stated	Not stated	1	2	Not stated	Best	Not stated	Arm use was permitted	Not stated	Not stated	Not stated
Gustavsson <i>et al</i> ²⁵	Vertical jump, single hop, drop jump, square hop, side hop	Controlled as stated	5 mins cycling plus squats, toe raises and jumps	Yes	three to 5	3	Un-involved then involved	Best	3 mins between tests, between trials not stated	Hands behind back	Landing held for 2 to 3 s	Heel	Yes
Hopper <i>et al</i> ²⁶	Timed hop, cross-over hop, stair hop, vertical hop	Randomised order	5 mins cycling and static stretching	Not stated	1	3	Randomised	Best	Not stated	Not stated	Not stated	Toe	Not stated
Järvelä <i>et al</i> ²⁷	Isokinetics, single hop	Order not stated	5 mins cycling	Not stated	Not stated	3	Not stated	Mean	Not stated	Not stated	Not stated	Not stated	Not stated
Kockum <i>et al</i> ²⁸	Vertical jump, single hop, side hop, single leg squat jump, knee power test	Order not stated	10 mins cycling, toe raises, tuck jumps	Not stated	three to 5	3	Right first and then alternating	Mean	Not stated	Not stated	Controlled landing	Not stated	Not stated
Logerstedt <i>et al</i> ²⁹	Single hop, cross-over hop, triple-hop, timed hop	Order not stated	Not stated	Not stated	Not stated	2	Un-involved then involved	Mean	Not stated	Not stated	Landing was stable, on one limb, under control, no additional hops forward	Heel	Not stated
Mumro and Herrington ³⁰	Agility t-test, single hop, triple-hop, cross-over hop, timed hop	Order not stated	Not stated	Not stated	Not stated	3	Not stated	Mean	30 s between trials and 2 mins between tests	Arm use was permitted	Landing held for 2 s	Heel	Not stated
Nawasreh <i>et al</i> ³¹	Isometric quad strength, single hop, cross-over hop, triple-hop, timed hop	Controlled as stated	Not stated	Not stated	2	2	Un-involved then involved	Mean	Not stated	Not stated	Maintained balance without touching the ground	Not stated	Not stated
Noyes <i>et al</i> ³²	Isokinetics, single hop, timed hop, triple-hop, cross-over hop	Order not stated	Not stated	Not stated	Not stated	2	Not stated	Mean	Not stated	Not stated	Not stated	Not stated	Not stated
Petschnig <i>et al</i> ³³	Vertical jump, single leg hop, triple-hop, isokinetics	Order not stated	Not stated	Not stated	Not stated	3	Not stated	Best (SLCMJ) and mean (hops)	2 mins between tests, between trials not stated	Hands behind back	Not stated	Not stated	Not stated
Reid <i>et al</i> ³⁴	Single hop, timed hop, triple-hop, cross-over hop	Controlled as stated	None other than practice trials	Yes	1	2	Un-involved then involved	Mean	30 s between trials and 2 mins between tests	Arm use was permitted	Landing held for 2 s	Toe	Not stated

Continued

Table 2 Continued

Study	Tests included	Testing order	Warm up activities	Familiarisation session	# Practice trials	# trials on each limb	Limb order	Best or mean score	Rest period	Hand placement	Landing requirements	Point of measurement	Checks for systematic bias
Ross <i>et al</i> ³⁵	Single hop, triple-hop, cross-over hop, timed hop	Randomised order	5 mins cycling and static stretching	Not stated	3	3	One limb only - randomly selected	Best	30 s between trials	Arm use was permitted	Not stated	Heel	Not stated

Procedural review applied to hop tests only.

Green, aligned with current evidence; yellow, reported but not aligned with current evidence; red, not stated.

sample is divided in half, and the limb order is reversed for each group to remove this confounding factor. Clinicians may wish to randomise the limb order for each hop performed during the test battery so that the uninvolved limb is not always first.

Rest periods

No consistent prescription of rest periods between tests/trials was applied across the studies included in the prior synthesis of the literature.⁶ Protocols ranged from 30 s to 3 min, with many studies not stating how long participants rested. Previous recommendations have indicated work to rest ratios of at least 1:5 during plyometric exercise.⁴³ Thus, counterbalancing limb order and 30 s rest between trials should provide adequate recovery and control for order and fatigue effects.

Reporting the best versus the mean score

Many studies reported either the best trial (three studies) or mean score (four studies), while the remaining 10 studies did not state which data were used for reporting. There are no clear recommendations for the preferred method (best vs mean score) to report hop test performance, and this is often based on individual preference. For countermovement jump tests, the mean score is more sensitive than the best recorded trial to detect performance changes and monitoring neuromuscular fatigue.⁴⁴ Similarly, systematic bias was shown between test sessions attended a week apart for the best but not mean asymmetry score during an isometric squat test, with lower test retest reliability.⁴⁵ Given that a limb symmetry index is a composite score derived from the performance of individual limbs, average scores may capture some of the inconsistencies between trials and the innate variability of asymmetry.⁴⁵ Thus, it is possible that the mean score is preferable when measuring hop performance following ACL reconstruction.

Including task constraints to avoid movement compensation

Absence of task constraints can affect scores obtained during hop testing due to movement compensations. Only six studies mentioned any landing requirements with descriptions varying from 'stable', 'controlled', 'without losing balance' and either 'hold for 2 s', or 'hold for 2–3 s'.

It may also be prudent when performing hop tests where the goal is to assess lower extremity power, to consider and control contributions from the extremities. The inclusion of an arm swing during jumping tasks has been shown to augment performance due to an increase in lower extremity work performed.⁴⁶ Therefore, when comparing results across different studies, arm placement should be controlled, or at least communicated. No study described the role of arm movement during testing. In the studies they cited (table 2), inconsistency was present with two requesting athletes place their hands behind the back, while four permitted arm use. Proponents of arm use suggest it is more 'functional', but it could be argued that hop testing itself is not truly 'functional' and lacks ecological validity for most sports. This was acknowledged by Noyes *et al*³² who stated 'the four hop tests used in this study were not sensitive enough to detect their functional limitations. This could be due to the nature of the tests themselves; they are performed in a safe environment instead of the playing field, there are no opponents or objects to undermine the participant's concentration, and the activities involve simple hopping with no cutting or twisting motions'.

Ageberg and Cronström⁴⁷ examined single leg hop performance in participants with lower extremity injury under two conditions: (1) with arms free and (2) arms behind back. Hop

distance was shorter on their involved versus uninvolved leg when the arms were placed behind their back, and nearly twice as many participants displayed 'abnormal' LSI (<90%). Agreement between the two conditions was also poor. These findings highlight the importance of limiting arm use, and that a stricter protocol should be adopted to avoid overestimating knee function.

Familiarisation and practice trials

Table 1 shows that no studies included a separate familiarisation session where athletes were provided with opportunities to practice the tests prior to data collection. There was also inconsistency in the number of practice trials, ranging from 3–5, 2–3, 1 and 'a few'. This affects data interpretation, as systematic learning will occur between trials and test sessions in both 'healthy' recreational athletes²⁴ and patients following ACL reconstruction.^{25–34} Significant differences in hop performance have been reported between test sessions 1 and 2, but not days 2 and 3 in previous research.^{25–34} These findings indicate a learning effect, supporting the need for a separate familiarisation session prior to testing.

The number of practice trials provided should allow for adequate familiarisation. Munro and Herrington³⁰ showed that learning affects were present in all four hop tests, where scores improved across trials. Single and triple-hop for distance test scores stabilised after three trials; whereas, cross-over hop scores stabilised after four trials. The timed hop stabilised after four trials in women and three in men. Therefore, practice trials should be provided to ensure a more reliable test outcome. Specifically, three practice trials are recommended for the single and triple-hop tests, with an additional trial included for the cross-over hop and timed hop.

Warm up procedures

A warm up is common practice in athletic endeavours. Table 1 shows 12 of the included studies did not state what/if any warm up procedures were applied, while five studies included either cycling, walking or running, with durations ranging from 5 to 10 min. Additional warm up activities included static stretching (one study) and dynamic exercises such as squats and toe raises (one study).

An active warm up can improve performance⁴⁸ and test outcomes might vary with such a large variation in physiological readiness. It has been shown that practice trials alone are insufficient to elicit maximal strength and jumping performance, which are positively related to muscle temperature.^{49–50} It is also recommended to avoid static stretching immediately prior to testing as jump performance has been shown to reduce compared with dynamic warm up protocols.⁵¹

Selecting the optimal warm up for athletes will depend on a variety of factors. We recommend a protocol consisting of general cardiovascular activity (eg, stationary cycling or jogging performed at approximately 60% of maximum perceived effort) for 5 min and task-specific activities (squats, lunges, practice jumps / hops, etc) to increase muscle temperature and movement pattern sequencing.

Other considerations

Results might be affected by a range of other factors beyond the scope of this review including the use of a knee brace, shoe-surface interaction and the athlete's state of readiness. These aspects should be considered in the study design and reported within the methods section. In addition, test administrators

should be well trained and have a thorough understanding of all protocols and procedures. Examples applied to hop tests have been outlined below.

Point of measurement

Only one study reported this procedure, where measurement was to the heel. Of the cited studies, four measured the distance to the heel and two to the toe. Measuring the distance hopped to the toe does not account for the wide variation in foot length, which is irrespective of the horizontal hop distance achieved during the task. Differences in the point of measurement (heel vs toe) can also affect the LSI score. Practitioners are encouraged to measure the distance hopped from the start line to the participant's heel. It is also prudent to report the absolute hop distances, and scores relative to leg length or body height to account for different anthropometric profiles, enabling comparisons of performance across different studies and athletes of the same sport/playing level.

Reporting details of the test raters

In cases where more than one person is conducting RTS testing, inter-rater reliability should be examined prior to data collection and the relevant statistics should be reported. To illustrate this point, consider the timed hop and the many inherent sources of error. The timer starts when the athlete's heel leaves the ground at the beginning of the test and stops when the athlete completes the 6-m distance. Thus, clinicians encounter four potential sources of error (1) heel raise; (2) hit start; (3) visually observe the athlete complete the 6 m distance and (4) hit stop). Ensuring appropriate consistency between raters prior to testing is therefore essential to ensure accurate data collection.

Check for systematic bias between trials

Raters should check the scores of each individual trial during hop testing to control for systematic bias. An athlete's score can progressively improve during the test (due to learning, increased confidence or warm up effects) or get worse (maybe from fatigue or wavering motivation). In the studies reviewed, only three stated that they checked for systematic bias. The most frequent protocols were 'if subjects increased their hop performance in all three trials, additional hops were performed until no increase was seen'.^{11–18} Beischer *et al.*⁸ stated 'if the test administrator felt that an even better result could be achieved, one or two additional hops were allowed'. An alternative is to use values greater than the minimal detectable change (MDC) to indicate the minimal amount of change required to determine if the observed performance increases or decreases between trials are 'real', accounting for the measurement error in the test. MDC values for the four hop tests have been reported in ACL patients (single hop: 8.09%; triple-hop: 10.02%; cross-over hop: 12.25%; timed hop: 12.96%; overall combination of hop tests: 7.05%).³⁴ To be confident a 'real' systematic increase in performance has been observed, a general guideline of ~10% may be applied, although this may increase for the cross-over and timed hop tests.

Description of hop test procedures in research studies

Several important methodological procedures were often not stated in the studies reviewed (table 1). Even when citing the methods of previous research (table 2), it was not possible to determine which procedures had been used. An example of this is the study of Kyritsis *et al.*² Text from their manuscript states: 'Single and triple hop distance tests were used'.^{24–26} The authors cited three manuscripts^{26–34–35} indicating further procedural

information. However, there were differences between these studies in test administration (table 2) and the original manuscript² did not specify which procedures from each of these studies were employed. Inadequate reporting makes it harder to replicate methodological procedures and threatens not only the external validity (ie, generalisability or applicability) of experimental studies but also those of subsequent systematic reviews. A previous systematic review examined the measurement properties of the hop tests and their relationships with future knee injury.⁵² The authors reported the methods and terminology varied greatly across studies and stated that a lack of standardisation limits the generality of the findings. We believe the heterogeneity in how data are collected and reported could at least in part, account for the equivocal results found within the synthesised literature⁶ and other systematic reviews in this area of research,⁴ which might influence current practice recommendations. Furthermore, the results of hop testing will almost certainly vary across patient groups (athletic vs non-athletic, older vs younger, presence of comorbidities vs no comorbidities, etc). A clear description of study population characteristics can aid clinicians in their interpretation of the data and translation of results into clinical practice.

CONCLUSION

Even small alterations to hop test procedures can affect performance outcomes and we propose the following:

- ▶ Increased rigour of test methods and reporting standards to enhance the quality and reproducibility of future research that examines RTS outcomes following ACL reconstruction.
- ▶ A standardised approach is needed with specific protocol instructions to obtain accurate data, heighten test sensitivity, and avoid overestimating or underestimating knee function.
- ▶ The procedural review guidelines provided in tables 1 and 2 could form part of a reporting checklist for future research.

What is already known?

- ▶ There is a lack of consistency in the ability of anterior cruciate ligament (ACL) return to sport (RTS) tests to predict who has a successful clinical outcome following ACL rehabilitation and who has a serious reinjury.
- ▶ We and other clinicians suspect there is insufficient rigour in how the hop tests are administered during an RTS battery.

What are the new findings?

- ▶ Authors reporting procedures used in RTS testing frequently did not detail how they administered the hop tests.
- ▶ We found substantial variation in RTS test administration, scoring and interpretation in different studies that reported hop testing post-ACL reconstruction
- ▶ Small alterations to hop test procedures can affect performance outcomes, making it difficult to compare the findings of the respective studies; a more standardised approach is required.
- ▶ There is a need for better quality in the reporting of hop test methods to allow practitioners and researchers to compare the findings of different studies. This is an important prerequisite to testing the utility (or not) of RTS batteries (that include the hop test).

Twitter Paul Read @paulread1010, Sean Mc Auliffe @Seaniemc89 and Gregory D Myer @gregmyer11

Contributors The lead author and one other author conceptualised the article contents. The lead author wrote the original draft of the manuscript. All authors contributed substantially to editing the manuscript in preparation to submit a final draft as well as subsequent revisions. All others read and approved the final copy of the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Paul Read <http://orcid.org/0000-0002-1508-8602>

Sean Mc Auliffe <http://orcid.org/0000-0002-8716-5005>

Mathew G Wilson <http://orcid.org/0000-0002-6317-0168>

Gregory D Myer <http://orcid.org/0000-0001-9829-813X>

REFERENCES

- 1 Grindem H, Snyder-Mackler L, Moksnes H, *et al*. Simple decision rules reduce injury risk after anterior cruciate ligament reconstruction: the Delaware-Orlando cohort study. *Br J Sports Med* 2016;50:804–8.
- 2 Kyritsis P, Bahr R, Landreau P, *et al*. Likelihood of ACL graft rupture: not meeting six clinical discharge criteria before return to sport is associated with a four times greater risk of rupture. *Br J Sports Med* 2016;50:946–51.
- 3 Nawasreh Z, Logerstedt D, Cummer K, *et al*. Functional performance 6 months after ACL reconstruction can predict return to participation in the same preinjury activity level 12 and 24 months after surgery. *Br J Sports Med* 2018;52:375–83.
- 4 Losciale JM, Zdeb RM, Ledbetter L, *et al*. The association between passing return-to-sport criteria and second anterior cruciate ligament injury risk: a systematic review with meta-analysis. *J Orthop Sports Phys Ther* 2019;49:43–54.
- 5 Toole AR, Ithurburn MP, Rauh MJ, *et al*. Young athletes cleared for sports participation after anterior cruciate ligament reconstruction: how many actually meet recommended return-to-sport criterion cutoffs? *J Orthop Sports Phys Ther* 2017;47:825–33.
- 6 Webster KE, Hewett TE. What is the evidence for and validity of Return-to-Sport testing after anterior cruciate ligament reconstruction surgery? A systematic review and meta-analysis. *Sports Med* 2019;49:917–29.
- 7 Capin JJ, Snyder-Mackler L, Risberg MA, *et al*. Keep calm and carry on testing: a substantive reanalysis and critique of 'what is the evidence for and validity of return-to-sport testing after anterior cruciate ligament reconstruction surgery? A systematic review and meta-analysis'. *Br J Sports Med* 2019;53:1444–6.
- 8 Beischer S, Hamrin Senorski E, Thomeé C, *et al*. Young athletes return too early to knee-strenuous sport, without acceptable knee function after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2018;26:1966–74.
- 9 Di Stasi SL, Logerstedt D, Gardinier ES, *et al*. Gait patterns differ between ACL-reconstructed athletes who pass return-to-sport criteria and those who fail. *Am J Sports Med* 2013;41:1310–8.
- 10 Ebert JR, Edwards P, Yi L, *et al*. Strength and functional symmetry is associated with post-operative rehabilitation in patients following anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2018;26:2353–61.
- 11 Fältström A, Häggglund M, Kvist J. Functional performance among active female soccer players after unilateral primary anterior cruciate ligament reconstruction compared with knee-healthy controls. *Am J Sports Med* 2017;45:377–85.
- 12 Gokeler A, Welling W, Zaffagnini S, *et al*. Development of a test battery to enhance safe return to sports after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2017;25:192–9.
- 13 Graziano J, Chiaia T, de Mille P, *et al*. Return to sport for skeletally immature athletes after ACL reconstruction: preventing a second injury using a quality of movement assessment and quantitative measures to address modifiable risk factors. *Orthop J Sports Med* 2017;5:232596711770059–10.
- 14 Krych AJ, Woodcock JA, Morgan JA, *et al*. Factors associated with excellent 6-month functional and isokinetic test results following ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2015;23:1053–9.
- 15 Logerstedt D, Di Stasi S, Grindem H, *et al*. Self-Reported knee function can identify athletes who fail return-to-activity criteria up to 1 year after anterior cruciate ligament reconstruction: a delaware-oslo ACL cohort study. *J Orthop Sports Phys Ther* 2014;44:914–23.

- 16 Luo TD, Ashraf A, Dahm DL, *et al.* Femoral nerve block is associated with persistent strength deficits at 6 months after anterior cruciate ligament reconstruction in pediatric and adolescent patients. *Am J Sports Med* 2015;43:331–6.
- 17 Sousa PL, Krych AJ, Cates RA, *et al.* Return to sport: does excellent 6-month strength and function following ACL reconstruction predict midterm outcomes? *Knee Surg Sports Traumatol Arthrosc* 2017;25:1356–63.
- 18 Thomeé R, Neeter C, Gustavsson A, *et al.* Variability in leg muscle power and hop performance after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2012;20:1143–51.
- 19 Welling W, Benjaminse A, Seil R, *et al.* Low rates of patients meeting return to sport criteria 9 months after anterior cruciate ligament reconstruction: a prospective longitudinal study. *Knee Surg Sports Traumatol Arthrosc* 2018;26:3636–44.
- 20 Wellsandt E, Failla MJ, Snyder-Mackler L. Limb symmetry indexes can overestimate knee function after anterior cruciate ligament injury. *J Orthop Sports Phys Ther* 2017;47:334–8.
- 21 Barber SD, Noyes FR, Mangine RE, *et al.* Quantitative assessment of functional limitations in normal and anterior cruciate ligament-deficient knees. *Clin Orthop Relat Res* 1990;255:204–14.
- 22 Bolgla LA, Keskula DR. Reliability of lower extremity functional performance tests. *J Orthop Sports Phys Ther* 1997;26:138–42.
- 23 Fitzgerald GK, Axe MJ, Snyder-Mackler L. A decision-making scheme for returning patients to high-level activity with nonoperative treatment after anterior cruciate ligament rupture. *Knee Surg Sports Traumatol Arthrosc* 2000;8:76–82.
- 24 Grindem H, Løgerstedt D, Eitzen I, *et al.* Single-legged hop tests as predictors of self-reported knee function in nonoperatively treated individuals with anterior cruciate ligament injury. *Am J Sports Med* 2011;39:2347–54.
- 25 Gustavsson A, Neeter C, Thomeé R, *et al.* A test battery for evaluating hop performance in patients with an ACL injury and patients who have undergone ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2006;14:778–88.
- 26 Hopper DM, Goh SC, Wentworth LA, *et al.* Test–retest reliability of knee rating scales and functional hop tests one year following anterior cruciate ligament reconstruction. *Physical Therapy in Sport* 2002;3:10–18.
- 27 Järvelä T, Kannus P, Latvala K, *et al.* Simple measurements in assessing muscle performance after an ACL reconstruction. *Int J Sports Med* 2002;23:196–201.
- 28 Kockum B, Heijne AI-LM. Hop performance and leg muscle power in athletes: reliability of a test battery. *Phys Ther Sport* 2015;16:222–7.
- 29 Løgerstedt D, Grindem H, Lynch A, *et al.* Single-legged hop tests as predictors of self-reported knee function after anterior cruciate ligament reconstruction: the Delaware-Oslo ACL cohort study. *Am J Sports Med* 2012;40:2348–56.
- 30 Munro AG, Herrington LC. Between-session reliability of four hop tests and the agility t-test. *J Strength Cond Res* 2011;25:1470–7.
- 31 Nawasreh Z, Løgerstedt D, Cummer K, *et al.* Do patients failing return-to-activity criteria at 6 months after anterior cruciate ligament reconstruction continue demonstrating deficits at 2 years? *Am J Sports Med* 2017;45:1037–48.
- 32 Noyes FR, Barber SD, Mangine RE. Abnormal lower limb symmetry determined by function hop tests after anterior cruciate ligament rupture. *Am J Sports Med* 1991;19:513–8.
- 33 Petschnig R, Baron R, Albrecht M. The relationship between isokinetic quadriceps strength test and hop tests for distance and one-legged vertical jump test following anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther* 1998;28:23–31.
- 34 Reid A, Birmingham TB, Stratford PW, *et al.* Hop testing provides a reliable and valid outcome measure during rehabilitation after anterior cruciate ligament reconstruction. *Phys Ther* 2007;87:337–49.
- 35 Ross MD, Langford B, Whelan PJ. Test-Retest reliability of 4 single-leg horizontal hop tests. *J Strength Cond Res* 2002;16:617–22.
- 36 Harman E. Principles of test selection and administration. In: Baechle TR, Earle RW, eds. *Essentials of strength training and conditioning*. 3rd edn. Champaign IL, USA: Human Kinetics, 2008: 245.
- 37 Kilduff LP, Owen N, Bevan H, *et al.* Influence of recovery time on post-activation potentiation in professional rugby players. *J Sports Sci* 2008;26:795–802.
- 38 Hodgson M, Docherty D, Robbins D. Post-Activation potentiation. *Sports Medicine* 2005;35:585–95.
- 39 Deusch M, Lloyd R. Effect of order of exercise on performance during a complex training session in rugby players. *J Sports Sci* 2008;26:803–9.
- 40 Masamoto N, Larson R, Gates T, *et al.* Acute effects of plyometric exercise on maximum squat performance in male athletes. *J Strength Cond Res* 2003;17:68–71.
- 41 Seitz LB, Haff GG. Factors modulating post-activation potentiation of jump, sprint, throw, and Upper-Body ballistic performances: a systematic review with meta-analysis. *Sports Med* 2016;46:231–40.
- 42 Augustsson J, Thomeé R, Karlsson J. Ability of a new hop test to determine functional deficits after anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2004;12:350–6.
- 43 Potach DH, Chu DA. Plyometric training. In: *The essentials of strength training and conditioning*. Baechle TR and Earle RW. Champaign, IL: Human Kinetics, 2008: 413–27.
- 44 Claudino JG, Cronin J, Mezêncio B, *et al.* The countermovement jump to monitor neuromuscular status: a meta-analysis. *J Sci Med Sport* 2017;20:397–402.
- 45 Bishop C, Read P, Chavda S, *et al.* Using unilateral strength, power and reactive strength tests to detect the magnitude and direction of asymmetry: a test-retest design. *Sports* 2019;7:58–14.
- 46 Hara M, Shibayama A, Takeshita D, *et al.* A comparison of the mechanical effect of arm swing and countermovement on the lower extremities in vertical jumping. *Hum Mov Sci* 2008;27:636–48.
- 47 Ageberg E, Cronström A. Agreement between test procedures for the single-leg hop for distance and the single-leg mini squat as measures of lower extremity function. *BMC Sports Sci Med Rehabil* 2018;10:1–7.
- 48 Bishop D, Up W II. Warm up II: performance changes following active warm up and how to structure the warm up. *Sports Med* 2003;33:483–98.
- 49 Vetter RE. Effects of six warm-up protocols on sprint and jump performance. *J Strength Cond Res* 2007;21:819–23.
- 50 Pyke FS. The effect of preliminary activity on maximal motor performance. *Res Q* 1968;39:1069–76.
- 51 Holt BW, Lambourne K. The impact of different warm-up protocols on vertical jump performance in male collegiate athletes. *J Strength Cond Res* 2008;22:226–9.
- 52 Hegedus EJ, McDonough S, Bleakley C, *et al.* Clinician-friendly lower extremity physical performance measures in athletes: a systematic review of measurement properties and correlation with injury, part 1. The tests for knee function including the hop tests. *Br J Sports Med* 2015;49:642–8.