

Postoperative cerebral infarction risk is related to lobectomy site in lung cancer: a retrospective cohort study of nationwide data in Japan

Natsumasa Nishizawa ,¹ Makoto Okawara,² Masataka Mori,¹ Yoshihisa Fujino,² Shinya Matsuda,³ Kiyohide Fushimi,⁴ Fumihiko Tanaka¹

To cite: Nishizawa N, Okawara M, Mori M, *et al*. Postoperative cerebral infarction risk is related to lobectomy site in lung cancer: a retrospective cohort study of nationwide data in Japan. *BMJ Open Resp Res* 2022;**9**:e001327. doi:10.1136/bmjresp-2022-001327

Received 9 June 2022
Accepted 16 July 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Second Department of Surgery, University of Occupational and Environmental Health Japan, Kitakyushu, Fukuoka, Japan
²Department of Environmental Epidemiology, Institute of Industrial Ecological Sciences, University of Occupational and Environmental Health Japan, Kitakyushu, Fukuoka, Japan
³Department of Public Health, University of Occupational and Environmental Health Japan, Kitakyushu, Fukuoka, Japan
⁴Department of Health Policy and Informatics, Tokyo Medical and Dental University Graduate School of Medical and Dental Sciences, Bunkyo-ku, Tokyo, Japan

Correspondence to
Dr Natsumasa Nishizawa;
nmasamed@med.uoeh-u.ac.jp

ABSTRACT

Objectives To determine the incidence of cerebral infarction after lobectomy at different sites using inpatient data from a diagnosis procedure combination database.

Design Retrospective cohort study.

Setting Data were retrieved from the Japanese diagnosis procedure combination inpatient database for patients who underwent lobectomies for lung cancer between April 2018 and March 2020.

Participants The analysis included 37 352 patients from 556 institutions who underwent lobectomies for lung cancer.

Main outcomes and measures The occurrence of cerebral infarction after lobectomy during hospitalisation was estimated using multilevel logistic regression models adjusted for sex, age, body mass index, smoking history, activity of daily living, surgical approach (thoracotomy or video-assisted), clinical cancer stage, comorbidities and hospital-level factors to describe the association between cerebral infarction and different lobectomy sites.

Results Overall cerebral infarction after lobectomy occurred in 99 patients (0.27%): 29 with left upper lobectomy (0.39%), 19 with left lower lobectomy (0.34%), 32 with right upper lobectomy (0.24%), 6 with right middle lobectomy (0.21%) and 13 with right lower lobectomy (0.16%). The multilevel multivariate logistic regression analysis revealed high ORs for the left upper lobectomy and left lower lobectomy groups. In both univariate and multivariate analyses, left upper lobectomy had the highest OR for the occurrence of cerebral infarction compared with lobectomies at other sites.

Conclusions and relevance Left upper lobectomy had the highest OR for the occurrence of cerebral infarction after lung cancer lobectomy during hospitalisation. There is an urgent need to investigate the specific mechanisms underlying postoperative cerebral infarction after left upper lobectomy and to establish preventive measures such as altering surgical methods, using radiological examinations for early detection and better use of anticoagulants.

INTRODUCTION

In Japan, lung cancer has the third highest morbidity and the highest mortality rate among all malignancies.¹ Depending on

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Cerebral infarction after lobectomy is an irreversible and sometimes life-threatening complication.
- ⇒ Previous studies have reported that left upper lobectomy contributes to a higher risk of developing cerebral infarction compared with lobectomies at other sites, but these studies have been case reports or small-scale case-control studies.
- ⇒ The relationship between the site of lobectomy and incidence of cerebral infarction after surgery still requires confirmation with large-scale clinical data.

WHAT THIS STUDY ADDS

- ⇒ Among all sites of lobectomies, the left upper lobectomy was found to have the highest OR for postoperative cerebral infarction.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ The findings of this study highlight the urgent need for new interventional research to define the specific mechanisms underlying postoperative cerebral infarction after left upper lobectomy and to establish preventive measures.

the tumour node metastasis (TNM) stage, different therapies are used as treatment, including surgery, radiation and chemotherapy. In general, patients with stage 0–3A lung cancer undergo surgery, with lobectomy being the standard procedure. The reported postoperative complications are bronchopleural fistula, arrhythmias, including atrial fibrillation and more serious complications such as acute exacerbation of interstitial pneumonia, infections, including pneumonia and fistulous empyema and thromboembolic disease such as cerebral infarction.²

Cerebral infarction after lobectomy is an irreversible and sometimes life-threatening complication. The risk factors for cerebral infarction after lung resection are cardiac thromboembolism due to postoperative atrial

fibrillation,³ cancer-associated hypercoagulability (Trousseau syndrome),⁴ postoperative dehydration⁵ and other common risks such as diabetes, smoking and prolonged bed rest.⁶ Previous studies have reported that left upper lobectomy contributes to a higher risk of developing cerebral infarction compared with lobectomies at other sites.^{7–13} However, because cerebral infarction is such a rare complication of pulmonary resection, with rates as low as 0.2%–0.8%, almost all of these studies were case reports or small-scale case-control studies.^{14 15} The relationship between the site of lobectomy and incidence of cerebral infarction after surgery still requires confirmation with large-scale clinical data.

The aim of this study was to identify the incidence of cerebral infarction after lobectomies performed at different sites by using a Japanese nationwide database from the diagnosis procedure combination (DPC) system. We hypothesised that left upper lobectomy would have the highest risk of causing postoperative cerebral infarction compared with lobectomies at other sites.

MATERIALS AND METHODS

Study design

This retrospective cohort study was conducted using data from the Japanese DPC inpatient database from April 2018 to March 2020. The DPC system was introduced in 2003 in Japanese medical hospitals authorised to treat patients enrolled in a health insurance plan. As of 2020 in Japan, over 1600 hospitals have adopted the DPC system.¹⁶ The system determines the per-diem hospital reimbursement based on patient diagnosis and the procedures made available to them during their hospital stay to regulate diagnostic and therapeutic approaches. DPC data consist of medical procedures such as surgeries and treatments performed, medications and disease names with the dates of onset or which are indexed in the original Japanese codes.^{16 17} Of the over 1600 hospitals participating in DPC, we constructed our database by analysing DPC data from 1291 hospitals, which covers approximately 90% of all acute care beds in Japan. We analysed 16 001 612 inpatient cases from these hospitals between 1 April 2018 and 31 March 2020.

Case selection

We identified patients who had undergone lobectomies based on the codes of the International Statistical Classification of Diseases and Related Health Problems version 10 (ICD-10 code) and the Japanese medical intervention classification master codes (treatment code).¹⁸ These were then used to unify information entered into the DPC database. Hospitalised patients who had undergone a lobectomy for primary lung cancer (ICD-10 code: C34) and patients who had undergone a lobectomy (treatment codes: K5143, 514-23) were included in the study. Exclusion criteria included missing information (body mass index (BMI), activity of daily living (ADL), smoking history and clinical TNM classification), failure to identify




Figure 1 Flow chart of inclusion and exclusion criteria for the study. From the data of 16 001 612 hospitalised patients, 37 352 cases of lobectomy for lung cancer were extracted.

the resected lobe or more than once surgical procedure for lung cancer (treatment code; K514: lung malignant tumour surgery, or K514-2: video-assisted lung malignant tumour surgery) during the same hospitalisation period. A flow chart of the case selection process is presented in figure 1.

Evaluation of the resected lobe

Each patient's resected lobe was labelled with the appropriate ICD-10 code (C341: upper lobe, C342: middle lobe, C343: lower lobe) and by the surgical information in the DPC database as right-sided or left-sided.

Evaluation of postoperative cerebral infarctions

From the DPC records of ICD-10 codes for postoperative complications during hospitalisation, we identified 'I63' as a postoperative cerebral infarction.¹⁶

Covariates variables

The following confounding factors were extracted from the information contained in the DPC data: patients' characteristics (age,¹⁹ sex, BMI, smoking history and ADL⁶), clinical cancer TNM stage,^{4 20} surgical approach (thoracotomy or video-assisted) and underlying comorbidities as risks for thrombosis. BMI was divided into the following quaternary groups: <18.5 (underweight), 18.5–25 (normal range) and ≥25 (obese), according to the guidelines of the Japan Society for the Study of Obesity.²¹ Smoking history was categorised into 'never' and 'current or former', and ADL was categorised into fully ambulatory, use of a cane, wheelchair users

and total assistance. We classified cancer TNM stage as follows: stage 0 (T0–Tis, N0, M0), stage 1 (T1–T2a, N0, M0), stage 2 (T2b–T3, N0, M0 or T1–T2, N1, M0), stage 3 (T4, N0, M0 or T3–T4, N1, M0 or any T, N2–N3, M0) and stage 4 (any T, any N, M1), according to WHO standards.²² Underlying comorbidities used as covariates included atrial fibrillation,³ diabetes, hypertension,²³ history of cerebral infarction,^{24 25} heart failure,²⁶ hypertrophic cardiomyopathy,^{27 28} chronic kidney disease (stage 3–5)²⁹ and vascular disease (history of myocardial infarction,³⁰ arterial plaque³¹ and venous thrombosis³²) according to previous studies and clinical experience.

Statistical analysis

The ORs with 95% CIs were estimated using a multilevel logistic model nested in each hospital. In the multivariate analysis, the model included age, sex, BMI, smoking history, ADL, clinical cancer TNM stage, surgical approach, operative time, underlying comorbidities and number of surgeries for lobectomy as the hospital-level factor, dividing them into quartiles. $P < 0.05$ was considered statistically significant. Statistical analyses were performed using commercial software (Stata Statistical Software: Release 16; StataCorp, College Station, Texas, USA).

RESULTS

Patient characteristics

This study included 37 352 patients from 556 institutions who underwent lobectomies for lung cancer. In this cohort, 7462 (20.0%) underwent left upper lobectomies, 5593 (15.0%) underwent left lower lobectomies, 13 075 (35.0%) underwent right upper lobectomies, 2912 (7.8%) underwent right middle lobectomies and 8310 (22.3%) underwent right lower lobectomies (table 1).

ORs for the occurrence of cerebral infarction

In this population, cerebral infarction after lobectomy occurred in 99 of 37 352 patients (0.27%); 29 with left upper lobectomies (0.39%), 19 with left lower lobectomies (0.34%), 32 with right upper lobectomies (0.24%), 6 with right middle lobectomies (0.21%) and 13 with right lower lobectomies (0.16%) (table 2). Multivariate analysis identified the highest ORs for the left upper lobectomy (OR 2.66, 95% CI 1.37 to 5.16, $p = 0.004$) and left lower lobectomy (OR 2.20, 95% CI 1.08 to 4.49, $p = 0.03$) groups.

In the post hoc analysis, we found that the existence of postoperative atrial fibrillation was significantly associated with cerebral infarction (adjusted OR 7.69, 95% CI 4.14 to 14.29, $p < 0.001$), but there was less change in ORs in the resected lobe for the occurrence of cerebral

infarction after adding postoperative atrial fibrillation as a covariate in the statistical model.

DISCUSSION

We examined the incidence of cerebral infarction in relation to different lobectomy sites. We found that left upper and left lower lobectomies were risk factors for cerebral infarction occurrence, with left upper lobectomies showing the highest OR among all other types of lobectomies.

We demonstrated that left upper lobectomy for cancer is a significant risk factor for postoperative cerebral infarction using a large, real-world dataset as opposed to case-control studies of smaller populations. Similar to the findings of our study, the incidence of postoperative cerebral infarction was reported to be higher in patients undergoing left upper lobectomy in a retrospective case-control study that collected data from 1670 patients' medical records between 2008 and 2015.¹¹ Resection of the left upper lobe was also reported to increase the risk of cerebral infarction in a multicentre, retrospective case-control study that used data of 610 patients collected from 153 institutions of the Japanese Association for Chest Surgery between 2004 and 2013.¹⁰

We suggest one major possible mechanism underlying the occurrence of postoperative cerebral infarction after left upper lobectomy. It is thrombosis in the long pulmonary vein stump, which has been reported previously.^{7–9 12 13} Long pulmonary vein stumps are more likely to remain after a left upper lobectomy, which may cause thrombosis in the left atrium after lung resection, as shown using contrast-enhanced CT.^{7 8 13} The median length of the pulmonary vein stump after left upper lobectomy was reported to be significantly longer than that of the other pulmonary vein stumps.⁸ Furthermore, blood flow evaluation using intraoperative ultrasonography showing the presence of spontaneous echo contrast in the left superior pulmonary vein stump has been reported to be predictive of thrombosis that could cause arterial infarction.⁹ In addition, the mechanism of thrombus formation in the vein stump was explainable by Virchow's triad (ie, blood stasis, endothelial injury and hypercoagulability).¹² These reports revealed congestion and blood flow turbulence in the long pulmonary vein stump, which may contribute to cerebral infarction after a left upper lobectomy.

On the contrary, we suggest atrial fibrillation as one minor possible mechanism underlying the occurrence of postoperative cerebral infarction after left upper lobectomy. Although only one article analysing the relationship between lobectomy and atrial fibrillation was found in our search, left lobectomy (including lower lobectomy) was suggested as a risk factor for atrial fibrillation after lobectomy in a single-centre retrospective cohort study that used data of 186 patients between 2005 and 2010,³ citing vagal stimulation associated with lobectomy

Table 1 Basic characteristics of the patients

	Total N=37 352	Left upper lobectomy N=7462	Left lower lobectomy N=5593	Right upper lobectomy N=13 075	Right middle lobectomy N=2912	Right lower lobectomy N=8310
		20.0%	15.0%	35.0%	7.80%	22.3%
Variables						
Age, mean (SD)	70.0 (9.1)	69.6 (9.0)	70.3 (9.1)	69.9 (9.0)	69.8 (10.0)	70.4 (8.9)
Sex, men	22 377 (59.9%)	4495 (60.2%)	3276 (58.6%)	7946 (60.8%)	1487 (51.1%)	5173 (62.3%)
Body mass index						
<18 kg/m ²	1443 (3.9%)	257 (3.4%)	210 (3.8%)	492 (3.8%)	132 (4.5%)	352 (4.2%)
18–24.9 kg/m ²	24 416 (65.4%)	4874 (65.3%)	3696 (66.1%)	8502 (65.0%)	1888 (64.8%)	5456 (65.7%)
>25 kg/m ²	11 493 (30.8%)	2331 (31.2%)	1687 (30.2%)	4081 (31.2%)	892 (30.6%)	2502 (30.1%)
Smoking history (current or former)	22 604 (60.5%)	4512 (60.5%)	3325 (59.4%)	8011 (61.3%)	1459 (50.1%)	5297 (63.7%)
Activity of daily living						
Self-reliant	36 727 (98.3%)	7345 (98.4%)	5503 (98.4%)	12 875 (98.5%)	2847 (97.8%)	8157 (98.2%)
Use of a cane	377 (1.0%)	63 (0.8%)	60 (1.1%)	125 (1.0%)	42 (1.4%)	87 (1.0%)
Wheelchair user	144 (0.4%)	33 (0.4%)	17 (0.3%)	38 (0.3%)	13 (0.4%)	43 (0.5%)
Total assistance	104 (0.3%)	21 (0.3%)	13 (0.2%)	37 (0.3%)	10 (0.3%)	23 (0.3%)
Cancer TNM stage						
0	1757 (4.7%)	302 (4.0%)	210 (3.8%)	703 (5.4%)	205 (7.0%)	337 (4.1%)
I	26 765 (71.7%)	5332 (71.5%)	3908 (69.9%)	9749 (74.6%)	2254 (77.4%)	5522 (66.5%)
II	5303 (14.2%)	1067 (14.3%)	955 (17.1%)	1582 (12.1%)	255 (8.8%)	1444 (17.4%)
III	3188 (8.5%)	691 (9.3%)	454 (8.1%)	943 (7.2%)	169 (5.8%)	931 (11.2%)
IV	339 (0.9%)	70 (0.9%)	66 (1.2%)	98 (0.7%)	29 (1.0%)	76 (0.9%)
Comorbidities						
Any of the following applies	16 183 (43.3%)	3240 (43.4%)	2377 (42.5%)	5649 (43.2%)	1215 (41.7%)	3702 (44.5%)
Chronic kidney disease	276 (0.7%)	39 (0.5%)	46 (0.8%)	110 (0.8%)	21 (0.7%)	60 (0.7%)
Vascular disease	1034 (2.8%)	230 (3.1%)	181 (3.2%)	326 (2.5%)	64 (2.2%)	233 (2.8%)
Hypertrophic cardiomyopathy	47 (0.1%)	7 (0.1%)	8 (0.1%)	19 (0.1%)	7 (0.2%)	6 (0.1%)
Heart failure	1409 (3.8%)	289 (3.9%)	208 (3.7%)	482 (3.7%)	102 (3.5%)	328 (3.9%)
Hypertension	9664 (25.9%)	1963 (26.3%)	1374 (24.6%)	3440 (26.3%)	761 (26.1%)	2126 (25.6%)
Diabetes	7380 (19.8%)	1438 (19.3%)	1097 (19.6%)	2530 (19.3%)	504 (17.3%)	1811 (21.8%)
Cerebral infarction or haemorrhage	394 (1.1%)	72 (1.0%)	56 (1.0%)	140 (1.1%)	34 (1.2%)	92 (1.1%)
Atrial fibrillation	1473 (3.9%)	300 (4.0%)	209 (3.7%)	554 (4.2%)	127 (4.4%)	283 (3.4%)

Continued

Table 1 Continued

	Total N=37 352	Left upper lobectomy N=7462	Left lower lobectomy N=5593	Right upper lobectomy N=13 075	Right middle lobectomy N=2912	Right lower lobectomy N=8310
Surgical approach		20.0%	15.0%	35.0%	7.80%	22.3%
Thoracotomy	5625 (15.1%)	1292 (17.3%)	828 (14.8%)	1696 (13.0%)	388 (13.3%)	1421 (17.1%)
Video-assisted	31 727 (84.9%)	6170 (82.7%)	4765 (85.2%)	11 379 (87.0%)	2524 (86.7%)	6889 (82.9%)
Hospital-level factor: number of lobectomies in the observation period						
1–72	9436 (25.3%)	1778 (23.8%)	1453 (26.0%)	3313 (25.3%)	761 (26.1%)	2131 (25.6%)
73–117	9266 (24.8%)	1915 (25.7%)	1405 (25.1%)	3235 (24.7%)	711 (24.4%)	2000 (24.1%)
120–182	9714 (26.0%)	1966 (26.3%)	1440 (25.7%)	3335 (25.5%)	749 (25.7%)	2224 (26.8%)
184–482	8936 (23.9%)	1803 (24.2%)	1295 (23.2%)	3192 (24.4%)	691 (23.7%)	1955 (23.5%)
TNM, tumour node metastasis.						

and cutting vagal cardiac branches during the superior mediastinal lymph node dissection. Atrial fibrillation is a well-known cause of thrombosis in the left atrium,³³ and this study provides additional evidence for the frequency of postoperative cerebral infarction after lobectomy. In addition, it may be one possible reason for the high OR for left lower lobectomy in this study. However, this was not demonstrated to be a strong mechanism in the present cohort, as the post hoc analysis including postoperative atrial fibrillation showed less OR variation. This suggestion should be clarified in subsequent interventional studies.

It is important, but challenging, to establish methods to decrease the risk of postoperative cerebral infarction for patients undergoing a left upper lobectomy for lung cancer. One suggested solution is ensuring that the pulmonary vein stumps are as short as possible during the lobectomy process, because the pulmonary vein stump after left upper lobectomy remains long in many cases.^{7–9 12 13} Although almost all pulmonary veins are resected with linear staplers, care should be taken to move the linear stapler to the mediastinal side when the pulmonary veins are resected to ensure that the vein stumps are short. In addition, radiological examinations, especially contrast-enhanced CT in the early postoperative period, should be considered for the early detection of blood clots. The reported median postoperative period interval before the detection of thrombus is reported to be 3 months (range, 2–19 months),⁸ but the onset was within 24–48 hours in some cases.³⁴ As for medication, the use of anticoagulants (ie, heparin, warfarin or other direct oral anticoagulants) should be considered in accordance with the treatment of left atrial thrombosis. Indeed, several reports have stated that starting anticoagulant therapy immediately after surgery for pulmonary vein thrombosis led to the prevention of thrombosis and elimination of thrombus.^{12 35 36} The dosage and duration of treatment need to be adjusted for each patient, but the length of treatment is unknown, and the risk of bleeding needs to be considered. We hope that further research will reveal a reliable method to prevent this complication.

There are inherent limitations in the present study. First, owing to the nature of the DPC system, only data during hospitalisation could be analysed, which made it difficult to track cerebral infarctions that were related to lobectomies but occurred after discharge from the hospital. Second, data that could not be entered into the DPC database (eg, laboratory data and detailed radiological findings) and preoperative treatment for comorbidities before admission (eg, surgery or other procedures for arrhythmias such as pacemaker implantation and atrial Maze procedure) were not included in this analysis. Although some of this information can be inferred to some extent from the codes that were added when abnormal values are reported, more specific content that could be used as variables by extraction methods from sources other than the DPC system would improve the overall analysis. Third, this study is an epidemiological

**Table 2** OR for the occurrence of cerebral infarction

% of events	Number of events (n=99)	Patients (n=37 352)	Sites of lobectomy	Univariate			Multivariate*		
				OR	95% CI	P value	Adjusted OR	95% CI	P value
Left upper lobectomy	7462	29	0.39%	2.49	1.29 to 4.80	0.006	2.66	1.37 to 5.16	0.004
Left lower lobectomy	5593	19	0.34%	2.17	1.07 to 4.41	0.032	2.20	1.08 to 4.49	0.030
Right upper lobectomy	13 075	32	0.24%	1.57	0.82 to 3.00	0.171	1.60	0.83 to 3.07	0.158
Right middle lobectomy	2912	6	0.21%	1.33	0.50 to 3.52	0.562	1.29	0.48 to 3.45	0.607
Right lower lobectomy	8310	13	0.16%	Reference			Reference		

*Adjusted for sex, age, body mass index, smoking, activity of daily living, surgical approach (thoracotomy or video-assisted), stage, comorbidities and hospital-level factors.

study and has not led to specific verification of the previously mentioned causes and preventive measures. These mentions of causes and measures were only supported by case reports or small-scale case-control studies in previous reports. Thus, it is necessary to conduct further interventional research to detect certain causes and construct treatment strategies.

CONCLUSIONS

In this nationwide DPC study conducted in Japan, we identified the incidence of cerebral infarction associated with each type of lobectomy used to surgically treat patients with lung cancer. We found that the left upper lobectomy had the highest OR for postoperative cerebral infarction among all sites of lobectomies. The findings of this study have demonstrated the urgent need for new studies to investigate the specific mechanisms underlying postoperative cerebral infarction after left upper lobectomy and to establish preventive measures.

Acknowledgements The authors are grateful to the members of the Department of Environmental Epidemiology and Department of Public Health at the University of Occupational and Environmental Health, and the DPC Research Institute for their support with data collection and management.

Contributors NN, MM and YF conceived and designed the study. MO, YF, SM and KF acquired, linked and cleaned the data. MO and YF were responsible for statistical analysis. All authors interpreted the results, prepared the manuscript and approved the final version. YF, SM, KF and FT are the guarantors. The corresponding author attests that all listed authors meet the authorship criteria and that no others meeting the criteria have been omitted.

Funding This study was funded by the Ministry of Health, Labour, and Welfare, Japan (grant number: 20AA2005).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study was approved by the Ethics Committee of Medical Care and Research at the University of Occupational and Environmental Health, Japan (approval no. R2-007).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. Data for this study were derived from the database maintained by the Ministry of Health, Labour, and Welfare, Japan.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Natsumasa Nishizawa <http://orcid.org/0000-0002-0749-8871>

REFERENCES

- 1 National Cancer Center Japan. Cancer information Japan. Available: <https://ganjoho.jp/public/index.html> [Accessed 09 Jun 2022].
- 2 Shimizu H, Okada M, *et al*. Thoracic and cardiovascular surgeries in Japan during 2017 : Annual report by the Japanese Association for Thoracic Surgery. *Gen Thorac Cardiovasc Surg* 2020;68:414–49.
- 3 Xin Y, Hida Y, Kaga K, *et al*. Left lobectomy might be a risk factor for atrial fibrillation following pulmonary lobectomy. *Eur J Cardiothorac Surg* 2014;45:247–50.
- 4 Trouseau A. Phlegmatia alba dolens. In: *Clinique Médicale de L'hôtel-dieu de Paris*. 2nd edn. Paris: J.-B. Baillièere et fils, 1865: 3. 654–712.
- 5 Tsai Y-H, Yang J-L, Lee I-N, *et al*. Effects of dehydration on brain perfusion and infarct core after acute middle cerebral artery occlusion in rats: evidence from high-field magnetic resonance imaging. *Front Neurol* 2018;9:786.
- 6 Zorowitz RD, Tietjen GE. Medical complications after stroke. *J Stroke Cerebrovasc Dis* 1999;8:192–6.
- 7 Ohtaka K, Hida Y, Kaga K, *et al*. Pulmonary vein thrombosis after video-assisted thoracoscopic left upper lobectomy. *J Thorac Cardiovasc Surg* 2012;143:e3–5.
- 8 Ohtaka K, Hida Y, Kaga K, *et al*. Thrombosis in the pulmonary vein stump after left upper lobectomy as a possible cause of cerebral infarction. *Ann Thorac Surg* 2013;95:1924–8.
- 9 Ohtaka K, Takahashi Y, Uemura S, *et al*. Blood stasis may cause thrombosis in the left superior pulmonary vein stump after left upper lobectomy. *J Cardiothorac Surg* 2014;9:159.
- 10 Matsumoto K, Sato S, Okumura M, *et al*. Left upper lobectomy is a risk factor for cerebral infarction after pulmonary resection: a multicentre, retrospective, case-control study in Japan. *Surg Today* 2020;50:1383–92.
- 11 Hattori A, Takamochi K, Kitamura Y, *et al*. Risk factor analysis of cerebral infarction and clinicopathological characteristics of left upper pulmonary vein stump thrombus after lobectomy. *Gen Thorac Cardiovasc Surg* 2019;67:247–53.

- 12 Asai K, Mochizuki T, Iizuka S, *et al.* Pulmonary vein stump thrombus: an early complication following upper division segmentectomy of the left lung. *Gen Thorac Cardiovasc Surg* 2014;62:244–7.
- 13 Yamamoto T, Suzuki H, Nagato K, *et al.* Is left upper lobectomy for lung cancer a risk factor for cerebral infarction? *Surg Today* 2016;46:780–4.
- 14 Matsumoto K, Sato S, Okumura M, *et al.* Frequency of cerebral infarction after pulmonary resection: a multicenter, retrospective study in Japan. *Surg Today* 2018;48:571–2.
- 15 Nojiri T, Inoue M, Takeuchi Y, *et al.* Impact of cardiopulmonary complications of lung cancer surgery on long-term outcomes. *Surg Today* 2015;45:740–5.
- 16 The Ministry of Health, Labour, and Welfare, Japan. Survey on impact assessment of DPC introduction: aggregate results. Available: <https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000049343.html> [Accessed 09 Jun 2022].
- 17 Yasunaga H, Horiguchi H, Kuwabara K, *et al.* Clinical features of bowel anisakiasis in Japan. *Am J Trop Med Hyg* 2010;83:104–5.
- 18 The Ministry of Health, Labour, and Welfare, Japan. Various information of medical fee. Available: <https://shinryohoshu.mhlw.go.jp/> [Accessed 09 Jun 2022].
- 19 Suzuki S, Yamashita T, Okumura K, *et al.* Incidence of ischemic stroke in Japanese patients with atrial fibrillation not receiving anticoagulation therapy--pooled analysis of the Shinken Database, J-RHYTHM Registry, and Fushimi AF Registry. *Circ J* 2015;79:432–8.
- 20 Grisold W, Oberndorfer S, Struhal W. Stroke and cancer: a review. *Acta Neurol Scand* 2009;119:1–16.
- 21 The Japan Society for the Study of Obesity. *Guidelines for obesity treatment 2016*. Tokyo: Life Science Publishing, 2016.
- 22 Rami-Porta R, Bolejack V, Crowley J, *et al.* The IASLC lung cancer staging project: proposals for the revisions of the T descriptors in the forthcoming eighth edition of the TNM classification for lung cancer. *J Thorac Oncol* 2015;10:990–1003.
- 23 D'Agostino RB, Wolf PA, Belanger AJ, *et al.* Stroke risk profile: adjustment for antihypertensive medication. The Framingham study. *Stroke* 1994;25:40–3.
- 24 Pearce LA, Hart RG, Halperin JL. Assessment of three schemes for stratifying stroke risk in patients with nonvalvular atrial fibrillation. *Am J Med* 2000;109:45–51.
- 25 EAFT (European Atrial Fibrillation Trial) Study Group. Secondary prevention in non-rheumatic atrial fibrillation after transient ischaemic attack or minor stroke. *Lancet* 1993;342:1255–62.
- 26 Gage BF, Waterman AD, Shannon W, *et al.* Validation of clinical classification schemes for predicting stroke: results from the National Registry of atrial fibrillation. *JAMA* 2001;285:2864–70.
- 27 Nozawa T, Inoue H, Hirai T, *et al.* D-Dimer level influences thromboembolic events in patients with atrial fibrillation. *Int J Cardiol* 2006;109:59–65.
- 28 Tomita F, Kohya T, Sakurai M, *et al.* Prevalence and clinical characteristics of patients with atrial fibrillation: analysis of 20,000 cases in Japan. *Jpn Circ J* 2000;64:653–8.
- 29 Kodani E, Atarashi H, Inoue H, *et al.* Impact of creatinine clearance on outcomes in patients with non-valvular atrial fibrillation: a subanalysis of the J-RHYTHM registry. *Eur Heart J Qual Care Clin Outcomes* 2018;4:59–68.
- 30 Investigators AF. Risk factors for stroke and efficacy of antithrombotic therapy in atrial fibrillation. Analysis of pooled data from five randomized controlled trials. *Arch Intern Med* 1994;154:1449–57.
- 31 Chan KL. Transesophageal echocardiographic correlates of thromboembolism in high-risk patients with nonvalvular atrial fibrillation. The stroke prevention in atrial fibrillation Investigators Committee on echocardiography. *Ann Intern Med* 1998;128:639–47.
- 32 European Heart Rhythm Association, European Association for Cardio-Thoracic Surgery, Camm AJ, *et al.* Guidelines for the management of atrial fibrillation: the task force for the management of atrial fibrillation of the European Society of cardiology (ESC). *Eur Heart J* 2010;31:2369–429.
- 33 Lip GYH, Nieuwlaat R, Pisters R, *et al.* Refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation using a novel risk factor-based approach: the Euro heart survey on atrial fibrillation. *Chest* 2010;137:263–72.
- 34 Koga T, Mori T, Shibata H, *et al.* Two cases of organ infarction due to thrombus in pulmonary vein stump after left lung lobectomy. *Jpn J Chest Surg* 2016;30:621–7.
- 35 Ichimura H, Ozawa Y, Nishina H, *et al.* Thrombus formation in the pulmonary vein stump after left upper lobectomy: a report of four cases. *Ann Thorac Cardiovasc Surg* 2014;20 Suppl:613–6.
- 36 Manabe S, Oshima Y, Nakano M, *et al.* Renal infarction in a patient with pulmonary vein thrombosis after left upper lobectomy. *Case Rep Nephrol Urol* 2014;4:103–8.