



Review

The Intersection of Rural Residence and Minority Race/Ethnicity in Cancer Disparities in the United States

Whitney E. Zahnd^{1,2,*}, Cathryn Murphy^{1,3}, Marie Knoll⁴, Gabriel A. Benavidez^{1,4}, Kelsey R. Day⁵,
Radhika Ranganathan^{1,4}, Parthenia Luke⁶, Anja Zgodic^{1,4}, Kwei Shi¹, Melinda A. Merrell¹,
Elizabeth L. Crouch^{1,7}, Heather M. Brandt⁸ and Jan M. Eberth^{1,2,4}

- ¹ Rural & Minority Health Research Center, Arnold School of Public Health, University of South Carolina, Columbia, SC 29210, USA; cathrynm@email.sc.edu (C.M.); BENAVIDG@email.sc.edu (G.A.B.); radhika@email.sc.edu (R.R.); azgodic@email.sc.edu (A.Z.); kwei.shi@yale.edu (K.S.); mmerrell@mailbox.sc.edu (M.A.M.); crouchel@mailbox.sc.edu (E.L.C.); jmeberth@mailbox.sc.edu (J.M.E.)
 - ² Big Data Health Science Center, Arnold School of Public Health, University of South Carolina, Columbia, SC 29208, USA
 - ³ College of Arts and Sciences, University of South Carolina, Columbia, SC 29208, USA
 - ⁴ Department of Epidemiology & Biostatistics, Arnold School of Public Health, University of South Carolina, Columbia, SC 29208, USA; mknoll@email.sc.edu
 - ⁵ Department of Exercise Science, Arnold School of Public Health, University of South Carolina, Columbia, SC 29208, USA; Krday@email.sc.edu
 - ⁶ College of Social Work, University of South Carolina, Columbia, SC 29208, USA; lukep@email.sc.edu
 - ⁷ Department of Health Services Policy and Management, Arnold School of Public Health, University of South Carolina, Columbia, SC 29208, USA
 - ⁸ St. Jude Children's Research Hospital, Memphis, TN 38105, USA; heather.brandt@stjude.org
- * Correspondence: zahnd@mailbox.sc.edu; Tel.: +1-803-576-6057



Citation: Zahnd, W.E.; Murphy, C.; Knoll, M.; Benavidez, G.A.; Day, K.R.; Ranganathan, R.; Luke, P.; Zgodic, A.; Shi, K.; Merrell, M.A.; et al. The Intersection of Rural Residence and Minority Race/Ethnicity in Cancer Disparities in the United States. *Int. J. Environ. Res. Public Health* **2021**, *18*, 1384. <https://doi.org/10.3390/ijerph18041384>

Academic Editor: Paul B. Tchounwou
Received: 18 December 2020
Accepted: 30 January 2021
Published: 3 February 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: One in every twenty-five persons in America is a racial/ethnic minority who lives in a rural area. Our objective was to summarize how racism and, subsequently, the social determinants of health disproportionately affect rural racial/ethnic minority populations, provide a review of the cancer disparities experienced by rural racial/ethnic minority groups, and recommend policy, research, and intervention approaches to reduce these disparities. We found that rural Black and American Indian/Alaska Native populations experience greater poverty and lack of access to care, which expose them to greater risk of developing cancer and experiencing poorer cancer outcomes in treatment and ultimately survival. There is a critical need for additional research to understand the disparities experienced by all rural racial/ethnic minority populations. We propose that policies aim to increase access to care and healthcare resources for these communities. Further, that observational and interventional research should more effectively address the intersections of rurality and race/ethnicity through reduced structural and interpersonal biases in cancer care, increased data access, more research on newer cancer screening and treatment modalities, and continued intervention and implementation research to understand how evidence-based practices can most effectively reduce disparities among these populations.

Keywords: rural; racial/ethnic minorities; cancer disparities; access to care; social determinants of health; cancer surveillance; cancer outcomes

1. Introduction

Rural populations in the United States comprise as much as 20% of the total population (i.e., as many as 59 million persons) [1]. Defining “rural” is objectively and subjectively challenging, as it is often driven by administrative geographic units (e.g., counties) for which there may be considerable variation in geographic and population size [1]. Regardless of the formal definition of “rural”, a “rural” area is typically characterized by small population density or size and notable distance to a metropolitan/urban area [1]. Three in

four rural persons in the United States live in the South or Midwest. However, the most isolated rural areas of the country are in the West, and even the population-dense Northeast includes sizable pockets of rural populations. Rural populations face a myriad of structural, socioeconomic, economic, environmental, and access-to-care barriers that put them at greater risk for higher prevalence of poor health behaviors (e.g., smoking, sedentary behavior), lower utilization of cancer-relevant preventive services, lower odds of receiving guideline-concordant treatment, and poorer cancer outcomes from incidence to survivorship [2–8].

Definitions of race and ethnicity are also socially and administratively driven. In this article, we categorize racial/ethnic groups as non-Hispanic White (the majority group), non-Hispanic Black, Hispanic, American Indian/Alaska Native, Asian, Native Hawaiian, and Other Pacific Islander. These are racial/ethnic groupings used by the Office of Management and Budget and the National Institute on Minority Health and Health Disparities [9,10]. Tangibly speaking, race and ethnicity are individual-level indicators of structural, institutional, and/or interpersonal disadvantage (or, conversely, advantage) that systematically and differentially affect population subgroups. Racial/ethnic minority groups are affected by both unique and similar historical and contemporary racism (both interpersonal and systemic) and other structural barriers to optimal health. Non-Hispanic Black populations have been affected by centuries of enslavement and brutality and have been particularly affected by segregationist policies (e.g., Jim Crow laws, redlining, exclusionary zoning) that have long-lasting social and health impacts beyond political, judicial, and legislative remedies (e.g., reconstruction, the Brown vs. the Board of Education decision, and the Fair Housing Act of 1968) [11,12]. American Indian and Alaska Native populations have similarly experienced sustained ill-health effects due to colonialism, forced migration, and racist policies (e.g., forced sterilization, delayed voting rights) [11]. Hispanic populations, comprising the largest current and fastest-growing rural population in the U.S., have been particularly affected by restrictive and xenophobic immigration policies that may affect health and access to healthcare services regardless of immigration status [12–14]. Asian and Native Hawaiian and Other Pacific Islander populations have been affected by discriminatory legislation and policies (e.g., exploitation of Chinese laborers in the late 19th century, exclusionary immigration policies, internment of Japanese Americans in World War II) [15,16]. These historic traumas, in addition to continued discriminatory policies and interpersonal biases, have limited minority populations' access to healthcare services, put them at greater odds of engaging in poorer health behaviors or being exposed to environmental carcinogens, make them less likely to receive optimal cancer treatment, and contribute to their greater cancer burden across the continuum [17–23].

The intersection of rurality and minority race/ethnicity in the United States is critical to our understanding of health disparities broadly, and cancer disparities specifically, but is often overlooked [24–27]. Overall, one in every 25 people in the U.S. is a non-White rural person (i.e., 22% of the U.S. rural population) [28]. Although most rural persons are White, 9% and 8% belong are Hispanic and Black, respectively. Rural, non-Hispanic Black and Hispanic populations reside primarily in the South (Figure 1), whereas rural American Indian/Alaska Native populations primarily reside in Alaska, the northern plains, and the Southwest. Although rural populations overall have declined because of outmigration and urban sprawl, rural Hispanic populations have grown in recent decades [29]. Furthermore, there are many counties throughout the country in which notable proportions of the populations are non-White (i.e., smaller proportions of racial/ethnic minority populations combine for a notable proportion of the total population). The intersection of the aforementioned structural barriers experienced by rural and racial/ethnic minority populations will be the focus of this paper. We begin our review by describing structural and contextual factors that disproportionately affect rural racial/ethnic minority populations and contribute to cancer disparities. Next, we summarize the literature from the past 20 years on disparities experienced by rural, racial/ethnic minority populations across

the cancer control continuum. Finally, we discuss the opportunities for observational and interventional research to mitigate these disparities.




Figure 1. Racial/ethnic composition across rural counties, 2018. Rural and urban counties are defined by the non-metro and metro designations from the United States Department of Agriculture [30]. Racial/ethnic composition by county is based on the 2018 Census Bureau Population Estimates Program [31].

2. A Conceptual Framework

Structural, social, and environmental factors at the macro and supramacro levels (e.g., policies, area-level poverty, area-level distance to care), in addition to micro-level factors (e.g., insurance status, age, behaviors), affect cancer across the continuum from etiology to survivorship. Supramacro factors, such as state-level policies, may have both an effect on lower level macro and micro factors and, subsequently, cancer outcomes across the continuum. For example, state-level Medicaid expansion, which is less common in largely rural states with higher proportions of Black residents, may affect the macro-level factor of potential access to care, the micro-level factor of individual insurance status and, subsequently, may affect cancer screening uptake, cancer staging, and cancer treatment [32–34]. To better contextualize the interplay of race/ethnicity and rurality, here we also adapt a framework from our previous work to depict how the intersection of rurality and race/ethnicity, combined with these multilevel factors, affects cancer outcomes across the continuum and demonstrate how racism at different levels may affect cancer outcomes by drawing from Gee and Ford, as well as Williams (Figure 2) [35–37]. Our adapted framework notes that historical racism (also referred to as “intergenerational drag”, as characterized by Gee and Ford) can affect different levels—the micro/individual level (interpersonal racism), the macro-level (racial/ethnic segregation, cultural racism), and the supramacro level (systemic racism)—also noting that racism has effects between levels as well. We place systemic racism and segregation at different levels, as often higher-level factors (e.g., state, federal policies that are systemic) have the greatest impact and/or are more effectively quantitatively measured at lower levels (e.g., county- or neighborhood-level residential segregation). The cancer continuum can be considered discretely within primary, secondary,

or tertiary prevention based on how public health and healthcare systems can intervene to address each area. As in our previous work, we here integrate both the National Cancer Institute’s Cancer Control Continuum and Wingo’s Framework for Cancer Surveillance to characterize rural cancer disparities within this continuum [37,38]. Areas of primary prevention include etiology and prevention, secondary prevention includes screening, early detection, and diagnosis, and tertiary prevention includes treatment, survivorship, and mortality.




Figure 2. A conceptual framework of the multilevel influence of racism on rural cancer disparities across the continuum. The role of racism across levels is noted in bold.

3. Social Determinants of Health and Their Role in Rural and Racial/Ethnic Disparities in Cancer

Rural racial/ethnic minority populations often fare worse than their non-minority rural and urban minority counterparts with regard to the social determinants of health and health outcomes, including cancer outcomes [39–41]. Here, we summarize how several social and structural determinants of health adversely affect rural racial/ethnic minority populations: structural racism, residential segregation, access to healthcare services, socioeconomic status, educational attainment, infrastructure, and environmental exposures.

3.1. Structural Racism and Residential Segregation

Interpersonal discrimination is recognized as an important social determinant of health that affects cancer outcomes [42,43]. However, both historical and systemic racism, such as residential segregation, also play a major role in observed racial and ethnic geographic health disparities [44,45]. Studies have shown that women, particularly Black women, born in the era of Jim Crow were more likely to be diagnosed with the breast cancer subtypes that have poorer prognoses [46]. Jim Crow laws were state and local regulations commonly found in the South that codified racial segregation in transportation, employment, restaurants, and other settings [46]. Similarly, in urban areas, living in a historically “redlined” area is associated with later stage at diagnosis and poorer cancer survival [47]. “Redlining” was another form of government-sponsored segregation that essentially prevented individuals in Black communities from successfully obtaining mortgages [47]. Segregation contributes to health disparities because it “isolates a minority

group from amenities, opportunities, and resources that affect social and economic well-being" [48]. One qualitative study that focused on Black residents of the rural Deep South, for example, identified several determinants of health at the structural level, including lack of opportunity for physical activities; lack of access to healthy foods; cronyism and nepotism in workplaces that favor White residents, a determinant of health given the association between income and health; persistent stress from poverty and institutional racism, a physiological cause of health risk [49]. Not only are Black people who live in segregated areas more likely to be diagnosed with breast and lung cancer at a later stage, but racially isolated areas also have less access to diagnostic cancer technologies [21,50]. Prior studies have also revealed that structural racism is harmful to residents of segregated areas regardless of race. Both Black and White breast cancer patients living in areas with high levels of Black segregation regardless of area-level poverty were less likely to receive appropriate care, and cancer survivors in similar areas had lower health-related quality of life even after accounting for area-level poverty, suggesting a systemic problem [51,52]. Although the intersection of race and economic class is important to consider, a national study found lung cancer mortality to be the highest among Black people living in the most segregated areas of the country, regardless of socioeconomic status or health insurance [21,53]. However, a study by Moss and colleagues found that screening rates were higher in more segregated areas regardless of race/ethnicity [54]. The intersection of residential segregation and rurality, as well as their compounding effects on health outcomes, is a critical area for additional research, especially as most rural minority populations live in the U.S. South, where the long-term effects of racist policies may be more prevalent, and interpersonal racism remains common [55].

3.2. Access to Healthcare Services

Healthcare access is often defined along five specific dimensions: availability, accessibility, accommodation, affordability, and acceptability [56]. Historical unethical treatment of racial/ethnic minority populations within the U.S. healthcare system (e.g., medical experimentation, forced sterilization, denial of needed medical treatment), as well as recent changes in rural healthcare systems, undermines access to healthcare [42]. Policies that segregated care among Black and White patients are embedded within the healthcare policy of the last century. The Hill-Burton Act passed in 1946 provided millions of dollars to increase the healthcare and public health infrastructure throughout the country but also allowed for racial segregation healthcare facilities [57]. Legal challenges and the Civil Rights Act led to the desegregation of hospitals. However, this also led to Black hospitals ultimately becoming financially inviable. This, coupled with cost slashing federal policies and changes to hospital payment structures in the 1970s and 1980s, led to a surge of hospital closures, particularly in the rural South [58]. In the context of both historical and contemporary policies, it is critical to examine these dimensions through the lenses of rurality, race/ethnicity, and cancer disparities.

Availability, accessibility, and affordability of healthcare services have been more comprehensively studied than other aspects of access. These dimensions are more readily quantified from publicly available health care workforce and insurance status data. Availability of services has decreased because of a surge in rural hospital closures in the past ten years, which has disproportionately affected rural minority populations [59]. Similarly, rural minority populations, particularly American Indian/Alaska Native populations, tend to live farther from healthcare services (i.e., less accessibility), including cancer care services such as National Cancer Institute (NCI)-designated cancer centers and cancer care specialists [60,61]. Affordability is affected by the inability of rural minority populations to access health insurance, either as a result of limited Medicaid expansion in states with large rural minority populations or because of the Affordable Care Act regulations associated with immigration status [62]. Furthermore, compared to their White counterparts, racial/ethnic minority populations are more likely to face financial hardship in paying for cancer care [63].

Acceptability is influenced by both racial/ethnic minority populations' expectations of, and reality with, experiencing discrimination in healthcare systems and their mistrust of healthcare providers as a result of historical unethical and criminal behaviors, such as unauthorized treatments and experiments performed on Black people, and present-day biases (e.g., pain perception) [64–66]. Acceptability may be facilitated by racial–ethnic concordance between patients and providers [67]. However, the closure of many Black medical schools in the 20th century may have reduced the number of Black physicians by ~35,000 roughly over the last century. In 2019, only 6.2% of medical school graduates were Black, less than half the proportion of the Black population nationwide [68]. This is particularly concerning in rural areas with large minority populations, as these areas already have a shortage of physicians.

Relatedly, racial/ethnic minority populations may face stigma accessing key health-care infrastructure, which led the American College of Physicians to issue a recent call to reduce such inequities [69]. Lack of accommodation in healthcare access may include limited hours of clinic operation or lack of translation services. These limited hours may have a disproportionate effect on rural and racial/ethnic minority populations who may be more likely to work for small businesses with less flexible work-leave policies or who are more likely to work in manufacturing and service professions for which there are multiple shifts [70,71]. Furthermore, rural racial/ethnic minority persons whose primary language is not English may face additional challenges with language barriers in health-care settings, which is especially critical with the growing population of immigrants from Spanish-speaking countries who now reside in rural America [14].

3.3. Socioeconomic Status, Educational Attainment, Housing, and Infrastructure

Generally, rural populations have higher rates of poverty, lower educational attainment, and more limited or aging community infrastructure than their urban counterparts [28,72]. Racial/ethnic minority populations, especially those living in rural areas, are disproportionately affected by these social and structural factors [73]. Historical racism and its long-term effects may play a role in these contextual factors [74]. Residential segregation also has led to concentrated Black poverty, as is seen by the clusters of persistent poverty along the Lower Mississippi River [74,75]. Furthermore, the impact of historical racism has had long-term effects on educational attainment with segregation shown to be associated with high school and college graduation among Black students, but not White students [74,76]. Poverty and educational attainment are both worse among rural racial/ethnic minority residents in comparison to their White counterparts. Poverty is higher among rural American Indian/Alaska Native (29%), non-Hispanic Black (24%), and Hispanic (21%) residents compared to White (10%) and Asian (9%) rural persons (Figure 3) [77–80]. Among rural residents, the highest rates of those with less than a high school diploma were among Hispanic (35%), Black (21%), and American Indian/Alaska Native (19.7%) populations compared to White (10%) and Asian (12%) populations [77–80]. Major housing inadequacies, such as plumbing issues, occur in a higher proportion of rural areas and decrease as areas become more metropolitan. Infrastructure issues, including lower access to safe roads, public transportation, and broadband, also exist in rural areas [77–81]. These social and structural determinants contribute to disparities in cancer outcomes among rural and racial/ethnic minority populations by inducing persistent stress, enabling or promoting poorer health behaviors, reducing access to cancer care, and affecting guideline-concordant treatment and cancer survivorship [73,82–86].




Figure 3. Social determinants of health across rural racial/ethnic groups (findings reported by Probst et al. based on American Community Survey data) [77–80].

3.4. Environmental Factors

Rural areas, particularly those with a higher proportion of residents reliant on well water and agricultural or mining industries, may be at an increased risk of exposure to carcinogens such as arsenic, radon, and nitrate [87]. Rural populations may be at greater risk of exposure to radon because of lower rates of home testing and, therefore, subsequent mitigation [88]. However, many of the previous studies that focused on well water, agricultural exposures, and radon testing and mitigation have been concentrated in the Northeast and Midwest, where rural populations are largely White [89–91]. Historical and contemporary environmental racism may affect rural, racial/ethnic minorities' exposures to carcinogens. Examples include the placement of landfills for carcinogenic polychlorinated biphenyl (PCB) in rural Warren County North Carolina in 1982, petrochemical exposures in “Cancer Alley” in rural, largely Black Louisiana, and exposures to carcinogens among tribal populations due to mining near reservations [91]. “Cancer Alley” is a 100-mile stretch between Baton Rouge and New Orleans, Louisiana, that includes the largely rural St. John the Baptist and St. James' counties. One-quarter of the nation's petrochemical production occurs in this area and includes a population that is 40% Black [92]. These kinds of exposures can lead to a greater risk of cancer incidence and poorer cancer outcomes. For example, recent studies from North Carolina showed that land quality, as indicated by agricultural activity, pesticide usage, and toxic release and priority cleanup sites, was associated with later stage at diagnosis for breast cancer among rural women [93]. Identifying the independent causal association between environmental exposures and cancer incidence is difficult at a population level, but it is critical to understanding and mitigating rural and racial/ethnic disparities in cancer [94,95].

4. Rural and Racial/Ethnic Disparities in Cancer Across the Continuum

4.1. Risk Factors and Primary Prevention

Significant differences in health behaviors have been observed across rural–urban and racial/ethnic subgroups, including tobacco use, alcohol consumption, diet, physical activity, and human papillomavirus (HPV) vaccination. Rural populations tend to have higher rates of tobacco use, diets higher in red/processed meat and lower in fruit/vegetable consumption, lower levels of physical activity, and lower uptake of HPV vaccination, but also tend to have lower alcohol consumption compared to urban areas [2–4]. Cancer-relevant health behaviors vary uniquely by race/ethnicity (e.g., smoking is higher among American Indian/Alaska Native populations, but Black populations have lower rates of physical activity) [55].

Smoking and smokeless tobacco use vary among rural racial/ethnic minority groups. Historically, American Indians and Alaskan Natives have been identified in national surveys, such as the Behavioral Risk Factor Surveillance System (BRFSS), as having the highest rates of current smoking among rural persons (36.7%), with rates higher than those

seen in other racial/ethnic minority groups: non-Hispanic Black (23.2%), Hispanic (17.0%), and Asian/Pacific Islander/Native Hawaiians (10.9%) [55]. Studies have also shown intrarural differences in smoking among American Indian/Alaska Native populations, with those living on tribal lands having lower current smoking prevalence compared to those living in non-tribal rural areas [96]. There are some regional and tribal variations in smoking prevalence among American Indian/Alaska Native populations, even within a specific state. For example, smoking prevalence among rural Alaska Natives varies notably across regions of the state [97]. It is important to note, however, that smoking in ceremonial settings is an important part of many tribal cultures [98]. This may affect the interpretation of national studies and the representation of smoking prevalence among these populations. Additionally, national surveys have shown that smoking rates vary by rurality and racial/ethnicity, e.g., among rural Black persons, smoking rates are lower than among urban Black persons [99]. Smokeless tobacco use tends to be most prevalent among rural White populations when a dichotomous analysis of race/ethnicity is considered (i.e., White vs. non-White) [100]. However, studies have shown that rural Alaskans, particularly Native Alaskans, have a higher prevalence of smokeless tobacco use compared to their White rural counterparts [101]. A rural North Carolina study showed that Black and Native American persons were more likely to use “snuff”, whereas Native Americans were more likely to use chewing tobacco compared to their White counterparts [102]. Another study showed that among a rural cohort of adolescents in California, non-Hispanic White teens were more likely to use smokeless tobacco compared to their rural Hispanic counterparts [103].

Alcohol use, which is associated with multiple cancer types, including forms of the liver, mouth, and esophagus, differs across rural–urban and racial/ethnic minority populations [104]. Binge drinking rates are lower among rural Black and Asian/Pacific Islander populations compared to rural White adults, but there was no difference between White adults and American Indian/Alaska Native or Hispanic adults [55]. Other recent studies focused on adolescent alcohol use have yielded similar findings [105]. Within the rural Hispanic farmworker population in North Carolina, the risk of alcohol abuse was found to be positively associated with years of residence in the United States [106]. This finding underscores the importance of considering the role of assimilation in cancer risk among rural Hispanic populations, as some rural areas are home to both newly immigrated and long-residing Hispanic populations.

Diet is a complex cancer-relevant health behavior that, depending on the foods consumed, can either decrease (e.g., fruits and vegetables) or increase (e.g., red and processed meats) risk for developing cancer [107]. A comprehensive study of consumption habits is critical in rural communities, given the strong overlap between food deserts and rural areas [108,109]. This intersection of rurality and diet is directly visible among different racial/ethnic groups. For example, in the Navajo Nation, 57% of residents in the Community Outreach and Patient Empowerment Program study stated they did not consume enough fruits and vegetables. Of these individuals, 61% cited expense as a reason, and over 51% reported needing to travel over an hour to obtain most of their food (which may have discouraged the purchase of fresh, perishable items) [110]. Similarly, a study in rural North Carolina found that rural Black older adults had a better Total Health Eating Index-2005 score compared to their White and American Indian counterparts, but the groups varied in their consumption of different types of healthy foods. However, no group had an overall high-quality diet [111]. Studies based in other states (Mississippi) have shown that rural Black persons have poorer diets than their rural White counterparts and the nation as a whole, suggesting that there may be geographic variation in these disparities [112]. The influence of these cultural and geographic-related factors on healthy diets among rural racial/ethnic minority groups demonstrates the need for tailored interventions for different populations to address the cancer burden, paying particular attention to the traditional food choices of particular regions and/or cultures.

In a similar vein, there is a connection between high body mass index and lack of physical activity and many types of cancer (e.g., colorectal cancer, postmenopausal breast cancer, pancreatic cancer). Studies have shown that rural populations have higher rates of obesity and lower levels of physical activity compared to their urban counterparts and that obesity/physical activity varies by race/ethnicity among rural populations [3,55]. Among rural populations, Black, Hispanic, and American Indian/Alaska Native populations had higher rates of obesity than their rural White peers, although roughly a third or more of all these groups were obese (ranging from 32.0% among rural White adults to 45.9% of rural Black adults) [55]. Rural Asian/Native Hawaiian/Other Pacific Islander populations had the lowest rate of obesity (15.5%) [55]. Other studies have shown that rates of obesity are higher among rural Black and White populations compared to their respective urban counterparts [113]. These higher rates of both obesity and physical inactivity may be due, in part, to obesogenic built environments that may be more prevalent in areas with larger racial/ethnic minority populations (i.e., the South) [114,115].

In addition to health behaviors that may put rural racial/ethnic minority populations at greater risk for cancer, it is important to assess the prevalence of health-promoting behaviors such as HPV vaccination, which can protect against several cancer types (e.g., cervical cancer, oropharyngeal cancer) [116]. Both rural and Black populations have higher incidence and mortality rates of these preventable, HPV-associated cancers [5,117]. HPV vaccination initiation rates are consistently lower among rural adolescents overall and among rural White and Hispanic adolescents specifically [118]. There has been no statistically significant difference in HPV vaccination initiation among Black adolescents across rural–urban designations [118]. Another recent study showed that vaccination rates were higher among Black and Hispanic teens compared to their White counterparts regardless of rural–urban residence [4]. Although vaccination rates have been more favorable among rural racial/ethnic minority populations, studies have shown that lower knowledge of the role of HPV vaccination in cancer prevention and financial and psychosocial barriers may play a role in lower rates of HPV vaccination in rural areas and among some rural racial/ethnic minority populations, such as newly immigrated Hispanics [119,120].

Racial segregation may play a role in many of these health behaviors that are more common among rural racial/ethnic minority populations. For example, studies show that Black persons in more segregated areas have higher smoking rates and have less access to healthy foods [121,122]. However, these types of studies examining the relationship between residential segregation and cancer-related health behaviors have primarily been performed in urban areas. Examining rural residential segregation is an important area of needed research to help explain and address cancer-related health behaviors [123].

4.2. Cancer Screening

For many cancer types (e.g., breast, colorectal, cervical, lung), receipt of screening at the recommended ages and intervals can detect cancer at an earlier, more treatable stage, improving outcomes and survival. Lower utilization of screening services in rural areas and areas with less geographic access to screening services have been well documented [124–126]. However, there have been mixed findings on racial/ethnic differences overall and at the intersection of race/ethnicity and rurality. Some studies have found that in rural communities, ethnicity (Hispanic vs. non-Hispanic) was not associated with having a guideline-concordant mammogram [127]. Disparities in race were seen in other studies that showed that rural Hispanic women were less likely to have had a mammogram within the past year compared with urban Hispanic women [128]. Rural Black women in persistent-poverty counties were less likely to report a recent mammogram compared to their urban counterparts [125]. However, another study found that for breast and cervical cancers, Black women had favorable odds of receipt of service regardless of rurality [129]. Multiple studies have shown that rural American Indian populations have lower rates of cancer screening compared to their urban counterparts, with lower rates found among those living in the most remote areas [128,130]. Rural Black persons had a

lower probability of reporting colorectal cancer screening than urban Black persons (44.8% vs. 51.8%) [7]. Similarly, rural Hispanics/Latinos had a lower predicted probability of reporting colorectal cancer screening than urban Hispanics/Latinos (40.8% vs. 43.7%) [7]. Since its recommendation by the United States Preventive Services Task Force (USPSTF) in 2013, lung cancer screening uptake at the intersection of race/ethnicity and rurality has been difficult to assess because of limited data availability and small sample sizes [131,132]. However, initial analyses of national surveys have shown that there are no rural–urban or racial/ethnic differences in screening uptake [132–134]. Studies have shown that lack of recommendation from a provider is a key barrier to being up to date with screening among rural patients [135]. Furthermore, low quality of provider information has been identified as a barrier to screening among rural Black individuals [136]. Provider biases may play a role in lack of provider recommendations and/or low-quality recommendations. Further research is needed to examine the role of biases in cancer screening recommendations and referrals among rural, racial/ethnic minority patients.

4.3. Cancer Incidence and Staging

We provide a summary of age-adjusted incidence rates for all cancers combined and several common cancer types across rural/urban designations and racial/ethnic groups for the most recent 5-year period of available data (2013–2017) using SEER 21 data (Table 1).

Generally speaking, rural, racial/ethnic minority populations have had higher incidence rates for lung and colorectal cancers. Rural Black populations had the highest rates of colorectal, female breast, prostate, and cervical cancer across rural groups. Previous studies utilizing more representative national data have shown that overall cancer incidence is higher among urban populations, driven in large part by higher rates of breast and prostate cancer, for which access to and utilization of screening may be a driving factor [5,6]. This holds across racial strata, except for American Indian/Alaska Native populations, among whom the all-cancer incidence rate was higher among rural populations, and Asian/Pacific Islanders, among whom there was no rural–urban difference [6]. However, for cancers with modifiable risk factors (e.g., tobacco-associated and HPV-associated cancers) or preventive screening opportunities (e.g., colorectal cancer), incidence rates are higher in rural compared to urban populations, with greater disparities among rural non-Hispanic Black persons, although rural Hispanic persons tended to experience lower incidence compared to their urban peers [5,6]. Early-onset colorectal cancer (i.e., colorectal cancers in those aged 20–49) has increased in rural populations since 2000, with persistently high rates among rural Black populations, and with the highest recent rate seen among rural American Indian/Alaska Native populations [138]. For HPV-associated cancers broadly and cervical cancer specifically, rates are highest for rural Black persons [5]. For cervical cancer, rural Black, White, and American Indian women have the higher rates compared to their urban counterparts [139]. Black men tended to have higher lung cancer incidence rates across all levels of rurality and for all histology types except small cell [140]. Although urban populations tend to have higher rates of testicular cancer overall, rural American Indian/Alaska Native and Asian/Pacific Islanders have higher rates compared to their urban counterparts, whereas other racial/ethnic groups experienced no rural–urban differences or showed more favorable rates in rural populations [141]. Studies have also tended to show that rural populations, regardless of race/ethnicity, are diagnosed with cancer at a more advanced stage, particularly for cancers with primary and secondary preventive opportunities, such as lung, cervical, and colorectal cancers [142,143].

Table 1. Rural and urban age-adjusted cancer incidence rates across racial/ethnic groups, 2013–2017.

Racial/Ethnic Group	Age-Adjusted Rural Incidence Rate per 100,000	Age-Adjusted Urban Incidence Rate per 100,000
All Cancers		
Non-Hispanic White	471.9	470.6
Non-Hispanic Black	455.9	454.1
American Indian/Alaska Native	338.5	337.8
Asian/Pacific Islander	320.3 *	305.9
Hispanic	322.9	349.3 *
Lung Cancer		
Non-Hispanic White	71.6 *	58.5
Non-Hispanic Black	68.1 *	56.1
American Indian/Alaska Native	35.7	41.2 *
Asian/Pacific Islander	40.5 *	36.4
Hispanic	31.3 *	28.9
Colorectal Cancer		
Non-Hispanic White	43.6 *	37.7
Non-Hispanic Black	52.7 *	44.6
American Indian/Alaska Native	42.3 *	33.1
Asian/Pacific Islander	34.8	32.2
Hispanic	35.3	33.7
Prostate Cancer		
Non-Hispanic White	99.7	106.0 *
Non-Hispanic Black	166.7	181.9 *
American Indian/Alaska Native	64.3	64.0
Asian/Pacific Islander	61.2	57.2
Hispanic	69.6	93.0 *
Female Breast Cancer		
Non-Hispanic White	120.1	139.7 *
Non-Hispanic Black	124.2	128.9
American Indian/Alaska Native	80.4	94.1 *
Asian/Pacific Islander	109.4	104.3
Hispanic	89.7	99.4 *
Cervical Cancer		
Non-Hispanic White	8.6 *	6.5
Non-Hispanic Black	10.8 *	8.8
American Indian/Alaska Native	10.8	7.9
Asian/Pacific Islander	6.6	6.6
Hispanic	8.4	9.2

Note: Rates are age-adjusted per the 2000 U.S. Standard Population and calculated from the Surveillance Epidemiology and End Results (SEER) 21 April 2020 data release [137]. Data from the Alaska and Hawaii Registries are not included as rural–urban indicators are not available. * indicates a statistically significant higher rate among a racial/ethnic group compared to their geographic counterpart.

4.4. Cancer Treatment

Although studies have shown that rural and racial/ethnic minorities are less likely to receive certain cancer treatments or receive them in accordance with guidelines, few studies have examined the intersection of rurality and race/ethnicity in their effects on cancer treatment [144,145]. Those studies that have explored this intersection have identified notable disparities among rural racial/ethnic minority cancer patients. For example, in the treatment of early-stage non-small cell lung cancer (NSCLC), complete surgical resection is considered the best treatment option to increase survival. However, in an analysis of 3481 Alabama Medicare beneficiaries diagnosed with NSCLC, the proportion of rural Black patients (10.7%) receiving surgery at any point was significantly less than

the proportion of rural White patients (28.7%) [146]. A study examining surgical treatment for endometrial cancer found that the proportion of Black women who did not undergo a lymphadenectomy, a procedure to remove and examine lymph nodes to help determine cancer progression, was significantly higher among those in rural compared to urban areas (40.9% vs. 30.3%) [147]. Multivariable analysis of national disparities in laparoscopic procedures for colon cancer treatment found similar results: the odds of undergoing less-invasive laparoscopic procedures were significantly higher among Black (OR: 2.2; 95% CI: 1.9, 2.5) and Hispanic (OR: 2.2; 95% CI: 1.9, 2.5) patients that were cared for in urban areas compared to their rural counterparts [148]. Several other studies have demonstrated that, when examined independently, racial/ethnic minorities and persons living in rural or non-metro areas have lower proportions and/or odds of receiving higher-quality cancer treatment across multiple modalities for a variety of cancers [149–154]. Furthermore, studies have shown that rural Black and Hispanic cancer patients were less likely to report easily getting care compared to White patients [155]. Some studies, although not specifically evaluating disparities in treatment, provide insight into why certain racial/ethnic disparities are likely to be present among rural populations. In a study of breast cancer patients from multiple U.S. states, Black (OR=1.93; 95% CI: 1.17, 3.18) and Hispanic (OR= 2.01; 95% CI: 1.21, 3.35) women were found to have significantly higher odds of being treated at low-volume hospitals [156]. Low-volume hospitals are much more prevalent in rural areas and, according to some studies, have been shown to offer lower-quality care in comparison to higher-volume hospitals [157,158].

Rural, racial/ethnic minority cancer patients are less likely to receive guideline-concordant care for their cancer, particularly when it comes to cancer surgery. This may be due in part to these populations being more frequently treated at low-volume hospitals. More research is needed to examine cancer treatment disparities across cancer sites and treatment modalities and to determine the independent contributions of contextual and treatment location factors. Furthermore, it is imperative to understand the dynamics between patient preference, provider biases, and provider–patient communication [159]. Studies have shown that perceived racism, distrust of the healthcare system, and providers' spurious perceptions of treatment efficacy play a role in the receipt of cancer treatment [159–162]. However, studies have yet to examine if and to what magnitude these factors may play a role in the treatment disparities experienced by rural racial/ethnic minority patients.

4.5. Cancer Survivorship

Survivorship comprises the health and wellbeing of a person with cancer from diagnosis until the end of life, including physical, mental, social, emotional, and financial effects of cancer that start at diagnosis and continue beyond treatment [163]. Rural and racial/ethnic disparities in survivorship issues (e.g., long-term follow-up care/surveillance, quality of life, financial toxicity) have been identified [164–169]. Rural cancer survivors have the highest rates of poor self-reported health, physical distress, and activity limitations as compared to their urban counterparts, as well as reported lower receipt of guidance about cancer follow-up care [167]. In addition, rural cancer survivors had poorer mental health outcomes compared to urban cancer survivors [165,168]. Studies have shown that rural cancer survivors report higher likelihood of financial toxicity compared to their urban counterparts, but this association is attenuated upon adjustment for other factors [170,171]. Studies have also identified cancer survivorship disparities among racial/ethnic minorities. Black and Hispanic breast cancer survivors were less likely to have surveillance mammography compared to their White counterparts. Other studies have shown that Black and Hispanic cancer survivors were more likely to forego prescription medications and dental care because of cost or experienced greater material and psychological financial hardship than White cancer survivors [172,173]. However, the intersection of rurality and race/ethnicity and their effect on cancer survivorship has been only minimally explored. A study among breast cancer survivors in the Carolina Breast Cancer Study found that rural survivors were

most financially affected by their diagnosis, but the magnitude of this impact was greatest among rural Black survivors [174]. Another study in South Carolina found that Black breast cancer survivors, particularly those in rural areas, had lower medication possession ratios, indicating greater nonadherence to endocrine therapy [85]. Opportunities exist to further quantify the intersection of race/ethnicity, rurality, and cancer survivorship, particularly among understudied American Indian/Alaska Native populations and the role that palliative care plays in cancer survivorship [175].

4.6. Cancer Mortality and Survival

In Table 2, we provide cancer mortality rates across racial/ethnic groups in both rural and urban areas.

Table 2. Rural and urban age-adjusted cancer mortality rates across racial/ethnic groups, 2013–2017.

Racial/Ethnic Group	Age-Adjusted Rural Mortality Rate per 100,000	Age-Adjusted Urban Mortality Rate per 100,000
All Cancers		
Non-Hispanic White	176.0 *	160.1
Non-Hispanic Black	199.1 *	184.4
American Indian/Alaska Native	161.2 *	120.8
Asian/Pacific Islander	104.7	98.4
Hispanic	109.1	111.0
Lung Cancer		
Non-Hispanic White	50.0 *	41.8
Non-Hispanic Black	49.2 *	42.6
American Indian/Alaska Native	40.0 *	29.8
Asian/Pacific Islander	24.6 *	22.0
Hispanic	48.5 *	38.4
Colorectal Cancer		
Non-Hispanic White	15.8 *	13.3
Non-Hispanic Black	22.0 *	18.5
American Indian/Alaska Native	18.6 *	11.8
Asian/Pacific Islander	9.9	9.5
Hispanic	16.0 *	13.4
Prostate Cancer		
Non-Hispanic White	18.6 *	17.8
Non-Hispanic Black	40.7 *	37.9
American Indian/Alaska Native	20.7 *	14.2
Asian/Pacific Islander	8.9	8.5
Hispanic	13.0	15.7 *
Female Breast Cancer		
Non-Hispanic White	20.5	20.4
Non-Hispanic Black	28.6	28.5
American Indian/Alaska Native	15.9	14.2
Asian/Pacific Islander	12.5	11.4
Hispanic	20.7	20.3
Cervical Cancer		
Non-Hispanic White	2.5 *	2.0
Non-Hispanic Black	4.6 *	3.4
American Indian/Alaska Native	3.4 *	1.9
Asian/Pacific Islander	1.8	1.8
Hispanic	2.7 *	2.2

Note: Rates are age-adjusted per the 2000 U.S. Standard Population and calculated from the CDC Wonder Dataset [176]. * indicates a statistically significant higher rate among a racial/ethnic group compared to their geographic counterpart.

Rural Black populations have the highest cancer mortality rate, and cancer mortality rates among rural White and Black populations exceed that of their urban counterparts.

Lung cancer mortality rates are higher among all rural racial/ethnic groups compared to urban. For colorectal and prostate cancer, rural Black populations have the highest mortality rate with rural White, Black, and American Indian/Alaska Native populations having higher mortality rates than their urban peers. Rural racial/ethnic disparities in cervical cancer mortality exist for all groups except Asian/Pacific Islander. Past studies have consistently shown that rural racial/ethnic minority patients have the poorest cancer-specific survival and mortality rates. Among all population groups, the rural Black population has the highest cancer mortality rate, but rural American Indian/Alaska Native populations have the largest disparity compared to their urban counterparts (a 33% higher rate) [41]. Studies using multiple categories of rurality have found that White, Black, and American Indian/Alaska Native populations have the highest cancer mortality burden [177,178]. Furthermore, despite overall improvement in cancer mortality trends, low-income Black patients in non-metropolitan areas experienced two to three times higher premature mortality risks than affluent Black and White patients in metropolitan areas. These associations were true for specific cancers as well. Studies have reported that living in rural areas increased the risk of mortality from lung cancer for Black patients overall by 54% and for Black women by 29% [179,180]. In colorectal cancer, Higginbotham et al. found that rural Black women had a 30% higher mortality rate than their urban counterparts, although Hines and colleagues found that the interaction term of Black race and rural residence was not statistically significant in models [180,181]. For breast cancer, one study found that in the urban setting of their study (Chicago communities), 17 out of 20 communities with the highest breast cancer mortality rates were predominantly Black, and that in the rural setting of their study (rural Washington state), Hispanic women had higher breast cancer mortality compared to non-Hispanic White women [182]. For cervical cancer, two studies found that Black women had higher mortality rates than White women regardless of geography, despite overall improvements among rural and urban women [139,183]. However, prostate cancer mortality rates in non-metropolitan areas compared to metropolitan areas were 12% higher in Black men and only 4% higher in White men [184]. Finally, another study provided results showing that urban American Indian and Alaska Native patients had a significantly higher risk of all-cancer mortality than urban White patients [185]. In children and adolescents in the state of Tennessee, Lindley and colleagues found that rural areas were more likely to be cancer mortality clusters and that the odds of living near a mortality cluster were nearly three times as high for Black children and adolescents (OR: 2.92, 95% CI: 1.13, 7.54) as their White counterparts [186].

To a lesser extent, cancer survival has been explored as a cancer outcome at the intersection of rurality and race/ethnicity, although some studies have examined rurality and race/ethnicity independently [147,187]. Generally, rural Black and American Indian/Alaska Native populations had the poorest survival outcomes. Five-year survival rates for cervical cancer among Black women in rural areas were 10% lower than Black women in metropolitan areas and 20% lower than White women in metropolitan areas [139]. In children and adolescents, American Indians/Alaska Natives residing in non-metro areas had more than twice the increased risk of cancer death than White residents in metro areas [188]. An analysis of National Cancer Database data showed that rural Black head and neck cancer patients had the shortest median survival time (35.1 months compared to 67 months among White urban patients) [189]. In rural–urban stratified analyses, Black men had poorer colorectal cancer survival than their White counterparts, with the greatest disparity among rural men; rural Asian men also had poorer survival outcomes [190].

5. Implications/Recommendations

We examined the extant research on cancer disparities among rural and racial/ethnic minority populations, extending from the social determinants of health that affect cancer disparities across the continuum. We summarize key findings from observational studies in the literature that explored disparities in primary prevention activities through sur-

ivorship among rural racial/ethnic minority populations. Our review notes several key opportunities for further policy changes and new areas for observational and interventional research to improve these identified disparities. Policy changes are critical to ensure that physical, social, and healthcare environments are optimized for health equity, particularly for those rural racial/ethnic minority populations that experience that greatest cancer burden: Black and American Indian/Alaska Native populations. Although we summarize many studies that have characterized the disparities at the intersection of rurality and race/ethnicity, continued research is needed to further elucidate and determine the appropriate actions to address these disparities. Furthermore, more studies are needed among American Indian/Alaska Native populations to address the particular burden among those individuals, and with the changes in availability and updated USPSTF recommendations, more research is needed to address disparities in lung and colorectal cancer among Black populations. Finally, there is a need for evidence-based interventions to be tested, implemented, and adopted in a culturally competent manner to improve cancer prevention and control activities at the community and clinic levels in rural settings.

To improve the social contextual and healthcare factors that affect cancer disparities across the continuum for rural racial/ethnic minority populations, it is important to develop and implement policies—at all levels from the federal government to local healthcare systems—that not only reduce disparities and biases but also promote equity. Our review notes that centuries of discriminatory policies (e.g., Jim Crow, redlining) have long-lasting effects on cancer outcomes that disproportionately affect racial/ethnic minority groups and that provider biases may affect screening and treatment disparities. It is critical to continue to examine the effects of these policies and biases long-term but also to implement policies that promote equity, including ensuring diversity in the healthcare workforce, enhancing community partnerships, and addressing institutional racism, as supported by the American Society of Clinical Oncology [191]. Diversity in the rural cancer care workforce, in particular, may be an effective means of addressing these disparities. Historically Black Colleges and Universities and student pipeline programs can play a key role in educating Black students and future healthcare professionals (e.g., Morehouse College's IMHOTEP internship program) [192]. In August 2020, Oklahoma State University and the Cherokee Nation opened an osteopathic medical school on Cherokee tribal lands with a goal of training future physicians, particularly those who are American Indian, to later practice in rural areas [193]. Improving representation among racial/ethnic minorities may help reduce provider biases and improve cancer care.

It is also important to consider the role of region in modifying the relationships observed between rurality, race/ethnicity, and cancer outcomes. The stark differences in regions' demographic and sociopolitical environments and physical access to healthcare-related services necessitates that researchers aim to determine whether observed rural racial/ethnic disparities hold uniformly across the U.S. or exist only in particular regions/localities, thereby requiring a different policy approach. Many states with high rural racial/ethnic minority populations in the South have not expanded access to Medicaid, which would afford many low-income Black and Hispanic populations greater access to primary and oncology care and has been shown to improve the use of cancer prevention services and cancer outcomes [194–196]. Much of the existing research on rural cancer disparities has focused on Appalachia, which experiences significant cancer burden; however, the Mississippi Delta and Black Belt regions of the country experience an even greater cancer mortality burden, as well as lower rates of cancer-prevention behaviors (e.g., HPV vaccination) and reduced access to care [117,197–199]. Much like Appalachia, the Delta Region (i.e., the Delta Regional Authority) is a federally designated region for socioeconomic development and receives federal funding for economic development and some health programming as well, as it is a designated area for loan repayment programs and visa waivers to increase access to healthcare providers. However, it is under-resourced, with the fiscal year 2020 budget allocating 45 cents for every dollar per capita directly to the Delta Regional Authority compared to the Appalachian Regional Commission (not in-

cluding regionally targeted funding from other federal agencies) [200,201]. More resources need to be targeted to this region, which has a high proportion of rural Black residents who experience less access to care and greater cancer burden.

Although a growing number of studies are exploring urban–rural disparities in health behaviors and outcomes across the cancer continuum and adding to the larger existing racial/ethnic disparities literature, fewer studies have focused on the intersection of race/ethnicity and rurality. To overcome this gap, there is first a collective need for increased awareness of the compounding effects of racial/ethnic minority status and rural residence in the U.S. Second, the effect of data collection and privacy rules on researchers' ability to study this important issue must be considered. Many population-based datasets have placed restrictions or moratoriums on accessing geographic-specific identifiers, including urban–rural status; however, in recent years, widely used datasets such as the National Health Interview Survey and BRFSS have begun incorporating such variables into their public-use files with minimal missingness. Unfortunately, many datasets are only available to access at federal Research Data Centers (RDCs), with costly data application/analysis fees, and some data remain masked because of privacy concerns (e.g., Medical Expenditures Panel Survey, Medicare data) [202,203]. Finally, because of administrative constraints and sampling approaches used in many population-based surveys, it is difficult to reach the sample sizes necessary to look at combinations of social strata (i.e., racial/ethnicity by rurality). Addressing this issue will likely require additional funding and/or oversampling to ensure both representative samples and large enough subgroup samples are obtained to make comparisons. The progress of observational research at the intersection of race/ethnicity and rurality hinges on dealing with these challenges.

As shown from our narrative review above, more research is needed on our smallest population groups, particularly rural American Indian/Alaska Natives, who experience a high cancer burden. As a result of small sample sizes, restrictive access to Indian Health Services data files, and insufficient payor categories (e.g., public vs. private), researchers know very little about whether American Indian/Alaska Native populations are receiving timely and high-quality care across the cancer continuum. Similarly, the challenge of small rural sample sizes in population-based surveys and other national data is exacerbated when rural American Indian/Alaska Natives are considered [204]. This is a particularly critical issue, as multiple studies have shown that American Indian/Alaska Native populations have the farthest travel burden to many cancer specialists and NCI-designated cancer centers and have the greatest rural–urban cancer mortality of any racial/ethnic group in the U.S. [41,60,61].

Furthermore, as new screening, diagnostic, treatment approaches emerge, it is important to understand how access to and utilization of these services vary based on rurality and race/ethnicity. For example, low-dose CT screening for lung cancer was initially recommended by the USPSTF in 2013, but critics were concerned that the high pack-year history requirements meant that fewer Black smokers may be eligible for screening [205,206]. Subsequently, the USPSTF released a draft statement that, if implemented, would lower the screening age and pack-year eligibility requirement [207]. Thus, this change may enable more rural Black residents, who experience a notable lung cancer burden, to be eligible for screening [140]. Although early national, population-based analyses of lung cancer screening utilization have not identified rural and/or racial disparities in screening, this is an important area for future research [132,134]. As our review notes, some research has examined disparities in the receipt and quality of cancer surgery, chemotherapy, and radiation treatments for cancer. In recent years, costly treatments such as immunotherapies and targeted therapies have been increasingly prescribed and are thus an important area for disparities research to ensure that innovations increase equity rather than intensify disparities [208–210].

Recent reports by the U.S. Department of Health and Human Services have outlined the critical need to focus research activities in rural communities to address disparities in the form of a rural health action plan and the need for social and behavioral intervention

research [211]. The historical scarcity of federally funded cancer research in rural settings has led to an increased focus on cancer prevention in these settings, especially in light of documented disparities in cancer outcomes [212–215]. There has been a limited focus on cancer prevention and control intervention research in rural settings [216]. When examining NCI's Evidence-Based Cancer Control Programs, only 40 (~25%) of 202 evidence-based interventions reported the community type as including rural with other settings, such as rural with suburban and urban/inner-city settings [217]. Of those 40, only 13 focused exclusively on rural communities. It is unclear how many of these evidence-based interventions in rural settings encompassed a focus on the intersection of rural life with myriad other factors. There is a need for clear guidance on developing interventions to accommodate the multiple and simultaneous forms of identity in rural communities. One without the other results in insufficient context to inform adaptation from one rural setting to the next and likely ineffective interventions on cancer-related outcomes.

Moreover, existing reporting guidelines aimed at promoting consistency in intervention reporting and replication and translating evidence into practice could promote more detailed geographic descriptions in their requirements to better understand rural context and intersectional identity among participants. Across existing frameworks and applications, the need to address intersectional identity matters. The AIMD framework considers the Aims, Ingredients, Mechanism, and Delivery of an intervention, which could provide context about the rural setting, rural-residing populations, and logistical and practical considerations of working in rural communities [218]. Similarly, the Standards for Reporting Implementation Studies (StaRI) checklist includes the opportunity to describe the context and targeted sites of an intervention and any contextual changes that may have affected the intervention's outcomes [219]. The Template for Intervention Description and Replication (TIDieR) also includes a focus on where the intervention occurred, including infrastructure or other relevant features [220]. Lastly, the equity extension to Consolidated Standards of Reporting Trials (CONSORT) calls for reporting who the study participants are, plus their relationship with their setting, given the interconnectedness of person and place, as well as the context and relationship of the study settings and locations to health inequity and generalizability [221]. The CONSORT equity extension demonstrates a promising approach to fully describing and understanding the importance of intersectional identity, especially in rural settings.

6. Conclusions

We provide a comprehensive review of the geographic distribution of rural racial/ethnic populations, provide a framework for and describe how the social determinants of health may disproportionately increase the risk of cancer disparities in these populations across the continuum, and detail a summary of the disparities these populations experience from cancer-relevant preventive behaviors to mortality. We identified that rural Black and American Indian/Alaska Native populations often experience the greatest cancer burden. As such, policies, as well as observational, interventional, and implementation research, must address the intersection of rurality and race/ethnicity to ensure that disparities are reduced, and equity is promoted.

Author Contributions: Conceptualization, W.E.Z.; methodology, W.E.Z.; formal analysis, W.E.Z.; data curation, W.E.Z.; writing—original draft preparation, W.E.Z., C.M., M.K., G.A.B., K.R.D., R.R., P.L., A.Z., K.S., M.A.M., E.L.C., H.M.B., J.M.E.; writing—review and editing, W.E.Z., C.M., M.K., G.A.B., K.R.D., R.R., P.L., A.Z., K.S., M.A.M., E.L.C., H.M.B., J.M.E.; visualization, W.E.Z.; supervision, J.M.E.; project administration, W.E.Z. and J.M.E.; funding acquisition, J.M.E., E.L.C., G.A.B., K.R.D., H.M.B. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by the Federal Office of Rural Health Policy (Health Resources and Services Administration, US Department of Health and Human Services) under cooperative agreement U1CRH30539 (W.E.Z., E.L.C., H.M.B., J.M.E.). The information, conclusions, and opinions expressed in this article are those of the authors, and no endorsement by the Federal Office of Rural Health Policy, the Health Resources and Services Administration, or the US Department of Health

and Human Services is intended or should be inferred. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. This publication was made possible in part by Grant Number T32-GM081740 from National Institutes of Health-National Institute of General Medical Sciences (NIH-NIGMS) (G.A.B. and K.R.D.). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIGMS or NIH. Research reported in this publication was supported by the National Cancer Institute of the National Institutes of Health under Award Number P30CA021765 and the American Lebanese and Syrian Associated Charities (ALSAC) of St. Jude Children’s Research Hospital (H.M.B.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Publicly available datasets were analyzed in this study. These data sources are cited in the footnotes below Tables 1 and 2 and in a corresponding position within the references.

Acknowledgments: The authors wish to thank Erica Sercy, MSPH, for her assistance with editing and formatting.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

- Bennett, K.J.; Borders, T.F.; Holmes, G.M.; Kozhimannil, K.B.; Ziller, E. What Is Rural? Challenges And Implications Of Definitions That Inadequately Encompass Rural People And Places. *Health Aff. (Proj. Hope)* **2019**, *38*, 1985–1992. [CrossRef] [PubMed]
- Doogan, N.J.; Roberts, M.E.; Wewers, M.E.; Stanton, C.A.; Keith, D.R.; Gaalema, D.E.; Kurti, A.N.; Redner, R.; Cepeda-Benito, A.; Bunn, J.Y.; et al. A growing geographic disparity: Rural and urban cigarette smoking trends in the United States. *Prev. Med.* **2017**, *104*, 79–85. [CrossRef] [PubMed]
- Matthews, K.A.; Croft, J.B.; Liu, Y.; Lu, H.; Kanny, D.; Wheaton, A.G.; Cunningham, T.J.; Khan, L.K.; Caraballo, R.S.; Holt, J.B.; et al. Health-Related Behaviors by Urban-Rural County Classification—United States, 2013. *Mmwr. Surveill Summ.* **2017**, *66*, 1–8. [CrossRef] [PubMed]
- Williams, C.L.; Walker, T.Y.; Elam-Evans, L.D.; Yankey, D.; Fredua, B.; Saraiya, M.; Stokley, S. Factors associated with not receiving HPV vaccine among adolescents by metropolitan statistical area status, United States, National Immunization Survey–Teen, 2016–2017. *Hum. Vaccines Immunother.* **2020**, *16*, 562–572. [CrossRef]
- Zahnd, W.E.; James, A.S.; Jenkins, W.D.; Izadi, S.R.; Fogleman, A.J.; Steward, D.E.; Colditz, G.A.; Brard, L. Rural-Urban Differences in Cancer Incidence and Trends in the United States. *Cancer Epidemiol. Biomark. Prev.* **2018**, *27*, 1265–1274. [CrossRef]
- Henley, S.J.; Anderson, R.N.; Thomas, C.C.; Massetti, G.M.; Peaker, B.; Richardson, L.C. Invasive Cancer Incidence, 2004–2013, and Deaths, 2006–2015, in Nonmetropolitan and Metropolitan Counties—United States. *Mmwr. Surveill Summ.* **2017**, *66*, 1–13. [CrossRef]
- Cole, A.M.; Jackson, J.E.; Doescher, M. Colorectal cancer screening disparities for rural minorities in the United States. *J. Prim. Care Community Health* **2013**, *4*, 106–111. [CrossRef]
- Atkins, G.T.; Kim, T.; Munson, J. Residence in Rural Areas of the United States and Lung Cancer Mortality. Disease Incidence, Treatment Disparities, and Stage-Specific Survival. *Ann. Am. Thorac. Soc.* **2017**, *14*, 403–411. [CrossRef]
- Alvidrez, J.; Castille, D.; Laude-Sharp, M.; Rosario, A.; Tabor, D. The National Institute on Minority Health and Health Disparities Research Framework. *Am. J. Public Health* **2019**, *109*, S16–S20. [CrossRef]
- Revisions to the Standards for the Classification of Federal Data on Race and Ethnicity | The White House. Available online: https://obamawhitehouse.archives.gov/omb/fedreg_1997standards (accessed on 27 September 2020).
- Jones, D.S. The persistence of American Indian health disparities. *Am. J. Public Health* **2006**, *96*, 2122–2134. [CrossRef]
- Philbin, M.M.; Flake, M.; Hatzenbuehler, M.L.; Hirsch, J.S. State-level immigration and immigrant-focused policies as drivers of Latino health disparities in the United States. *Soc. Sci. Med.* **2018**, *199*, 29–38. [CrossRef] [PubMed]
- Findling, M.G.; Bleich, S.N.; Casey, L.S.; Blendon, R.J.; Benson, J.M.; Sayde, J.M.; Miller, C. Discrimination in the United States: Experiences of Latinos. *Health Serv. Res.* **2019**, *54*, 1409–1418. [CrossRef] [PubMed]
- Lichter, D.T.; Johnson, K.M. A Demographic Lifeline? Immigration and Hispanic Population Growth in Rural America. *Popul. Res. Policy Rev.* **2020**, *39*, 785–803. [CrossRef]
- Gee, G.C.; Ro, A.; Shariff-Marco, S.; Chae, D. Racial discrimination and health among asian americans: Evidence, assessment, and directions for future research. *Epidemiol. Rev.* **2009**, *31*, 130–151. [CrossRef]

16. McMurtry, C.L.; Findling, M.G.; Casey, L.S.; Blendon, R.J.; Benson, J.M.; Sayde, J.M.; Miller, C. Discrimination in the United States: Experiences of Asian Americans. *Health Serv. Res.* **2019**, *54*, 1419–1430. [[CrossRef](#)]
17. Caldwell, J.T.; Ford, C.L.; Wallace, S.P.; Wang, M.C.; Takahashi, L.M. Racial and ethnic residential segregation and access to health care in rural areas. *Health Place* **2017**, *43*, 104–112. [[CrossRef](#)]
18. Shariff-Marco, S.; Klassen, A.C.; Bowie, J.V. Racial/ethnic differences in self-reported racism and its association with cancer-related health behaviors. *Am. J. Public Health* **2010**, *100*, 364–374. [[CrossRef](#)]
19. Morello-Frosch, R.; Jesdale, B.M. Separate and unequal: Residential segregation and estimated cancer risks associated with ambient air toxins in U.S. metropolitan areas. *Environ. Health Perspect.* **2006**, *114*, 386–393. [[CrossRef](#)]
20. Fang, P.; He, W.; Gomez, D.; Hoffman, K.E.; Smith, B.D.; Giordano, S.H.; Jagsi, R.; Smith, G.L. Racial disparities in guideline-concordant cancer care and mortality in the United States. *Adv. Radiat Oncol.* **2018**, *3*, 221–229. [[CrossRef](#)]
21. Landrine, H.; Corral, I.; Lee, J.G.L.; Eford, J.; Hall, M.B.; Bess, J.J. Residential Segregation and Racial Cancer Disparities: A Systematic Review. *J. Racial Ethn. Health Disparities* **2017**, *4*, 1195–1205. [[CrossRef](#)]
22. Pruitt, S.L.; Lee, S.J.; Tiro, J.A.; Xuan, L.; Ruiz, J.M.; Inrig, S. Residential racial segregation and mortality among black, white, and Hispanic urban breast cancer patients in Texas, 1995 to 2009. *Cancer* **2015**, *121*, 1845–1855. [[CrossRef](#)] [[PubMed](#)]
23. Warner, E.T.; Gomez, S.L. Impact of neighborhood racial composition and metropolitan residential segregation on disparities in breast cancer stage at diagnosis and survival between black and white women in California. *J. Community Health* **2010**, *35*, 398–408. [[CrossRef](#)] [[PubMed](#)]
24. Williams, D.R.; Kontos, E.Z.; Viswanath, K.; Haas, J.S.; Lathan, C.S.; MacConaill, L.E.; Chen, J.; Ayanian, J.Z. Integrating Multiple Social Statuses in Health Disparities Research: The Case of Lung Cancer. *Health Serv. Res.* **2012**, *47*, 1255–1277. [[CrossRef](#)] [[PubMed](#)]
25. Zavala, V.A.; Bracci, P.M.; Carethers, J.M.; Carvajal-Carmona, L.; Coggins, N.B.; Cruz-Correa, M.R.; Davis, M.; de Smith, A.J.; Dutil, J.; Figueiredo, J.C.; et al. Cancer health disparities in racial/ethnic minorities in the United States. *Br. J. Cancer* **2020**, *124*, 315–332. [[CrossRef](#)] [[PubMed](#)]
26. Caldwell, J.T.; Ford, C.L.; Wallace, S.P.; Wang, M.C.; Takahashi, L.M. Intersection of Living in a Rural Versus Urban Area and Race/Ethnicity in Explaining Access to Health Care in the United States. *Am. J. Public Health* **2016**, *106*, 1463–1469. [[CrossRef](#)]
27. Green, M.A.; Evans, C.R.; Subramanian, S.V. Can intersectionality theory enrich population health research? *Soc. Sci. Med.* **2017**, *178*, 214–216. [[CrossRef](#)]
28. Pender, J.; Hertz, T.; Cromartie, J.; Farrigan, T. Rural America at a Glance, 2019 Edition. Economic Information Bulletin No. (EIB-212). Economic Research Service, United States Department of Agriculture. Available online: <https://www.ers.usda.gov/publications/pub-details/?pubid=95340> (accessed on 9 December 2020).
29. Lichter, D.T. Immigration and the New Racial Diversity in Rural America. *Rural Sociol.* **2012**, *77*, 3–35. [[CrossRef](#)]
30. USDA ERS—Rural Classifications. Available online: <https://www.ers.usda.gov/topics/rural-economy-population/rural-classifications/> (accessed on 9 December 2020).
31. U.S. Census Bureau. Population Estimates Program. Available online: <https://www2.census.gov/programs-surveys/popest/datasets/2010-2018/> (accessed on 9 December 2020).
32. Foutz, J.; Artiga, S.; Garfield, R. The Role of Medicaid in Rural America. 25 April 2017. Available online: <https://www.kff.org/medicaid/issue-brief/the-role-of-medicare-in-rural-america/> (accessed on 9 December 2020).
33. Hendryx, M.; Luo, J. Increased Cancer Screening for Low-income Adults Under the Affordable Care Act Medicaid Expansion. *Med. Care* **2018**, *56*, 944–949. [[CrossRef](#)]
34. Sineshaw, H.M.; Ellis, M.A.; Yabroff, K.R.; Han, X.; Jemal, A.; Day, T.A.; Graboyes, E.M. Association of Medicaid Expansion Under the Affordable Care Act With Stage at Diagnosis and Time to Treatment Initiation for Patients With Head and Neck Squamous Cell Carcinoma. *JAMA Otolaryngol. Head Neck Surg.* **2020**, *146*, 247–255. [[CrossRef](#)]
35. Gee, G.C.; Ford, C.L. Structural racism and health inequities: Old Issues, New Directions. *Du Bois Rev.* **2011**, *8*, 115–132. [[CrossRef](#)]
36. Williams, D.R.; Lawrence, J.A.; Davis, B.A. Racism and Health: Evidence and Needed Research. *Annu. Rev. Public Health* **2019**, *40*, 105–125. [[CrossRef](#)] [[PubMed](#)]
37. Zahnd, W.E.; McLafferty, S.L.; Eberth, J.M. Multilevel analysis in rural cancer control: A conceptual framework and methodological implications. *Prev. Med.* **2019**, *129*, 105835. [[CrossRef](#)] [[PubMed](#)]
38. Wingo, P.A.; Howe, H.L.; Thun, M.J.; Ballard-Barbash, R.; Ward, E.; Brown, M.L.; Sylvester, J.; Friedell, G.H.; Alley, L.; Rowland, J.H.; et al. A national framework for cancer surveillance in the United States. *Cancer Causes Control.* **2005**, *16*, 151–170. [[CrossRef](#)] [[PubMed](#)]
39. Probst, J.C.; Bellinger, J.D.; Walsemann, K.M.; Hardin, J.; Glover, S.H. Higher risk of death in rural blacks and whites than urbanites is related to lower incomes, education, and health coverage. *Health Aff.* **2011**, *30*, 1872–1879. [[CrossRef](#)]
40. Probst, J.C.; Moore, C.G.; Glover, S.H.; Samuels, M.E. Person and place: The compounding effects of race/ethnicity and rurality on health. *Am. J. Public Health* **2004**, *94*, 1695–1703. [[CrossRef](#)]
41. Probst, J.C.; Zahnd, W.E.; Hung, P.; Eberth, J.M.; Crouch, E.L.; Merrell, M.A. Rural-Urban Mortality Disparities: Variations Across Causes of Death and Race/Ethnicity, 2013–2017. *Am. J. Public Health* **2020**, *110*, 1325–1327. [[CrossRef](#)]
42. Feagin, J.; Bennefield, Z. Systemic racism and U.S. health care. *Soc. Sci. Med.* **2014**, *103*, 7–14. [[CrossRef](#)]
43. Williams, D.R.; Neighbors, H.W.; Jackson, J.S. Racial/ethnic discrimination and health: Findings from community studies. *Am. J. Public Health* **2003**, *93*, 200–208. [[CrossRef](#)]

44. Acevedo-Garcia, D.L.K. Residential Segregation and Health. In *Neighborhoods and Health*; Kawachi, I., Berkman, L.F., Eds.; Oxford University Press: New York, NY, USA, 2003.
45. Acevedo-Garcia, D.; Lochner, K.A.; Osypuk, T.L.; Subramanian, S.V. Future directions in residential segregation and health research: A multilevel approach. *Am. J. Public Health* **2003**, *93*, 215–221. [[CrossRef](#)]
46. Krieger, N.; Jahn, J.L.; Waterman, P.D.; Chen, J.T. Breast Cancer Estrogen Receptor Status According to Biological Generation: US Black and White Women Born 1915–1979. *Am. J. Epidemiol.* **2018**, *187*, 960–970. [[CrossRef](#)]
47. Krieger, N.; Wright, E.; Chen, J.T.; Waterman, P.D.; Huntley, E.R.; Arcaya, M. Cancer Stage at Diagnosis, Historical Redlining, and Current Neighborhood Characteristics: Breast, Cervical, Lung, and Colorectal Cancers, Massachusetts, 2001–2015. *Am. J. Epidemiol.* **2020**, *189*, 1065–1075. [[CrossRef](#)] [[PubMed](#)]
48. Massey, D.S.; Denton, N.A. Hypersegregation in U.S. Metropolitan Areas: Black and Hispanic Segregation along Five Dimensions. *Demography* **1989**, *26*, 373–391. [[CrossRef](#)] [[PubMed](#)]
49. Scott, A.; Wilson, R.F. Social determinants of health among African Americans in a rural community in the Deep South: An ecological exploration. *Rural Remote Health* **2011**, *11*, 196. [[CrossRef](#)]
50. Menon, N.M.; Leslie, T.F.; Frankenfeld, C.L. Cancer-related diagnostic and treatment capabilities of hospitals in the context of racial residential segregation. *Public Health* **2020**, *182*, 95–101. [[CrossRef](#)] [[PubMed](#)]
51. Haas, J.S.; Earle, C.C.; Orav, J.E.; Brawarsky, P.; Neville, B.A.; Williams, D.R. Racial segregation and disparities in cancer stage for seniors. *J. Gen. Intern. Med.* **2008**, *23*, 699–705. [[CrossRef](#)] [[PubMed](#)]
52. Hao, Y.; Landrine, H.; Smith, T.; Kaw, C.; Corral, I.; Stein, K. Residential Segregation and Disparities in Health-Related Quality of Life Among Black and White Cancer Survivors. *Health Psychol.* **2011**, *30*, 137–144. [[CrossRef](#)]
53. Hayanga, A.J.; Zeliadt, S.B.; Backhus, L.M. Residential segregation and lung cancer mortality in the United States. *Arch. Surg.* **2013**, *148*, 37–42. [[CrossRef](#)]
54. Moss, J.L.; Ehrenkranz, R.; Perez, L.G.; Hair, B.Y.; Julian, A.K. Geographic disparities in cancer screening and fatalism among a nationally representative sample of US adults. *J. Epidemiol. Community Health* **2019**, *73*, 1128–1135. [[CrossRef](#)]
55. James, C.V.; Moonesinghe, R.; Wilson-Frederick, S.M.; Hall, J.E.; Penman-Aguilar, A.; Bouye, K. Racial/ethnic health disparities among rural adults—United States, 2012–2015. *Mmwr. Surveill. Summ.* **2017**, *66*, 1–9. [[CrossRef](#)]
56. Penchansky, R.; Thomas, J.W. The concept of access: Definition and relationship to consumer satisfaction. *Med. Care* **1981**, *19*, 127–140. [[CrossRef](#)]
57. Largent, E.A. Public Health, Racism, and the Lasting Impact of Hospital Segregation. *Public Health Rep.* **2018**, *133*, 715–720. [[CrossRef](#)] [[PubMed](#)]
58. Government Accountability Office. Rural Hospitals: Federal Efforts Should Target Areas Where Closures Would Threaten Access to Care. Available online: <https://www.gao.gov/assets/160/150195.pdf> (accessed on 29 January 2021).
59. Thomas, S.R.; Holmes, G.M.; Pink, G.H. To What Extent do Community Characteristics Explain Differences in Closure among Financially Distressed Rural Hospitals? *J. Health Care Poor Underserved* **2016**, *27*, 194–203. [[CrossRef](#)] [[PubMed](#)]
60. Hung, P.; Deng, S.; Zahnd, W.E.; Adams, S.A.; Olatosi, B.; Crouch, E.L.; Eberth, J.M. Geographic disparities in residential proximity to colorectal and cervical cancer care providers. *Cancer* **2019**, *126*, 1068–1076. [[CrossRef](#)] [[PubMed](#)]
61. Onega, T.; Alford-Teaster, J.; Wang, F. Population-based geographic access to parent and satellite National Cancer Institute Cancer Center Facilities. *Cancer* **2017**, *123*, 3305–3311. [[CrossRef](#)] [[PubMed](#)]
62. Grogan, C.M.; Park, S.E. The racial divide in state medicaid expansions. *J. Health Politics Policy Law* **2017**, *42*, 539–572. [[CrossRef](#)] [[PubMed](#)]
63. Hastert, T.A.; Banegas, M.P.; Hamel, L.M.; Reed, A.R.; Baird, T.; Beebe-Dimmer, J.L.; Schwartz, A.G. Race, financial hardship, and limiting care due to cost in a diverse cohort of cancer survivors. *J. Cancer Surviv.* **2019**, *13*, 429–437. [[CrossRef](#)]
64. Prather, C.; Fuller, T.R.; Jeffries, W.L.; Marshall, K.J.; Howell, A.V.; Belyue-Umole, A.; King, W. Racism, African American Women, and Their Sexual and Reproductive Health: A Review of Historical and Contemporary Evidence and Implications for Health Equity. *Health Equity* **2018**, *2*, 249–259. [[CrossRef](#)] [[PubMed](#)]
65. Jacobs, E.A.; Mendenhall, E.; McAlearney, A.S.; Rolle, I.; Whitaker, E.E.; Warnecke, R.; Ferrans, C.E. An exploratory study of how trust in health care institutions varies across African American, Hispanic and white populations. *Commun. Med.* **2011**, *8*, 89–98. [[CrossRef](#)]
66. Hoffman, K.M.; Trawalter, S.; Axt, J.R.; Oliver, M.N. Racial bias in pain assessment and treatment recommendations, and false beliefs about biological differences between blacks and whites. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, 4296–4301. [[CrossRef](#)]
67. Takeshita, J.; Wang, S.; Loren, A.W.; Mitra, N.; Shults, J.; Shin, D.B.; Sawinski, D.L. Association of Racial/Ethnic and Gender Concordance Between Patients and Physicians With Patient Experience Ratings. *JAMA Netw. Open* **2020**, *3*, e2024583. [[CrossRef](#)]
68. American Association of Medical Colleges. Diversity in Medicine: Facts and Figures 2019. Available online: <https://www.aamc.org/data-reports/workforce/interactive-data/figure-13-percentage-us-medical-school-graduates-race-ethnicity-alone-academic-year-2018-2019> (accessed on 26 January 2021).
69. Butkus, R.; Rapp, K.; Cooney, T.G.; Engel, L.S. Envisioning a Better U.S. Health Care System for All: Reducing Barriers to Care and Addressing Social Determinants of Health. *Ann. Intern. Med.* **2020**, *172*, S50–S59. [[CrossRef](#)] [[PubMed](#)]
70. Labor Force Characteristics by Race and Ethnicity, 2018: BLS Reports: U.S. Bureau of Labor Statistics. Available online: <https://www.bls.gov/opub/reports/race-and-ethnicity/2018/home.htm> (accessed on 28 November 2020).

71. Charlton, M.; Schlichting, J.; Chioreso, C.; Ward, M.; Vikas, P. Challenges of Rural Cancer Care in the United States. *Oncol. (Williston Park)* **2015**, *29*, 633–640.
72. Probst, J.; Eberth, J.M.; Crouch, E. Structural Urbanism Contributes to Poorer Health Outcomes For Rural America. *Health Aff. (Proj. Hope)* **2019**, *38*, 1976–1984. [[CrossRef](#)] [[PubMed](#)]
73. Singh, G.K.; Daus, G.P.; Allender, M.; Ramey, C.T.; Martin, E.K.; Perry, C.; Reyes, A.A.L.; Vedamuthu, I.P. Social Determinants of Health in the United States: Addressing Major Health Inequality Trends for the Nation, 1935–2016. *Int. J. Mch. Aids* **2017**, *6*, 139–164. [[CrossRef](#)] [[PubMed](#)]
74. Hicken, M.T.; Burnside, L.; Edwards, D.L.; Lee, H. Black–White Health Inequalities by Intentional Design: The Lasting Health Impact of Racial Residential Segregation. In *Racism: Science & Tools for the Public Health Professional*; Ford, C., Ed.; APHA Press: Washington, DC, USA, 2019; pp. 117–132.
75. Parisi, D.; Grice, S.M.; Taquino, M.; Gill, D.A. Community Concentration of Poverty and Its Consequences on Nonmetro persistence of poverty in Mississippi. *Sociol. Spectrum*. **2005**, *5*, 469–483. [[CrossRef](#)]
76. Quillian, L. Does Segregation Create Winners and Losers? Residential Segregation and Inequality in Educational Attainment. *Soc. Probl.* **2014**, *61*, 402–426. [[CrossRef](#)]
77. Probst, J.C.; Ajmal, F. Social Determinants of Health among Rural Asian and Pacific Islander Populations. Policy Brief. July 2019. Available online: <https://www.ruralhealthresearch.org/publications/1271> (accessed on 9 December 2020).
78. Probst, J.C.; Ajmal, F. Social Determinants of Health among the Rural Hispanic Population. Policy Brief. July 2019. Available online: <https://www.ruralhealthresearch.org/publications/1269> (accessed on 9 December 2020).
79. Probst, J.C.; Ajmal, F. Social Determinants of Health among the Rural African American population. Policy Brief. July 2019. Available online: <https://www.ruralhealthresearch.org/publications/1268> (accessed on 9 December 2020).
80. Probst, J.C.; Ajmal, F. Social Determinants of Health among Rural American Indian and Alaska Native Populations. Policy Brief. July 2019. Available online: <https://www.ruralhealthresearch.org/publications/1270> (accessed on 9 December 2020).
81. Greenberg-Worisek, A.J.; Kurani, S.; Finney Rutten, L.J.; Blake, K.D.; Moser, R.P.; Hesse, B.W. Tracking healthy people 2020 Internet, broadband, and mobile device access goals: An update using data from the health information national trends survey. *J. Med. Internet Res.* **2019**, *21*, e13300. [[CrossRef](#)]
82. Beydoun, M.A.; Beydoun, H.A.; Mode, N.; Dore, G.A.; Canas, J.A.; Eid, S.M.; Zonderman, A.B. Racial disparities in adult all-cause and cause-specific mortality among us adults: Mediating and moderating factors. *BMC Public Health* **2016**, *16*, 1113. [[CrossRef](#)]
83. Fairfield, K.M.; Black, A.W.; Lucas, F.L.; Murray, K.; Ziller, E.; Korsen, N.; Waterston, L.B.; Han, P.K.J. Association between Rurality and Lung Cancer Treatment Characteristics and Timeliness. *J. Rural Health* **2019**, *35*, 560–565. [[CrossRef](#)]
84. Boscoe, F.P.; Henry, K.A.; Sherman, R.L.; Johnson, C.J. The relationship between cancer incidence, stage, and poverty in the United States. *Int. J. Cancer* **2016**, *139*, 607–612. [[CrossRef](#)]
85. Heiney, S.P.; Truman, S.; Babatunde, O.A.; Felder, T.M.; Eberth, J.M.; Crouch, E.; Wickersham, K.E.; Adams, S.A. Racial and Geographic Disparities in Endocrine Therapy Adherence among Younger Breast Cancer Survivors. *Am. J. Clin. Oncol.* **2020**, *43*, 504–509. [[CrossRef](#)] [[PubMed](#)]
86. Desmond, R.A.; Jackson, B.E.; Waterbor, J.W. Disparities in Cancer Survivorship Indicators in the Deep South Based on BRFSS Data: Recommendations for Survivorship Care Plans. *South. Med. J.* **2017**, *110*, 181–187. [[CrossRef](#)] [[PubMed](#)]
87. Strosnider, H.; Kennedy, C.; Monti, M.; Yip, F. Rural and Urban Differences in Air Quality, 2008–2012, and Community Drinking Water Quality, 2010–2015—United States. *Mmwr. Surveill. Summ.* **2017**, *66*, 1–10. [[CrossRef](#)] [[PubMed](#)]
88. Zahnd, W.E.; Mueller-Luckey, G.S.; Ratnapradipa, K.; Smith, T. Predictors and Spatial Variation of Radon Testing in Illinois, 2005–2012. *J. Public Health Manag. Pr.* **2018**, *24*, e1–e9. [[CrossRef](#)] [[PubMed](#)]
89. Baris, D.; Waddell, R.; Beane Freeman, L.E.; Schwenn, M.; Colt, J.S.; Ayotte, J.D.; Ward, M.H.; Nuckols, J.; Schned, A.; Jackson, B.; et al. Elevated Bladder Cancer in Northern New England: The Role of Drinking Water and Arsenic. *J. Natl. Cancer Inst.* **2016**, *108*, djw099. [[CrossRef](#)]
90. National Institutes of Health. About | Agricultural Health Study. Available online: <https://aghealth.nih.gov/about/index.html> (accessed on 3 October 2020).
91. Tyson, F.L.; Midwife, K.C.; Gavin, J.; Gaylord, C.E.; Lee, C.; Setlow, V.P.; Wilson, S. Cancer, the environment, and environmental justice? *Cancer* **2000**, *83*, 1784–1792. [[CrossRef](#)]
92. James, W.; Jia, C.; Kedia, S. Uneven magnitude of disparities in cancer risks from air toxics. *Int. J. Environ. Res. Public Health* **2012**, *9*, 4365–4385. [[CrossRef](#)]
93. Gearhart-Serna, L.M.; Hoffman, K.; Devi, G.R. Environmental Quality and Invasive Breast Cancer. *Cancer Epidemiol. Biomark. Prev.* **2020**, *29*, 1920–1928. [[CrossRef](#)]
94. Madia, F.; Worth, A.; Whelan, M.; Corvi, R. Carcinogenicity assessment: Addressing the challenges of cancer and chemicals in the environment. *Environ. Int.* **2019**, *128*, 417–429. [[CrossRef](#)]
95. Patel, C.J.; Kerr, J.; Thomas, D.C.; Mukherjee, B.; Ritz, B.; Chatterjee, N.; Jankowska, M.; Madan, J.; Karagas, M.R.; Mcallister, K.A.; et al. Opportunities and challenges for environmental exposure assessment in population-based studies. *Cancer Epidemiol. Biomark. Prev.* **2017**, *26*, 1370–1380. [[CrossRef](#)]
96. Cunningham, J.K.; Ritchey, J.; Solomon, T.A.; Cordova, F.M. Cigarette Use Among American Indians and Alaska Natives in Metropolitan Areas, Rural Areas, and Tribal Lands. *J. Public Health Manag. Pract.* **2019**, *25*, S11–S19. [[CrossRef](#)] [[PubMed](#)]

97. Koller, K.R.; Flanagan, C.A.; Day, G.E.; Patten, C.; Umans, J.G.; Austin, M.A.; Hopkins, S.E.; Raindl, C.; Boyer, B.B. High tobacco use prevalence with significant regional and sex differences in smokeless tobacco use among western alaska native people: The WATCH study. *Int. J. Circumpolar Health* **2017**, *76*, 1398009. [CrossRef] [PubMed]
98. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. Health Reports of the Surgeon General. In *The Health Consequences of Smoking-50 Years of Progress: A Report of the Surgeon General*; Centers for Disease Control and Prevention (US): Atlanta, GA, USA, 2014.
99. Doescher, M.P.; Jackson, J.E.; Jerant, A.; Gary Hart, L. Prevalence and trends in smoking: A national rural study. *J. Rural Health* **2006**, *22*, 112–118. [CrossRef] [PubMed]
100. Chang, J.T.; Levy, D.T.; Meza, R. Trends and factors related to smokeless Tobacco use in the United States. *Nicotine Tob. Res.* **2016**, *18*, 1740–1748. [CrossRef] [PubMed]
101. Dille, J.A.; Peterson, E.; Bobo, M.; Pickle, K.E.; Rohde, K. Tobacco use prevalence—Disentangling associations between Alaska Native race, low socio-economic status and rural disparities. *Int. J. Circumpolar Health* **2013**, *72*, 21582. [CrossRef] [PubMed]
102. Bell, R.A.; Arcury, T.A.; Chen, H.; Anderson, A.M.; Savoca, M.R.; Kohrman, T.; Quandt, S.A. Use of tobacco products among rural older adults: Prevalence of ever use and cumulative lifetime use. *Addict. Behav.* **2009**, *34*, 662–667. [CrossRef]
103. Chaffee, B.W.; Couch, E.T.; Urata, J.; Gansky, S.A.; Essex, G.; Cheng, J. Predictors of Smokeless Tobacco Susceptibility, Initiation, and Progression Over Time Among Adolescents in a Rural Cohort. *Subst. Use Misuse* **2019**, *54*, 1154–1166. [CrossRef]
104. Boffetta, P.; Hashibe, M. Alcohol and cancer. *Lancet Oncol.* **2006**, *7*, 149–156. [CrossRef]
105. Dixon, M.A.; Chartier, K.G. Alcohol use patterns among urban and rural residents: Demographic and social influences. *Alcohol Res. Curr. Rev.* **2016**, *38*, 69–77.
106. Arcury, T.A.; Talton, J.W.; Summers, P.; Chen, H.; Laurienti, P.J.; Quandt, S.A. Alcohol Consumption and Risk for Dependence Among Male Latino Migrant Farmworkers Compared to Latino Nonfarmworkers in North Carolina. *Alcohol. Clin. Exp. Res.* **2016**, *40*, 377–384. [CrossRef]
107. Steck, S.E.; Murphy, E.A. Dietary patterns and cancer risk. *Nat. Rev. Cancer* **2020**, *20*, 125–138. [CrossRef] [PubMed]
108. Dutko, P.; Ver Ploeg, M.; Farrigan, T. Characteristics and Influential Factors of Food Deserts. United States Department of Agriculture, Economic Research Service, Economic Research Report Number 140; August 2012. Available online: https://www.ers.usda.gov/webdocs/publications/45014/30940_err140.pdf (accessed on 9 December 2020).
109. Quandt, S.A.; Groeschel-Johnson, A.; Kinzer, H.T.; Jensen, A.; Miles, K.; O'Hara, H.M.; Chen, H.; Arcury, T.A. Migrant Farmworker Nutritional Strategies: Implications for Diabetes Management. *J. Agromedicine* **2018**, *23*, 347–354. [CrossRef] [PubMed]
110. Sundberg, M.A.; Warren, A.C.; Vanwassenhove-Paetzold, J.; George, C.; Carroll, D.S.; Becenti, L.J.; Martinez, A.; Jones, B.; Bachman-Carter, K.; Begay, M.G.; et al. Implementation of the Navajo fruit and vegetable prescription programme to improve access to healthy foods in a rural food desert. *Public Health Nutr.* **2020**, *23*, 2199–2210. [CrossRef] [PubMed]
111. Savoca, M.R.; Arcury, T.A.; Leng, X.; Bell, R.A.; Chen, H.; Anderson, A.; Kohrman, T.; Quandt, S.A. The Diet Quality of Rural Older Adults in the South as Measured by Healthy Eating Index-2005 Varies by Ethnicity. *J. Am. Diet. Assoc.* **2009**, *109*, 2063–2067. [CrossRef] [PubMed]
112. Champagne, C.M.; Bogle, M.L.; McGee, B.B.; Yadrick, K.; Allen, H.R.; Kramer, T.R.; Simpson, P.; Gossett, J.; Weber, J. Dietary intake in the lower Mississippi delta region: Results from the foods of our delta study. *J. Am. Diet. Assoc.* **2004**, *104*, 199–207. [CrossRef] [PubMed]
113. Befort, C.A.; Nazir, N.; Perri, M.G. Prevalence of obesity among adults from rural and urban areas of the United States: Findings from NHANES (2005–2008). *J. Rural Health* **2012**, *28*, 392–397. [CrossRef] [PubMed]
114. Wen, M.; Zhang, X.; Harris, C.D.; Holt, J.B.; Croft, J.B. Spatial Disparities in the Distribution of Parks and Green Spaces in the USA. *Ann. Behav. Med.* **2013**, *45*, 18–27. [CrossRef]
115. Kaczynski, A.T.; Eberth, J.M.; Stowe, E.W.; Wende, M.E.; Liese, A.D.; McLain, A.C.; Breneman, C.B.; Josey, M.J. Development of a national childhood obesogenic environment index in the United States: Differences by region and rurality. *Int. J. Behav. Nutr. Phys. Act.* **2020**, *17*, 83. [CrossRef]
116. Chaturvedi, A.K.; Graubard, B.I.; Broutian, T.; Pickard, R.K.L.; Tong, Z.Y.; Xiao, W.; Kahle, L.; Gillison, M.L. Effect of prophylactic human papillomavirus (HPV) vaccination on oral HPV infections among young adults in the United States. *J. Clin. Oncol.* **2018**, *36*, 262–267. [CrossRef]
117. Zahnd, W.E.; Jenkins, W.D.; Mueller-Luckey, G.S. Cancer mortality in the mississippi delta region: Descriptive epidemiology and needed future research and interventions. *J. Health Care Poor Underserved* **2017**, *28*, 315–328. [CrossRef]
118. Walker, T.Y.; Elam-Evans, L.D.; Williams, C.L.; Fredua, B.; Yankey, D.; Markowitz, L.E.; Stokley, S. Trends in human papillomavirus (HPV) vaccination initiation among adolescents aged 13–17 by metropolitan statistical area (MSA) status, National Immunization Survey–Teen, 2013–2017. *Hum. Vaccines Immunother.* **2020**, *16*, 554–561. [CrossRef] [PubMed]
119. Luque, J.S.; Raychowdhury, S.; Weaver, M. Health care provider challenges for reaching Hispanic immigrants with HPV vaccination in rural Georgia. *Rural Remote Health* **2012**, *12*, 168.
120. Cates, J.R.; Brewer, N.T.; Fazekas, K.I.; Mitchell, C.E.; Smith, J.S. Racial differences in HPV knowledge, HPV vaccine acceptability, and related beliefs among rural, Southern women. *J. Rural Health* **2009**, *25*, 93–97. [CrossRef] [PubMed]
121. Landrine, H.; Klonoff, E.A. Racial segregation and cigarette smoking among Blacks: Findings at the individual level. *J. Health Psychol.* **2000**, *5*, 211–219. [CrossRef]

122. Havewala, F. The dynamics between the food environment and residential segregation: An analysis of metropolitan areas. *Food Policy* **2020**. [CrossRef]
123. Lichter, D.T.; Parisi, D.; Grice, S.M.; Taquino, M.C. National estimates of racial segregation in rural and small-town America. *Demography* **2007**, *44*, 563–581. [CrossRef]
124. Anderson, A.E.; Henry, K.A.; Samadder, N.J.; Merrill, R.M.; Kinney, A.Y. Rural vs urban residence affects risk-appropriate colorectal cancer screening. *Clin. Gastroenterol. Hepatol.* **2013**, *11*, 526–533. [CrossRef]
125. Bennett, K.J.; Pumkam, C.; Bellinger, J.D.; Probst, J.C. Cancer screening delivery in persistent poverty rural counties. *J. Prim. Care Community Health* **2011**, *2*, 240–249. [CrossRef]
126. Khan-Gates, J.A.; Ersek, J.L.; Eberth, J.M.; Adams, S.A.; Pruitt, S.L. Geographic Access to Mammography and Its Relationship to Breast Cancer Screening and Stage at Diagnosis: A Systematic Review. *Womens Health Issues* **2015**, *25*, 482–493. [CrossRef]
127. Duggan, C.; Molina, Y.; Carosso, E.; Ibarra, G.; Thompson, B. County of residence and screening practices among Latinas and non-Latina whites in two rural communities. *Ethn. Dis.* **2019**, *29*, 31–38. [CrossRef]
128. Nuno, T.; Gerald, J.K.; Harris, R.; Martinez, M.E.; Estrada, A.; Garcia, F.; Nuño, T.; Gerald, J.K.; Harris, R.; Martinez, M.E.; et al. Comparison of breast and cervical cancer screening utilization among rural and urban Hispanic and American Indian women in the Southwestern United States. *Cancer Causes Control.* **2012**, *23*, 1333–1341. [CrossRef] [PubMed]
129. Bennett, K.J.; Probst, J.C.; Bellinger, J.D. Receipt of cancer screening services: Surprising results for some rural minorities. *J. Rural Health* **2012**, *28*, 63–72. [CrossRef] [PubMed]
130. Towne, S.D., Jr.; Smith, M.L.; Ory, M.G. Geographic variations in access and utilization of cancer screening services: Examining disparities among American Indian and Alaska Native Elders. *Int. J. Health Geogr.* **2014**, *13*, 18. [CrossRef] [PubMed]
131. Rai, A.; Doria-Rose, V.P.; Silvestri, G.A.; Yabroff, K.R. Evaluating Lung Cancer Screening Uptake, Outcomes, and Costs in the United States: Challenges With Existing Data and Recommendations for Improvement. *JNCI J. Natl. Cancer Inst.* **2019**, *111*, 342–349. [CrossRef] [PubMed]
132. Odahowski, C.L.; Zahnd, W.E.; Eberth, M.J. Challenges and Opportunities for Lung Cancer Screening in Rural America. *J. Am. Coll. Radiol.* **2019**, *16*, 590–595. [CrossRef] [PubMed]
133. Jemal, A.; Fedewa, S.A. Lung Cancer Screening With Low-Dose Computed Tomography in the United States-2010 to 2015. *Jama Oncol.* **2017**, *3*, 1278–1281. [CrossRef]
134. Zgodic, A.; Zahnd, W.E.; Miller, D.P.; Studts, J.L.; Eberth, J.M. Predictors of Lung Cancer Screening Utilization in a Population-Based Survey. *J. Am. Coll. Radiol.* **2020**. [CrossRef]
135. Wang, H.; Roy, S.; Kim, J.; Farazi, P.A.; Siashpush, M.; Su, D. Barriers of colorectal cancer screening in rural USA: A systematic review. *Rural Remote Health* **2019**, *19*, 5181. [CrossRef]
136. Katz, M.L.; James, A.S.; Pignone, M.P.; Hudson, M.A.; Jackson, E.; Oates, V.; Campbell, M.K. Colorectal cancer screening among African American church members: A qualitative and quantitative study of patient-provider communication. *BMC Public Health* **2004**, *4*, 62. [CrossRef]
137. Surveillance Epidemiology and End Results (SEER) Program (www.seer.cancer.gov) SEER*Stat Database: Incidence-SEER Research Limited Field Data. 21 Registries. Nov 2019 Sub (2000–2017)-Linked to County Attributes-Time Dependent (1990–2017)/Income-Rurality, 1969–2018 Counties, National Cancer Institute. DCCPS, Surveillance Research Program, Released April 2020. Based upon the November 2019 Submission. Available online: <https://seer.cancer.gov/data/citation.html> (accessed on 3 February 2021).
138. Zahnd, W.E.; Gomez, S.L.; Steck, S.E.; Brown, M.J.; Ganai, S.; Zhang, J.; Arp Adams, S.; Berger, F.G.; Eberth, J.M. Rural-urban and racial/ethnic trends and disparities in early-onset and average-onset colorectal cancer. *Cancer* **2020**. [CrossRef]
139. Singh, G.K. Rural-urban trends and patterns in cervical cancer mortality, incidence, stage, and survival in the United States, 1950–2008. *J. Community Health* **2012**, *37*, 217–223. [CrossRef] [PubMed]
140. Houston, K.A.; Mitchell, K.A.; King, J.; White, A.; Ryan, B.M. Histologic Lung Cancer Incidence Rates and Trends Vary by Race/Ethnicity and Residential County. *J. Thorac. Oncol.* **2018**, *13*, 497–509. [CrossRef] [PubMed]
141. Rovito, M.J.; Taylor, S.; Lockwood, R.; Adams, W.B.; Craycraft, M. Testicular Cancer Incidence and Mortality Within Rural and Urban Regions. *J. Adolesc. Young Adult Oncol.* **2020**, *9*, 202–207. [CrossRef] [PubMed]
142. Zahnd, W.E.; Fogleman, A.J.; Jenkins, W.D. Rural-Urban Disparities in Stage of Diagnosis Among Cancers With Preventive Opportunities. *Am. J. Prev. Med.* **2018**, *54*, 688–698. [CrossRef] [PubMed]
143. Yu, L.; Sabatino, S.A.; White, M.C. Rural-urban and racial/ethnic disparities in invasive cervical cancer incidence in the United States, 2010–2014. *Prev. Chronic Dis.* **2019**, *16*, E70. [CrossRef] [PubMed]
144. Camacho, F.T.; Tan, X.; Alcalá, H.E.; Shah, S.; Anderson, R.T.; Balkrishnan, R. Impact of patient race and geographical factors on initiation and adherence to adjuvant endocrine therapy in medicare breast cancer survivors. *Medicine (Baltim.)* **2017**, *96*, e7147. [CrossRef] [PubMed]
145. Chen, L.; Li, C.I. Racial disparities in breast cancer diagnosis and treatment by hormone receptor and HER2 status. *Cancer Epidemiol Biomark. Prev.* **2015**, *24*, 1666–1672. [CrossRef]
146. Steele, C.B.; Pisu, M.; Richardson, L.C. Urban/rural patterns in receipt of treatment for non-small cell lung cancer among black and white medicare beneficiaries, 2000–2003. *J. Natl. Med. Assoc.* **2011**, *103*, 711–718. [CrossRef]

147. Zahnd, W.E.; Hyon, K.S.; Diaz-Sylvester, P.; Izadi, S.R.; Colditz, G.A.; Brard, L. Rural-urban differences in surgical treatment, regional lymph node examination, and survival in endometrial cancer patients. *Cancer Causes Control*. **2018**, *29*, 221–232. [CrossRef]
148. AlNasser, M.; Schneider, E.B.; Gearhart, S.L.; Wick, E.C.; Fang, S.H.; Haider, A.H.; Efron, J.E. National disparities in laparoscopic colorectal procedures for colon cancer. *Surg. Endosc.* **2014**, *28*, 49–57. [CrossRef]
149. Thomas, P.S.; Class, C.A.; Gandhi, T.R.; Bambhroliya, A.; Do, K.A.; Brewster, A.M. Demographic, clinical, and geographical factors associated with lack of receipt of physician recommended chemotherapy in women with breast cancer in Texas. *Cancer Causes Control*. **2019**, *30*, 409–415. [CrossRef] [PubMed]
150. Des Bordes, J.K.A.; Lopez, D.S.; Swartz, M.D.; Volk, R.J. Sociodemographic Disparities in Cure-Intended Treatment in Localized Prostate Cancer. *J. Racial Ethn. Health Disparities* **2018**, *5*, 104–110. [CrossRef] [PubMed]
151. Bregar, A.J.; Melamed, A.; Diver, E.; Clemmer, J.T.; Uppal, S.; Schorge, J.O.; Rice, L.W.; del Carmen, M.G.; Rauh-Hain, J.A. Minimally Invasive Staging Surgery in Women with Early-Stage Endometrial Cancer: Analysis of the National Cancer Data Base. *Ann. Surg. Oncol.* **2017**, *24*, 1677–1687. [CrossRef] [PubMed]
152. Gabriel, E.; Thirunavukarasu, P.; Al-Sukhni, E.; Attwood, K.; Nurkin, S.J. National disparities in minimally invasive surgery for rectal cancer. *Surg. Endosc.* **2016**, *30*, 1060–1067. [CrossRef] [PubMed]
153. Arsoniadis, E.G.; Fan, Y.; Jarosek, S.; Gaertner, W.B.; Melton, G.B.; Madoff, R.D.; Kwaan, M.R. Decreased Use of Sphincter-Preserving Procedures Among African Americans with Rectal Cancer. *Ann. Surg. Oncol.* **2018**, *25*, 720–728. [CrossRef] [PubMed]
154. Nguyen, C.; Lairson, D.R.; Swartz, M.D.; Du, X.L. Racial, Socioeconomic, and Geographic Disparities in the Receipt, Timing to Initiation, and Duration of Adjuvant Androgen Deprivation Therapy in Men with Prostate Cancer. *J. Racial Ethn. Health Disparities* **2019**, *6*, 133–142. [CrossRef]
155. Mollica, M.A.; Weaver, K.E.; McNeel, T.S.; Kent, E.E. Examining urban and rural differences in perceived timeliness of care among cancer patients: A SEER-CAHPS study. *Cancer* **2018**, *124*, 3257–3265. [CrossRef]
156. Kong, A.L.; Yen, T.W.F.; Pezzin, L.E.; Miao, H.; Sparapani, R.A.; Laud, P.W.; Nattinger, A.B. Socioeconomic and racial differences in treatment for breast cancer at a low-volume hospital. *Ann. Surg. Oncol.* **2011**, *18*, 3220–3227. [CrossRef]
157. Hughes, R.G.; Hunt, S.S.; Luft, H.S. Effects of surgeon volume and hospital volume on quality of care in hospitals. *Med. Care* **1987**, *25*, 489–503. [CrossRef]
158. Whitaker, R.G.; Holmes, G.M.; Pink, G.H. The Impact of the Low Volume Hospital (LVH) Program on the Viability of Small, Rural Hospitals. NC Rural Health Research Program, Findings Brief. October 2016. Available online: https://www.shepscenter.unc.edu/wp-content/uploads/dlm_uploads/2016/10/Impact-of-LVH.pdf (accessed on 9 December 2020).
159. Esnaola, N.F.; Ford, M.E. Racial Differences and Disparities in Cancer Care and Outcomes: Where’s the Rub? *Surg. Oncol. Clin. N. Am.* **2012**, *21*, 417. [CrossRef]
160. Jessup, J.M.; Stewart, A.; Greene, F.L.; Minsky, B.D. Adjuvant chemotherapy for stage III colon cancer: Implications of race/ethnicity, age, and differentiation. *JAMA* **2005**, *29*, 2703–2711. [CrossRef] [PubMed]
161. Mandelblatt, J.S.; Kerner, J.F.; Hadley, J.; Hwang, Y.T.; Eggert, L.; Johnson, L.E.; Gold, K.; OPTIONS. Variations in breast carcinoma treatment in older medicare beneficiaries: Is it black or white. *Cancer* **2002**, *95*, 1401–1414. [CrossRef] [PubMed]
162. Cykert, S.; Dilworth-Anderson, P.; Monroe, M.H.; Walker, P.; McGuire, F.R.; Corbie-Smith, G.; Edwards, L.J.; Bunton, A.J. Factors associated with decisions to undergo surgery among patients with newly diagnosed early-stage lung cancer. *JAMA* **2010**, *303*, 2368–2376. [CrossRef] [PubMed]
163. National Cancer Institute. Definition of survivorship—NCI Dictionary of Cancer Terms—National Cancer Institute. Available online: <https://www.cancer.gov/publications/dictionaries/cancer-terms/def/survivorship> (accessed on 15 October 2020).
164. Khan, S.; Hicks, V.; Rancilio, D.; Langston, M.; Richardson, K.; Drake, B.F. Predictors of Follow-Up Visits Post Radical Prostatectomy. *Am. J. Men’s Health* **2018**, *12*, 760–765. [CrossRef]
165. Weaver, K.E.; Geiger, A.M.; Lu, L.; Case, L.D. Rural-urban disparities in health status among US cancer survivors. *Cancer* **2013**, *119*, 1050–1057. [CrossRef]
166. Spencer, J.C.; Reeve, B.B.; Troester, M.A.; Wheeler, S.B. Factors Associated with Endocrine Therapy Non-Adherence in Breast Cancer Survivors. *Psycho Oncol.* **2020**, *29*, 647–654. [CrossRef]
167. Schootman, M.; Homan, S.; Weaver, K.E.; Jeffe, D.B.; Yun, S. The health and welfare of rural and urban cancer survivors in Missouri. *Prev. Chronic Dis.* **2013**, *10*, E152. [CrossRef]
168. Andrykowski, M.A.; Steffens, R.F.; Bush, H.M.; Tucker, T.C. Disparities in mental health outcomes among lung cancer survivors associated with ruralness of residence. *Psychooncology* **2014**, *23*, 428–436. [CrossRef]
169. Weaver, K.E.; Palmer, N.; Lu, L.; Case, L.D.; Geiger, A.M. Rural-urban differences in health behaviors and implications for health status among US cancer survivors. *Cancer Causes Control*. **2013**, *24*, 1481–1490. [CrossRef]
170. Odahowski, C.L.; Zahnd, W.E.; Zgodic, A.; Edward, J.S.; Hill, L.N.; Davis, M.M.; Perry, C.K.; Shannon, J.; Wheeler, S.B.; Vanderpool, R.C.; et al. Financial hardship among rural cancer survivors: An analysis of the Medical Expenditure Panel Survey. *Prev. Med.* **2019**, 105881. [CrossRef]
171. Zahnd, W.E.; Davis, M.M.; Rotter, J.S.; Vanderpool, R.C.; Perry, C.K.; Shannon, J.; Ko, L.K.; Wheeler, S.B.; Odahowski, C.L.; Farris, P.E.; et al. Rural-urban differences in financial burden among cancer survivors: An analysis of a nationally representative survey. *Supportive Care Cancer* **2019**, *27*, 4779–4786. [CrossRef] [PubMed]

172. Weaver, K.E.; Rowland, J.H.; Bellizzi, K.M.; Aziz, N.M. Forgoing medical care because of cost: Assessing disparities in healthcare access among cancer survivors living in the United States. *Cancer* **2010**, *116*, 3493–3504. [CrossRef] [PubMed]
173. Yabroff, K.R.; Dowling, E.C.; Guy, G.P., Jr.; Banegas, M.P.; Davidoff, A.; Han, X.; Virgo, K.S.; McNeel, T.S.; Chawla, N.; Blanch-Hartigan, D.; et al. Financial Hardship Associated With Cancer in the United States: Findings From a Population-Based Sample of Adult Cancer Survivors. *J. Clin. Oncol.* **2016**, *34*, 259–267. [CrossRef] [PubMed]
174. Spencer, J.C.; Rotter, J.S.; Eberth, J.M.; Zahnd, W.E.; Vanderpool, R.C.; Ko, L.K.; Davis, M.M.; Troester, M.A.; Olshan, A.F.; Wheeler, S.B. Employment Changes Following Breast Cancer Diagnosis: The Effects of Race and Place. *J. Natl. Cancer Inst.* **2020**, *112*, 647–650. [CrossRef] [PubMed]
175. Guadagnolo, B.A.; Petereit, D.G.; Coleman, C.N. Cancer Care Access and Outcomes for American Indian Populations in the United States: Challenges and Models for Progress. *Semin. Radiat. Oncol.* **2017**, *27*, 143–149. [CrossRef] [PubMed]
176. Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death “ ” 1999–2019 on CDC WONDER Online Database, released in 2020. Data are from the Multiple Cause of Death Files, 1999–2019, as “ ” compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Available online: <http://wonder.cdc.gov/ucd-icd10.html> (accessed on 25 January 2021).
177. Singh, G.K.; Williams, S.D.; Siahpush, M.; Mulhollen, A. Socioeconomic, Rural-Urban, and Racial Inequalities in US Cancer Mortality: Part I-All Cancers and Lung Cancer and Part II-Colorectal, Prostate, Breast, and Cervical Cancers. *J. Cancer Epidemiol.* **2011**, *2011*, 107497. [CrossRef]
178. Singh, G.K.; Siahpush, M. Widening rural-urban disparities in all-cause mortality and mortality from major causes of death in the USA, 1969–2009. *J. Urban. Health* **2014**, *91*, 272–292. [CrossRef]
179. Johnson, A.M.; Johnson, A.; Hines, R.B.; Mohammadi, R. Neighborhood context and non-small cell lung cancer outcomes in Florida non-elderly patients by race/ethnicity. *Lung Cancer* **2020**, *142*, 20–27. [CrossRef]
180. Higginbotham, J.C.; Moulder, J.; Currier, M. Rural v. urban aspects of cancer: First-year data from the Mississippi Central Cancer Registry. *Fam. Community Health* **2001**, *24*, 1–9. [CrossRef]
181. Hines, R.; Markossian, T.; Johnson, A.; Dong, F.; Bayakly, R. Geographic residency status and census tract socioeconomic status as determinants of colorectal cancer outcomes. *Am. J. Public Health* **2014**, *104*, e63–e71. [CrossRef]
182. Thompson, B.; Hohl, S.D.; Molina, Y.; Paskett, E.D.; Fisher, J.L.; Baltic, R.D.; Washington, C.M. Breast Cancer Disparities Among Women in Underserved Communities in the USA. *Curr. Breast Cancer Rep.* **2018**, *10*, 131–141. [CrossRef] [PubMed]
183. Puechl, A.M.; Chino, F.; Havrilesky, L.J.; Davidson, B.A.; Chino, J.P. Place of death by region and urbanization among gynecologic cancer patients: 2006–2016. *Gynecol. Oncol.* **2019**, *155*, 98–104. [CrossRef] [PubMed]
184. Jemal, A.; Ward, E.; Wu, X.; Martin, H.J.; McLaughlin, C.C.; Thun, M.J. Geographic patterns of prostate cancer mortality and variations in access to medical care in the United States. *Cancer Epidemiol. Biomark. Prev.* **2005**, *14*, 590–595. [CrossRef]
185. Jacobs-Wingo, J.L.; Espey, D.K.; Groom, A.V.; Phillips, L.E.; Haverkamp, D.S.; Stanley, S.L. Causes and disparities in death rates among Urban American Indian and Alaska native populations, 1999–2009. *Am. J. Public Health* **2016**, *106*, 906–914. [CrossRef] [PubMed]
186. Lindley, L.C.; Oyana, T.J. Geographic Variation in Mortality Among Children and Adolescents Diagnosed With Cancer in Tennessee: Does Race Matter? *J. Pediatric Oncol. Nurs.* **2016**, *33*, 129–136. [CrossRef] [PubMed]
187. Hines, R.B.; Markossian, T.W. Differences in Late-Stage Diagnosis, Treatment, and Colorectal Cancer-Related Death Between Rural and Urban African Americans and Whites in Georgia. *J. Rural Health* **2012**, *28*, 296–305. [CrossRef]
188. Delavar, A.; Feng, Q.; Johnson, K.J. Rural/urban residence and childhood and adolescent cancer survival in the United States. *Cancer* **2019**, *125*, 261–268. [CrossRef]
189. Clarke, J.A.; Despotis, A.M.; Ramirez, R.J.; Zevallos, J.P.; Mazul, A.L. Head and Neck Cancer Survival Disparities by Race and Rural–Urban Context. *Cancer Epidemiol. Biomark. Prev.* **2020**, *29*, 1955–1961. [CrossRef]
190. Rogers, C.R.; Blackburn, B.E.; Huntington, M.; Curtin, K.; Thorpe, R.J.; Rowe, K.; Snyder, J.; Deshmukh, V.; Newman, M.; Fraser, A.; et al. Rural–urban disparities in colorectal cancer survival and risk among men in Utah: A statewide population-based study. *Cancer Causes Control.* **2020**, *31*, 241–253. [CrossRef]
191. Patel, M.I.; Lopez, A.M.; Blackstock, W.; Reeder-Hayes, K.; Allyn Moushey, E.; Phillips, J.; Tap, W. Cancer disparities and health equity: A policy statement from the American Society of Clinical Oncology. *J. Clin. Oncol.* **2020**, *38*, 3439–3448. [CrossRef]
192. Bouye, K.E.; McCleary, K.J.; Williams, K.B. Increasing Diversity in the Health Professions: Reflections on Student Pipeline Programs. *J. Healthc. Sci. Humanit* **2016**, *6*, 67–79. [PubMed]
193. Oklahoma State University. OSU Ceremony Opens First Tribally Affiliated Medical School. Available online: <https://news.okstate.edu/articles/communications/2020/osu-ceremony-opens-first-tribally-affiliated-medical-school.html/> (accessed on 25 January 2021).
194. Davidoff, A.J.; Hill, S.C.; Bernard, D.; Yabroff, K.R. The Affordable Care Act and Expanded Insurance Eligibility among Nonelderly Adult Cancer Survivors. *Jnci J. Natl. Cancer Inst.* **2018**, *107*, djv181. [CrossRef] [PubMed]
195. Gan, T.; Sinner, H.F.; Walling, S.C.; Chen, Q.; Huang, B.; Tucker, T.C.; Patel, J.A.; Evers, B.M.; Bhakta, A.S. Impact of the Affordable Care Act on Colorectal Cancer Screening, Incidence, and Survival in Kentucky. *J. Am. Coll. Surg.* **2019**, *228*, 342–353.e1. [CrossRef] [PubMed]
196. Yankey, D.; Elam-Evans, L.D.; Bish, C.L.; Stokley, S.K. Human papillomavirus vaccination estimates among adolescents in the Mississippi delta region: National immunization survey-teen, 2015–2017. *Prev. Chronic Dis.* **2020**, *17*, E31. [CrossRef] [PubMed]

197. Gennuso, K.P.; Jovaag, A.; Catlin, B.B.; Rodock, M.; Park, H. Assessment of Factors Contributing to Health Outcomes in the Eight States of the Mississippi Delta Region. *Prev. Chronic Dis.* **2016**, *13*, E33. [CrossRef] [PubMed]
198. Mokdad, A.H.; Dwyer-Lindgren, L.; Fitzmaurice, C.; Stubbs, R.W.; Bertozzi-Villa, A.; Morozoff, C.; Charara, R.; Allen, C.; Naghavi, M.; Murray, C.J. Trends and Patterns of Disparities in Cancer Mortality Among US Counties, 1980–2014. *JAMA* **2017**, *317*, 388–406. [CrossRef]
199. Siegel, R.L.; Sahar, L.; Robbins, A.; Jemal, A. Where can colorectal cancer screening interventions have the most impact? *Cancer Epidemiol. Biomark. Prev.* **2015**, *24*, 1151–1156. [CrossRef]
200. Grants and Contracts—Appalachian Regional Commission. Available online: <https://www.arc.gov/grants-and-contracts/> (accessed on 1 December 2020).
201. DRA: FY 2021 Close-Out Budget Justification. 2019. Available online: https://dra.gov/images/uploads/content_files/FY_2021_Close_Out_Justification_FINAL.pdf (accessed on 1 December 2020).
202. Blewett, L.A.; Call, K.T.; Turner, J.; Hest, R. Data Resources for Conducting Health Services and Policy Research. *Annu. Rev. Public Health* **2018**, *39*, 437–452. [CrossRef]
203. Zahnd, W.E.; Askelson, N.; Vanderpool, R.C.; Stradtman, L.; Edward, J.; Farris, P.E.; Petermann, V.; Eberth, J.M. Challenges of using nationally representative, population-based surveys to assess rural cancer disparities. *Prev. Med.* **2019**, *129*, 105812. [CrossRef]
204. Becker, T.; Babey, S.H.; Shimkhada, R.; Scheitler, A.J.; Ponce, N.A. Limited access to health data on American Indian and Alaska Natives impedes population health insights. Policy Brief. 16 November 2020. Available online: <https://healthpolicy.ucla.edu/publications/search/pages/detail.aspx?PubID=2004> (accessed on 9 December 2020).
205. Haddad, D.N.; Sandler, K.L.; Henderson, L.M.; Rivera, M.P.; Aldrich, M.C. Disparities in lung cancer screening: A review. *Ann. Am. Thorac. Soc.* **2020**, *17*, 399–405. [CrossRef]
206. Aldrich, M.C.; Mercaldo, S.F.; Sandler, K.L.; Blot, W.J.; Grogan, E.L.; Blume, J.D. Evaluation of USPSTF Lung Cancer Screening Guidelines among African American Adult Smokers. *JAMA Oncol.* **2019**, *5*, 1318–1324. [CrossRef] [PubMed]
207. United States Preventive Services Taskforce. Recommendation: Lung Cancer: Screening. Available online: <https://www.uspreventiveservicestaskforce.org/uspstf/draft-update-summary/lung-cancer-screening1> (accessed on 1 December 2020).
208. Huey, R.W.; Hawk, E.; Offodile, A.C. Mind the Gap: Precision Oncology and Its Potential to Widen Disparities. *J. Oncol. Pract.* **2019**, *15*, 301–304. [CrossRef] [PubMed]
209. Shih, Y.C.T.; Smieliauskas, F.; Geynisman, D.M.; Kelly, R.J.; Smith, T.J. Trends in the cost and use of targeted cancer therapies for the privately insured nonelderly: 2001 to 2011. *J. Clin. Oncol.* **2015**, *33*, 2190–2196. [CrossRef] [PubMed]
210. Haslam, A.; Prasad, V. Estimation of the Percentage of US Patients With Cancer Who Are Eligible for and Respond to Checkpoint Inhibitor Immunotherapy Drugs. *JAMA Netw. Open* **2019**, *2*, e192535. [CrossRef] [PubMed]
211. National Institutes of Health. RFA-CA-20-051: Social and Behavioral Intervention Research to Address Modifiable Risk Factors for Cancer in Rural Populations (R01 Clinical Trial Required). Available online: <https://grants.nih.gov/grants/guide/rfa-files/RFA-CA-20-051.html> (accessed on 1 December 2020).
212. Moonshot Blue Ribbon Panel Implementation Science Working Group. Accelerating Implementation of Evidence-Base Cancer Prevention and Screening Strategies. Available online: <https://www.cancer.gov/research/key-initiatives/moonshot-cancer-initiative/blue-ribbon-panel/implementation-science-working-group-report.pdf> (accessed on 9 December 2020).
213. Weaver, S.J.; Blake, K.D.; Vanderpool, R.C.; Gardner, B.; Croyle, R.T.; Srinivasan, S. Advancing Rural Cancer Control Research: National Cancer Institute Efforts to Identify Gaps and Opportunities. *Cancer Epidemiol. Biomark. Prev.* **2020**, *29*, 1515–1518. [CrossRef]
214. Blake, K.D.; Moss, J.L.; Gaysynsky, A.; Srinivasan, S.; Croyle, R.T. Making the Case for Investment in Rural Cancer Control: An Analysis of Rural Cancer Incidence, Mortality, and Funding Trends. *Cancer Epidemiol. Biomark. Prev.* **2017**, *26*, 992–997. [CrossRef]
215. National Cancer Institute, Division of Cancer Control and Population Sciences. Rural Cancer Control. Available online: <https://cancercontrol.cancer.gov/research-emphasis/rural> (accessed on 9 December 2020).
216. Vanderpool, R.C.; Stradtman, L.R.; Brandt, H.M. Policy opportunities to increase HPV vaccination in rural communities. *Hum. Vaccines Immunother.* **2019**, *15*, 1527–1532. [CrossRef]
217. HPV Vaccination Evidence-Based Programs Listing | Evidence-Based Cancer Control Programs (EBCCP). Available online: <https://ebccp.cancercontrol.cancer.gov/topicPrograms.do?topicId=22626661&choice=default> (accessed on 1 December 2020).
218. Bragge, P.; Grimshaw, J.M.; Lokker, C.; Colquhoun, H.; Albrecht, L.; Baron, J.; Dadich, A.; Damschroder, L.; Danko, K.; Fernandez, M.E.; et al. AIMD—A validated, simplified framework of interventions to promote and integrate evidence into health practices, systems, and policies. *BMC Med. Res. Methodol.* **2017**, *17*, 38. [CrossRef]
219. Pinnock, H.; Barwick, M.; Carpenter, C.R.; Eldridge, S.; Grandes, G.; Griffiths, C.J.; Rycroft-Malone, J.; Meissner, P.; Murray, E.; Patel, A.; et al. Standards for Reporting Implementation Studies (StaRI) Statement. *BMJ* **2017**, *356*. [CrossRef]
220. Hoffmann, T.C.; Glasziou, P.P.; Boutron, I.; Milne, R.; Perera, R.; Moher, D.; Altman, D.G.; Barbour, V.; Macdonald, H.; Johnston, M.; et al. Better reporting of interventions: Template for intervention description and replication (TIDieR) checklist and guide. *BMJ* **2014**, *348*. [CrossRef] [PubMed]
221. Welch, V.A.; Norheim, O.F.; Jull, J.; Cookson, R.; Sommerfelt, H.; Tugwell, P. CONSORT-Equity 2017 extension and elaboration for better reporting of health equity in randomised trials. *BMJ Clin. Res.* **2017**, *359*, j5085. [CrossRef] [PubMed]