



Lysyl oxidase family gene polymorphisms and risk of aneurysmal subarachnoid hemorrhage: a case-control study

Chun Luo^{1#^}, Bingyang Li^{1,2#^}, Chao Liu³, Rui Dong^{1^}, Chongyu Hu^{4^}, Junyu Liu^{5^}, Liming Hu^{1^}, Xin Liao^{6^}, Jilin Zhou^{5^}, Lu Xu^{5^}, Songlin Liu^{5^}, Dun Yuan^{5^}, Weixi Jiang^{5^}, Junxia Yan^{1,7^}, Yifeng Li^{5^}

¹Department of Epidemiology and Health Statistics, Xiangya School of Public Health, Central South University, Changsha, China; ²Department of Information Statistics, Changsha Hospital of Traditional Chinese Medicine (Changsha Eighth Hospital), Changsha, China; ³Department of Neurosurgery, Central Hospital of Zhuzhou, Zhuzhou, China; ⁴Department of Neurology, Hunan People's Hospital, Changsha, China; ⁵Department of Neurosurgery, Xiangya Hospital, Central South University, Changsha, China; ⁶Department of Biomedical Sciences, Faculty of Health Sciences, University of Macau, Macau, China; ⁷The Hunan Provincial Key Laboratory of Clinical Epidemiology, Xiangya School of Public Health, Central South University, Changsha, China

Contributions: (I) Conception and design: C Luo, B Li, Y Li; (II) Administrative support: W Jiang, J Yan, Y Li; (III) Provision of study materials or patients: C Liu, S Liu, C Hu, J Liu, D Yuan; (IV) Collection and assembly of data: C Luo, B Li, L Hu, X Liao, J Zhou, L Xu; (V) Data analysis and interpretation: C Luo, B Li, C Liu, R Dong; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

[#]These authors contributed equally to this work.

Correspondence to: Yifeng Li. Department of Neurosurgery, Xiangya Hospital, Central South University, Changsha 410008, China. Email: liyifeng87@csu.edu.cn.

Background: Aneurysmal subarachnoid hemorrhage (aSAH) is a devastating disease caused by intracranial aneurysm (IA) rupture. Lysyl oxidase (*LOX*) family genes (*LOX-like [LOXL] 1-4*) have roles in collagen cross-linking in the extracellular matrix (ECM) and may be associated with IA rupture. We aimed to explore the association between *LOX* polymorphisms and the risk of aSAH.

Methods: This case-control study included 2 cohorts: 133 single ruptured and 115 unruptured IA patients, and 65 multiple ruptured and 71 unruptured IA patients. Genotyping of 27 single nucleotide polymorphisms (SNPs) in *LOX* was performed. Logistic regression analysis was performed to calculate the odds ratios (ORs) and 95% confidence intervals (CIs) of the SNPs of *LOX* and the risk of aSAH.

Results: *LOX* rs1800449 and *LOXL4* rs3793692 were positively associated with the risk of single IA rupture in the recessive model (OR =5.66, 2.06; 95% CI =1.22–26.24, 1.11–3.82, respectively) and *LOX* rs10519694 demonstrated a protective effect on single IA rupture (dominant model: OR =0.42, 95% CI =0.21–0.83; recessive model: OR =0.16, 95% CI =0.04–0.65; additive model: OR =0.46, 95% CI =0.28–0.78). *LOXL1* rs2165241, *LOXL2* rs1063582, and *LOXL3* rs17010021 showed risk effects on multiple IAs rupture. *LOXL3* rs17010022 showed a protective effect on multiple IAs ruptures (dominant model: OR =0.41, 95% CI =0.21–0.82; additive model: OR =0.51, 95% CI =0.30–0.85).

Conclusions: *LOX* and *LOXL4* may be susceptibility genes for single IA rupture, whereas *LOXL1-3* may have a role in susceptibility to multiple IAs ruptures in the Chinese population, suggesting that *LOX* family genes may be associated with aSAH.

Keywords: Intracranial aneurysms (IAs); subarachnoid hemorrhage; aneurysmal subarachnoid hemorrhage (aSAH); *LOXL2*; *LOX*

Submitted Jun 20, 2022. Accepted for publication Aug 08, 2022.

doi: 10.21037/atm-22-3484

View this article at: <https://dx.doi.org/10.21037/atm-22-3484>

[^] ORCID: Chun Luo, 0000-0001-9935-9467; Bingyang Li, 0000-0003-0138-773X; Rui Dong, 0000-0002-5701-7669; Chongyu Hu, 0000-0002-4086-0916; Junyu Liu, 0000-0002-3803-4068; Liming Hu, 0000-0002-0166-8585; Xin Liao, 0000-0002-9640-8638; Jilin Zhou, 0000-0003-2091-980X; Lu Xu, 0000-0002-1905-4557; Songlin Liu, 0000-0002-2985-0266; Dun Yuan, 0000-0002-9665-4222; Weixi Jiang, 0000-0003-1570-0854; Junxia Yan, 0000-0001-6067-1613; Yifeng Li, 0000-0002-7593-8453.

Introduction

Intracranial aneurysm (IA) is a complex disease characterized by incomplete integrity of the artery wall, which is typically induced by pathological expansion and swelling of the weak artery wall prone to rupture (1). IA affects 3–5% of the global population and approximately 7% of the Chinese population aged 35–75 years (2–4). IA can be classified as single (i.e., 1 aneurysm) or multiple (i.e., equal to or more than 2 aneurysms) events, which accounts for approximately 20–30% of cases (5). Most IAs are asymptomatic before rupture; however, rupture can result in an aneurysmal subarachnoid hemorrhage (aSAH), which is a devastating condition with a poor prognosis (6). Thus, to manage IA more effectively, it is necessary to identify the risk factors associated with aSAH as early as possible.

Although the etiology of IA rupture is not entirely clear, both environmental and genetic factors have been recognized to possibly lead to IA rupture (7,8). For instance, a study has revealed that smoking and hypertension are predictors of IA rupture (9). Furthermore, first-degree relatives of aSAH patients are more likely to be diagnosed with IA or aSAH, which indicates a familial tendency (10). Histopathologically, extracellular matrix (ECM) plays an important role in the structural support of cerebral artery blood vessels (11). Some IA remain stable over time, and the walls of unruptured IAs exhibit with ECM defects and tissue thrombosis, but in others mural cell die, the ECM degenerates too fragile to resist intravascular hemodynamic pressure, the IA will rupture, causing aSAH (1,12). Elastin and collagens are abundant matrix proteins in the ECM, while the lysyl oxidase (*LOX*) family of genes in the extracellular copper-containing enzyme family initiate cross-linking of collagen and elastin by oxidative deamination of lysine residues (13). Therefore, *LOX* family genes may play roles in aSAH, and the alteration of matrix proteins may cause vasculature-related pathological changes in aSAH.

Although numerous studies have explored the genetic susceptibility of aSAH, there are few studies on the association between *LOX* family gene polymorphisms and aSAH. Our previous study found that *LOX* was associated with the risk of single IA, while *LOX*-like 2 (*LOXL2*) was associated with the risk of multiple IAs (14). *LOXL2*, which belongs to the *LOX* family of genes, was found to be associated with susceptibility to familial IA (FIA) in Chinese and Japanese populations (15,16); In Korea, Hong *et al.* discovered that *LOX* was associated with IA

formation and rupture using candidate gene association analysis (17). However, since the pathophysiology of IA rupture is somewhat different from the process of IA formation (12), whether *LOX* family genes are also associated with aSAH in Chinese population remains unclear. Since there is high homology among the subtypes of the *LOX* family genes, other members of this family may also be associated with aSAH. Therefore, we aimed to explore whether *LOX* family gene polymorphisms are associated with aSAH by comparing ruptured and unruptured IA in a Chinese sample and provide a reference for the etiological study of aSAH. We present the following article in accordance with the MDAR reporting checklist (available at <https://atm.amegroups.com/article/view/10.21037/atm-22-3484/rc>).

Methods

Study population

A total of 384 patients with IA were collected from 2 third-class hospitals in Hunan province (Xiangya Hospital of Central South University and Hunan People's Hospital) from July 2018 to December 2020. This case-control study included 248 single IA (133 ruptured and 115 unruptured) and 136 multiple IA (65 ruptured, 158 aneurysms; 71 unruptured, 183 aneurysms) patients. IA patients were confirmed by cerebral angiography (computed tomography, magnetic resonance angiography, and digital subtraction angiography) or detected during surgery. IA patients with autosomal dominant polycystic nephropathy, Marfan's syndrome, other autosomal dominant hereditary diseases, other cerebrovascular diseases, and first- or second-degree relatives diagnosed with IA or aSAH disease were excluded. Moreover, patients with IA who had ruptured aneurysms were classified into the ruptured group. The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). The current project was approved by the Ethics Committee of Central South University (permit No. CTXY-150002-1), and the other hospital (Hunan People's Hospital) was informed and agreed the study. All patients provided informed consent.

Single nucleotide polymorphism (SNP) selection and genotyping

SNPs were selected based on tag SNPs and the Genome Variation Server 150 (<http://gvs.gs.washington.edu/>)

Table 1 General demographic and clinical characteristics of the subjects

Variables	Single IA			Multiple IAs		
	Ruptured (n=133)	Unruptured (n=115)	P	Ruptured (n=65)	Unruptured (n=71)	P
Age (years), mean \pm SD	58.31 \pm 10.8	56.18 \pm 9.8	0.109	55.42 \pm 11.23	57.74 \pm 10.64	0.218
Female, n (%)	95 (71.4)	71 (61.7)	0.106	52 (80.0)	49 (69.0)	0.143
Smoking, n (%)	23 (17.3)	19 (16.5)	0.872	7 (10.8)	10 (14.1)	0.559
Drinking, n (%)	16 (12.0)	8 (7.0)	0.178	5 (7.7)	7 (9.9)	0.656
Hypertension, n (%)	64 (48.1)	63 (54.8)	0.295	39 (60.0)	42 (59.2)	0.920
Diabetes, n (%)	6 (4.5)	9 (7.8)	0.275	3 (4.6)	6 (8.5)	0.580
Hyperlipidemia, n (%)	4 (3.0)	9 (7.8)	0.090	3 (4.6)	6 (8.5)	0.580
Intracranial aneurysm, n	133	115		158	183	
Shape of the aneurysm, n (%)			<0.001*			<0.001*
Regular	112 (84.2)	112 (97.4)		128 (81.0)	175 (95.6)	
Irregular	21 (15.8)	3 (2.6)		30 (19.0)	8 (4.4)	
Location, n (%)			0.001*			0.055
Internal carotid artery	53 (39.8)	60 (52.2)		78 (49.4)	96 (52.5)	
Anterior cerebral artery	9 (6.8)	6 (5.2)		14 (8.9)	10 (5.5)	
Middle cerebral artery	21 (15.8)	24 (20.9)		34 (21.5)	35 (19.1)	
Posterior cerebral artery	1 (0.8)	1 (0.9)		1 (0.6)	12 (6.6)	
Anterior communicating artery	34 (25.6)	7 (6.1)		11 (7.0)	9 (4.9)	
Posterior communicating artery	11 (8.3)	7 (6.1)		9 (5.7)	7 (3.8)	
Vertebral basilar artery	4 (3.0)	10 (8.7)		5 (3.2)	11 (6.0)	
Others	0	0		6 (3.8)	3 (1.6)	

*, P<0.05. SD, standard deviation.

GVS150/index.jsp). After screening SNPs by name, preference was given to mutations associated with IA, SNPs located in the functional exon region, or those covering many other loci. Finally, we selected 27 SNPs from the *LOX* family genes to be included (Table S1).

Fasting blood samples (5–10 mL) of each participant were obtained in the morning before the treatment, placed in EDTAK2 anticoagulant tubes (10 mL), and refrigerated at 4 °C. A blood genomic DNA extraction kit (Tiangen Biochemical Technology Co. Ltd., Beijing, China) was used for DNA extraction and refrigerated at –80 °C. Genotyping was conducted using the MassARRAY iPLEX platform (Agena Bioscience Inc., San Diego, CA, USA). The primers were designed using Assay Design 3.1 software (Sequenom, San

Diego, CA, USA), as detailed in Table 1. The mixture for the polymerase chain reaction (PCR) amplification reaction included dddH₂O (1.8 μ L), 10 \times PCR buffer solution (0.5 μ L), MgCl₂ (0.4 μ L), deoxynucleoside triphosphate (0.1 μ L), Taq polymerase (0.2 μ L), PCR primer (1 μ L), and DNA sample (1 μ L). The PCR amplification was conducted in the following steps: pre-denaturation for 2 min at 95 °C, followed by 45 cycles of 30 s at 9 °C, 30 s at 56 °C, 60 s at 72 °C, and finally extension for 5 min at 72 °C. The final products were stored at 25 °C until further use. After shrimp alkaline phosphatase, single nucleotide extension, resin desalination steps, and the matrix assisted laser desorption ionization time-of flight mass spectrometry reaction, MassArray TYPER 4.0 software (Sequenom, San Diego,

CA, USA) was used to interpret the genotype of each sample target site.

Statistical analysis

Statistical analysis was performed using SPSS (version 23.0; IBM, Armonk, NY, USA). Data following a normal distribution were described using mean \pm standard deviation. Normally distributed continuous variables were compared using *t*-tests, and non-normally distributed variables were compared using Mann-Whitney U tests. Categorical variables were compared between the two groups using chi-square or Fisher's exact tests. The association between *LOX* family gene polymorphisms and the risk of IA rupture was evaluated by odds ratios (ORs) and 95% confidence intervals (CIs) using logistic regression in additive, recessive, and dominant models. Differences were considered statistically significant at $P < 0.05$.

Results

Characteristics of the participants

The basic characteristics of the participants are listed in *Table 1*. We included 248 single IA patients (133 ruptured and 115 unruptured) and 136 patients with multiple IAs (65 ruptured, 158 aneurysms; 71 unruptured, 183 aneurysms). Among the single and multiple IAs patients, there were no differences in age, sex, smoking status, drinking status, hypertension, diabetes, and hyperlipidemia between the ruptured and unruptured groups ($P > 0.05$), but there were differences in the morphological distribution of IA ($P < 0.05$). The distribution of aneurysm location was different in the single IA ruptured and unruptured groups ($P < 0.05$), but there was no difference in multiple IAs patients ($P > 0.05$).

Associations between *LOX* family gene polymorphisms and the risk of single IA rupture

Univariate analysis revealed that *LOX* rs1800449, *LOX* rs10519694, and *LOXL4* rs3793692 were associated with single IA rupture, which remained significant after adjusting for the shape and location of IA (*Table S2*). *LOX* rs1800449 was associated with the risk of a single IA rupture (recessive model: OR = 5.66, 95% CI = 1.22–26.24, $P = 0.027$). Nevertheless, *LOX* rs10519694 demonstrated a protective effect on single IA rupture under all 3 genetic models (dominant: OR = 0.42, 95% CI = 0.21–0.83, $P = 0.013$;

recessive: OR = 0.16, 95% CI = 0.04–0.65, $P = 0.010$; additive: OR = 0.46, 95% CI = 0.28–0.78, $P = 0.004$). *LOXL4* rs3793692 was associated with single IA rupture in the recessive model (OR = 2.06, 95% CI = 1.11–3.82, $P = 0.022$). These results are shown in *Table 2*.

Associations between *LOX* family gene polymorphisms and the risk of multiple IAs ruptures

Since every patient with multiple IAs had 2 or more IAs, we were unable to adjust for morphological confounders in the multivariate analysis. The univariate analysis results indicated that *LOXL1* rs2165241 was associated with multiple IAs ruptures (dominant model: OR = 2.99, 95% CI = 1.32–6.78, $P = 0.009$; additive model: OR = 2.53, 95% CI = 1.16–5.56, $P = 0.020$). Moreover, *LOXL2* rs1063582 was associated with the risk of multiple IAs ruptures in the recessive model (OR = 4.12, 95% CI = 1.08–15.71, $P = 0.038$). We found that 2 sites in *LOXL3* were significantly associated with multiple IAs ruptures, but the directionality of the function was different. Furthermore, rs17010021 was associated with the risk of multiple IAs ruptures (additive model: OR = 1.72, 95% CI = 1.03–2.89, $P = 0.039$), but rs17010022 may be an effective factor for reducing the risk of multiple IA ruptures (dominant model: OR = 0.41, 95% CI = 0.21–0.82, $P = 0.011$; additive model: OR = 0.51, 95% CI = 0.30–0.85, $P = 0.010$; *Table 3*).

Discussion

The present study extensively explored the associations between *LOX* family gene polymorphisms and the risk of aSAH. We demonstrated that *LOX* and *LOXL4* polymorphisms were associated with single IA rupture, whereas *LOXL1-3* polymorphisms were associated with multiple IAs ruptures, suggesting that members of the *LOX* family may have roles in aSAH.

The *LOX* family can be classified into two groups based on the structure of their N-terminal domains: *LOX* and *LOXL1* have a propeptide at their N-terminal, whereas *LOXL2*, *LOXL3*, and *LOXL4* have 4 scavenger receptor cysteine-rich domains (18). The *LOX* family gene subtypes (*LOX*, *LOXL1-4*) are all amine oxidases and contain a highly conserved C-terminal binding domain that forms a special lysine tyrosylquinone cofactor-moiety after binding to the copper ion cofactor (19). These family genes are critical enzymes that regulate the crosslinking of elastin and collagen and have a regulatory role in ECM assembly

Table 2 Multivariate analysis of the association between *LOX* family gene polymorphisms and single IA rupture

Gene	SNP	Genotype [†]		Dominant model		Recessive model		Additive model	
		Ruptured (n)	Unruptured (n)	OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P
<i>LOX</i>	rs1800449(C>T)	86/36/11	77/36/2	1.11 (0.65–1.91)	0.706	5.66 (1.22–26.24)	0.027*	1.33 (0.85–2.06)	0.212
	rs2956540(G>C)	75/41/17	62/46/7	0.87 (0.52–1.47)	0.610	2.22 (0.87–5.70)	0.096	1.08 (0.73–1.59)	0.713
	rs10519694(C>T)	115/15/3	86/17/12	0.42 (0.21–0.83)	0.013*	0.16 (0.04–0.65)	0.010*	0.46 (0.28–0.78)	0.004*
	rs2303656(G>T)	122/11/0	111/4/0	2.38 (0.71–7.99)	0.160	–	–	2.38 (0.71–7.99)	0.160
	rs763497(A>G)	100/32/1	77/32/6	0.66 (0.37–1.16)	0.148	0.19 (0.02–1.63)	0.130	0.63 (0.38–1.05)	0.076
	rs3900446(A>G)	110/19/4	93/21/1	0.78 (0.40–1.54)	0.475	3.44 (0.36–32.79)	0.283	0.93 (0.52–1.66)	0.793
<i>LOXL1</i>	rs2165241(C>T)	108/23/2	93/20/2	0.87 (0.44–1.70)	0.674	0.92 (0.13–6.72)	0.933	0.89 (0.49–1.60)	0.694
	rs3825942(G>A)	103/26/4	83/30/2	0.76 (0.42–1.38)	0.367	1.90 (0.32–11.37)	0.482	0.86 (0.51–1.45)	0.564
	rs2304721(C>A)	77/45/11	64/39/12	1.09 (0.65–1.84)	0.749	0.78 (0.32–1.89)	0.588	1.00 (0.68–1.48)	0.998
	rs12441130(T>C)	63/54/16	53/43/19	1.03 (0.61–1.73)	0.919	0.73 (0.35–1.54)	0.412	0.94 (0.65–1.35)	0.738
<i>LOXL2</i>	rs2294128(C>T)	105/26/2	93/2/20	1.24 (0.65–2.35)	0.510	0.34 (0.03–3.51)	0.366	1.12 (0.62–2.01)	0.708
	rs7818494(A>G)	84/40/9	74/34/7	0.99 (0.58–1.69)	0.961	1.14 (0.40–3.30)	0.805	1.01 (0.66–1.55)	0.951
	rs4323477(A>G)	32/67/34	26/64/25	1.00 (0.54–1.86)	0.989	1.21 (0.66–2.23)	0.543	1.08 (0.74–1.57)	0.699
	rs7818416(G>A)	44/66/23	33/59/23	0.79 (0.45–1.39)	0.418	0.90 (0.47–1.74)	0.761	0.87 (0.60–1.27)	0.479
	rs1063582(G>T)	82/47/4	72/38/5	1.10 (0.64–1.86)	0.738	0.57 (0.13–2.49)	0.455	1.01 (0.64–1.61)	0.961
	rs2280936(C>G)	86/43/4	67/42/6	0.75 (0.44–1.27)	0.286	0.39 (0.10–1.59)	0.188	0.73 (0.46–1.15)	0.171
	rs2294133(C>T)	81/39/13	67/35/13	0.86 (0.51–1.46)	0.579	0.86 (0.37–2.00)	0.725	0.90 (0.61–1.31)	0.575
	rs2280935(A>C)	53/57/23	41/61/13	0.81 (0.48–1.38)	0.445	1.87 (0.88–3.96)	0.102	1.06 (0.73–1.54)	0.754
	rs1010156(T>C)	45/63/25	30/55/30	0.66 (0.38–1.17)	0.154	0.64 (0.35–1.20)	0.166	0.73 (0.51–1.05)	0.088
	rs142252012(G>A)	129/4/0	112/3/0	1.04 (0.21–5.14)	0.958	–	–	1.04 (0.21–5.14)	0.958
<i>LOXL3</i>	rs715407(T>G)	90/41/2	71/38/6	0.85 (0.49–1.45)	0.540	0.31 (0.06–1.58)	0.158	0.79 (0.49–1.25)	0.310
	rs6707302(C>T)	94/38/1	74/35/6	0.81 (0.47–1.40)	0.446	0.16 (0.02–1.34)	0.091	0.73 (0.45–1.19)	0.205
	rs17010021(T>A)	57/63/13	54/49/12	1.00 (0.60–1.68)	0.999	0.70 (0.29–1.69)	0.422	0.93 (0.62–1.39)	0.717
	rs17010022(C>G)	57/63/13	53/49/13	1.25 (0.74–2.10)	0.406	0.89 (0.39–2.05)	0.780	1.10 (0.75–1.64)	0.621
<i>LOXL4</i>	rs3793692(G>A)	25/69/39	26/68/21	1.29 (0.68–2.46)	0.432	2.06 (1.11–3.82)	0.022*	1.48 (1.00–2.19)	0.051
	rs1983864(G>T)	40/73/20	32/55/28	0.93 (0.53–1.65)	0.806	0.58 (0.30–1.13)	0.108	0.81 (0.56–1.18)	0.274
	rs7077266(G>T)	95/37/1	75/36/4	0.70 (0.40–1.22)	0.208	0.26 (0.03–2.38)	0.233	0.68 (0.41–1.13)	0.137

[†], genotype presented as wild type/heterozygous/homozygous; *, P<0.05. IA, intracranial aneurysm; SNP, single nucleotide polymorphism; OR, odds ratio; CI, confidence interval; –, not available.

(20,21), while the dysregulation of ECM may disrupt the function or structure of the arterial wall, and may be a risk factor in the pathogenesis of aSAH (22,23). Therefore, they are plausible functional candidates for exploring the associations with aSAH.

The *LOX* gene is located on chromosome 5q23.3–31.2.

Being a copper amine oxidase, *LOX* initiates the covalent cross-linking of collagen and elastin by condensing the oxidized peptidyl α -amino adipic- δ -semialdehyde with neighboring peptidyl aldehydes, thereby consolidating the collagen and elastin fibers of the ECM (13,24). Genetic mouse models for *LOX* have also demonstrated its

Table 3 Univariate analysis of the association between LOX family gene polymorphisms and multiple IAs ruptures

Gene	SNP	Genotype [†]		Dominant model		Recessive model		Additive model	
		Ruptured (n)	Unruptured (n)	OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P
LOX	rs1800449(C>T)	35/25/5	39/28/4	1.05 (0.53–2.05)	0.899	1.40 (0.36–5.44)	0.631	1.09 (0.63–1.87)	0.767
	rs2956540(G>C)	27/27/11	31/33/7	1.09 (0.55–2.15)	0.803	1.86 (0.68–5.14)	0.23	1.22 (0.74–1.99)	0.437
	rs10519694(C>T)	53/9/3	59/9/3	1.11 (0.46–2.69)	0.812	1.10 (0.21–5.64)	0.912	1.08 (0.56–2.08)	0.824
	rs2303656(G>T)	55/10/0	60/11/0	0.99 (0.39–2.52)	0.986	–	–	0.99 (0.39–2.52)	0.986
	rs763497(A>G)	45/16/4	51/17/3	1.13 (0.54–2.37)	0.740	1.49 (0.32–6.91)	0.613	1.15 (0.64–2.06)	0.646
	rs3900446(A>G)	51/14/0	51/17/2	0.70 (0.32–1.54)	0.374	–	–	0.65 (0.31–1.35)	0.248
LOXL1	rs2165241(C>T)	43/23/0	60/10/1	2.99 (1.32–6.78)	0.009*	–	–	2.53 (1.16–5.56)	0.020*
	rs3825942(G>A)	46/18/1	52/16/3	1.13 (0.53–2.39)	0.749	0.35 (0.04–3.49)	0.374	0.99 (0.52–1.89)	0.981
	rs2304721(C>A)	43/21/1	39/28/4	0.62 (0.31–1.25)	0.183	0.26 (0.03–2.41)	0.236	0.61 (0.33–1.13)	0.117
	rs12441130(T>C)	28/31/6	36/26/9	1.36 (0.69–2.67)	0.374	0.70 (2.24–2.09)	0.523	1.10 (0.67–1.81)	0.717
LOXL2	rs2294128(C>T)	47/17/1	53/17/1	1.13 (0.53–2.42)	0.757	1.09 (0.07–17.85)	0.950	1.11 (0.55–2.24)	0.765
	rs7818494(A>G)	44/19/2	39/28/4	0.58 (0.29–1.17)	0.129	0.53 (0.09–3.01)	0.475	0.63 (0.34–1.14)	0.126
	rs4323477(A>G)	21/30/14	18/36/17	0.71 (0.34–1.50)	0.371	0.87 (0.39–1.95)	0.738	0.83 (0.52–1.34)	0.446
	rs7818416(G>A)	21/28/16	21/37/13	0.88 (0.43–1.82)	0.731	1.46 (0.64–3.32)	0.371	1.07 (0.67–1.72)	0.771
	rs1063582(G>T)	39/16/10	43/25/3	1.02 (0.52–2.04)	0.947	4.12 (1.08–15.71)	0.038*	1.31 (0.78–2.18)	0.306
	rs2280936(C>G)	40/20/5	48/20/3	1.30 (0.65–2.64)	0.460	1.89 (0.43–8.24)	0.397	1.30 (0.74–2.29)	0.356
	rs2294133(C>T)	46/17/2	42/25/4	0.60 (0.29–1.22)	0.158	0.53 (0.09–3.01)	0.475	0.64 (0.35–1.18)	0.153
	rs2280935(A>C)	33/27/5	28/32/11	0.63 (0.32–1.25)	0.185	0.46 (1.15–1.39)	0.166	0.65 (0.39–1.09)	0.102
LOXL3	rs1010156(T>C)	11/41/13	17/38/16	1.55 (0.66–3.61)	0.314	0.86 (0.38–1.96)	0.719	1.11 (0.66–1.87)	0.687
	rs142252012(G>A)	64/1/0	68/3/0	0.35 (0.04–3.49)	0.374	–	–	0.35 (0.04–3.49)	0.374
	rs715407(T>G)	44/20/1	47/22/2	0.94 (0.46–1.91)	0.853	0.54 (0.05–6.09)	0.617	0.90 (0.47–1.72)	0.757
	rs6707302(C>T)	46/18/1	51/18/2	1.05 (0.50–2.22)	0.891	0.54 (0.05–6.09)	0.617	0.99 (0.51–1.92)	0.980
	rs17010021(T>A)	22/31/12	33/33/5	1.70 (0.85–3.40)	0.135	2.99 (0.99–9.02)	0.052	1.72 (1.03–2.89)	0.039*
LOXL4	rs17010022(C>G)	38/22/5	26/33/12	0.41 (0.21–0.82)	0.011*	0.41 (0.14–1.24)	0.113	0.51 (0.30–0.85)	0.010*
	rs3793692(G>A)	18/31/16	17/38/16	0.82 (0.38–1.78)	0.618	1.12 (0.51–2.48)	0.775	0.97 (0.60–1.56)	0.890
	rs1983864(G>T)	27/31/7	27/33/11	0.86 (0.43–1.72)	0.676	0.66 (0.24–1.82)	0.419	0.84 (0.51–1.38)	0.480
	rs7077266(G>T)	48/14/3	53/17/1	1.04 (0.48–2.25)	0.915	3.39 (0.34–33.41)	0.296	1.16 (0.60–2.24)	0.649

[†], genotype presented as wild type/heterozygous/homozygous; *, P<0.05. IA, intracranial aneurysm; SNP, single nucleotide polymorphism; OR, odds ratio; CI, confidence interval; –, not available.

significant contribution to the cardiovascular system (25,26). In the present study, significant associations between LOX (rs1800449 and rs10519694) and single IA rupture were detected. Similar to our previous study, it was found that LOX was associated with IA susceptibility (14), but these results are inconsistent with those of a previous study

by Hong *et al.*, who conducted a case-control study with 41 ruptured and 39 unruptured IA patients in a Korean population and showed that LOX may not be a susceptibility gene for IA rupture (17). We found that population heterogeneity may be the reason for the discordance between these 2 countries, and minor allele frequency in the

2 sites was discrepant between these 2 populations.

The *LOXL1* gene is located on chromosome 15q24.1. The homogeneity of *LOX* and *LOXL1* has been found to be as high as 88%, so their functions are similar (27). The pro-sequence contained by *LOX* and *LOXL1* can directly interact with the ECM to direct these enzyme deposits on the elastic tissues (28). The distinction of *LOXL1* from *LOX* is that *LOXL1* specifically locates at the elastic formation site and interacts with fibulin-5. Mice deficient in *LOXL1* did not deposit normal elastic fibers postpartum, thus demonstrating their specific role in elastogenesis (29). Recent studies have indicated that *LOXL1* may also have a role in type II collagen formation and suppression, as well as the promotion of tumorigenesis (30,31). *LOXL1* deficiency has been associated with pseudoexfoliation syndrome, idiopathic pulmonary fibrosis, and aneurysms (28,32). Our present study demonstrated that *LOXL1* rs2165241 was associated with multiple IAs ruptures. This is different from our previous study, which did not find an association between *LOXL1* and IAs susceptibility (14). This may be due to the fact that the pathobiology leading to IAs formation and its rupture are not exactly the same, causing the existing IA rupture (aSAH) is a separate process from an IA formation (12). Therefore, the association between *LOXL1* polymorphisms and aSAH and its mechanism needs to be further explored.

LOXL2 is located on chromosome 8p21.3 and its protein products are helpful in maintaining the integrity and stability of the vascular wall. Thus, *LOXL2* may play a role in susceptibility to IA rupture (33). The unbiased proteomic analysis demonstrated that *LOXL2* could accelerate vascular sclerosis by promoting matrix stiffness and vascular smooth muscle stiffness and contractility (34), and an additional study has identified that *LOXL2* polymorphisms are associated with blood pressure (33). Increased vascular stiffness and high blood pressure are independent risk factors for cardiovascular diseases, such as stroke and subarachnoid hemorrhage (35). Akagawa *et al.* conducted an association study to systematically screen the *LOX* family genes in 402 IA patients and 462 controls from a Japanese population and found that *LOXL2* rs1010156 was associated with FIA (15). Using whole-exome sequencing, a significant association was also found with *LOXL2* in FIA patients from a Chinese population (16). Our previous research also found that *LOXL2* is associated with IA (14). Similarly, our present results also demonstrated that *LOXL2* is associated with IA rupture but with multiple IAs, not total IA or single IA rupture. If the same gene has different roles in the

process of single and multiple IAs ruptures, this may be due to the higher rupture risk in patients with multiple IAs than in patients with a single IA (36); however, the mechanism of *LOXL2* in IA rupture is unclear, and further studies are required.

The *LOXL3* gene is located on chromosome 2p13.1 and its expression level has been found to be high in the heart, spleen, lung, aorta, and coronary arteries (37). *LOXL3* showed beta-aminopropionitrile inhibition of amine oxidase activity towards elastin and collagen. The highest activity was observed for type VIII collagen, which is a network collagen mainly expressed in vascular endothelial cells and smooth muscle cells, possibly having a role in the maintenance of vessel wall integrity (38). Mouse models have also described the oxidative effect of *LOXL3* on ECM fibronectin (39). In the present study, we found that *LOXL3* was associated with multiple IAs ruptures, suggesting that a variant of *LOXL3* may have a role in aSAH, but the mechanism of function needs to be further studied.

The *LOXL4* gene is located on chromosome 10q24.2, and contains an additional 13 amino acid inserts that differ from *LOXL2* and *LOXL3*. *LOXL4* is present in multiple human tissues, including the lung, liver, heart, brain, and colon (40). It has been found to be abnormally expressed in several tumors, and the potential biological function of *LOXL4* has been extended to the remodeling of the vascular ECM (41). Although our current results suggest that *LOXL4* may have a role in single IA rupture, whether it leads to IA rupture by affecting the remodeling of ECM or other methodologies is unclear; therefore, future studies are needed.

Our study had several limitations. First, the sample size was relatively small, which may have contributed to false associations due to limited statistical power; therefore, it is important to use larger studies to further verify the association between *LOX* family genes and aSAH. Second, we could not modify the morphological factors for multivariate analysis due to patients with 2 or more aneurysms in the multiple IA group; however, irregular aneurysms are more likely to rupture than regular aneurysms, and irregular aneurysms were more common in the ruptured group. Hence, we suggest that the univariate analysis results of multiple IA ruptures may provide a reference for multiple IA etiological research. Third, functional studies on susceptibility genes of IA rupture were not conducted, and we did not explore the specific mechanisms of aSAH; therefore, further research is needed to clarify the mechanism of function in the future. Despite

the above limitations, our present work provides evidence of the association between *LOX* family gene polymorphisms and aSAH. This may provide a basis for management and treatment of aSAH.

In summary, after exploring the association between *LOX* family genes and single and multiple IAs ruptures, we found that *LOX* and *LOXL4* may be associated with single IA rupture, while *LOXLI-3* were associated with multiple IAs ruptures in this Chinese sample. This suggests that the expression of *LOX* family genes may be associated with aSAH, which should be further studied and explored.

Acknowledgments

The authors thank all participants in this study.

Funding: This work was supported by the National Nature Science Foundation, China (grant No. 81502881) and the Hunan Province Nature Science Foundation, China (Nos. 2021JJ30911, 2021JJ31077 and 2021JJ31121).

Footnote

Reporting Checklist: The authors have completed the MDAR reporting checklist. Available at <https://atm.amegroups.com/article/view/10.21037/atm-22-3484/rc>

Data Sharing Statement: Available at <https://atm.amegroups.com/article/view/10.21037/atm-22-3484/dss>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://atm.amegroups.com/article/view/10.21037/atm-22-3484/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. This study was reviewed and approved by The Ethics Committee at Central South University (permit No. CTXY-150002-1), and the other hospital (Hunan People's Hospital) was informed and agreed the study. The patients/participants provided their written informed consent to participate in this study, and the study was conducted in accordance with the Declaration of Helsinki (as revised in 2013).

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons

Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Jung KH. New Pathophysiological Considerations on Cerebral Aneurysms. *Neurointervention* 2018;13:73-83.
- Jeon TY, Jeon P, Kim KH. Prevalence of Unruptured Intracranial Aneurysm on MR Angiography. *Korean J Radiol* 2011;12:547-53.
- Li M-H, Chen S-W, Li Y-D, et al. Prevalence of unruptured cerebral aneurysms in Chinese adults aged 35 to 75 years: a cross-sectional study. *Ann Intern Med* 2013;159:514-21.
- Vlak MH, Algra A, Brandenburg R, et al. Prevalence of unruptured intracranial aneurysms, with emphasis on sex, age, comorbidity, country, and time period: a systematic review and meta-analysis. *Lancet Neurol* 2011;10:626-36.
- McDowell MM, Zhao Y, Kellner CP, et al. Demographic and clinical predictors of multiple intracranial aneurysms in patients with subarachnoid hemorrhage. *J Neurosurg* 2018;128:961-8.
- Roquer J, Cuadrado-Godia E, Guimaraens L, et al. Short- and long-term outcome of patients with aneurysmal subarachnoid hemorrhage. *Neurology* 2020;95:e1819-e29.
- Greving JP, Wermer MJ, Brown RD, Jr., et al. Development of the PHASES score for prediction of risk of rupture of intracranial aneurysms: a pooled analysis of six prospective cohort studies. *Lancet Neurol* 2014;13:59-66.
- Theodotou CB, Snelling BM, Sur S, et al. Genetic associations of intracranial aneurysm formation and sub-arachnoid hemorrhage. *Asian J Neurosurg* 2017;12:374-81.
- Kang H, Peng T, Qian Z, et al. Impact of hypertension and smoking on the rupture of intracranial aneurysms and their joint effect. *Neurol Neurochir Pol* 2015;49:121-5.
- Zuurbier C, Greving JP, Rinkel G, et al. Higher risk of intracranial aneurysms and subarachnoid haemorrhage in siblings of families with intracranial aneurysms. *Eur Stroke J* 2020;5:73-7.
- Ma Z, Mao C, Jia Y, et al. Extracellular matrix dynamics in vascular remodeling. *Am J Physiol Cell Physiol*

- 2020;319:C481-c99.
12. Juhana F, Riikka T, Anders P, et al. Saccular intracranial aneurysm: pathology and mechanisms. *Acta Neuropathologica* 2012;123:773-86.
 13. Csiszar K. Lysyl oxidases: a novel multifunctional amine oxidase family. *Prog Nucl Acid Res Mol Biol* 2001;70:1-32.
 14. Luo C, Hu C, Li B, et al. Polymorphisms in Lysyl Oxidase Family Genes Are Associated With Intracranial Aneurysm Susceptibility in a Chinese Population. *Front Endocrinol (Lausanne)* 2021;12:642698.
 15. Akagawa H, Narita A, Yamada H. Systematic screening of lysyl oxidase-like (LOXL) family genes demonstrates that LOXL2 is a susceptibility gene to intracranial aneurysms. *Hum Genet* 2007;121:377-87.
 16. Wu YQ, Li Z, Shi Y, et al. Exome Sequencing identifies LOXL2 mutation as a cause of familial intracranial aneurysm. *World Neurosurg* 2017;109:e812-e8.
 17. Hong EP, Pyeong JJ, Sung-Eun K, et al. A Novel Association between Lysyl Oxidase Gene Polymorphism and Intracranial Aneurysm in Koreans. *Yonsei Med J* 2017;58:1006-11.
 18. Finney J, Moon HJ, Ronnebaum T, et al. Human copper-dependent amine oxidases. *Arch Biochem Biophys* 2014;546:19-32.
 19. Setargew YFI, Wyllie K, Grant RD, et al. Targeting Lysyl Oxidase Family Mediated Matrix Cross-Linking as an Anti-Stromal Therapy in Solid Tumours. *Cancers (Basel)* 2021;13:491.
 20. Yang N, Cao DF, Yin XX, et al. Lysyl oxidases: Emerging biomarkers and therapeutic targets for various diseases. *Biomed Pharmacother* 2020;131:110791.
 21. Nguyen XX, Nishimoto T, Takihara T, et al. Lysyl oxidase directly contributes to extracellular matrix production and fibrosis in systemic sclerosis. *Am J Physiol Lung Cell Mol Physiol* 2021;320:L29-L40.
 22. Ruigrok YM, Rinkel GJE, Slot RVT, et al. Evidence in favor of the contribution of genes involved in the maintenance of the extracellular matrix of the arterial wall to the development of intracranial aneurysms. *Hum Mol Genet* 2006;15:3361-8.
 23. Rempe RG, Hartz AMS, Bauer B. Matrix metalloproteinases in the brain and blood-brain barrier: Versatile breakers and makers. *J Cereb Blood Flow Metab* 2016;36:1481-507.
 24. Kagan HM, Li W. Lysyl oxidase: properties, specificity, and biological roles inside and outside of the cell. *J Cell Biochem* 2003;88:660-72.
 25. Martinez-Gonzalez J, Varona S, Canes L, et al. Emerging Roles of Lysyl Oxidases in the Cardiovascular System: New Concepts and Therapeutic Challenges. *Biomolecules* 2019;9:610.
 26. Joni MM, Juha R, Hilkka T, et al. Inactivation of the Lysyl Oxidase Gene *Lox* Leads to Aortic Aneurysms, Cardiovascular Dysfunction, and Perinatal Death in Mice. *Circulation* 2002;106:2503-9.
 27. Al-U'datt D, Allen BG, Nattel S. Role of the lysyl oxidase enzyme family in cardiac function and disease. *Cardiovasc Res* 2019;115:1820-37.
 28. Greene AG, Eivers SB, Dervan EWJ, et al. Lysyl Oxidase Like 1: Biological roles and regulation. *Exp Eye Res* 2020;193:107975.
 29. Liu X, Zhao Y, Gao J, et al. Elastic fiber homeostasis requires lysyl oxidase like 1 protein. *Nat Genet* 2004;36:178-82.
 30. Alsofi L, Daley E, Hornstra I, et al. Sex-Linked Skeletal Phenotype of Lysyl Oxidase Like-1 Mutant Mice. *Calcif Tissue Int* 2016;98:172-85.
 31. Zeltz C, Pasko E, Cox TR, et al. LOXL1 Is Regulated by Integrin $\alpha 11$ and Promotes Non-Small Cell Lung Cancer Tumorigenicity. *Cancers (Basel)* 2019;11:705.
 32. Li X, He J, Sun J. LOXL1 gene polymorphisms are associated with exfoliation syndrome/exfoliation glaucoma risk: An updated meta-analysis. *PLoS One* 2021;16:e0250772.
 33. Gao Y, Qi GX, Jia ZM, et al. Prediction of marker genes associated with hypertension by bioinformatics analyses. *Int J Mol Med* 2017;40:137-45.
 34. Steppan J, Wang H, Bergman Y, et al. Lysyl oxidase-like 2 depletion is protective in age-associated vascular stiffening. *Am J Physiol Heart Circ Physiol* 2019;317:H49-h59.
 35. Backes D, Rinkel GJ, Laban KG, et al. Patient- and Aneurysm-Specific Risk Factors for Intracranial Aneurysm Growth: A Systematic Review and Meta-Analysis. *Stroke* 2016;47:951-7.
 36. Sato H, Kamide T, Kikkawa Y, et al. Clinical Characteristics of Ruptured Intracranial Aneurysm in Patients with Multiple Intracranial Aneurysms. *World Neurosurg* 2021;149:e935-e41.
 37. Dudakova L, Sasaki T, Liskova P, et al. The presence of lysyl oxidase-like enzymes in human control and keratoconic corneas. *Histol Histopathol* 2016;31:63-71.
 38. Lee JE, Kim Y. A tissue-specific variant of the human lysyl oxidase-like protein 3 (LOXL3) functions as an amine oxidase with substrate specificity. *J Biol Chem*

- 2006;281:37282-90.
39. Kraft-Sheleg O, Zaffryar-Eilot S, Genin O, et al. Localized LoxL3-Dependent Fibronectin Oxidation Regulates Myofiber Stretch and Integrin-Mediated Adhesion. *Dev Cell* 2016;36:550-61.
 40. Asuncion L, Fogelgren B, Fong KS, et al. A novel human lysyl oxidase-like gene (LOXL4) on chromosome 10q24 has an altered scavenger receptor cysteine rich domain. *Matrix Biol* 2001;20:487-91.
 41. Busnadiago O, González-Santamaría J, Lagares D, et al. LOXL4 is induced by transforming growth factor β 1 through Smad and JunB/Fra2 and contributes to vascular matrix remodeling. *Mol Cell Biol* 2013;33:2388-401.

(English Language Editor: C. Betlazar-Maseh)

Cite this article as: Luo C, Li B, Liu C, Dong R, Hu C, Liu J, Hu L, Liao X, Zhou J, Xu L, Liu S, Yuan D, Jiang W, Yan J, Li Y. Lysyl oxidase family gene polymorphisms and risk of aneurysmal subarachnoid hemorrhage: a case-control study. *Ann Transl Med* 2022;10(17):925. doi: 10.21037/atm-22-3484