

BRCA1/2 Mutation Status Impact on Autophagy and Immune Response: Unheralded Target

Susan Morand, BS,¹ Laura Stanbery, PhD,² Adam Walter, MD,³ Rodney P. Rocconi , MD,⁴ John Nemunaitis , MD^{2,*}

¹Department of Internal Medicine, University of Toledo, Toledo, OH, USA; ²Gradalis, Inc, Carrollton, TX, USA; ³Promedica Health System, Toledo, OH, USA and ⁴University of South Alabama - Mitchell Cancer Institute, Mobile, AL, USA

*Correspondence to: John Nemunaitis, MD, Gradalis, Inc, 2545 Golden Bear Carrollton, TX 75006, USA (e-mail: Johnnemunaitis@gmail.com).

Abstract

BRCA1 and possibly BRCA2 proteins may relate to the regulation of autophagy. Autophagy plays a key role in immune response from both a tumor and immune effector cell standpoint. In cells with BRCA mutations, increased autophagy leads to elevated expression of major histocompatibility complex class II but may cause subclonal neoantigen presentation, which may impair the immune response related to clonal neoantigen visibility. We review evidence of BRCA1/2 regulation of autophagy, immune response, and antigen presentation.

Autophagy refers to the process by which a cell consumes its own constituents (1). It is increasingly evident that basal levels of autophagy may play a more important role than previously thought regarding sensitivity and/or resistance to immunotherapy, maintaining homeostasis by recycling cytosolic material, regulating metabolism, and eliminating harmful free radicals (2). Autophagy also plays a key role as a tumor suppressor (3-6). In the absence of autophagy, proto(oncoproteins) accumulate, exerting effects on cell growth, progression through the cell cycle, or angiogenesis, among other hallmarks (7-12). Similarly, impaired autophagy allows defective organelles to accumulate. Particularly, the accumulation of mitochondria in the absence of mitophagy results in increased reactive oxygen species and further damages DNA (13-15). Several studies demonstrate increased autophagy in cancer stem cells, during the Warburg effect, in anoikis and metastasis, and in resistance to chemotherapy (16-19). However, autophagy has a key role in major histocompatibility complex (MHC) processing (20-22) and permits the mounting of intracellular material onto MHC class II (MHC-II), which is traditionally thought of as the bearer of extracellular threats (23). Furthermore, autophagy communicates closely with the exosomal and endosomal pathway, thereby modulating the intercellular exchange of material, including tumor antigens and a further relationship to MHC-I display along with other immunomodulatory molecule expression (1,24). Together, these actions have important consequences on antitumor immunity.

Autophagy Pathway

Autophagy begins with 3 principal steps: initiation, expansion, and formation of the autophagosome (1). During initiation, a phagophore is derived from the endoplasmic reticulum. In expansion, the phagophore approaches cytosolic material and expands. Finally, the phagophore completely envelopes the cytosolic contents to form the autophagosome. To complete autophagy, the autophagosome may fuse with a lysosome, which subsequently degrades the material and makes it available for processing via MHC-II. Alternatively, the autophagosome may fuse with an endosome to form an amphisome, which fuses with the lysosome to complete the process (1). Otherwise, it may engage in exosome transfer, in which it secretes its material to the extracellular matrix (1,24-26). Depending on the derivate cell, these exosomes may be referred to as tumor exosomes (TEXs) or dendritic cell exosomes (1).

The machinery for autophagy and exosome production overlap considerably. In the total absence of one process, the other process is incapable of occurring (24,26). Similarly, when conditions including hypoxia, chemotherapy, and endoplasmic reticulum stress occur, both autophagy and exosomal production are increased in response (24). Homeostasis exists between the 2 pathways, which share endosomes as a substrate (24,26). With activation of autophagy, intravesicular bodies (IVBs) preferentially fuse with autophagosomes to undergo degradation by

Received: 10 June 2020; Revised: 6 August 2020; Accepted: 10 August 2020

© The Author(s) 2020. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

lysosomes, thereby decreasing flux down the exosomal pathway. In contrast, when autophagy is inhibited, intravesicular bodies are shunted toward the exosomal pathway, where they are secreted as TEXs (1,24,26).

Autophagy Role in Immune Response

Tumor cells predominately express MHC-I; however, they can express MHC-II even when the tissue of origin does not (27). Elevated expression of MHC-II by both breast and ovarian tumors, which are typically associated with BRCA1/2 mutation, has been correlated with better prognosis and overall survival advantage (28-30). Autophagy increases the tumor antigen supply of likely subclonal neoantigens for MHC-II presentation while decreasing expression of MHC-I by tumor cells (1,25,31). MHC-II expression of tumor neoantigens is a well-established immune therapeutic development direction (32). However, clonal neoantigen display is critical toward effective anticancer immune response (33). Overdisplay of subclonal neoantigens may dilute focused immune response to clonal neoantigens and even activate immune editing of clonal neoantigen effector cells (34,35). Despite autophagy's role in refreshing the pool of available peptides to be presented by MHC-I (31), autophagy appears to recycle MHC-I more rapidly, resulting in decreased expression duration. Indeed, inhibition of autophagy related to gamma interferon expression augmented MHC-I expression in melanoma (36-38).

Molecular Signaling and BRCA1/2 Function Related to Autophagy Regulation

Several molecular pathways overlap to control autophagy, particularly mTOR and Beclin1/ATG5. Type I PI3K/AKT/mTOR inhibits autophagy, whereas type III PI3K/vps34/Beclin1 activates autophagy (39). New evidence suggests that wild-type BRCA1 and BRCA2 proteins negatively regulate autophagy, with increased autophagy noted in mutant BRCA1/2 gene samples as well as in BRCA1/2 gene silencing studies (40-43). Coimmunoprecipitation studies of BRCA1 with Beclin1 show that when this complex is interrupted via a BRCA1 mutation, the canonical pathway is enhanced because of the free action of Beclin1, and autophagy levels are increased (40). Additionally, in the role that BRCA1 functions as an antioxidant, the absence of functional BRCA1 increases reactive oxygen species, and autophagy increases in response to oxidative stress (41). Finally, BRCA1 interacts with PTEN, an inhibitor of the type I PI3K/AKT/mTOR pathway (44).

BRCA2 has been less thoroughly studied in relation to autophagy. Nonetheless, evidence supports a negative regulatory role of wild-type BRCA2 on autophagy (42,43). Knockdown of BRCA2 via RNA interference in tumors with BRCA1 allelic loss enhanced autophagy and mitophagy, a derivative of autophagy that recycles mitochondria (42). This same study suggested that PARP inhibitors, particularly olaparib whose mechanism depends on interrupted BRCA1/2 function, have enhanced function in the context of increased autophagy (42). It remains unknown if autophagy is similarly upregulated regardless of BRCA gene variant or in tumors that are homologous recombination deficient. It is known that autophagy is required for homologous recombination to take place, and in the absence of autophagy, DNA damage accumulates, resulting in cell death (45).

BRCA1/2 Mutation Role in Autophagy and Cancer Immunity

Autophagy and exosome transport are multifaceted processes, and their role in cancer development, progression, and immunity relates to BRCA1/2 expression and genetic stability (31). BRCA1/2 wild-type expression decreases autophagy, whereas BRCA1/2 mutation enhances autophagy activity (1,40). Tumors with increased autophagy due to defective BRCA1/2 demonstrate increased MHC-II expression of tumor antigens (1,40,42). However, tumors with BRCA1/2 mutation also exhibit increased estimated glomerular filtration rate (EGFR) expression, which decreases MHC expression compared with wild-type tumors (46,47). EGFR also directly regulates autophagy in a context-dependent manner, which may be depend on localization (48). Through association with Beclin1, inactive EGFR can decrease autophagy; however, when localized to the endosome, EGFR can also initiate autophagy (49,50). The complex correlation between autophagy, EGFR, and MHC expression warrants further research.

It is hypothetically possible that germline vs somatic BRCA mutation status may affect immune response differently. In patients with germline mutations, it is presumed that immune and cancer cells would be impacted by altered BRCA functioning, potentially resulting in increased autophagy across the board. It has been shown that BRCA heterozygous mice have decreased white blood cells and lymphocytes. The same study also showed that heterozygous BRCA1 carriers are more at increased risk for chemotherapy-associated hematopoietic complications (51). Although this could be due to increased DNA damage in these cells, it may be linked to autophagy. Further investigation would be needed to validate this hypothesis. Of interest, immune cells with increased autophagy express both increased MHC-I and MHC-II (1,31,52,53). This is explained by increased dendritic cell participation in cross-presentation (1,25,54). Complementary to autophagy, cross-presentation allows extracellular material to be presented by MHC-I (autophagy places intracellular material on MHC-II) (1,25). Moreover, cross-presentation allows DCs to accept tumor antigens from TEXs and to present them on MHC-I and II on the DC surface (1,25). Because DCs express costimulatory molecules, cross-presentation allows antigens from TEXs to elicit a more effective immune response than the same antigen could in the absence of DCs (1,54). In contrast, we presume that patients with somatic BRCA1/2 mutations demonstrate the effect of autophagy induction exclusively in the tumor cells containing the BRCA1/2 mutation.

BRCA1/2 activity on autophagy, however, is complicated. For example, increased autophagy in BRCA1/2 mutant or dysfunctional tumor cells destabilizes the immunological synapse between cytolytic immune cells and their targets, consequently disrupting transfer of cytotoxic molecules (55). In particular, autophagy increases degradation of granzyme B and connexin-43, which are 2 important molecules in the tumor-killing pathway enacted by cytotoxic T lymphocytes (CTLs) and natural killer (NK) cells (55-59). Moreover, inhibition of autophagy restores cytolytic activity in cell populations that are resistant to CTL- and NK-mediated killing (57,60,61). Conversely, upregulation of autophagy decreases CTL- and NK-mediated killing, particularly during the endothelial-mesenchymal transition, which is vital for metastasis (62).

Autophagy also suppresses the immune response indirectly via its effects on the exosomal pathway. Presumably, expressive BRCA wild-type tumor cells have increased exosome secretion

as opposed to BRCA-mutation tumor cells. With low levels of autophagy, homeostasis shifts toward the exosomal pathway, with subsequently increased release of TEXs (24,26). Although TEXs represent a vehicle for transporting tumor antigens to distant antigen presenting cells (APCs), they also have a major role in immunosuppression of the anticancer immune response (54). This is accomplished through expression of immunosuppressive proteins including TGF- β , IL-6, and PGE2 (63-66). Furthermore, TEXs frequently carry miRNAs that suppress expression of immunostimulatory genes in target cells (54,67,68). Other immunosuppressive effects include redirection of myeloid precursors toward myeloid-derived suppressor cells, along with inhibited differentiation of DCs from precursors (63,69). Along these same lines, TEXs prevent the maturation of immature DCs, alter pattern recognition receptors to impair antigen recognition, and reduce DC production of immunostimulatory cytokines including TNF- α and IL-12 (66,70). Furthermore, TEXs induce the CD14⁺ HLA-DR^{-low} monocyte subtype, which suppresses T-cell proliferation and cytotoxicity (64). Beyond monocytes, TEXs negatively impact the development of NK cells and CTLs (71-76). Finally, TEXs induce the regulatory subtypes of T and B cells (77-80). In sum, TEXs have a complicated impact on anticancer immunity, and they are increased in settings with low autophagy. This includes wild-type expression of BRCA1/2 (40-43).

Clinical Relevance

Tumor expression of PD-L1, an immune checkpoint, is inversely related to autophagy (31). With activation of autophagy, PD-L1 levels are decreased. In contrast, inhibition of autophagy results in increased expression of PD-L1 (31). Similarly, approximately 90% of human lung cancer samples with increased PD-L1 expression showed inhibition of autophagy, whereas 83% of tumors with negative PD-L1 had increased autophagy (81). Likewise, blocking PD-L1 disinhibited autophagy, whereas increasing PD-L1 signaling inhibited autophagy (82). Thus, it is expected that BRCA wild-type patients would have increased PD-L1 signaling, whereas BRCA mutant would have decreased PD-L1 signaling. Therefore, BRCA mutation status may relate to clinical benefit of immune checkpoint inhibitor (ICI) therapy.

In the BRCA1/2 wild-type population, for example, patients would likely derive benefit from therapies that introduce tumor antigens to the immune system via an alternative mechanism, for example, use of Vigil, a personalized neoantigen-educating immunotherapy as opposed to checkpoint-inhibitor therapy, which does not modulate neoantigen expression and would have limited activity in lower clonal neoantigen expressive tumors (83-86). Vigil is an experimental therapeutic that educates the immune system via an autologous tumor vaccine along with plasmid DNA-encoding upregulators of the immune system (granulocyte-macrophage colony-stimulating factor) and blocking downregulators of the immune system (furin, which activates TGF- β , a major immunomodulatory molecule) (87). In clinical trials involving melanoma, Ewing sarcoma, ovarian cancer, and other solid tumors, Vigil was well tolerated (83,87-91). In solid tumors, improved clinical outcomes related to Vigil treatment were correlated with γ -IFN-ELISPOT positive response, indicating that Vigil is able to activate a durable immune response. Combination of Vigil followed by immune checkpoint inhibitor may sensitize patients to checkpoint inhibitors by priming T cells to the relevant clonal tumor neoantigens.

Conclusion

Increased tumor cell autophagy (as in BRCA mutant) likely leads to enhanced presentation of subclonal neoantigens to the immune system but impaired cytotoxic killing. Conversely, inhibited tumor autophagy (as in BRCA wild-type) likely causes lower tumor antigen presentation but may preserve clonal neoantigen display supporting increased target-directed susceptibility to cell-mediated destruction. Consequently, BRCA status may impact a patient's response to certain immunotherapies. Going forward, research involving clinical therapeutic measures that involve and/or modulate BRCA1/2 signaling as involved in autophagy control and relationship to antigen presentation is justified.

Funding

The authors have no funding to disclose.

Notes

Role of the funder: Not applicable.

Author contributions: Susan Morand: Contributed to the research and writing of the manuscript. Laura Stanbery: Contributed to the research and writing of the manuscript. Adam Walter: Contributed to the writing and editing of the manuscript. Rodney Rocconi: Contributed to the writing and editing of the manuscript. John Nemunaitis: Initiated review design, contributed to article design, writing, and editing.

Disclosure: The authors have no conflicts of interest to disclose.

Acknowledgments: The authors would like to acknowledge Brenda Marr for her competent and knowledgeable assistance in the preparation of the manuscript.

Prior presentations: None

Data availability

Not applicable.

References

1. You L, Mao L, Wei J, et al. The crosstalk between autophagic and endo-/exosomal pathways in antigen processing for MHC presentation in anticancer T cell immune responses. *J Hematol Oncol*. 2017;10(1):165.
2. Mizushima N, Levine B, Cuervo AM, et al. Autophagy fights disease through cellular self-digestion. *Nature*. 2008;451(7182):1069-1075.
3. Gozuacik D, Kimchi A. Autophagy as a cell death and tumor suppressor mechanism. *Oncogene*. 2004;23(16):2891-2906.
4. Yue Z, Jin S, Yang C, et al. Beclin 1, an autophagy gene essential for early embryonic development, is a haploinsufficient tumor suppressor. *Proc Natl Acad Sci USA*. 2003;100(25):15077-15082.
5. Takamura A, Komatsu M, Hara T, et al. Autophagy-deficient mice develop multiple liver tumors. *Genes Dev*. 2011;25(8):795-800.
6. Marino G, Salvador-Montoliu N, Fueyo A, et al. Tissue-specific autophagy alterations and increased tumorigenesis in mice deficient in Atg4C/autophagin-3. *J Biol Chem*. 2007;282(25):18573-18583.
7. Goussetis DJ, Gounaris E, Wu EJ, et al. Autophagic degradation of the BCR-ABL oncoprotein and generation of antileukemic responses by arsenic trioxide. *Blood*. 2012;120(17):3555-3562.
8. Isakson P, Bjoras M, Boe SO, et al. Autophagy contributes to therapy-induced degradation of the PML/RARA oncoprotein. *Blood*. 2010;116(13):2324-2331.
9. Larrue C, Saland E, Boutzen H, et al. Proteasome inhibitors induce FLT3-ITD degradation through autophagy in AML cells. *Blood*. 2016;127(7):882-892.
10. Li P, Ji M, Lu F, et al. Degradation of AF1Q by chaperone-mediated autophagy. *Exp Cell Res*. 2014;327(1):48-56.
11. Tse W, Deeg HJ, Stirewalt D, et al. Increased AF1q gene expression in high-risk myelodysplastic syndrome. *Br J Haematol*. 2005;128(2):218-220.
12. Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell*. 2000;100(1):57-70.

13. Frank M, Duvezin-Caubet S, Koob S, et al. Mitophagy is triggered by mild oxidative stress in a mitochondrial fission dependent manner. *Biochim Biophys Acta*. 2012;1823(12):2297-2310.
14. Klionsky DJ, Cuervo AM, Dunn WA Jr, et al. How shall I eat thee? *Autophagy*. 2007;3(5):413-416.
15. Lemasters JJ. Selective mitochondrial autophagy, or mitophagy, as a targeted defense against oxidative stress, mitochondrial dysfunction, and aging. *Rejuvenation Res*. 2005;8(1):3-5.
16. Komatsu M, Waguri S, Ueno T, et al. Impairment of starvation-induced and constitutive autophagy in Atg7-deficient mice. *J Cell Biol*. 2005;169(3):425-434.
17. Avivar-Valderas A, Salas E, Bobrovnikova-Marjon E, et al. PERK integrates autophagy and oxidative stress responses to promote survival during extracellular matrix detachment. *Mol Cell Biol*. 2011;31(17):3616-3629.
18. Peng Y-F, Shi Y-H, Ding Z-B, et al. Autophagy inhibition suppresses pulmonary metastasis of HCC in mice via impairing anoikis resistance and colonization of HCC cells. *Autophagy*. 2013;9(12):2056-2068.
19. Kim D-G, Jung KH, Lee D-G, et al. 20 (S)-Ginsenoside Rg3 is a novel inhibitor of autophagy and sensitizes hepatocellular carcinoma to doxorubicin. *Oncotarget*. 2014;5(12):4438-4451.
20. You L, Jin S, Zhu L, et al. Autophagy, autophagy-associated adaptive immune responses and its role in hematologic malignancies. *Oncotarget*. 2017;8(7):12374-12388.
21. Deretic V, Kimura T, Timmins G, et al. Immunologic manifestations of autophagy. *J Clin Invest*. 2015;125(1):75-84.
22. Schmid D, Pypaert M, Münz C. Antigen-loading compartments for major histocompatibility complex class II molecules continuously receive input from autophagosomes. *Immunity*. 2007;26(1):79-92.
23. Li Y, Wang LX, Yang G, et al. Efficient cross-presentation depends on autophagy in tumor cells. *Cancer Res*. 2008;68(17):6889-6895.
24. Xu J, Camfield R, Gorski SM. The interplay between exosomes and autophagy - partners in crime. *J Cell Sci*. 2018;131(15):jcs215210.
25. Blum JS, Wearsch PA, Cresswell P. Pathways of antigen processing. *Annu Rev Immunol*. 2013;31(1):443-473.
26. Gudbergsson JM, Johnsen KB. Exosomes and autophagy: rekindling the vesicular waste hypothesis. *J Cell Commun Signal*. 2019;13(4):443-450.
27. Axelrod ML, Cook RS, Johnson DB, et al. Biological consequences of MHC-II expression by tumor cells in cancer. *Clin Cancer Res*. 2019;25(8):2392-2402.
28. Oldford SA, Robb JD, Codner D, et al. Tumor cell expression of HLA-DM associates with a Th1 profile and predicts improved survival in breast carcinoma patients. *Int Immunol*. 2006;18(11):1591-1602.
29. Forero A, Li Y, Chen D, et al. Expression of the MHC class II pathway in triple-negative breast cancer tumor cells is associated with a good prognosis and infiltrating lymphocytes. *Cancer Immunol Res*. 2016;4(5):390-399.
30. Callahan MJ, Nagymanyoki Z, Bonome T, et al. Increased HLA-DMB expression in the tumor epithelium is associated with increased CTL infiltration and improved prognosis in advanced-stage serous ovarian cancer. *Clin Cancer Res*. 2008;14(23):7667-7673.
31. Folkerts H, Hilgendorf S, Vellenga E, et al. The multifaceted role of autophagy in cancer and the microenvironment. *Med Res Rev*. 2019;39(2):517-560.
32. Thibodeau J, Bourgeois-Daigneault MC, Lapointe R. Targeting the MHC class II antigen presentation pathway in cancer immunotherapy. *Oncimmunology*. 2012;1(6):908-916.
33. McGranahan N, Furness AJ, Rosenthal R, et al. Clonal neoantigens elicit T cell immunoreactivity and sensitivity to immune checkpoint blockade. *Science*. 2016;351(6280):1463-1469.
34. Keenan TE, Burke KP, Van Allen EM. Genomic correlates of response to immune checkpoint blockade. *Nat Med*. 2019;25(3):389-402.
35. Milo I, Bedora-Faure M, Garcia Z, et al. The immune system profoundly restricts intratumor genetic heterogeneity. *Sci Immunol*. 2018;3(29):eaat1435.
36. Li B, Lei Z, Lichty BD, et al. Autophagy facilitates major histocompatibility complex class I expression induced by IFN-gamma in B16 melanoma cells. *Cancer Immunol Immunother*. 2010;59(2):313-321.
37. Loi M, Muller A, Steinbach K, et al. Macroautophagy proteins control MHC class I levels on dendritic cells and shape anti-viral CD8(+) T cell responses. *Cell Rep*. 2016;15(5):1076-1087.
38. Parekh VV, Wu L, Boyd KL, et al. Impaired autophagy, defective T cell homeostasis, and a wasting syndrome in mice with a T cell-specific deletion of Vps34. *J Immunol*. 2013;190(10):5086-5101.
39. Young AR, Chan EY, Hu XW, et al. Starvation and ULK1-dependent cycling of mammalian Atg9 between the TGN and endosomes. *J Cell Sci*. 2006;119(18):3888-3900.
40. Tang MK, Kwong A, Tam KF, et al. BRCA1 deficiency induces protective autophagy to mitigate stress and provides a mechanism for BRCA1 haploinsufficiency in tumorigenesis. *Cancer Lett*. 2014;346(1):139-147.
41. Esteve JM, Armengod ME, Knecht E. BRCA1 negatively regulates formation of autophagic vacuoles in MCF-7 breast cancer cells. *Exp Cell Res*. 2010;316(16):2618-2629.
42. Arun B, Akar U, Gutierrez-Barrera AM, et al. The PARP inhibitor AZD2281 (Olaparib) induces autophagy/mitophagy in BRCA1 and BRCA2 mutant breast cancer cells. *Int J Oncol*. 2015;47(1):262-268.
43. Wan B, Dai L, Wang L, et al. Knockdown of BRCA2 enhances cisplatin and cisplatin-induced autophagy in ovarian cancer cells. *Endocr Relat Cancer*. 2018;25(1):69-82.
44. Minami A, Nakanishi A, Ogura Y, et al. Connection between tumor suppressor BRCA1 and PTEN in damaged DNA repair. *Front Oncol*. 2014;4:318.
45. Gillespie DA, Ryan KM. Autophagy is critically required for DNA repair by homologous recombination. *Mol Cell Oncol*. 2016;3(1):e1030538.
46. Arnes JB, Bégin LR, Stefansson I, et al. Expression of epidermal growth factor receptor in relation to BRCA1 status, basal-like markers and prognosis in breast cancer. *J Clin Pathol*. 2009;62(2):139-146.
47. Toyama T, Yamashita H, Kondo N, et al. Frequently increased epidermal growth factor receptor (EGFR) copy numbers and decreased BRCA1 mRNA expression in Japanese triple-negative breast cancers. *BMC Cancer*. 2008;8(1):309.
48. Li H, You L, Xie J, et al. The roles of subcellularly located EGFR in autophagy. *Cell Signal*. 2017;35:223-230.
49. Tan X, Thapa N, Sun Y, et al. A kinase-independent role for EGF receptor in autophagy initiation. *Cell*. 2015;160(1-2):145-160.
50. Wei Y, Zou Z, Becker N, et al. EGFR-mediated Beclin 1 phosphorylation in autophagy suppression, tumor progression, and tumor chemoresistance. *Cell*. 2013;154(6):1269-1284.
51. Mgbemena VE, Signer RAJ, Wijayatunge R, et al. Distinct BRCA1 mutations differentially reduce hematopoietic stem cell function. *Cell Rep*. 2017;18(4):947-960.
52. Dengel J, Schoor O, Fischer R, et al. Autophagy promotes MHC class II presentation of peptides from intracellular source proteins. *Proc Natl Acad Sci USA*. 2005;102(22):7922-7927.
53. Li Y, Wang L-X, Yang G, et al. Efficient cross-presentation depends on autophagy in tumor cells. *Cancer Res*. 2008;68(17):6889-6895.
54. Liu Y, Gu Y, Cao X. The exosomes in tumor immunity. *Oncimmunology*. 2015;4(9):e1027472.
55. Tittarelli A, Mendoza-Naranjo A, Farias M, et al. Gap junction intercellular communications regulate NK cell activation and modulate NK cytotoxic capacity. *J Immunol*. 2014;192(3):1313-1319.
56. Tittarelli A, Janji B, Van Moer K, et al. The selective degradation of synaptic Connexin 43 protein by hypoxia-induced autophagy impairs natural killer cell-mediated tumor cell killing. *J Biol Chem*. 2015;290(39):23670-23679.
57. Baginska J, Viry E, Berchem G, et al. Granzyme B degradation by autophagy decreases tumor cell susceptibility to natural killer-mediated lysis under hypoxia. *Proc Natl Acad Sci USA*. 2013;110(43):17450-17455.
58. Noman MZ, Janji B, Kaminska B, et al. Blocking hypoxia-induced autophagy in tumors restores cytotoxic T-cell activity and promotes regression. *Cancer Res*. 2011;71(18):5976-5986.
59. Noman MZ, Buart S, Van Pelt J, et al. The cooperative induction of hypoxia-inducible factor-1 alpha and STAT3 during hypoxia induced an impairment of tumor susceptibility to CTL-mediated cell lysis. *J Immunol*. 2009;182(6):3510-3521.
60. Messai Y, Noman MZ, Janji B, et al. The autophagy sensor ITPR1 protects renal carcinoma cells from NK-mediated killing. *Autophagy*. 2015;00-1080/15548627.2015.1017194.0.
61. Khazen R, Muller S, Gaudenzio N, et al. Melanoma cell lysosome secretory burst neutralizes the CTL-mediated cytotoxicity at the lytic synapse. *Nat Commun*. 2016;7(1):10823.
62. Akalay I, Janji B, Hasmim M, et al. Epithelial-to-mesenchymal transition and autophagy induction in breast carcinoma promote escape from T-cell-mediated lysis. *Cancer Res*. 2013;73(8):2418-2427.
63. Liu Y, Xiang X, Zhuang X, et al. Contribution of MyD88 to the tumor exosome-mediated induction of myeloid derived suppressor cells. *Am J Pathol*. 2010;176(5):2490-2499.
64. Valenti R, Huber V, Filipazzi P, et al. Human tumor-released microvesicles promote the differentiation of myeloid cells with transforming growth factor-beta-mediated suppressive activity on T lymphocytes. *Cancer Res*. 2006;66(18):9290-9298.
65. Yu S, Liu C, Su K, et al. Tumor exosomes inhibit differentiation of bone marrow dendritic cells. *J Immunol*. 2007;178(11):6867-6875.
66. Yang C, Kim SH, Bianco NR, et al. Tumor-derived exosomes confer antigen-specific immunosuppression in a murine delayed-type hypersensitivity model. *PLoS One*. 2011;6(8):e22517.
67. Kogure T, Lin WL, Yan IK, et al. Intercellular nanovesicle-mediated microRNA transfer: a mechanism of environmental modulation of hepatocellular cancer cell growth. *Hepatology*. 2011;54(4):1237-1248.
68. Ohshima K, Inoue K, Fujiwara A, et al. Let-7 microRNA family is selectively secreted into the extracellular environment via exosomes in a metastatic gastric cancer cell line. *PLoS One*. 2010;5(10):e13247.
69. Condamine T, Gabrilovich DI. Molecular mechanisms regulating myeloid-derived suppressor cell differentiation and function. *Trends Immunol*. 2011;32(1):19-25.
70. Zhou M, Chen J, Zhou L, et al. Pancreatic cancer derived exosomes regulate the expression of TLR4 in dendritic cells via miR-203. *Cell Immunol*. 2014;292(1-2):65-69.
71. Groh V, Wu J, Yee C, et al. Tumour-derived soluble MIC ligands impair expression of NKG2D and T-cell activation. *Nature*. 2002;419(6908):734-738.
72. Clayton A, Mitchell JP, Court J, et al. Human tumor-derived exosomes down-modulate NKG2D expression. *J Immunol*. 2008;180(11):7249-7258.
73. Szczepanski MJ, Szajnik M, Welsh A, et al. Blast-derived microvesicles in sera from patients with acute myeloid leukemia suppress natural killer cell function via membrane-associated transforming growth factor-beta1. *Haematologica*. 2011;96(9):1302-1309.

74. Hedlund M, Nagaeva O, Kargl D, et al. Thermal- and oxidative stress causes enhanced release of NKG2D ligand-bearing immunosuppressive exosomes in leukemia/lymphoma T and B cells. *PLoS One*. 2011;6(2):e16899.
75. Klibi J, Niki T, Riedel A, et al. Blood diffusion and Th1-suppressive effects of galectin-9-containing exosomes released by Epstein-Barr virus-infected nasopharyngeal carcinoma cells. *Blood*. 2009;113(9):1957-1966.
76. Soderberg A, Barral AM, Soderstrom M, et al. Redox-signaling transmitted in trans to neighboring cells by melanoma-derived TNF-containing exosomes. *Free Radic Biol Med*. 2007;43(1):90-99.
77. Wieckowski EU, Visus C, Szajnik M, et al. Tumor-derived microvesicles promote regulatory T cell expansion and induce apoptosis in tumor-reactive activated CD8+ T lymphocytes. *J Immunol*. 2009;183(6):3720-3730.
78. Szajnik M, Czystowska M, Szczepanski MJ, et al. Tumor-derived microvesicles induce, expand and up-regulate biological activities of human regulatory T cells (Treg). *PLoS One*. 2010;5(7):e11469.
79. Yin Y, Cai X, Chen X, et al. Tumor-secreted miR-214 induces regulatory T cells: a major link between immune evasion and tumor growth. *Cell Res*. 2014;24(10):1164-1180.
80. Yang C, Chalasani G, Ng YH, et al. Exosomes released from mycoplasma infected tumor cells activate inhibitory B cells. *PLoS One*. 2012;7(4):e36138.
81. Lastwika KJ, Wilson W III, Li QK, et al. Control of PD-L1 expression by oncogenic activation of the AKT-mTOR pathway in non-small cell lung cancer. *Cancer Res*. 2016;76(2):227-238.
82. Clark CA, Gupta HB, Sareddy G, et al. Tumor-intrinsic PD-L1 signals regulate cell growth, pathogenesis, and autophagy in ovarian cancer and melanoma. *Cancer Res*. 2016;76(23):6964-6974.
83. Senzer N, Barve M, Kuhn J, et al. Phase I trial of "bi-shRNAi(furin)/GMCSF DNA/autologous tumor cell" vaccine (FANG) in advanced cancer. *Mol Ther*. 2012;20(3):679-686.
84. Oh J, Barve M, Matthews CM, et al. Phase II study of Vigil® DNA engineered immunotherapy as maintenance in advanced stage ovarian cancer. *Gynecol Oncol*. 2016;143(3):504-510.
85. Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer*. 2012;12(4):252-264.
86. Alsaab HO, Sau S, Alzhrani R, et al. PD-1 and PD-L1 checkpoint signaling inhibition for cancer immunotherapy: mechanism, combinations, and clinical outcome. *Front Pharmacol*. 2017;8(561):1-15.
87. Oh J, Barve M, Matthews CM, et al. Phase II study of Vigil(R) DNA engineered immunotherapy as maintenance in advanced stage ovarian cancer. *Gynecol Oncol*. 2016;143(3):504-510.
88. Barve M, Kuhn J, Lamont J, et al. Follow-up of bi-shRNA furin/GM-CSF engineered autologous tumor cell (EATC) immunotherapy Vigil® in patients with advanced melanoma. *Biomed Genet Genomics*. 2016;1(4):81-86.
89. Senzer N, Barve M, Nemunaitis J, et al. Long term follow up: phase I trial of "bi-shRNA furin/GMCSF DNA/autologous tumor cell" immunotherapy (FANG) in advanced cancer. *J Vaccines Vaccination*. 2013;4(8):1-8.
90. Ghisoli M, Barve M, Schneider R, et al. Pilot trial of FANG immunotherapy in Ewing's Sarcoma. *Mol Ther*. 2015;23(6):1103-1109.
91. Ghisoli M, Barve M, Mennel R, et al. Three-year follow up of GMCSF/bi-shRNA(furin) DNA-transfected autologous tumor immunotherapy (Vigil) in metastatic advanced Ewing's Sarcoma. *Mol Ther*. 2016;24(8):1478-1483.