

Better living through chemistry: Addressing emerging antibiotic resistance

Nathan P Coussens¹, Ashley L Molinaro², Kayla J Culbertson², Tyler Peryea¹, Gergely Zahoránszky-Köhalmi¹, Matthew D Hall¹ and Dayle A Daines²

¹National Center for Advancing Translational Sciences, Division of Pre-Clinical Innovation, Rockville, MD 20850, USA; ²Department of Biological Sciences, Old Dominion University, Norfolk, VA 23529, USA

Corresponding author: Dayle A Daines. Email: ddaines@odu.edu

Impact statement

The alarming increase in antibiotic-resistant microorganisms is a rapidly emerging threat to human health throughout the world. Historically, small molecule drugs have played a major role in controlling bacterial infections and they continue to offer tremendous potential in countering resistant organisms. This minireview provides a broad overview of the relevant issues, including the diversity of FDA-approved small molecule drugs and mechanisms of drug resistance, unintended consequences of antibiotic use, the current state of development for small molecule antibacterials and financial challenges that impact progress towards novel therapies. The content will be informative to diverse stakeholders, including clinicians, basic scientists, translational scientists and policy makers, and may be used as a bridge between these key players to advance the development of much-needed therapeutics.

Abstract

The increasing emergence of multidrug-resistant bacteria is recognized as a major threat to human health worldwide. While the use of small molecule antibiotics has enabled many modern medical advances, it has also facilitated the development of resistant organisms. This minireview provides an overview of current small molecule drugs approved by the US Food and Drug Administration (FDA) for use in humans, the unintended consequences of antibiotic use, and the mechanisms that underlie the development of drug resistance. Promising new approaches and strategies to counter antibiotic-resistant bacteria with small molecules are highlighted. However, continued public investment in this area is critical to maintain an edge in our evolutionary “arms race” against antibiotic-resistant microorganisms.

Keywords: Bacteria, drugs, mechanisms, pharmacology, resistance, small molecule inhibitors

Experimental Biology and Medicine 2018; 243: 538–553. DOI: 10.1177/1535370218755659

Introduction

Antibiotic-resistant bacteria are a major public health concern, and the treatment of drug-resistant infections is increasingly problematic. More than 150 small molecule drugs have been approved by the US Food and Drug Administration (FDA) over eight decades for the treatment of bacterial infections. In the context of this article, we will define a “small molecule” based on the FDA’s assessment of a drug substance as a “chemical” according to ISO IDMP 11238.¹ Additionally, we include mixtures, as specified by the FDA, provided that each component of the mixture is

classified as a “chemical”. Both the misuse and overuse of antibiotics have been cited as the major driving forces for the development of resistance among human pathogens. Globalization and international travel increases the likelihood that an easily-transmissible antibiotic-resistant infection could result in a pandemic. The lack of effective antibiotic therapies jeopardizes many modern medical practices including transplantation, cancer chemotherapy, surgeries, and care of the critically ill or premature infants.² Understanding the mechanisms of antibiotic resistance is critical for developing new and innovative approaches for

better and more effective treatments.^{3–5} This article provides some context and reviews the recent literature describing the development of small molecule therapies to circumvent antibiotic resistance.

US Food and Drug Administration-approved small molecule antibacterial drugs and their targets

Many antibacterial drugs are natural products that were isolated from microbes in the soil.⁶ Therefore, elucidating the antimicrobial mechanisms of action has been a subject of intense interest, and numerous investigations have led to the categorization of these drugs by their effects on the bacterial cell. The vast majority of antibiotics target the disruption of either the bacterial cell wall or membrane, or interfere with essential processes involved in metabolism and replication (Figure 1). While the mechanisms of action for many antibiotics have been described, there are still classes of antimicrobials for which specific binding sites or target(s) remain unknown. These systems continue to be scrutinized for insight into potential areas of exploitation for promising drugs. Studies suggest that the activities of antibacterial drugs are extended beyond the initial drug/target interaction to metabolic and homeostatic networks so that the efficacy of a drug depends on its mode of action in a given cellular context.⁷ Table 1 lists the small molecule antibiotics approved by the FDA along with the approval years. Figure 2 shows the number of small molecule antibacterial drugs approved for human use between 1939 and 2017 grouped according to the antibiotic class.

Cell wall synthesis

Peptidoglycan provides the bacterial cell with its fundamental structural integrity and essential reactions can be inhibited when antibiotics bind to penicillin binding proteins (PBPs) that are crucial for forming peptidoglycan. Upon the arrest of peptidoglycan production, the cell wall loses its integrity, causing lysis of the bacterium.⁸ Antibiotics that include the β -lactams and glycopeptides are effective against bacteria due to their ability to interfere with bacterial peptidoglycan synthesis.⁹ Subclasses of the β -lactams include penicillins, cephalosporins, monobactams, carbapenems, and clavams. Vancomycin is a glycopeptide class antibiotic that has been FDA-approved. Peptidoglycan synthesis is also inhibited by the oxazolidinone class antibiotic cycloserine and the polypeptide class antibiotic bacitracin.

Cell membrane function

The lipopeptide class antibiotic daptomycin targets the cell wall in Gram-positive microorganisms. While the exact mechanism responsible for the membrane disruption remains unknown, it has been shown that the presence of both phosphatidylglycerol (PG) and calcium are necessary for daptomycin to be effective. Comparisons among membranes that contain PG and those made entirely of phosphatidylcholine have shown that daptomycin is less effective on membranes without PG. Calcium levels are

also important, as the concentration of daptomycin needed to penetrate the cell membrane decreases by as much as 50-fold in the presence of calcium ions.¹⁰ The polypeptide class antibiotic polymyxin B is active against Gram-negative organisms, destabilizing both the outer and inner membranes.¹¹

Protein synthesis

Several classes of antibiotics target ribosomes in order to disrupt protein synthesis in bacteria. These classes of antibiotics include the phenicols, macrolides, lincosamides, and streptogramins which bind to the 50S ribosomal subunit and hinder translation. It is thought that the structures of lincosamides might resemble the 3'-ends of L-Pro-Met-tRNA and deacylated-tRNA, allowing them to bind to and subsequently inhibit the ribosome.¹² The oxazolidinone class antibiotics linezolid and tedizolid also inhibit protein synthesis by binding the 50S ribosomal subunit.

In the bacterial ribosome, the 30S subunit is a target for other classes of antibiotics such as the tetracyclines and aminoglycosides. Examples of aminoglycosides include kanamycin, gentamicin, neomycin, and streptomycin. These drugs bind to the 16S ribosomal RNA within the 30S subunit and do not allow the binding of charged aminoacyl-tRNA to the ribosome, thereby arresting protein synthesis. While the exact site of tetracycline binding has yet to be identified, a recent study suggested that the ribosomal RNA structure is more important than the actual ribonucleotide sequence for binding, as tetracyclines can bind to random sequences of double-stranded RNA.¹³

Nucleic acid synthesis

Bacterial enzymes are targets of antibiotics, including those that play roles in nucleic acid synthesis, DNA replication and regulation. Sulfonamide drugs were the first class to be FDA-approved beginning in 1939 and act as competitive inhibitors of the enzyme dihydropteroate synthase, which is required for the synthesis of tetrahydrofolic acid and nucleotides. The diaminopyrimidine trimethoprim targets the same pathway by inhibiting dihydrofolate reductase. Enzymes like DNA gyrase, type IIA topoisomerases, and topoisomerase IV serve as targets of the quinolone class of antibiotics. Topoisomerases form covalent bonds with DNA and catalyze DNA breakage and re-ligation. The quinolones inhibit these topoisomerases, resulting in the formation of ternary complexes that cause strand breakage and block DNA re-ligation. Quinolones were also shown to lower the threshold of magnesium ions required for topoisomerases to initiate strand breakage, allowing for increased rates of cleavage and the formation of ternary complexes.¹⁴ The rifamycin class of antibiotics inhibit RNA polymerase and prevent RNA synthesis.¹⁵

Recent FDA approvals

There were no new approvals of small molecule antibacterial drugs by the FDA in 2016; however, three were approved in 2017 as of 10 October 2017. Secnidazole (Symbiomix Therapeutics, approved 15 September 2017)




Figure 1. Anatomy of Gram-negative and Gram-positive bacteria and their susceptibility to antibiotics. (a) The cell wall of Gram-negative bacteria includes outer and inner (cytoplasmic) membranes, which form the barriers of the periplasmic space that contains a thin layer of peptidoglycan. The outer leaflet of the outer membrane contains lipopolysaccharide (LPS) consisting of lipid A and O-polysaccharide. The outer membrane contains porins, which provide entrance to the periplasm for some molecules. (b) The outer wall of Gram-positive bacteria contains a thick layer of peptidoglycan that retains the crystal violet stain used in the Gram stain test. Teichoic acids, which are not found in Gram-negative bacteria, are called lipoteichoic acids (LTA) when anchored to the cytoplasmic membrane. Gram-positive bacteria lack an outer membrane and thus are more susceptible to antibiotics. Both Gram-negative and Gram-positive cell walls contain efflux pumps and other active pumps that can export intracellular molecules, including antibiotics. (c) Gram-negative bacteria are intrinsically resistant to many antibiotics due to the architecture of their cell walls. Nevertheless, a variety of antibiotics are effective by targeting the synthesis of protein, the cell wall or nucleic acids. Resistance to antibiotics can be mediated by increased efflux by pumps and reduced permeability through porins in addition to other mechanisms, including target mutation, overexpression or protection and drug inactivation.

is a nitroimidazole antibiotic reported as early as 1976, but only now approved for bacterial vaginosis in adult women.¹⁶ Also, a combination of the broad-spectrum β -lactam antibiotic meropenem with the new β -lactamase inhibitor vaborbactam (trade name Vabomere, Rempex Pharmaceuticals, approved 29 August 2017) was approved as an injection formulation for treating complicated urinary tract infections. Vaborbactam is a new chemotype

containing a boronic acid that covalently inhibits serine-containing carbapenemases at the active site.¹⁷ While vaborbactam inhibits by a new mechanism of action, it does not act against a new target class and it is an adjuvant to the antibiotic. Finally, delafloxacin is a fluoroquinolone (Melinta Therapeutics, approved 19 June 2017) used to treat patients with acute bacterial skin and skin structure infections (ABSSSI).¹⁸

Table 1. Small molecule antibacterials approved by the US Food and Drug Administration (FDA) for use in humans.a

Type	Class	Drug	FDA-Approval Year	Figure 3 ID	MCS Cluster	ISO 11238 Substance Type	Molecular Weight or Weight Range (Daltons)
β-Lactams							
	Penicillins						
		Penicillin G	1947	1	2	chemical	334.39
		Penicillin V	1954	2	2	chemical	350.39
		Pheneticillin	1959	3	2	chemical	364.42
		Methicillin	1960	4	2	chemical	380.42
		Oxacillin	1962	5	1	chemical	401.44
		Ampicillin	1963	6	2	chemical	349.41
		Nafcillin	1964	7	2	chemical	414.48
		Cloxacillin	1965	8	1	chemical	435.88
		Dicloxacillin	1968	10	1	chemical	470.33
		Carbenicillin	1972	13	2	chemical	378.40
		Amoxicillin	1974	15	2	chemical	365.40
		Ticarcillin	1976	19	2	chemical	384.43
		Bacampicillin	1980	24	5	chemical	465.52
		Mezlocillin	1981	26	2	chemical	539.58
		Piperacillin	1981	27	2	chemical	517.56
		Amdinocillin	1984	32	0	chemical	325.43
		Sulbactam	1986	41	11	chemical	233.24
	Cephalosporins						
		Cephaloridine	1968	9	0	chemical	415.49
		Cephaloglycin	1970	11	5	chemical	405.43
		Cephalexin	1971	12	5	chemical	347.39
		Cefazolin	1973	14	8	chemical	454.51
		Cephalothin	1974	16	8	chemical	396.44
		Cephradine	1974	17	8	chemical	349.41
		Cephapirin	1974	18	8	chemical	423.46
		Cefadroxil	1978	20	5	chemical	363.39
		Cefamandole	1978	21	6	chemical	462.50
		Cefoxitin	1978	22	10	chemical	427.45
		Cefaclor	1979	23	5	chemical	367.81
		Latamoxef	1981	25	0	chemical	520.47
		Cefotaxime	1981	28	3	chemical	455.47
		Cefoperazone	1982	29	5	chemical	645.67
		Ceftizoxime	1983	30	3	chemical	383.40
		Cefuroxime	1983	31	10	chemical	424.39
		Cefonicid	1984	35	7	chemical	542.57
		Ceforanide	1984	36	7	chemical	519.55
		Ceftriaxone	1984	34	3	chemical	554.58
		Ceftazidime	1985	38	3	chemical	545.57
		Cefotetan	1985	39	7	chemical	575.62
		Cefmenoxime	1987	42	3	chemical	511.56
		Cefotiam	1988	43	6	chemical	525.63
		Cefpiramide	1989	45	5	chemical	612.64
		Cefixime	1989	44	3	chemical	453.45
		Cefmetazole	1989	46	7	chemical	471.53
		Cefprozil	1991	48	5	mixture	389.43
		Loracarbef	1991	47	5	chemical	349.77
		Cefpodoxime	1992	49	4	chemical	427.46
		Tazobactam	1993	50	11	chemical	300.29
		Ceftibuten	1995	51	0	chemical	410.43
		Cefepime	1996	52	3	chemical	480.56
		Cefdinir	1997	54	3	chemical	395.41
		Cefditoren	2001	55	3	chemical	506.58
		Ceftaroline fosamil	2010	58	0	chemical	684.69
		Ceftolozane	2014	59	4	chemical	666.69
	Monobactams						
		Aztreonam	1986	40	0	chemical	435.43
	Carbapenems						
		Imipenem	1985	37	0	chemical	299.35
		Meropenem	1996	53	9	chemical	383.46
		Ertapenem	2001	56	9	chemical	475.52
		Doripenem	2007	57	9	chemical	420.50

(continued)

Table 1. Continued

Type	Class	Drug	FDA-Approval Year	Figure 3 ID	MCS Cluster	ISO 11238 Substance Type	Molecular Weight or Weight Range (Daltons)	
MLS family	Clavams	Clavulanic acid	1984	33	0	chemical	199.16	
		Lincosamides	Lincomycin					1964
		Clindamycin	1970					424.98
	Macrolides	Erythromycin	1952					733.93
		Troleandomycin	1956					813.97
		Clarithromycin	1991					747.95
		Azithromycin	1991					748.98
		Dirithromycin	1995					835.07
		Telithromycin	2004					812.00
	Streptogramins	Quinupristin	1999					1022.22
Dalfopristin		1999	690.85					
Other Antibiotics	Acylampicillin							
		Azlocillin	1982		chemical	461.49		
	Aminocyclitol							
		Spectinomycin	1971		chemical	332.35		
	Aminoglycosides	Streptomycin	1946	581.57				
		Neomycin	1951	322.36 – 614.64				
		Paromomycin	1960	615.63				
		Gentamicin	1966	449.54 – 477.60				
		Tobramycin	1975	467.51				
		Amikacin	1976	585.60				
		Kanamycin	1981	483.51 – 484.50				
		Netilmicin	1983	475.58				
	Aminosalicylates							
		Aminosalicylic acid	1948		chemical	153.14		
	Diaminopyrimidines							
		Trimethoprim	1973		chemical	290.32		
	Diarylquinolines							
		Bedaquiline	2012		chemical	555.51		
	Glycopeptides							
		Vancomycin	1958		chemical	1449.25		
	Glycylcyclines							
		Tigecycline	2005		chemical	585.65		
	Isonicotinic acid							
		Isoniazid	1952		chemical	137.14		
	Lipoglycopeptides	Telavancin	2009	1755.64				
		Oritavancin	2014	1793.10				
		Dalbavancin	2014	1802.67 – 1830.72				
Oritavancin		2014	1793.10					
Lipopeptides								
	Daptomycin	2003		chemical	1620.67			
Macrocyclics								
	Fidaxomicin	2011		chemical	1058.04			
Monoxycarboic acids								
	Mupirocin	1987		chemical	500.62			
N-substituted ethylenediamine	Ethambutol	1967		chemical	204.31			
	Nicotinamide derivative							
	Ethionamide	1965		chemical	166.24			
Nitrofurans	Nitrofurantoin	1953		chemical	238.16			
	Furazolidone	1955		chemical	225.16			

(continued)

Table 1. Continued

Type	Class	Drug	FDA-Approval Year	Figure 3 ID	MCS Cluster	ISO 11238 Substance Type	Molecular Weight or Weight Range (Daltons)
	Nitroimidazoles	Metronidazole	1963			chemical	171.15
		Tinidazole	2004			chemical	247.27
		Secnidazole	2017			chemical	185.18
	Non--lactam -lactamase inhibitors	Avibactam	2015			chemical	265.24
		Vaborbactam	2017			chemical	297.14
	Oxazolidinones	Cycloserine	1964			chemical	102.09
		Linezolid	2000			chemical	337.35
		Tedizolid	2014			chemical	370.34
	Phenicol	Chloramphenicol	1949			chemical	323.13
	Phosphonic acid derivative	Fosfomycin	1996			chemical	138.06
	Pleuromutilins	Retapamulin	2007			chemical	517.76
	Polymyxins	Colistin	1959			mixture	1155.43 – 1169.46
	Polypeptides	Bacitracin	1948			mixture	1408.67 – 1422.69
		Polymyxin B	1951			mixture	1189.45 – 1203.48
	Pyrazines	Pyrazinamide	1955			chemical	123.11
	Quinolones	Nalidixic acid	1964			chemical	232.24
		Oxolinic acid	1975			chemical	261.23
		Norfloxacin	1986			chemical	319.33
		Ciprofloxacin	1987			chemical	331.34
		Ofloxacin	1990			chemical	361.37
		Enoxacin	1991			chemical	320.32
		Lomefloxacin	1992			chemical	351.35
		Sparfloxacin	1996			chemical	392.40
		Levofloxacin	1996			chemical	361.37
		Grepafloxacin	1997			chemical	359.39
		Trovafloxacin	1997			chemical	416.35
		Moxifloxacin	1999			chemical	401.43
		Gatifloxacin	2003			chemical	375.39
		Gemifloxacin	2003			chemical	389.38
		Besifloxacin	2009			chemical	393.84
		Finafloxacin	2014			chemical	398.39
		Delafloxacin	2017			chemical	440.76
	Rifamycins	Rifampin	1971			chemical	822.94
		Rifabutin	1992			chemical	847.00
		Rifapentine	1998			chemical	877.03
		Rifaximin	2004			chemical	785.88
	Riminophenazine	Clofazimine	1986			chemical	473.40
	Sulfonamides	Sulfapyridine	1939			chemical	249.29
		Sulfadiazine	1941			chemical	250.28
		Sulfamerazine	1943			chemical	264.30
		Sulfathiazole	1945			chemical	255.32
		Sulfacetamide	1945			chemical	214.24
		Sulfoxone	1947			chemical	404.48
		Sulfisoxazole	1949			chemical	267.30
		Sulfamethizole	1953			chemical	270.33
		Sulfamethoxazole	1961			chemical	253.28

(continued)

Table 1. Continued

Type	Class	Drug	FDA-Approval Year	Figure 3 ID	MCS Cluster	ISO 11238 Substance Type	Molecular Weight or Weight Range (Daltons)
		Sulfanilamide	1965			chemical	172.21
		Mafenide	1969			chemical	186.23
		Sulfaphenazole	1974			chemical	314.36
		Sulfadoxine	1981			chemical	310.33
	Sulfones						
		Dapsone	1957			chemical	248.30
	Tetracyclines						
		Chlortetracycline	1950			chemical	478.88
		Oxytetracycline	1950			chemical	460.43
		Tetracycline	1953			chemical	444.43
		Demeclocycline	1960			chemical	464.85
		Methacycline	1966			chemical	442.42
		Doxycycline	1967			chemical	444.43
		Minocycline	1971			chemical	457.48
	Triazinanes						
		Methenamine	1967			chemical	140.19
	Tuberactinomycins						
		Viomycin	1953			chemical	685.69
		Capreomycin	1971			mixture	524.53 – 668.71

^aThe table is organized according to antibiotic type, class, drug name, FDA-approval year, identification number in Figure 3 and MCS cluster designation (β -lactam drugs only), ISO 11238 substance type, and molecular weight or weight range (for mixtures). The maximal common substructure (MCS) clustering analysis was performed with β -lactam drugs as described in the Figure 3 legend and all β -lactams were assigned as either singletons (designated as zero) or one of eleven MCS clusters as indicated in the table. The list of antibacterial drugs was constructed from the NCATS Inxight Drugs database (<https://drugs.ncats.io>) and eMedExpert (<http://www.emedexpert.com/lists/antibiotics.shtml#1>) followed by manual curation. Both FDA drug approval databases, Drugs@FDA and the FDA Orange Book, were used to confirm each approval status and date. Effort was made to exclude salts, esters and known pro-drug variants of other approved antibiotic substances. For any active moiety not associated with a definite approval date from FDA's databases, the approval date from the NCATS Inxight Drugs resource was used.




Figure 2. Small molecule antibacterial drugs approved by the US Food and Drug Administration (FDA) for use in humans. This plot shows the number of drugs approved within five-year periods from 1939 until 2017 (all drugs are listed in Table 1). The 157 drugs are grouped according to 20 classifications as shown by the legend. The greatest number of drugs were approved between 1981 and 1985, consisting of 19 antibacterials that span more than six classifications. During the past three decades, the number of approved small molecule antibacterial drugs has steadily decreased.

Unintended consequences of antibiotic use

Alexander Fleming's discovery of penicillin in 1928 and its release for use in 1942 marked the era of antibiotics and the foundation for the treatment of diseases that had

previously been lethal, such as staphylococcal and streptococcal infections.¹⁹ However, over the last several decades we have seen an overwhelming increase in resistance to various antimicrobials. Further, there are a number of undesirable outcomes that occur during and after antibiotic

treatment that can exacerbate the duration and severity of infectious diseases.

Alteration of the microbiome

The human microbiome plays a vital role in host health. It is comprised of trillions of microbial cells encompassing as many as 500 species,^{20,21} maintains a crucial symbiotic relationship with the host and plays a fundamental role in educating the immune system.²² The components of one's own microbiota are unique enough to be used for self-identification, and success in using the community structure of the human gut to distinguish individuals in a group has been demonstrated.²³ The commensal bacteria of the human microbiota maintain over two orders of magnitude more genes than their host,²⁴ including those that encode enzymes that facilitate host digestion, lipid absorption via bile acid conversion, vitamin production as well as the protection and maintenance of the intestinal mucosa.^{25–28} Along with providing essential functions to the host, commensals also compete with pathogenic microorganisms for nutrition and attachment sites and therefore help prevent infection. Bohnhoff and Miller first reported in 1962 what is now known as “colonization resistance” by determining that mice were significantly more susceptible to *Salmonella* infection after treatment with streptomycin, which decreased the normal gut microbiota.²⁹ Antibiotic treatment eliminates many commensal organisms in the gut, enabling enteric pathogens such as *Clostridium difficile*, vancomycin-resistant *Enterococcus faecalis*, and *Salmonella enterica* serovar Typhimurium to colonize and infect the host.^{30–32} Antibiotic-mediated disruption of the normal microbiota has also been implicated in liver disease and colorectal cancers.^{33–35} Further, these changes in the gut flora can lead to significantly elevated levels of resistance genes in the microbial population, the increased carriage of which might last for many years following a single course of antibiotic treatment.³⁶ Changes in the microbiota of the upper respiratory tract, skin and genitourinary tract can likewise affect susceptibility to infections. A recent systematic review of the potential for exposure to antibiotics to increase the risk of community-acquired infections found that the collateral damage to the patient's microbiome could be associated with upper respiratory, urinary tract and soft tissue infections as well as infectious diarrhea, and that this association decreased with increasing time since antibiotic exposure.³⁷

Activation of toxin–antitoxin systems

Another consequence of antibiotic use is the induction of growth arrest in bacteria subsequent to the activation of toxin–antitoxin (TA) modules. Five unique TA families have been identified (I–V); the toxin is always a protein but the antitoxin can be a protein or nucleic acid, and this characteristic determines to which family a module belongs. These gene pairs are found in hundreds of bacteria as well as in the archaea.³⁸ The type II TA systems have protein antitoxins and are the best characterized. Upon translation, the toxin and antitoxin interact to form a non-toxic complex. When bacteria experience stress in their

microenvironment, such as exposure to reactive oxidative species, nutrient limitation, temperature, pH or pressure changes and antibiotic therapy, induced proteases such as Lon and ClpXP degrade the labile antitoxin, freeing the more stable toxin to carry out its function.^{39,40} In the type II TA modules, the toxin is often a ribonuclease enzyme which when released from its cognate antitoxin cleaves bacterial mRNA, eliciting a state of reversible growth arrest. It is thought that many bacterial species in the environment have maintained these systems to enhance their ability to survive nutrient-poor and harsh conditions over hundreds (and, in some cases, thousands) of years.⁴¹ For bacterial pathogens, this state of stasis results in nonspecific tolerance to antimicrobials that target enzymes and pathways essential to growth and replication, allowing a subpopulation to survive antibiotic therapy and resume growth later when conditions improve.

Livestock use generating resistance

Antimicrobial resistance is a problem in veterinary medicine and the inappropriate use of these agents in agriculture continues to have a significant impact on human health. Antibiotics are routinely used in animal production to promote growth by preventing and treating infections.⁴² The preventative treatment of entire flocks or herds over time has facilitated the evolution of increased numbers of resistant pathogens. For example, there has been a marked rise in multidrug-resistant (MDR) *Escherichia coli*, *Salmonella*, *Campylobacter*, and methicillin-resistant *Staphylococcus aureus* (MRSA) emerging from commercial animal production across the globe.^{43–45} In 2014, approximately 9.7 million kilograms of medically-important antibiotics were sold for use in food-producing animals in the United States (including almost 7 million kilograms of the tetracycline drug class alone), compared with approximately 3.5 million kilograms of antibiotics sold for human use.⁴⁶

On large farms with poor living conditions, livestock are treated with high doses of various antibiotics to decrease disease until slaughter. This practice selects for resistant bacteria which can be spread to packaging plants and end up in the kitchens of consumers.⁴⁷ Improper cooking and poor hand hygiene can facilitate dissemination, resulting in difficult-to-treat infections. Farm wastewater contaminated with fecal matter harboring resistant pathogens can accumulate as runoff in nearby rivers, streams, and ground water.⁴⁸ These resistant pathogens can exchange DNA with the existing organisms of that environment and cause life-threatening bacterial infections in human hosts downstream.⁴⁹ Manure from treated animals used for fertilization results in the selection for increased antibiotic resistance in soil microorganisms, which are ultimately transferred to the crop at harvest and then on to consumers.^{48,50,51}

Establishment of environmental reservoirs

In many parts of the world, over-the-counter sales resulting in inappropriate and/or excessive use, payback mechanisms, and lack of oversight plague antibiotic production.⁵² A related concern is the potential for the release of drug-

containing effluent from pharmaceutical manufacturing centers into the environment, resulting in selection pressure that facilitates the evolution of antibiotic resistance. For example, an LC-MS analysis of samples from 28 different sites around a bulk drug producing area in Hyderabad, South India, resulted in the identification of high concentrations of three antimicrobials (and increased concentrations of eight others) from each site. Further, tandem microbiological studies found the presence of extended-spectrum β -lactamase (ESBL) and carbapenemase-producing *Enterobacteriaceae* and non-fermenters in 95% of the samples.⁵³ The release of drugs into the environment has global consequences in terms of increasing the potential for infections by pathogens that are resistant to the very antimicrobials that we are producing on an industrial scale.

Mechanisms of antibiotic resistance

Antibiotic resistance is a natural phenomenon that predates the discovery of antibiotics and the selective pressures of antibiotic use, as genes providing resistance to the β -lactam, tetracycline, and glycopeptide antibiotics have been identified from 30,000-year-old Beringian permafrost sediments.⁵⁴ This natural resistance can be a direct result of competition between microorganisms in the environment but can also stem from spontaneous mutations that confer an adaptive benefit.⁵⁵ The dissemination of resistance can occur horizontally, carried between species via mobile genetic elements such as prophages, conjugative plasmids, or insertion sequences.⁵⁶ It can also be inherited vertically by genetic mutations that are subsequently propagated in each daughter cell.⁵⁷ Bacteria can be resistant to certain antibiotics by virtue of their physiology as well. For instance, the low permeability barrier of the Gram-negative outer membrane (Figure 1(a)) serves as the first line of defense and provides intrinsic resistance to many antibiotics.⁵⁸ As an example, Gram-negative bacteria are not susceptible to daptomycin, likely due to a lower abundance of anionic phospholipids that are required for the Ca^{2+} -mediated insertion of the lipopeptide antibiotic.⁵⁹ Thus, daptomycin is limited to the treatment of Gram-positive organisms (Figure 1(b)). The widespread use of antibiotics is thought to have provided the selective pressure to accelerate development of antibiotic resistance and MDR in previously susceptible clinically-relevant microorganisms.⁶⁰ Acquired resistance to essentially all antibiotics can be achieved by genetic mutations, the acquisition of genes or modulation of gene expression, and new resistance mechanisms continue to be characterized.^{61,62} Generally, these mechanisms include lowering the antibiotic concentration in the cell (either by reducing entry or increasing efflux⁶³), protecting the target or altering the target structure (through mutations, enzymatic modifications, or protein binding⁶⁴), over-expression of the target⁶⁵ or modifications to inactivate the antibiotic⁶⁶ (Figure 1(c)). Resistance can also be due to multiple mechanisms that are not mutually exclusive. In addition to target mutations, other mutations that result in reduced drug accumulation by either increased efflux or decreased uptake, as well as horizontal transmission by plasmids

carrying genes for target protection proteins or drug modifying enzymes can give rise to resistance.⁶⁷

Porins

Hydrophilic antibiotics (such as the β -lactams) can cross Gram-negative and mycobacterial membranes through aqueous channels called porins that normally provide the organism with nutrients.⁶³ Porins are classified according to their structure, selectivity and expression regulation. Antibiotic uptake can be decreased by altering the type or level of porin expression or altering the selectivity/function of these channels.^{61,62} As an example, studies of *Klebsiella pneumoniae* clinical isolates revealed that porin integrity was more of a determinant for carbapenem resistance than the presence of ESBL.⁶⁸ The study did not detect carbapenemases among 61 Chilean clinical isolates of carbapenem-resistant *Enterobacteriaceae*, so resistance was attributed to a combination of porin loss or alteration and β -lactamase activity. Another study characterized clinical isolates of *Enterobacter aerogenes* from patients treated with the carbapenem drug imipenem and linked altered porin expression with the adaptive response of *E. aerogenes* to the antibiotic.⁶⁹

Efflux pumps

While alterations to porins can reduce the penetration of antibiotics, efflux pumps actively transport many antibiotics out of Gram-negative and Gram-positive organisms.⁷⁰ Among the first efflux pumps to be described was the system that pumps tetracyclines out of *E. coli*.⁷¹ The efflux pumps can have a narrow or broad range of specificity for antibiotic substrates and increased expression can enable higher levels of resistance to an antibiotic. Pumps able to transport different classes of antibiotics are associated with MDR. The five major families of efflux pumps include the ATP-binding cassette (ABC) superfamily, the major facilitator superfamily (MFS), the multidrug and toxic-compound extrusion (MATE) family, the small multidrug-resistance (SMR) family, and the resistance nodulation division (RND) family.⁷⁰ Classification into a family is based on the number of components comprising the pump, the number of transmembrane-spanning regions, the energy source for active transport and the substrate type. The efflux transporters with the highest clinical relevance from Gram-positive bacteria are members of the MFS, whereas for Gram-negative organisms the transporters are from the RND family, which are able to extrude a wide range of antibiotics.⁷⁰ Recently, genomic analysis of pre- and post-therapy MDR clinical isolates of *Salmonella typhimurium* were compared from a patient that failed antibacterial therapy and a G288D mutation was identified in the Na^+ regulatory domain (NRD) transporter acriflavine resistance protein (AcrB) of the AcrAB-TolC tripartite MDR efflux pump system.⁷² Characterization studies indicated that the mutation increased the efflux of the fluoroquinolone antibiotic ciprofloxacin but decreased the efflux of other drugs. While this study demonstrates how mutations to a transporter can alter bacterial susceptibility to antibiotic therapy, mutations can also alter the regulatory

network controlling the expression of efflux pumps to result in increased antibiotic resistance.⁶² Resistance to antibiotics has also been linked to a combination of efflux pump overexpression and porin downregulation in clinical isolates from *Pseudomonas aeruginosa*, *K. pneumoniae*, and *E. coli*.^{73,74}

Protecting the target

A general strategy to achieve antibiotic resistance is to prevent an antibiotic from binding its target, which can be accomplished by different means. One paradigm of this is a target-protection mechanism, whereby a protein can associate with the drug target to prevent association with the antibiotic. An example is given by the FusB fusidic acid resistance protein, which protects elongation factor G (EF-G) from the antibiotic fusidic acid (FA). FA is a steroid antibiotic that is effective against Gram-positive bacteria, including MRSA, and acts by inhibiting the release of EF-G from the ribosome, thereby halting protein synthesis and inducing growth arrest.⁷⁵ Recent structural studies of the FusB-EF-G complex by solution NMR revealed dynamic changes in EF-G following FusB binding that facilitates disassembly of the stalled post-translational complex to rescue protein synthesis.⁷⁶ Another mechanism to prevent target engagement by an antibiotic is genetic mutations to the target gene. An example of this is fluoroquinolone resistance, which can result from mutations to DNA gyrase and topoisomerase IV that are targets of the antibiotic.⁷⁷ Mutations to the topoisomerases alter their structures to reduce the binding efficiency of quinolones.

Altering the target structure

Alternatively, modifications to antibiotic target structures can be achieved enzymatically. For example, the chloramphenicol-florfenicol resistance (Cfr) ribosomal RNA (rRNA) methyltransferase of Gram-positive and Gram-negative bacteria modifies the 23S rRNA at position A2503, providing MDR to a range of antibiotics including: phenicols, lincosamides, oxazolidinones, pleuromutilins, and streptogramin A.⁷⁸

Altered gene expression

Resistance to antibiotics can be due to the modulation of gene expression. For example, resistance to methicillin in *S. aureus* results from the *mecA* gene that encodes PBP2a, a PBP that has low affinity for many β -lactams including penicillins, cephalosporins, and carbapenems.⁷⁹ While this is an example of replacing a target with another protein of similar biochemical function that is not susceptible to antibiotics, resistance has also been achieved by overexpression of a target to a concentration beyond what the antibiotic can inhibit. For instance, a clinically isolated strain of *E. coli* was shown to have mutations that resulted in overexpression of dihydrofolate reductase by several hundredfold, which resulted in a three-fold increase in the K_i for the antifolate drug trimethoprim.⁸⁰

Modifications to the antibiotic

Antibiotic resistance can also be achieved through biochemical modifications to the antibiotic, including hydrolysis, group transfer, and redox mechanisms.⁶⁶ The β -lactamase enzymes provide a classic example of a hydrolysis mechanism, as the first β -lactamase termed "penicillinase" was identified in, 1940 from *E. coli* prior to the clinical use of penicillin.^{81,82} Since that time, more than 850 β -lactamases have been identified, which are able to hydrolyze β -lactam antibiotics (including penicillins, cephalosporins, monobactams, and carbapenems).⁸³ Alternatively, antibiotics can be inactivated due to various chemical modifications, including acylation, phosphorylation, glycosylation, nucleotidylation, ribosylation, and thiol transfer.⁶⁶ For example, the aminoglycoside antibiotic kanamycin B can be enzymatically modified by acylation, phosphorylation, and nucleotidylation.⁸⁴ A well-studied example of the redox mechanism of antibiotic inactivation is the enzyme TetX, which catalyzes the monohydroxylation of tetracyclines at position 11a, disrupting the antibiotic Mg^{2+} -binding site which is required for antibacterial activity.⁶⁶

From a perspective of evolving drug resistance, it is interesting to consider how the structures of approved antibiotics have changed over time. Fifty-nine β -lactam drugs were approved by the FDA between 1947 and 2014 (Table 1). To capture the evolution of these molecules since penicillin G was approved, structural similarities were evaluated among all 59 β -lactam drugs. The method of maximal common substructure (MCS) clustering was applied to detect various chemotypes through the identification of common structural elements among similar molecules.^{85,86} The expected outcome from such an analysis is that molecules in a given cluster are more similar to each other than to molecules of another cluster. The resulting chemical space-time analysis is shown in Figure 3. The 2-dimensional transformation reveals closely related molecules and the overall structural diversity among the β -lactam antibacterial drugs. To highlight the evolution of β -lactam chemotypes, selected structures are shown. The plot clearly shows a trend towards new scaffolds over the course of time.

New approaches for small molecule antimicrobials

Due to the crisis of antibiotic resistance, it is necessary to explore alternate or adjunctive therapies to enhance our current defenses against infectious diseases. Analyzing the role of small molecules to advance our strategies against antibiotic-tolerant and -resistant bacterial infections is of great interest and has afforded a number of successful new approaches.

New chemotypes

As indicated in the previous section, the development of resistance against new antibiotics by pathogenic organisms is essentially inevitable due to a range of mechanisms. However, resistance is not anticipated to be easily developed against the recently discovered antibacterial molecule




Figure 3. Chemical space-time of β -lactam antibacterial drugs. The dots indicate structures of β -lactam antibacterial drugs in a 2-dimensional chemical space. The distance between dots reflects the extent of structural similarity, with similar structures in closer proximity. The color and size of the dots corresponds to the year of FDA-approval, with more recent approvals indicated by a color closer to the blue end of the spectrum and a smaller size. The maximal common substructure (MCS) clustering was performed with a similarity threshold of 0.60 using the StarDrop scientific software suite (version: 6.2.0, Optibrium Ltd, Cambridge, UK). The clustering resulted in 11 MCS clusters and eight singletons (Table 1). In order to characterize the diversity among β -lactam structures, they were embedded in a 2-dimensional space using the t-distributed stochastic neighbor embedding algorithm of StarDrop.⁸⁷

pseudouridimycin (PUM), which is a nucleoside analog.⁸⁸ Because PUM binds directly and specifically to the nucleotriphosphate-binding active site of bacterial RNA polymerase (RNAP), mutations that reduce PUM binding would likely also reduce or abolish enzymatic activity. The compound was identified from extracts of soil microbes in a screen for inhibitors of bacterial RNAP. PUM inhibited Gram-negative and Gram-positive bacterial growth, including drug resistant *Streptococcus* strains, at low micromolar concentrations and cleared infection in a mouse model of *Streptococcus pyogenes* peritonitis. The antibiotic rifampin binds RNAP at a distinct site and additive antibacterial activity was observed when PUM and rifampin were co-administered.⁸⁸

New chemotypes are required to stay ahead of the rapid evolution of antibiotic resistance. The fatty acid biosynthesis type 1 enzyme (FabI) has been known as an antibacterial target for many years, but several new inhibitors are currently in clinical development that appear to offer genuinely new chemotypes, including afabicin (Deniopharm International), and CG400549 (CrystalGenomics), both of which are in phase II clinical trials.⁸⁹ As well, the mycolic acid transporter MmpL3 in *Mycobacterium tuberculosis* has been targeted by novel diamine- and indolamide-based compounds, and these have been determined to be synergistic or additive with the activity of other antimicrobials, increasing their utility and effectiveness.⁹⁰

Chemotypes for broad-spectrum activity

A recent advancement in the understanding of ideal physicochemical properties for small molecule antibacterials was made by Richter and colleagues, who studied a diverse set of compounds and used structure-activity relationship studies with computational analyses to develop predictive guidelines for small molecule accumulation in Gram-negative bacteria.⁹¹ Notably, the results indicated that small molecules most likely to accumulate contain an amine, are both rigid and amphiphilic and have low globularity, which differed substantially from the properties thought to be important based on a retrospective analysis of known antibiotics, particularly polarity and molecular weight. The authors note that compounds with primary amines are rare in standard compound libraries used for high-throughput screening and the assembly of compound collections that meet the guidelines for accumulation might enable the discovery of novel chemical leads for the development of broad-spectrum antibiotics.

New targets

A potential target for new antibiotics is the membrane enzyme phospho-MurNac-pentapeptide translocase (MraY), which is essential for the synthesis of the bacterial envelope. The role of MraY is to catalyze the initial processes in peptidoglycan synthesis by generating uridine-monophosphate (UMP) and lipid 1. There are MraY

inhibitors currently in development, but some issues remain, not the least of which is the challenge of bacterial cell entry and the problem of the inhibitor reaching effective levels within the cell.⁹

Riboswitches, which are segments of mRNA that can bind to small molecules and metabolites, are also potential targets for antimicrobial molecules. Engineered small molecules can inhibit the function of riboswitches and some were shown to be effective in bacteria. Ribocil is a selective chemical modulator that binds to riboflavin riboswitches and prevents riboflavin production in bacteria. The use of ribocil was shown to lower the *E. coli* bacterial burden in a mouse model of septicemia by two to three orders of magnitude.⁹² While further research on riboswitches as drug targets is warranted, caveats include the potential for gain of mutations as well as the observed difficulty for some modulators to cross the bacterial cell wall. Nevertheless, these molecules represent unique targets and an opportunity for novel antimicrobials.⁹³

A promising approach for new antibacterials is antivirulence, which focuses on targeting pathogenic bacteria by binding to and inhibiting virulence factors such as toxins, particularly those with enzymatic activity.^{94,95} For example, a small molecule was identified *in silico* and confirmed *in vitro* and *in vivo* to target the enzymatic site of Shiga toxin from *Shigella dysenteriae*.⁹⁶ In theory, because these virulence factors are not necessary for bacterial survival outside the context of a host infection, this approach will not apply direct selective pressure and therefore resistance should not rapidly emerge. Antivirulence treatments are anticipated to supplement small molecule therapeutics and improve their efficacies, resulting in optimal clinical treatment regimens.

Drug repurposing

The approval of new antibiotics has declined during the past three decades (Figure 2) while resistance to the approved antibiotics continues to increase. Despite an urgent need for new antibiotics, most major pharmaceutical companies have moved away from the antibacterial market thereby drying up the pipeline for new treatments.² Therefore, repurposing of approved drugs to treat infectious diseases might provide an alternative approach for effective new therapies.⁹⁸ Approved drugs repurposed for a new indication can be rapidly advanced to phase II clinical trials without a phase I clinical trial due to the existing preclinical, human pharmacokinetics, and drug safety data. An example of repurposing a drug to treat bacterial infections is provided by the gold-containing compound auranofin. This drug is FDA-approved for the treatment of rheumatoid arthritis, but has also demonstrated antimicrobial activity against a range of Gram-positive and Gram-negative pathogens.^{98–105} The thiol ligand of auranofin can form irreversible adducts with selenol or thiol groups, and auranofin is a well-known inhibitor of reduction/oxidation enzymes including thioredoxin reductase (TrxR).⁹⁸ The antibacterial activity of auranofin has been linked to the disruption of selenium metabolism and interference with a number of selenoproteins including glycine reductase, proline reductase, and xanthine dehydrogenase.^{99–101}

Biochemical studies have demonstrated that auranofin is a potent inhibitor of bacterial TrxR, which is not targeted by other antibiotics and is involved in a variety of processes required for survival and proliferation, such as DNA synthesis and protein repair.¹⁰⁴ The ability of auranofin to interfere with multiple bacterial enzymes might reduce the development of resistance.

Synergistic combinations

Drug combinations have been approved for the treatment of bacterial infections, such as the combination of amoxicillin (a β -lactam) with clavulanic acid (a β -lactamase inhibitor).¹⁰⁶ Advantages of drug combinations include extending the spectrum of susceptible organisms, overcoming drug resistance, reducing the development of antibiotic resistance, and the potential for drug synergies.⁹⁷ New combinations of antibiotics have been successfully applied to treat drug-resistant pathogens. For example, a combination of three β -lactams (meropenem, piperacillin, tazobactam) was reported to act synergistically against MRSA and showed similar *in vivo* activity to the oxazolidinone antibiotic linezolid.¹⁰⁷ While Gram-negative bacteria are intrinsically resistant to the antibiotic vancomycin, a study showed strong synergies against *E. coli* when combining vancomycin with antibiotics such as trimethoprim or nitrofurantoin.¹⁰⁸ The drug fosfomycin, which has been registered in the United States for oral treatment of uncomplicated UTIs since 1995, has been tested as a participant in a cocktail with polymyxin B, tobramycin or ciprofloxacin to target *P. aeruginosa*, although issues with resistance remain.^{109,110} Other novel combinations of antibiotics have shown synergy against Gram-negative pathogens, including *Acinetobacter baumannii*.¹¹¹ Progress has also been made in identifying synergistic drug combinations that combine antibiotics with approved drugs not previously used to treat infections. As an example, the antibiotic penicillin G is primarily only active against Gram-positive organisms, but showed synergistic activity with the non-antibiotic drug promethazine against *E. coli*.¹¹² The opioid receptor agonist loperamide (Imodium) is an approved anti-diarrheal that was shown to synergize with tetracyclines due to its ability to dissipate bacterial membrane potential and enhance antibiotic uptake.¹¹³ Recently, a drug combination screen identified 17 synergistic three-drug combinations that were effective against *K. pneumoniae* at concentrations which are clinically achievable. Additionally, three sets of three-drug combinations, two of which contained auranofin, were active against a panel of 10 MDR clinical isolates, including *K. pneumoniae*, *A. baumannii*, *P. aeruginosa*, *Citrobacter freundii*, *Enterobacter cloacae*, and *E. coli*.¹⁰⁵ While the previous examples of combinations utilized antibiotics, an interesting strategy was recently applied to completely replace an antibiotic pair with non-antibiotic drugs. The aim was to identify synergistic pairs based on the antibiotics trimethoprim and sulfamethoxazole that have been used in combination for decades to treat bacterial infections. First, the antiviral drug azidothymidine (AZT) was found to synergize with trimethoprim or sulfamethoxazole against MDR *E. coli* and

K. pneumoniae. The development of a mechanistic understanding of the underlying interaction enabled the discovery of synergy between AZT and the cancer drug floxuridine, which reduced the median bacterial burden by 10,000-fold compared with the traditional pair of trimethoprim/sulfamethoxazole in a zebrafish model.¹¹⁴

The potential opportunities to repurpose drugs, although exciting, are not without challenges.^{115–117} There are safety concerns associated with utilizing drugs known to target host tissues, such as oncology medications.¹¹⁶ Additionally there are currently major economic barriers to repurposing because there are insufficient market incentives to advance drugs that are off-patent. It has been estimated that repurposing a drug costs around \$300 million and takes about 6.5 years.¹¹⁷ Other current challenges include intellectual property rights and other legal concerns, regulatory issues and timelines, return on investment, as well as the design and execution of proof-of-concept clinical trials.

Financial concerns

Another unsolved but important issue in the search for effective small molecule antibiotics is that of return on investment. Most pharmaceutical companies are in business to make a profit for their shareholders, and antimicrobial drug discovery is currently not profitable. It was recently estimated that the cost of a new drug with market approval is \$2.6 billion and takes more than a decade to develop.¹¹⁸ When a new antibiotic is finally available to the public, it is prescribed only transiently for infections and not needed by otherwise healthy people on a regular basis, unlike medications that lower cholesterol or treat high blood pressure. To address these concerns and incentivize antibacterial drug development, in 2012 the 112th United States Congress passed the Food and Drug Administration Safety and Innovation Act,¹¹⁹ which included as Title VIII the “Generating Antibiotic Incentives Now” (GAIN) Act. As a commercial inducement, the GAIN Act extended the market exclusivity period, which is the length of time that a pharmaceutical company is allowed to be the sole provider of an antimicrobial by law. At the same time, an FDA Task Force was launched to identify challenges and provide guidance for antibiotic drug resistance.

A novel solution to this problem has been put forward which consists of incentivizing antibacterial research and development by drug companies using a model that is either fully or partially delinked from drug sales through milestone payments and other financial encouragements.¹²⁰ The successful implementation of this model would require both multinational coordination and cooperation as well as regional and global banking institutional support. However, because the health threat posed by antibiotic-resistant organisms affects the population worldwide, this approach might have merit and should be subject to serious consideration.

Conclusions

It is not an exaggeration to describe our current situation as an evolutionary “arms race” against antibiotic-resistant microorganisms. There is not likely to be a single compound or class of drugs that will continue to be effective for the foreseeable future against bacterial infections. Rather, the prediction is that we will discover novel treatments (natural, man-made or new combinations) that will initially be valuable additions to our arsenal of antibiotics, but which will eventually lose efficacy due to bacterial resistance mechanisms. This is a clear indication that academia, industry and government must be called upon to invest in a concerted and unified effort to fill the drug development pipeline with new antimicrobials. The first antibiotic was released to the public more than 80 years ago, which means that there are people alive today that were born in the pre-antibiotic era. For those of us that were raised in a world with available and effective treatments, imagining a future without antibiotics is daunting indeed.

AUTHORS' CONTRIBUTIONS: All authors participated in the writing and review of the manuscript.

ACKNOWLEDGMENTS

Minireview space limitations prevented citation of all relevant literature. The authors wish to thank Noel Southall for helpful discussions.

DECLARATION OF CONFLICT OF INTEREST

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

FUNDING

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: in part by U01 DC014756 (DAD) and the Intramural Research Program of the National Institutes of Health, National Center for Advancing Translational Sciences (NPC, TP, GZ-K and MDH).

REFERENCES

1. FDIS I. 11238 Health informatics – identification of medicinal products – data elements and structures for the unique identification and exchange of regulated information on substances. Available at: <http://bit.ly/obxMhU> (accessed October 2011)
2. Infectious Diseases Society of A, Spellberg B, Blaser M, Guidos RJ, Boucher HW, Bradley JS, Eisenstein BI, Gerding D, Lynfield R, Reller LB, Rex J, Schwartz D, Septimus E, Tenover FC, Gilbert DN. Combating antimicrobial resistance: policy recommendations to save lives. *Clin Infect Dis* 2011;**52**:S397–428
3. Khan MZ, Bhaskar A, Upadhyay S, Kumari P, Rajmani RS, Jain P, Singh A, Kumar D, Bhavesh NS, Nandicoori VK. Protein kinase G confers survival advantage to *Mycobacterium tuberculosis* during latency like conditions. *J Biol Chem* 2017;**292**:16093–108
4. Sิริyong T, Srimanote P, Chusri S, Yingyongnarongkul BE, Suaisom C, Tipmanee V, Voravuthikunchai SP. Conessina as a novel inhibitor of

- multidrug efflux pump systems in *Pseudomonas aeruginosa*. *BMC Complement Altern Med* 2017;**17**:405
5. Eswari JS, Dhagat S, Kaser S, Tiwari A. Homology modeling and molecular docking studies of Bacillomycin and Iturin synthetases with novel ligands for the production of therapeutic lipopeptides. *Curr Drug Discov Technol*. Epub ahead of print 15 August 2017. doi: 10.2174/1570163814666170816112536.
 6. Forsberg KJ, Reyes A, Wang B, Selleck EM, Sommer MO, Dantas G. The shared antibiotic resistance of soil bacteria and human pathogens. *Science* 2012;**337**:1107–11
 7. Kohanski MA, Dwyer DJ, Collins JJ. How antibiotics kill bacteria: from targets to networks. *Nat Rev Microbiol*. 2010;**8**:423–35
 8. Cho H, Uehara T, Bernhardt TG. Beta-lactam antibiotics induce a lethal malfunctioning of the bacterial cell wall synthesis machinery. *Cell* 2014;**159**:1300–11
 9. Liu Y, Breukink E. The membrane steps of bacterial cell wall synthesis as antibiotic targets. *Antibiotics* 2016;**5**:28
 10. Miller WR, Bayer AS, Arias CA. Mechanism of action and resistance to daptomycin in *Staphylococcus aureus* and Enterococci. *Cold Spring Harb Perspect Med* 2016;**6**. doi: 10.1101/cshperspect.a026997.
 11. Velkov T, Roberts KD, Nation RL, Thompson PE, Li J. Pharmacology of polymyxins: new insights into an 'old' class of antibiotics. *Future Microbiol* 2013;**8**:711–24
 12. Spizek J, Rezanka T. Lincosamides: chemical structure, biosynthesis, mechanism of action, resistance, and applications. *Biochem Pharmacol* 2017;**133**:20–8
 13. Chukwudi CU. rRNA binding sites and the molecular mechanism of action of the tetracyclines. *Antimicrob Agents Chemother* 2016;**60**:4433–41
 14. Opegard LM, Schwanz HA, Towle TR, Kerns RJ, Hiasa H. Fluoroquinolones stimulate the DNA cleavage activity of topoisomerase IV by promoting the binding of Mg(2+) to the second metal binding site. *Biochim Biophys Acta* 2016;**1860**:569–75
 15. Calvori C, Frontali L, Leoni L, Tecce G. Effect of rifamycin on protein synthesis. *Nature* 1965;**207**:417–8
 16. Hillier SL, Nyirjesy P, Waldbaum AS, Schwabke JR, Morgan FG, Adetoro NA, Braun CJ. Secnidazole treatment of bacterial vaginosis: a randomized controlled trial. *Obstet Gynecol* 2017;**130**:379–86
 17. Bush K, Page MG. What we may expect from novel antibacterial agents in the pipeline with respect to resistance and pharmacodynamic principles. *J Pharmacokinet Pharmacodyn* 2017;**44**:113–32
 18. Candel FJ, Penuelas M. Delafloxacin: design, development and potential place in therapy. *Drug Des Dev Ther* 2017;**11**:881–91
 19. Fleming A. On the antibacterial action of cultures of a penicillium, with special reference to their use in the isolation of *B. influenzae* 1929. *Bull World Health Organ* 2001;**79**:780–90
 20. Li J, Jia H, Cai X, Zhong H, Feng Q, Sunagawa S, Arumugam M, Kultima JR, Prifti E, Nielsen T, Juncker AS, Manichanh C, Chen B, Zhang W, Levenez F, Wang J, Xu X, Xiao L, Liang S, Zhang D, Zhang Z, Chen W, Zhao H, Al-Aama JY, Edris S, Yang H, Hansen T, Nielsen HB, Brunak S, Kristiansen K, Guarner F, Pedersen O, Dore J, Ehrlich SD, Bork P. An integrated catalog of reference genes in the human gut microbiome. *Nat Biotechnol* 2014;**32**:834–41
 21. Eckburg PB, Bik EM, Bernstein CN, Purdom E, Dethlefsen L, Sargent M, Gill SR, Nelson KE, Relman DA. Diversity of the human intestinal microbial flora. *Science* 2005;**308**:1635–8
 22. Zeissig S, Blumberg RS. Life at the beginning: perturbation of the microbiota by antibiotics in early life and its role in health and disease. *Nat Immunol* 2014;**15**:307–10
 23. Franzosa EA, Huang K, Meadow JF, Gevers D, Lemon KP, Bohannan BJ, Huttenhower C. Identifying personal microbiomes using metagenomic codes. *Proc Natl Acad Sci U S A* 2015;**112**:E2930–8
 24. Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, Nielsen T, Pons N, Levenez F, Yamada T, Mende DR, Li J, Xu J, Li S, Li D, Cao J, Wang B, Liang H, Zheng H, Xie Y, Tap J, Lepage P, Bertalan M, Batto JM, Hansen T, Le Paslier D, Linneberg A, Nielsen HB, Pelletier E, Renault P, Sicheritz-Ponten T, Turner K, Zhu H, Yu C, Jian M, Zhou Y, Li Y, Zhang X, Qin N, Yang H, Wang J, Brunak S, Dore J, Guarner F, Kristiansen K, Pedersen O, Parkhill J, Weissenbach J, Bork P, Ehrlich SD. A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* 2010;**464**:59–65
 25. Cario E, Gerken G, Podolsky DK. Toll-like receptor 2 controls mucosal inflammation by regulating epithelial barrier function. *Gastroenterology* 2007;**132**:1359–74
 26. Long SL, Gahan CGM, Joyce SA. Interactions between gut bacteria and bile in health and disease. *Mol Aspects Med* 2017;**56**:54–65
 27. Hooper LV, Midtvedt T, Gordon JI. How host-microbial interactions shape the nutrient environment of the mammalian intestine. *Annu Rev Nutr* 2002;**22**:283–307
 28. Mikov MM, Stojančević MP, Bojić GM. Probiotics as a promising treatment for inflammatory bowel disease. *Hosp Pharmacol* 2014;**1**:52–60
 29. Bohnhoff M, Miller CP. Enhanced susceptibility to Salmonella infection in streptomycin-treated mice. *J Infect Dis* 1962;**111**:117–27
 30. Buffie CG, Jarchum I, Equinda M, Lipuma L, Gobburne A, Viale A, Ubeda C, Xavier J, Pamer EG. Profound alterations of intestinal microbiota following a single dose of clindamycin results in sustained susceptibility to *Clostridium difficile*-induced colitis. *Infect Immun* 2012;**80**:62–73
 31. Reeves AE, Theriot CM, Bergin IL, Huffnagle GB, Schloss PD, Young VB. The interplay between microbiome dynamics and pathogen dynamics in a murine model of *Clostridium difficile* Infection. *Gut Microbes* 2011;**2**:145–58
 32. Donskey CJ, Chowdhry TK, Hecker MT, Hoyen CK, Hanrahan JA, Hujer AM, Hutton-Thomas RA, Whalen CC, Bonomo RA, Rice LB. Effect of antibiotic therapy on the density of vancomycin-resistant enterococci in the stool of colonized patients. *N Engl J Med* 2000;**343**:1925–32
 33. Yang B, Hagberg KW, Chen J, Sahasrabudhe VV, Graubard BI, Jick S, McGlynn KA. Associations of antibiotic use with risk of primary liver cancer in the Clinical Practice Research Datalink. *Br J Cancer* 2016;**115**:85–9
 34. Dik VK, van Oijen MG, Smeets HM, Siersema PD. Frequent use of antibiotics is associated with colorectal cancer risk: results of a nested case-control study. *Dig Dis Sci* 2016;**61**:255–64
 35. Boursi B, Haynes K, Mamtani R, Yang YX. Impact of antibiotic exposure on the risk of colorectal cancer. *Pharmacoepidemiol Drug Saf* 2015;**24**:534–42
 36. Schuller A, Blaser MJ. Risks of antibiotic exposures early in life on the developing microbiome. *PLoS Pathog* 2015;**11**:e1004903
 37. Malik U, Armstrong D, Ashworth M, Dregan A, L'esperance V, McDonnell L, Molokhia M, White P. Association between prior antibiotic therapy and subsequent risk of community-acquired infections: a systematic review. *J Antimicrob Chemother* 2018;**73**:287–96
 38. Pandey DP, Gerdes K. Toxin-antitoxin loci are highly abundant in free-living but lost from host-associated prokaryotes. *Nucleic Acids Res* 2005;**33**:966–76
 39. Van Melderen L, Thi MH, Lecchi P, Gottesman S, Couturier M, Maurizi MR. ATP-dependent degradation of CcdA by Lon protease. *J Biol Chem* 1996;**271**:27730–8
 40. Aizenman E, Engelberg-Kulka H, Glaser G. An *Escherichia coli* chromosomal "addiction module" regulated by guanosine [corrected] 3',5'-bispyrophosphate: a model for programmed bacterial cell death. *Proc Natl Acad Sci U S A* 1996;**93**:6059–63
 41. Jorgensen BB, D'hondt S. Ecology. A starving majority deep beneath the seafloor. *Science* 2006;**314**:932–4
 42. Durso LM, Cook KL. Impacts of antibiotic use in agriculture: what are the benefits and risks?. *Curr Opin Microbiol* 2014;**19**:37–44
 43. Jiang HX, Lü DH, Chen ZL, Wang XM, Chen JR, Liu YH, Liao XP, Liu JH, Zeng ZL. High prevalence and widespread distribution of multi-resistant *Escherichia coli* isolates in pigs and poultry in China. *Vet J* 2011;**187**:99–103
 44. Marshall BM, Levy SB. Food animals and antimicrobials: impacts on human health. *Clin Microbiol Rev* 2011;**24**:718–33
 45. Hussain HI, Iqbal Z, Seleem MN, Huang D, Sattar A, Hao H, Yuan Z. Virulence and transcriptome profile of multidrug-resistant *Escherichia coli* from chicken. *Sci Rep* 2017;**7**:8335
 46. Spellberg B, Hansen GR, Kar A, Cordova CD, Price LB, Johnson JR. *Antibiotic resistance in humans and animals*. National Academy of

- Medicine, 2016. Available at: <https://nam.edu/antibiotic-resistance-in-humans-and-animals/>
47. Bélanger L, Garenaux A, Harel J, Boulianne M, Nadeau E, Dozois CM. *Escherichia coli* from animal reservoirs as a potential source of human extraintestinal pathogenic *E. coli*. *FEMS Immunol Med Microbiol* 2011;**62**:1–10
 48. He LY, Ying GG, Liu YS, Su HC, Chen J, Liu SS, Zhao JL. Discharge of swine wastes risks water quality and food safety: antibiotics and antibiotic resistance genes from swine sources to the receiving environments. *Environ Int* 2016;**92–93**:2109
 49. Jia S, Zhang XX, Miao Y, Zhao Y, Ye L, Li B, Zhang T. Fate of antibiotic resistance genes and their associations with bacterial community in livestock breeding wastewater and its receiving river water. *Water Res* 2017;**124**:259–68
 50. Marti R, Scott A, Tien YC, Murray R, Sabourin L, Zhang Y, Topp E. Impact of manure fertilization on the abundance of antibiotic-resistant bacteria and frequency of detection of antibiotic resistance genes in soil and on vegetables at harvest. *Appl Environ Microbiol* 2013;**79**:5701–9
 51. Casey JA, Curriero FC, Cosgrove SE, Nachman KE, Schwartz BS. High-density livestock operations, crop field application of manure, and risk of community-associated methicillin-resistant *Staphylococcus aureus* infection in Pennsylvania. *JAMA Intern Med* 2013;**173**:1980–90
 52. Morel CM, Edwards SE, Harbarth S. Preserving the ‘commons’: addressing the sustainable use of antibiotics through an economic lens. *Clin Microbiol Infect* 2017;**23**:718–22
 53. Lübbert C, Baars C, Dayakar A, Lippmann N, Rodloff AC, Kinzig M, Sörgel F. Environmental pollution with antimicrobial agents from bulk drug manufacturing industries in Hyderabad, South India, is associated with dissemination of extended-spectrum beta-lactamase and carbapenemase-producing pathogens. *Infection* 2017;**45**:479–91
 54. D’Costa VM, King CE, Kalan L, Morar M, Sung WW, Schwarz C, Froese D, Zazula G, Calmels F, Debruyne R, Golding GB, Poinar HN, Wright GD. Antibiotic resistance is ancient. *Nature* 2011;**477**:457–61
 55. Waglchner N, Wright GD. Antibiotic resistance: it’s bad, but why isn’t it worse?. *BMC Biol* 2017;**15**:84
 56. Porse A, Schonning K, Munck C, Sommer MO. Survival and evolution of a large multidrug resistance plasmid in new clinical bacterial hosts. *Mol Biol Evol* 2016;**33**:2860–73
 57. Jaidane N, Naas T, Mansour W, Radhia BB, Jerbi S, Boujaafar N, Bouallegue O, Bonnin RA. Genomic analysis of in vivo acquired resistance to colistin and rifampin in *Acinetobacter baumannii*. *Int J Antimicrob Agents*. Epub ahead of print 7 November 2017. doi: 10.1016/j.ijantimicag.2017.10.016.
 58. Zgurskaya HI, Lopez CA, Gnanakaran S. Permeability barrier of Gram-negative cell envelopes and approaches to bypass it. *ACS Infect Dis* 2015;**1**:512–22
 59. Randall CP, Mariner KR, Chopra I, O’neill AJ. The target of daptomycin is absent from *Escherichia coli* and other gram-negative pathogens. *Antimicrob Agents Chemother* 2013;**57**:637–9
 60. Boucher HW, Ambrose PG, Chambers HF, Ebricht RH, Jezek A, Murray BE, Newland JG, Ostrowsky B, Rex JH. America IDSo White Paper: developing antimicrobial drugs for resistant pathogens, narrow-spectrum indications, and unmet needs. *J Infect Dis* 2017;**216**:228–36
 61. Munita JM, Arias CA. Mechanisms of antibiotic resistance. *Microbiol Spectr* 2016;**4**. doi: 10.1128/microbiolspec.VMBF-0016-2015.
 62. Blair JM, Webber MA, Baylay AJ, Ogbolu DO, Piddock LJ. Molecular mechanisms of antibiotic resistance. *Nat Rev Microbiol* 2015;**13**:42–51
 63. Fernandez L, Hancock RE. Adaptive and mutational resistance: role of porins and efflux pumps in drug resistance. *Clin Microbiol Rev* 2012;**25**:661–81
 64. Lambert PA. Bacterial resistance to antibiotics: modified target sites. *Adv Drug Deliv Rev* 2005;**57**:1471–85
 65. Sandegren L, Andersson DI. Bacterial gene amplification: implications for the evolution of antibiotic resistance. *Nat Rev Microbiol* 2009;**7**:578–88
 66. Wright GD. Bacterial resistance to antibiotics: enzymatic degradation and modification. *Adv Drug Deliv Rev* 2005;**57**:1451–70
 67. Correia S, Poeta P, Hebraud M, Capelo JL, Igrejas G. Mechanisms of quinolone action and resistance: where do we stand?. *J Med Microbiol* 2017;**66**:551–9
 68. Wozniak A, Villagra NA, Undabarrena A, Gallardo N, Keller N, Moraga M, Roman JC, Mora GC, Garcia P. Porin alterations present in non-carbapenemase-producing Enterobacteriaceae with high and intermediate levels of carbapenem resistance in Chile. *J Med Microbiol* 2012;**61**:1270–9
 69. Lavigne JP, Sotto A, Nicolas-Chanoine MH, Bouziges N, Pages JM, Davin-Regli A. An adaptive response of *Enterobacter aerogenes* to imipenem: regulation of porin balance in clinical isolates. *Int J Antimicrob Agents* 2013;**41**:130–6
 70. Piddock LJ. Multidrug-resistance efflux pumps – not just for resistance. *Nat Rev Microbiol* 2006;**4**:629–36
 71. McMurry L, Petrucci RE, Jr, Levy SB. Active efflux of tetracycline encoded by four genetically different tetracycline resistance determinants in *Escherichia coli*. *Proc Natl Acad Sci U S A* 1980;**77**:3974
 72. Blair JM, Bavro VN, Ricci V, Modi N, Cacciotto P, Kleinekathfer U, Ruggerone P, Vargiu AV, Baylay AJ, Smith HE, Brandon Y, Galloway D, Piddock LJ. AcrB drug-binding pocket substitution confers clinically relevant resistance and altered substrate specificity. *Proc Natl Acad Sci U S A* 2015;**112**:3511–6
 73. Baroud M, Dandache I, Araj GF, Wakim R, Kanj S, Kanafani Z, Khairallah M, Sabra A, Shehab M, Dbaibo G, Matar GM. Underlying mechanisms of carbapenem resistance in extended-spectrum beta-lactamase-producing *Klebsiella pneumoniae* and *Escherichia coli* isolates at a tertiary care centre in Lebanon: role of OXA-48 and NDM-1 carbapenemases. *Int J Antimicrob Agents* 2013;**41**:75–9
 74. Xavier DE, Picao RC, Girardello R, Fehlberg LC, Gales AC. Efflux pumps expression and its association with porin down-regulation and beta-lactamase production among *Pseudomonas aeruginosa* causing bloodstream infections in Brazil. *BMC Microbiol* 2010;**10**:217
 75. Howden BP, Grayson ML. Dumb and dumber – the potential waste of a useful antistaphylococcal agent: emerging fusidic acid resistance in *Staphylococcus aureus*. *Clin Infect Dis* 2006;**42**:394–400
 76. Tomlinson JH, Thompson GS, Kalverda AP, Zhuravleva A, O’neill AJ. A target-protection mechanism of antibiotic resistance at atomic resolution: insights into FusB-type fusidic acid resistance. *Sci Rep* 2016;**6**:19524
 77. Redgrave LS, Sutton SB, Webber MA, Piddock LJ. Fluoroquinolone resistance: mechanisms, impact on bacteria, and role in evolutionary success. *Trends Microbiol* 2014;**22**:438–45
 78. Long KS, Poehlsgaard J, Kehrenberg C, Schwarz S, Vester B. The Cfr rRNA methyltransferase confers resistance to phenicols, lincosamides, oxazolidinones, pleuromutilins, and streptogramin A antibiotics. *Antimicrob Agents Chemother* 2006;**50**:2500–5
 79. Chambers HF, Deleo FR. Waves of resistance: *Staphylococcus aureus* in the antibiotic era. *Nat Rev Microbiol* 2009;**7**:629–41
 80. Flensburg J, Skold O. Massive overproduction of dihydrofolate reductase in bacteria as a response to the use of trimethoprim. *Eur J Biochem* 1987;**162**:473–6
 81. Abraham EP, Chain E. An enzyme from bacteria able to destroy penicillin. *Nature* 1940;**146**:837
 82. Abraham EP, Gardner AD, Chain E, Heatley NG, Fletcher CM, Jennings MA. Further observations on penicillin. *Lancet* 1941;**2**:177–89
 83. Drawz SM, Bonomo RA. Three decades of beta-lactamase inhibitors. *Clin Microbiol Rev* 2010;**23**:160–201
 84. Ramirez MS, Tolmasky ME. Aminoglycoside modifying enzymes. *Drug Resist Update* 2010;**13**:151–71
 85. Raymond JW. RASCAL: calculation of graph similarity using maximum common edge subgraphs. *Comput J* 2002;**45**:631–44
 86. Gardiner EJ, Gillet VJ, Willett P, Cosgrove DA. Representing clusters using a maximum common edge substructure algorithm applied to reduced graphs and molecular graphs. *J Chem Inf Model* 2007;**47**:354–66

87. Maaten L, Hinton G. Visualizing data using t-SNE. *J Mach Learn Res* 2008;**9** Nov: 2579–605
88. Maffioli SI, Zhang Y, Degen D, Carzaniga T, Del Gatto G, Serina S, Monciardini P, Mazzetti C, Guglielame P, Candiani G, Chiriack AI, Facchetti G, Kaltofen P, Sahl HG, Deho G, Donadio S, Ebricht RH. Antibacterial nucleoside-analog inhibitor of bacterial RNA polymerase. *Cell* 2017;**169**:1240–8 e23
89. Schiebel J, Chang A, Shah S, Lu Y, Liu L, Pan P, Hirschbeck MW, Tareilus M, Eltschkner S, Yu W, Cummings JE, Knudson SE, Bommineni GR, Walker SG, Slayden RA, Sotriffer CA, Tonge PJ, Kisker C. Rational design of broad spectrum antibacterial activity based on a clinically relevant enoyl-acyl carrier protein (ACP) reductase inhibitor. *J Biol Chem* 2014;**289**:15987–6005
90. Li W, Sanchez-Hidalgo A, Jones V, de Moura VC, North EJ, Jackson M. Synergistic interactions of MmpL3 inhibitors with antitubercular compounds in vitro. *Antimicrob Agents Chemother* 2017;**61**. doi: 10.1128/AAC.02399-1.
91. Richter MF, Drown BS, Riley AP, Garcia A, Shirai T, Svec RL, Hergenrother PJ. Predictive compound accumulation rules yield a broad-spectrum antibiotic. *Nature* 2017;**545**:299–304
92. Howe JA, Wang H, Fischmann TO, Balibar CJ, Xiao L, Galgoczi AM, Malinverni JC, Mayhood T, Villafania A, Nahvi A, Murgolo N, Barbieri CM, Mann PA, Carr D, Xia E, Zuck P, Riley D, Painter RE, Walker SS, Sherborne B, de Jesus R, Pan W, Plotkin MA, Wu J, Rindgen D, Cummings J, Garlisi CG, Zhang R, Sheth PR, Gill CJ, Tang H, Roemer T. Selective small-molecule inhibition of an RNA structural element. *Nature* 2015;**526**:672–7
93. Dersch P, Khan MA, Muhlen S, Gorke B. Roles of regulatory RNAs for antibiotic resistance in bacteria and their potential value as novel drug targets. *Front Microbiol* 2017;**8**:803
94. Coussens NP, Daines DA. Wake me when it's over – bacterial toxin-antitoxin proteins and induced dormancy. *Exp Biol Med* 2016;**241**:1332–42
95. Hauser AR, Mecas J, Moir DT. Beyond antibiotics: new therapeutic approaches for bacterial infections. *Clin Infect Dis* 2016;**63**:89–95
96. Chauhan V, Chaudhary D, Pathak U, Saxena N, Dhaked RK. In silico discovery and validation of amide based small molecule targeting the enzymatic site of shiga toxin. *J Med Chem* 2016;**59**:10763–73
97. Zheng W, Sun W, Simeonov A. Drug repurposing screens and synergistic drug-combinations for infectious diseases. *Br J Pharmacol* 2018;**175**:181–91
98. Roder C, Thomson MJ. Auranofin: repurposing an old drug for a golden new age. *Drugs R&D* 2015;**15**:13–20
99. Jackson-Rosario S, Self WT. Inhibition of selenium metabolism in the oral pathogen *Treponema denticola*. *J Bacteriol* 2009;**191**:4035–40
100. Jackson-Rosario S, Cowart D, Myers A, Tarrion R, Levine RL, Scott RA, Self WT. Auranofin disrupts selenium metabolism in *Clostridium difficile* by forming a stable Au-Se adduct. *J Biol Inorg Chem* 2009;**14**:507–19
101. Srivastava M, Mallard C, Barke T, Hancock LE, Self WT. A selenium-dependent xanthine dehydrogenase triggers biofilm proliferation in *Enterococcus faecalis* through oxidant production. *J Bacteriol* 2011;**193**:1643–52
102. Hokai Y, Jurkowicz B, Fernandez-Gallardo J, Zakirkhodjaev N, Sanau M, Muth TR, Contel M. Auranofin and related heterometallic gold(I)-thiolates as potent inhibitors of methicillin-resistant *Staphylococcus aureus* bacterial strains. *J Inorg Biochem* 2014;**138**:81–8
103. Aguinagalde L, Diez-Martinez R, Yuste J, Royo I, Gil C, Casa I, Martin-Fontecha M, Marin-Ramos NI, Ardanuy C, Linares J, Garcia P, Garcia E, Sanchez-Puelles JM. Auranofin efficacy against MDR *Streptococcus pneumoniae* and *Staphylococcus aureus* infections. *J Antimicrob Chemother* 2015;**70**:2608–17
104. Harbut MB, Vilcheze C, Luo X, Hensler ME, Guo H, Yang B, Chatterjee AK, Nizet V, Jacobs WR, Jr, Schultz PG, Wang F. Auranofin exerts broad-spectrum bactericidal activities by targeting thiol-redox homeostasis. *Proc Natl Acad Sci U S A* 2015;**112**:4453–8
105. Sun W, Weingarten RA, Xu M, Southall N, Dai S, Shinn P, Sanderson PE, Williamson PR, Frank KM, Zheng W. Rapid antimicrobial susceptibility test for identification of new therapeutics and drug combinations against multidrug-resistant bacteria. *Emerg Microbes Infect* 2016;**5**:e116
106. Smith GM, Slocombe B, Abbott KH, Mizen LW. Activity of amoxicillin-clavulanate against penicillin-resistant *Streptococcus pneumoniae* in an experimental respiratory infection model in rats. *Antimicrob Agents Chemother* 1998;**42**:813–7
107. Gonzales PR, Pesesky MW, Bouley R, Ballard A, Bidy BA, Suckow MA, Wolter WR, Schroeder VA, Burnham CA, Mobashery S, Chang M, Dantas G. Synergistic, collaterally sensitive beta-lactam combinations suppress resistance in MRSA. *Nat Chem Biol* 2015;**11**:855–61
108. Zhou A, Kang TM, Yuan J, Beppler C, Nguyen C, Mao Z, Nguyen MQ, Yeh P, Miller JH. Synergistic interactions of vancomycin with different antibiotics against *Escherichia coli*: trimethoprim and nitrofurantoin display strong synergies with vancomycin against wild-type E. coli. *Antimicrob Agents Chemother* 2015;**59**:276–81
109. Silver LL. Fosfomycin: mechanism and resistance. *Cold Spring Harb Perspect Med*. Epub ahead of print 6 January 2017. doi: 10.1101/cshperspect.a025262.
110. Walsh CC, Landersdorfer CB, McIntosh MP, Peleg AY, Hirsch EB, Kirkpatrick CM, Bergen PJ. Clinically relevant concentrations of fosfomycin combined with polymyxin B, tobramycin or ciprofloxacin enhance bacterial killing of *Pseudomonas aeruginosa*, but do not suppress the emergence of fosfomycin resistance. *J Antimicrob Chemother* 2016;**71**:2218–29
111. Hornsey M, Longshaw C, Phee L, Wareham DW. In vitro activity of telavancin in combination with colistin versus Gram-negative bacterial pathogens. *Antimicrob Agents Chemother* 2012;**56**:3080–5
112. Lehtinen J, Lilius EM. Promethazine renders *Escherichia coli* susceptible to penicillin G: real-time measurement of bacterial susceptibility by fluoro-luminometry. *Int J Antimicrob Agents* 2007;**30**:44–51
113. Ejim L, Farha MA, Falconer SB, Wildenhain J, Coombes BK, Tyers M, Brown ED, Wright GD. Combinations of antibiotics and nonantibiotic drugs enhance antimicrobial efficacy. *Nat Chem Biol* 2011;**7**:348–50
114. Wambaugh MA, Shakya VPS, Lewis AJ, Mulvey MA, Brown JCS. High-throughput identification and rational design of synergistic small-molecule pairs for combating and bypassing antibiotic resistance. *PLoS Biol* 2017;**15**:e2001644
115. Novac N. Challenges and opportunities of drug repositioning. *Trends Pharmacol Sci* 2013;**34**:267–72
116. Rangel-Vega A, Bernstein LR, Mandujano-Tinoco EA, Garcia-Contreras SJ, Garcia-Contreras R. Drug repurposing as an alternative for the treatment of recalcitrant bacterial infections. *Front Microbiol* 2015;**6**:282
117. Nosengo N. Can you teach old drugs new tricks?. *Nature* 2016;**534**:314–6
118. DiMasi JA, Grabowski HG, Hansen RW. Innovation in the pharmaceutical industry: new estimates of R&D costs. *J Health Econ* 2016;**47**:20–33
119. 112th Congress: Food and Drug Administration Safety and Innovation Act, 2012.
120. Årdal C, Baraldi E, Ciabuschi F, Outtersson K, Rex JH, Piddock LJV, Findlay D, Committee D-AS. To the G20: incentivising antibacterial research and development. *Lancet Infect Dis* 2017;**17**:799–801