



Published in final edited form as:

Nature. 2014 November 13; 515(7526): 228–233. doi:10.1038/nature13846.

Clathrin regenerates synaptic vesicles from endosomes

Shigeki Watanabe¹, Thorsten Trimbuch², Marcial Camacho-Pérez², Benjamin R. Rost^{2,3}, Bettina Brokowski², Berit Söhl-Kielczynski², Annegret Felies², M. Wayne Davis¹, Christian Rosenmund^{2,*}, and Erik M. Jorgensen^{1,*}

¹Department of Biology and Howard Hughes Medical Institute, University of Utah, Salt Lake City, UT 84112-0840, U.S.A.

²Neuroscience Research Center Charité Universitätsmedizin Berlin, Berlin, 10117, Germany

³German Center for Neurodegenerative Diseases (DZNE), 10117 Berlin, Germany

Summary

Ultrafast endocytosis can retrieve a single large endocytic vesicle as fast as 50-100 ms after synaptic vesicle fusion. However, the fate of the large endocytic vesicles is not known. Here we demonstrate that these vesicles transition to a synaptic endosome about one second after stimulation. The endosome is resolved into coated vesicles after 3 seconds, which in turn become small-diameter synaptic vesicles 5-6 seconds after stimulation. We disrupted clathrin function using RNAi and found that clathrin is not required for ultrafast endocytosis but is required to generate synaptic vesicles from the endosome. Ultrafast endocytosis fails when actin polymerization is disrupted, or when neurons are stimulated at room temperature instead of physiological temperature. In the absence of ultrafast endocytosis, synaptic vesicles are retrieved directly from the plasma membrane by clathrin-mediated endocytosis. These results explain in large part discrepancies among published experiments concerning the role of clathrin in synaptic vesicle endocytosis.

Introduction

Clathrin is thought to act at the plasma membrane of synapses to retrieve synaptic vesicle membrane and proteins. There is a very large literature supporting this conclusion. Classic ultrastructural studies of frog neuromuscular junctions revealed the presence of clathrin coats on the plasma membrane after stimulation, suggesting that synaptic vesicles are reconstituted at the surface¹. Biochemical purification of clathrin-coated vesicles from rat brain demonstrated that synaptic vesicle proteins co-purify with clathrin and AP2², and AP2

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

*Corresponding authors Contact: jorgensen@biology.utah.edu, christian.rosenmund@charite.de.

Author contributions

SW, CR, and EMJ conceived and designed experiments. SW and BS performed the freezing experiments. SW performed electron microscopy imaging and analysis. TT designed shRNA constructs. TT and BB generated lentivirus and performed biochemistry. TT and BS performed immunocytochemistry. TT, MC, and BRR performed electrophysiology. AF prepared cell cultures. WD designed the stimulation device. SW, TT, MC, BRR, CR, and EMJ wrote the manuscript. CR and EMJ provided funding, experiments were performed at the Charité, Berlin.

and clathrin are sufficient to generate vesicles from purified brain lipids^{3,4}. Transmembrane vesicle proteins, such as synaptotagmin⁵⁻⁷ and synaptobrevin^{8,9}, interact with adaptor proteins at the plasma membrane and these interactions are required for regenerating functional synaptic vesicles. These studies lend strong support to the idea that clathrin acts at the plasma membrane to regenerate functional synaptic vesicles. However, recent morphological studies^{10,11} suggest that endocytosis may be much faster than previously described for clathrin-mediated endocytosis.

‘Flash-and-freeze’ fixation combines optogenetic stimulation with high-pressure freezing to capture events at synapses milliseconds after stimulation. Our ultrastructural studies demonstrated that endocytic pits, which lacked stereotypical clathrin coats, appeared ~50 ms after stimulation at both *C. elegans*¹¹ and mouse synapses¹⁰. Ultrafast endocytosis is compensatory, that is, membrane retrieval is triggered by the membrane added by fusion¹⁰. Under our stimulation conditions this appears to be due to multiquantal release¹⁰, as has been previously observed at cultured hippocampal synapses¹². The large endocytic vesicles retrieved by ultrafast endocytosis are too big to be functional synaptic vesicles – they are typically equivalent to the surface area of 4 synaptic vesicles, and the fate of these vesicles was unknown.

Here, we examined events that occur up to 20 seconds after stimulation. We performed ‘flash-and-freeze’ experiments on mouse hippocampal neurons and analyzed the synaptic ultrastructure blind from ~200 synaptic profiles per time point. A total of 10,514 synaptic profiles and 361 tomograms were analyzed. We applied a single stimulus in the presence of 4 mM Ca²⁺ or 10 stimuli (20 Hz) in the presence of 2 mM Ca²⁺. At physiological temperatures, clathrin is not required for ultrafast endocytosis but is required to bud synaptic vesicles from synaptic endosomes (Extended Data Fig. 1). However, at room temperature, clathrin functions at the plasma membrane during endocytosis.

Results

Synaptic vesicles are regenerated from endosomes

Clathrin polyhedral coats on endocytic membranes are distinctive in electron micrographs (Fig. 1, Extended Data Fig 2). In our preparations, these coats are observed on pits on the cell body (Extended Data Fig. 2d) but rarely observed on the plasma membrane at synapses. To test the requirement of clathrin after stimulation, a single stimulus was applied to hippocampal cells expressing a variant of channelrhodopsin (ChetaTC)¹³. The experiments were performed in the presence of 4 mM Ca²⁺ in the external solution at 34°C (Fig. 1) and 37°C (Extended Data Fig. 2). After a single stimulus, endocytic pits that lack distinctive clathrin coats formed 50-100 ms after fusion (Fig. 1a, left panel; Extended Data Fig. 2a, left panel)¹⁰. These invaginations resolved into large vesicles about 80 nm in diameter at the lateral edges of the active zone (Fig. 1a; Extended Data Fig. 2a). The number of large endocytic vesicles adjacent to the plasma membrane peaked at 100 ms (Fig. 1d; Extended Data Fig. 2e). Thereafter, endosome-like structures began accumulating in the bouton and peaked at 1 s (Fig. 1b,d; Extended Data Fig. 2b,e). These organelles are larger in diameter (116.4 ± 2.5 nm, 8 synaptic vesicle equivalents) than the endocytic vesicles at the periphery (80.6 ± 0.7 nm; 4 synaptic vesicle equivalents; Extended Data Fig. 3a, *see Methods*),

suggesting that the large endocytic vesicles fuse to form a synaptic endosome¹⁴. Tomographic reconstructions of these endosomes demonstrated that they are not connected to the plasma membrane (Extended Data Fig. 2c and Supplementary Video 1). Clathrin-like coats were visible on some of these endosomes (Fig. 1b,c; Extended Data Fig. 2b,c,d), and budded endosomes peaked 3 s after stimulation (Fig. 1c,d; Extended Data Fig. 2b,c,e). The decline in endosomes was accompanied by an accumulation of clathrin-coated vesicles inside the bouton at 3 s (Fig. 1d; Extended Data Fig. 2e, f). By contrast, clathrin-coated pits on the plasma membrane were only observed in 0.4% of synaptic profiles (4/907) between 1 to 10 s after stimulation (1s 2/332; 3s 2/345; 10 s 0/330) (Fig. 1d; Extended Data Fig. 2e). These results suggest that clathrin does not regenerate synaptic vesicles via endocytosis at the plasma membrane but rather by budding vesicles from an endosome.

Ultrafast endocytosis is intact after clathrin knock-down

To test the role of clathrin in ultrafast endocytosis, we reduced expression of clathrin using RNA interference¹⁵. Cultures were infected with lentivirus expressing a short hairpin RNA (shRNA) targeted against clathrin heavy chain mRNA, or a scrambled shRNA control. Clathrin was significantly reduced seven days after infection as determined by Western blot (~80% reduction, n=3; Extended Data Fig. 4a) and immuno-staining (64% reduction; Extended Data Fig. 4b,d,e). Similarly, transferrin uptake was reduced by 66% (Extended Data Fig. 4c,f). Clathrin knock-down reduced release-ready vesicles but did not affect the exocytic machinery in electrophysiological recordings (Fig. 2a, Extended Data Fig. 5a-f, see Supplementary Information). Likewise, a smaller docked pool of vesicles was observed by electron microscopy, but the overall morphology of synapses was normal (Extended Data Fig. 5g,h, see Supplementary Information).

Clathrin knock-down did not disrupt ultrafast endocytosis, which remained proportional to vesicle fusion. Following a single stimulus (34°C, 4 mM Ca²⁺), endocytic pits and large endocytic vesicles appeared at the edges of the active zone both in the control and clathrin knock-down cells (Fig. 2b, c). The number of endocytic pits and vesicles was decreased by 41% (Fig. 2c) - closely matching the reduction in exocytosis of synaptic vesicles (Extended Data Fig. 5h). These results suggest that membrane retrieval via ultrafast endocytosis does not depend on clathrin, which is consistent with the lack of stereotypic clathrin coats on pits during ultrafast endocytosis (Fig. 1a; Extended Data Fig. 2a).

Clathrin is required to regenerate synaptic vesicles from endosomes

To determine whether clathrin is required to resolve endosomes into synaptic vesicles, control and clathrin knock-down cells were stimulated in the presence of cationized ferritin (4 mM Ca²⁺, 34°C). In control cultures 1 s after a single stimulus (Fig. 3a,b), ferritin molecules were found in endosomes but not in clathrin-coated vesicles. At 3 s endosomes were budding and ferritin began to appear in clathrin-coated vesicles. The size of endosomes was larger than endocytic vesicles, again suggesting that endocytic vesicles carrying ferritin fuse to form synaptic endosomes (Extended Data Fig. 3b,c). Sixteen ferritin-positive endosomes were reconstructed from tomograms, none of them extended a tubule to the plasma membrane (Extended Data Fig. 6a). However, half of those endosomes were not fully contained within the 200 nm tomogram. Therefore, we reconstructed 11 complete

synapses by assembling serial tomograms of synapses 3 s after stimulation. None of the 17 complete end-to-end endosomes were connected to the plasma membrane (6 of which contained ferritin), suggesting that they are true intracellular organelles, not extensions of the plasma membrane (Supplementary Videos 1-5). The decline in the number of endosomes is followed by an accumulation of ferritin-positive coated vesicles and synaptic vesicles (Fig. 3b). Some clathrin-coated vesicles were observed before 3 s after stimulation (1 s, 4/149 synaptic profiles and Fig. 1d), but they did not contain ferritin molecules, suggesting that they were derived from pre-existing endosomes. The total number of ferritin granules per synaptic profile does not increase suggesting that there is not an additional wave of endocytic events that occurs during these 20 s (Extended Data Fig. 6c). Some of the ferritin-filled small vesicles docked to the active zone (Fig. 3a, right panel; Extended Data Fig. 6b,d), suggesting that these vesicles are synaptic vesicles. These results indicate that recently endocytosed membrane passes through endosomes to regenerate synaptic vesicles within about 5-6 s after fusion.

In clathrin knock-down cultures, ferritin-filled endosomes increased after 1 s (Fig. 3d). However, these endosomes did not decline across the 3 s, 10 s or 20 s (Fig. 3c,d). These endosomes formed almost no coated buds and remained spherical (Fig. 3c), suggesting that membrane curvature of the buds requires clathrin. Together, these results suggest that clathrin is required at endosomes to regenerate synaptic vesicles following a single stimulus.

Clathrin acts on endosomes after high-frequency stimulation

At mammalian central synapses, action potentials often fire in bursts^{16,17} and could exhaust ultrafast endocytosis. To assay endocytosis after high-frequency stimulation, we delivered 10 stimuli (20 Hz for 500 ms in 2 mM Ca^{2+} , 34°C) and froze cultures 1 s, 3 s, 10 s, and 20 s after the end of the stimulus. Under these conditions, 90% of the cells fired action potentials for all 10 light pulses (Extended Data Fig. 7a). Using cationized ferritin, we followed the fate of the newly endocytosed vesicles (Fig. 4 and Extended Data Fig. 3 d,e). The results were similar to those after a single stimulus. Coated pits were not observed on the plasma membrane. Ferritin accumulation in endosomes peaked at 1 s; endosomes exhibited coated buds at 3 s, accompanied by the appearance of clathrin-coated vesicles and some uncoated synaptic vesicles. All vesicles were uncoated by 10 s (Fig. 4a,b; Extended Data Fig. 7b). Ferritin-filled vesicles were eventually recruited to the active zone, suggesting that these vesicles are likely synaptic vesicles (Extended Data Fig. 7c).

We then applied an identical 20 Hz stimulation to cultures treated with clathrin shRNA. In these cultures ferritin was taken up into large endosomes, again suggesting that membrane internalization after a high frequency burst does not require clathrin (Fig. 4c). However, ferritin-containing endosomes were abundant and were not resolved into vesicles even after 20 s (Fig. 4c,d), indicating that clathrin is essential for regenerating vesicles from endosomes after a burst of action potentials. The acute inhibition of clathrin function by Pitstop 2 (30 μM , 2 min) also blocked regeneration of synaptic vesicles from endosomes (Extended Data Fig. 8; see Supplementary Information). These results suggest that following a short burst of action potentials, clathrin acts on endosomes to regenerate synaptic vesicles.

To determine whether intense stimulation triggers clathrin-mediated endocytosis, we applied trains of 3, 10, 30, and 100 light pulses at 20 Hz to hippocampal neurons and froze them 10 s after stimulation (2 mM Ca^{2+} , 34°C). To arrest all endocytic intermediates at the plasma membrane, we applied 80 μM dynasore for 30 s before the stimulation. Dynasore is a drug that binds dynamin¹⁸; although it also has off-target effects¹⁹, it nevertheless effectively locks both clathrin-mediated endocytosis and ultrafast endocytosis at a late stage of vesicle scission^{10,18}. After dynasore treatment, no clathrin-coated pits were trapped on the plasma membrane under any condition. Instead, strings of uncoated vesicles attached to the plasma membrane accumulated at the edges of active zones (Fig. 4e,f). The diameter of vesicles in strings (53.3 ± 3.7 nm) was larger than synaptic vesicles (~40 nm) but smaller than large endocytic vesicles formed after a single stimulus (~80 nm), reflecting reduced fusion during repetitive stimulation. These results suggest ultrafast endocytosis is responsible for removing membrane from the surface even during high frequency bursts of action potentials and that clathrin functions on endosomes rather than at the plasma membrane to regenerate synaptic vesicles.

Clathrin acts at the plasma membrane when ultrafast endocytosis fails

Actin polymerization is required for ultrafast endocytosis¹⁰. To study synaptic vesicle recycling in the absence of ultrafast endocytosis, we blocked actin polymerization. We treated cultured neurons with 10 μM latrunculin-A for 30 s, stimulated once by a 10 ms light flash, and froze 3 or 10 s later (4 mM Ca^{2+} , 34°C). Large endocytic vesicles adjacent to the active zone were absent following the latrunculin-A application, indicating that ultrafast endocytosis was blocked (Fig. 5a-d). Under these conditions, clathrin-coated pits were observed on the plasma membrane 3 s after stimulation, followed by an accumulation of coated vesicles between 3 and 10 s, and the appearance of large vesicles at 10 s (Fig. 5c,d). The size of coated pits, coated vesicles and synaptic vesicles were similar (Extended Data Fig. 9a).

Actin polymerization is known to be less efficient at room temperature in mammalian cells^{20,21}. To test whether ultrafast endocytosis fails at low temperature, we incubated cells at room temperature (22°C) for 5 min prior to flash-and-freeze (Fig. 5e-h). In control cells maintained at 34°C, coated pits were not observed after stimulation (Fig. 5e-h, unstimulated 0/251 synaptic profiles, 3 s 0/248 synaptic profiles). However, at 22°C, clathrin-coated pits appeared on the plasma membrane adjacent to the active zone 3 s after a single stimulus (Fig. 5g, h). Like in the latrunculin-A treated cells, large vesicles, possibly endosomes, accumulated in the center of the bouton 10 s after stimulation^{22,23}, as observed previously in hippocampal cultures stimulated at room temperature^{22,23}. The diameter of coated pits and synaptic vesicles were similar (Extended Data Fig. 9b). These results suggest that failure of ultrafast endocytosis leads to clathrin-mediated endocytosis on the plasma membrane.

Discussion

In classic electron microscopy studies of stimulated synapses, clathrin-mediated endocytosis occurs about 20 s after stimulation^{24,25}. By contrast, clathrin-independent ultrafast endocytosis removes excess membrane 30-100 ms after stimulation^{10,11}, and as

demonstrated here, synaptic vesicles are regenerated via an endosome 5-6 s after stimulation (Extended Data Fig. 1). These contradictory results can be reconciled simply by considering temperature.

Temperature

Ultrafast endocytosis is only observed at physiological temperatures. When neurons are cooled to 22°, ultrafast endocytosis fails, and the slower clathrin-mediated process takes over. Actin polymerization is greatly reduced in cultured cells at non-physiological temperature^{26,27}. Because ultrafast endocytosis requires actin polymerization, it is likely that vesicle components accumulate on the plasma membrane at room temperature, and the clathrin machinery is recruited to the plasma membrane instead of the synaptic endosome. The shift to clathrin-mediated endocytosis may contribute to rapid synaptic depression that is observed at room temperature^{28,29}, perhaps because excess membrane in the active zone could block exocytosis.

Previous ultrastructural data have suggested the presence of a fast form of endocytosis at physiological temperature. When frog pectoral muscles were stimulated at 10°C in the original Heuser and Reese experiments, clathrin-coated pits accumulated on the plasma membrane¹; whereas when stimulated at room temperature, two forms of endocytosis were observed: a fast form 1 s after stimulation, and a slow form 20 s after stimulation²⁴. At snake neuromuscular junctions clathrin-coated pits accumulated when stimulated at 7°C³⁰, whereas at room temperature, membrane was retrieved in 1-2 s³¹. In rat hippocampal neurons endosomes (> 70 nm) were observed after stimulation at physiological temperature but not at room temperature³². Capacitance studies also indicate fast and slow mechanisms. In the Calyx of Held, fast endocytosis was observed at physiological temperature (35°C-37°C), but was abolished at room temperature³³. Fast endocytosis was also observed in fish retinal bipolar cells, membrane was recovered ~1s after a single stimulation^{34,35}.

Ultrastructure and capacitance analyses only measure membrane endocytosis; protein endocytosis can be measured by pHluorin assays. At room temperature, a slow form of endocytosis is observed in hippocampal neurons with a surface dwell time of 15 s³⁶. When pHluorin experiments are conducted at physiological temperatures, the time constant for endocytosis is significantly faster: 14 s at 24° and 10 s at 30°C and as fast as 6 s at 37°C^{37,38}. At first glance, a 6 s recovery for vesicle proteins agrees well with a 6 s recovery of morphologically defined synaptic vesicles. However, such a comparison is deceptive, since it is unlikely that proteins in a fusing vesicle could diffuse to the lateral edge of the active zone and into an endocytic vesicle within 50 ms. Nevertheless, temperature seems to be responsible for differences observed among ultrastructural studies.

Clathrin at endosomes

Endosomes are sorting compartments that receive cargo from transport vesicles generated by endocytosis from the plasma membrane. Heuser and Reese proposed in their original model that clathrin-mediated endocytosis generated a transport vesicle that fused to an endosome ('cisterna'), and synaptic vesicles were then regenerated from these endosomes¹. However, subsequent experiments demonstrated that these cisternae were likely to arise via bulk

endocytosis due to non-physiological stimulation²⁵, and the model was amended to propose that synaptic vesicles were generated directly from the plasma membrane via clathrin-mediated endocytosis, whereas bulk endocytosis removed excess membrane added during intense stimulation. Thus, the cisternae do not represent true endosomes since they do not arise via transport vesicles but rather from a simple pinching off from the plasma membrane. Moreover, it was observed that internal clathrin-coated buds were often, or even always, attached to the plasma membrane via long tubules^{1,4,39}, which led to the idea that the formation of clathrin-coated pits is limited to the plasma membrane. The different lipid compositions of endosomes and the plasma membrane support this notion. Endosomes are enriched for PI-(3)P, whereas the plasma membrane is enriched in PI-(4,5)P2 and therefore can recruit clathrin-AP2⁴⁰.

By contrast, our data suggest that clathrin regenerates synaptic vesicles from a synaptic endosome rather than from the surface. Tomographic reconstructions of endosomes failed to identify connecting tubules to the plasma membrane. Moreover, these compartments appear to be true endosomes, in that they are generated by transport vesicles. Because endosomes are twice as large as endocytic vesicles, the synaptic endosome probably forms by fusion of endocytic vesicles. The synaptic endosome is resolved into synaptic vesicles by clathrin-mediated budding, and for this reason it is unlikely that synaptic endosomes are typical endosomes. The presence of clathrin on these endosomes suggests they are enriched in PI-(4,5)P2. In this case, the membrane composition of “synaptic endosomes” is more like the plasma membrane than that of classical endosomes.

A growing body of molecular studies support the conclusion that synaptic vesicles may be regenerated from endosomes. Live imaging in hippocampal neurons indicates that recently endocytosed vesicle proteins are sorted in an endosome after stimulation²². Endosomal rab GTPases, such as Rab5 and Rab35, are abundant at synapse and can be co-purified with synaptic vesicles⁴¹, and inhibition of Rab5⁴² and skywalker⁴³ (required for Rab35 function) results in an accumulation of endosomes in synaptic boutons. Finally, accumulation of endosomes is also observed in the absence of AP2, clathrin, clathrin accessory proteins, or dynamin⁴⁴⁻⁴⁸.

In summary, our results resolve an important contradiction: most studies indicate that clathrin is essential for endocytosis at hippocampal synapses³⁶, whereas ultrafast endocytosis is clathrin independent. Here we demonstrate that even the ultrafast pathway requires clathrin to regenerate synaptic vesicles, not at the plasma membrane but rather at the synaptic endosome. It is likely that at some synapses and under some physiological conditions clathrin acts at the plasma membrane to regenerate synaptic vesicles. Whether ultrafast endocytosis is a general mechanism or a specialization for synapses with high turnover will require further study.

Methods online

Cell culture and lentivirus infection

Animals were handled according to the rules of Berlin authorities and the animal welfare committee of the Charité Berlin, Germany. Hippocampi were dissected from newborn C57/

BL6-N mice and cultured at 13×10^3 cells /cm² on 6 mm sapphire disks for ‘flash-and-freeze’ electron microscopy experiments and 25 mm coverslips for all other experiments as previously described¹⁰. Astrocytes were seeded a week in advance to generate a feeder layer. For autaptic cultures, microislands of astrocytes were prepared as previously described²⁸ and neurons were plated at 300 cells/cm². For biochemistry, 10×10^3 cells/cm² were cultured without the feeder layer.

Lentivirus was produced as described previously⁵⁰. The cells were infected with lentivirus expressing ChetaTC¹⁰ on DIV1 and clathrin or non-specific shRNA on DIV7. Infection of clathrin shRNA at DIV1 caused severe loss of neurons at DIV14, and thus, cells were only infected for 1 week with clathrin shRNA for all the experiments.

shRNA constructs

A mouse CHC-1 specific siRNA target sequence (5′ -GTT GGT GAC CGT TGT TAT G-3′) was obtained using Genscript siRNA Target Finder (<https://www.genscript.com/ssl-bin/app/rnai>) and cloned as shRNA into a lentiviral shuttle vector under the control of a U6 promoter. For the scrambled shRNA control sequence we adapted the scrambled siRNA sequence from Royel *et al.* (2005)¹⁵ and cloned it as shRNA (5′-TTC GCA CCC TAC TTC GTG G-3′) into a lentiviral shuttle vector. Both sequences were subject to a BLAST search to ensure that mouse CHC-1 shRNA was specific and that the scrambled shRNA did not match any sequence. To identify infected cells, the shuttle vector contained a human Synapsin-1 promoter, driving an expression of a nuclear-targeted red fluorescent protein (NLS-RFP).

Western blots and immunocytochemical staining

For detection of clathrin levels by western blots, protein lysates were obtained from astrocyte-free mass cultures of hippocampal neurons. Briefly, cells were lysed using 50 mM Tris/HCl (pH 7.9), 150 mM NaCl, 5 mM EDTA, 1% Triton-X-100, 1% Nonidet P-40, 1% sodium deoxycholate, and protease inhibitors (cOMplete protease inhibitor cocktail tablet, Roche Diagnostics GmbH, Mannheim, Germany). Protein concentration was determined by BCA assay. Proteins were separated by SDS-PAGE and transferred to nitrocellulose membranes. Membranes were then incubated with rabbit anti-CHC-1 (1:500, Abcam, ab21679) and mouse anti-tubulin III (1:2000, Sigma-Aldrich, T8660) antibodies overnight at 4°C. After incubation with corresponding horseradish peroxidase-conjugated goat secondary antibodies (1h at room temperature) and ECL Plus Western Blotting Detection Reagents (GE Healthcare Biosciences), chemiluminescent was imaged using a Vilber Lourmat Fusion FX7 detection system. Ratiometric quantification of signal intensities was measured with the supplied BIO-ID software package of the Fusion FX system. The signals are linear with the amount of protein lysate loaded (Extended Data Fig. 4a), suggesting that we can detect the clathrin signals in the knock-down without saturating the signals from the control.

Immunocytochemical staining was performed on autaptic cultures. One week after infection with the shRNA constructs, cells were washed once in PBS and fixed with 4% paraformaldehyde (PFA) for 10 min. After washing in PBS, cells were permeabilized with 0.1% PBS-Tween20 for 10 min and blocked in 5% NGS for 30 min. Subsequently cells

were incubated with rabbit anti-CHC-1 (1:1000, abcam, ab21679) and guinea pig anti-synaptophysin (1:1000, Synaptic System) antibodies overnight at 4 °C. Primary antibodies were labeled with anti-rabbit Alexa Fluor 647 and anti-guinea pig Alexa Fluor 488 (each 1:500, Jackson Immunoresearch Laboratories) for 1h at room temperature. After washing, cover slips were mounted with Mowiol 4-88 anti-fade medium (Polysciences Europe GmbH, Eppelheim, Germany). Transduced cells, visible by nuclear RFP expression, were imaged using an Olympus IX81 microscope equipped with a Princeton camera and controlled by Metamorph software, with same acquisition settings for all groups. Ratiometric quantification of clathrin signals over synaptophysin signals was conducted in the automated fashion using ImageJ (National Institute of Health, Bethesda, MD) with custom-written macros. In short, the program first subtracts uneven illumination using a “rolling ball” function with the radius set at 30 pixels (http://fiji.sc/Rolling_Ball_Background_Subtraction). Then, the synapses are defined by thresholding the synaptophysin signals at 15%. The binary image of synaptophysin was used as the regions of interest. The average intensities of fluorescence signals from clathrin heavy chains and synaptophysin were measured from those locations and were divided to obtain ratio between those two proteins. A total of 28 images from the control shRNA cells and 30 images from the clathrin knock-down cells (N = 2 cultures).

Electrophysiology

Whole cell patch-clamp and cell-attached voltage-clamp recordings were performed as previously described¹⁰. The extracellular solution contained 140 mM NaCl, 2.4 mM KCl, 10 mM Hepes, 10 mM Glucose (pH adjusted to 7.3 with NaOH, 300 mOsm). For a single light stimulus experiment, 4 mM CaCl₂, and 1 mM MgCl₂ were added to the solution while 2 mM CaCl₂ and 1 mM MgCl₂ were added for high-frequency stimulation experiments. All recordings were performed at 34°C.

For recordings in autapses, neurons at DIV15-17 were clamped at -70 mV with a Multiclamp 700B amplifier (Molecular Devices) under control of Clampex 9 (Molecular Devices). The extracellular solution contained 2 mM CaCl₂ and 4 mM MgCl₂. The patch pipette solution contained 136 mM KCl, 17.8 mM Hepes, 1 mM EGTA, 0.6 mM MgCl₂, 4 mM ATP-Mg, 0.3 mM GTP-Na, 12 mM phosphocreatine and 50 units/ml phosphocreatine kinase (300 mOsm, pH 7.4). EPSCs were evoked by a brief 2 ms somatic depolarization to 0 mV. RRP size was determined by measuring the charge transfer of the transient synaptic current induced by a pulsed 5 s application of hypertonic solution (500 mM sucrose in extracellular solution). Vesicular release probability was calculated as the ratio of the charge from the evoked EPSC (integrated for 1s) and the readily-releasable pool size.

Electron Microscopy

‘Flash-and-freeze’ experiments were performed as previously described¹⁰. Briefly, sapphire disks with cultured cells were mounted in the freezing chamber of the high-pressure freezer (HPM100, Leica), which was set at 34°C. 34°C was chosen to match the temperature of electrophysiological recordings. To minimize the exposure to room temperature, a petri dish containing the sapphire disk was placed on a 37°C cryopack while mounting, and the transparent polycarbonate sample cartridges were stored between cryopacks equilibrated to

37°C. Immediately after the sapphire disk was mounted on the sample holder, recording solution kept at 37°C was applied to the specimen and the cartridge was inserted into the freezing chamber and allowed to equilibrate to 34°C for 30 seconds. For experiments at 37°C (Extended Data Fig. 2), a temperature-controlled chamber (Leica) was placed around the specimen loading table of the high-pressure freezer. After transferring in a petri dish, the cells were allowed to recover in the 37°C chamber for 2 min and then mounted onto the specimen carriers for freezing. The freezing chamber was set at 37°C in these experiments. Using a home-built light stimulation controller, we applied either 1 or 10 light pulses (20 Hz) to the specimens. The controls for each experiment were always frozen on the same day from the same animal. Each light pulse was for 10 ms. Under these conditions we observe multivesicular release; we estimate that up to 4 synaptic vesicles are released per active zone (data not shown and¹⁰). We set the device so that the samples were frozen at 15, 30, 50, 100, 300, 1000, 3000, 10,000, or 20,000 ms after the initiation of the first stimulus¹⁰.

For ferritin-loading experiments, cationized ferritin (Sigma-Aldrich) was added in the recording solution at 0.25mg/ml. The calcium concentration was reduced to 1 mM to suppress spontaneous activity during the loading. The cells were incubated in the solution for 5 min at 37°C. For Pitstop 2 experiments, 30 μ M Pitstop 2 was additionally included in the solution, and the cells were only incubated for 2 min. After 2 min incubation with Pitstop 2, some cells were dead while performing patch-clamp experiments. After ferritin incubation, the cells were immersed in the recording solution containing either 4 mM Ca^{2+} for a single stimulus experiment or 2 mM Ca^{2+} for high-frequency stimulation. The change in calcium concentrations from 1 mM to 4 mM increases the rate of miniature EPSCs and thus may contribute to the background ferritin loading before the experiments. In our experiments, about 1% of synaptic profiles contained endosomes and synaptic vesicles that were ferritin-positive without stimulation. The absence of ferritin-positive coated pits on the plasma membrane in the unstimulated controls suggests that ferritin passes through endosomes even during spontaneous activity. 3 μ M NBQX and 30 μ M bicuculline were also included in the recording media to minimize the recurrent network activity¹⁰. Ferritin-positive synaptic vesicles were only found in 5-7% synaptic profiles scored before stimulation using this protocol.

Following high-pressure freezing, samples were transferred into a vial containing 1 % osmium tetroxide (EMS), 1% glutaraldehyde (EMS), 1% milliQ water, in anhydrous acetone (EMS). The freeze-substitution was performed in AFS2 (Leica) with the following program: -90°C for 5-7 hours, 5°C/hour to -20°C, 12 hours at -20°C, and 10°C/hour to 20°C. Following *en bloc* staining with 0.1% uranyl acetate, the samples were infiltrated and embedded into epon and cured for 48 hours in a 60°C oven. Serial 40-nm sections were cut using a microtome (Leica UCT) and collected onto formvar-coated single-slot grids. Sections were stained with 2.5% uranyl acetate prior to imaging. For ferritin experiments, sections were not stained after sectioning to improve contrast of ferritin molecules in our images – this, in turn, might have compromised our ability to distinguish clathrin-coated vesicles. Approximately 150 synaptic profiles were collected from a single section from each specimen, and the experiments were repeated with second cultures in each case (for detailed n values, see Supplementary Table 1). The sample size was chosen based on the

previous experiments that allowed us to acquire a sufficiently large set of data for statistical analysis¹⁰. The total numbers of synaptic profiles and tomograms analyzed for these experiments are 10,514 and 361, respectively. Note that there is some variability from experiment to experiment; this is likely due to the short lifetimes of these structures. The synaptic profiles were chosen randomly to sample unbiased populations. Active zones were defined as regions juxtaposed to a post-synaptic density. Docked vesicles are defined as those directly in contact with membrane. The large endocytic vesicles are defined as vesicles larger than ~50 nm by visual inspection and within 50 nm of active zone, measured in ImageJ. Endosomes are defined as membrane-bound organelles that are in the center of the bouton and larger than 100 nm by visual inspection. Vesicular compartments with coated buds were also categorized as endosomes, which appeared frequently in our 3 s time points. Vesicles are scored as clathrin-coated only if distinctive coats were visible which can lead to underscoring (Fig. 1c, and Extended Data Fig. 2c). Furthermore, the chance of capturing a coated vesicle in a given synaptic profile is low compared to an endosome due to its size. These factors will lead to an underestimate of the number of coated vesicles observed in profiles. The morphometry was performed blind using custom-written ImageJ macro and Matlab scripts (Watanabe, Davis, and Jorgensen).

For electron tomography, 100-200 nm thick sections were collected on pioloform-coated single-slot grids. Sections were post-stained with uranyl acetate as described for thin sections. 10-nm gold particles were sparsely applied to both sides of the grids by incubating the grids in drops of solution containing 5.7×10^{12} particles/ml (<http://microspheres-nanospheres.com/>) for 2 min. 17-50 tilt-series ($\pm 65^\circ$) were collected from each sample using FEI TF20 electron microscope. For serial tomograms, low magnification images acquired from each section on a grid were used to locate the same synapses in the serial sections. Typically, the entire synapses span 4-5 200-nm sections. The tomograms were generated from tilt-series using IMOD. The tomograms were segmented and reconstructed using Amira or TrakEM2.

Membrane calculations

The amount of membrane on large endocytic vesicles and endosomes was calculated to determine the number of synaptic vesicle equivalents contained in these structures. These structures were scored blind and were distinguished by position in the bouton, by size, and by shape. For large endocytic vesicles in single stimulus experiments (Fig. 1), the mean diameter was 80.6 ± 1.5 nm. The surface area of the large endocytic vesicles is therefore about 20,000 nm², since the surface area of a synaptic vesicle is 5,300 nm², large endocytic vesicles correspond to 4 synaptic vesicle equivalents. In the RNA interference experiments after a single stimulus, the size of large endocytic vesicles was 82.6 ± 4.5 nm (4 SVs) in the control and 74.4 ± 2.0 nm (3 SVs) in the clathrin knock-down. In the RNA interference experiments after high-frequency stimulation, the size of large endocytic vesicles was 85.5 ± 2.2 nm (4 SV) in the control and 70.1 ± 1.4 nm (3 SV) in the knock-down.


The surface area of endosomes was calculated by measuring the diameter and assuming a spherical shape. This calculation will lead to an underestimate of the surface area of endosomes because the membrane in budding endosomes is convoluted. To minimize this

error, endosomes with coated buds were not included in the calculations. After a single stimulus (Fig. 1), the mean diameter of endosomes was 116.4 ± 3.0 nm (8 SV). In the ferritin experiments with a single stimulus, the diameter of endosomes was 108.7 ± 4.2 nm (7 SV) in the control and 93.1 ± 2.8 nm (5 SV) in the clathrin knock-down. In the ferritin experiments with multiple stimuli, the diameter of endosomes was 111.8 ± 4.1 nm (8 SV) in the control and 100.9 ± 2.6 nm (6.5 SV) in the clathrin knock-down. The smaller size of endocytic structures in the clathrin knock-downs is likely due to the reduced neurotransmission in these cells. To confirm the estimates of surface area, we measured the surface area of complete endosomes in tomograms of wild-type synapses. Most endosomes were spherical with the diameter of 111.2 ± 5.3 nm (39,000 nm² or 7.3 SV equivalents). The surface area of budded endosomes was measured to be $\sim 41,000$ nm² or 7.7 synaptic vesicle equivalents.


Statistics

For detailed numbers and statistical analysis, *see* Supplementary Table 1.

Extended Data

**Extended Data Fig. 1.**


Ultrafast endocytosis regenerates synaptic vesicles in a two-step process. Ultrafast endocytosis removes membrane added by vesicle fusion at the lateral edge of the active zone. Large endocytic vesicles then fuse to endosomes. Endosomes are resolved into synaptic vesicles via a clathrin-dependent process. Newly formed synaptic vesicles can be recruited back to the active zone.



Extended Data Fig. 2.

Synaptic vesicles are regenerated from endosomes at 37°C. Hippocampal synapses were stimulated once and frozen at the indicated times. The experiments were performed at 37°C in the presence of 4 mM external Ca^{2+} . (a) Electron micrographs showing invaginations and large endocytic vesicles (arrowheads) recovered via ultrafast endocytosis. (b) Micrographs showing single coated buds (left, middle) or multiple buds (right) forming on an endosome. (c) Virtual section from an electron tomogram (left) and reconstruction (middle) showing a synaptic endosome with buds following a single stimulus. The average intensity of coated


buds from the top 20 nm (right, top) and the bottom 40 nm (right, bottom) is shown. Clathrin-cages can be preserved in our fixation and are visible in the tomogram. We found a total of 32 endosomes in these reconstructions (14 endosomes in the unstimulated control and 28 endosomes 3 seconds after stimulation). Of the total 32 endosomes, none were connected to the plasma membrane or showed evidence of a truncated tubule extending from the endosomal membrane. Of the 14 unstimulated endosomes, 8 were contained within single tomograms, and are therefore unambiguously closed on both ends. Of the 28 endosomes in stimulated synapses, 16 endosomes were contained within single tomograms so that it was clear that no attachment to the plasma membrane was possible. (d) Examples of a coated pit on the plasma membrane (top) and a coated bud on an endosome (bottom). Note that the morphology of the coats is similar. (e) Increase in the number of each endocytic structure per synaptic profile after a single stimulus at 37°C. The prevalence of large endocytic vesicles and endosomes is followed by an increase in the number of clathrin-coated vesicles. Coated pits were not observed on the plasma membrane (PM). (f) Frequency of profiles or tomograms that contain endosomal structures at 37°C. Roughly, 60% of unstimulated synapses contained one endosome. One second after stimulation, 60% of the synapses contained at least one endosome, and half of those synapses contained two endosomes. Three seconds after stimulation, ~30% of the synapses contained budded endosomes and clathrin-coated vesicles, suggesting that those synapses were active. The standard error of the mean is shown in each graph. For N values, detailed numbers and statistical analysis, *see* Supplementary Table 1.



Extended Data Fig. 3.


Large endocytic vesicles likely fuse to become synaptic endosomes. (a-e) Histograms (left) and cumulative plots (right), showing the size of large endocytic vesicles (red) and endosomes (orange) after one stimulus from control cells without ferritin (a), scrambled shRNA infected cells (b), and clathrin knock-down cells (c) both with ferritin. 10 stimuli were applied to scrambled shRNA (d) or clathrin knock-down cells (e). The large endocytic vesicles are defined as those that are within 50 nm of active zone and larger than a synaptic vesicle by visual inspection. Endosomes are defined as large vesicles greater than 50 nm

from the active zone (often in the center of the bouton) and are larger than ~100 nm. Any vesicular compartment that has coated buds in the center of the bouton is also categorized as an endosome. The numbers of endocytic structures in (a) represent all the structures scored from 100 ms and 300 ms time points. The number of large vesicles and endosomes in (b-e) represent all the ferritin-positive structures from later time points (3, 10, and 20 s). In the control shRNA (b), the number of large endocytic vesicles and endosomes has declined by these late time points, whereas the ferritin is trapped in large endocytic vesicles and synaptic endosomes in the clathrin knock-down experiments. Because ferritin passes from large endocytic vesicles to even larger budded endosomes, it is likely that the endocytic vesicles fuse with either each other or an existing compartment.

**Extended Data Fig. 4.**


Clathrin shRNA reduces clathrin expression. (a) Left, western blot showing clathrin levels after one-week expression of a scrambled shRNA control or clathrin heavy chain shRNA in cultured hippocampal neurons. Right, an 80% reduction was observed ($n=3$, $p < 0.001$, paired T-test). (b) Normalized ratio of clathrin heavy chain and synaptophysin fluorescence in control and clathrin knock-down (chc KD) cultures. The clathrin/synaptophysin ratio is reduced to 64% in the knock-down cells. (c) The mean fluorescence intensity (normalized to 30 min) representing the amount of transferrin uptake in control and knock-down cells.

Transferrin uptake was reduced by 66% in the knock-down cells. (d, e) Fluorescence images of immuno-cytochemical staining of hippocampal autaptic cultures using anti-synaptophysin (left), anti-clathrin heavy chain (middle), and merge in control (d) and clathrin knock-down cultures (e). (f) Example micrographs of hippocampal mass cultures showing transferrin uptake. The standard error of the mean is shown. *** indicates p-value of <0.001. For detailed numbers and statistical analysis, see Supplementary Table 1.



Extended Data Fig. 5.


Exocytic machinery is intact in the clathrin knock-down cells. Sample traces from cell-attached voltage clamp during light stimulation in control (a) and clathrin knock-down (b). Number of action potentials triggered during the 10 ms light pulse is shown to the right. (c) Readily-releasable pool (RRP) in control and clathrin knock-down cells, defined by brief application of 500 mM sucrose to autaptic neurons (control: 622 ± 56 pC, knockdown: 443 ± 52 pC, $p < 0.01$). (d) Vesicular release probability (Pvr) in these cells (control $5.4 \pm 0.4\%$, $n = 54$; knock-down $3.5 \pm 0.4\%$, $n = 48$; $p < 0.001$). (e, f) Average miniature EPSC (mEPSC) frequency (e) and amplitude (f). No change was observed in knockdown cells. (g) Average number of synaptic vesicles per synaptic profile ($n = 142$ synapses for control and 137 for knockdown). (h) Average number of docked vesicles in active zones per synaptic profile (control: no stimulation, 1.5 ± 0.1 , $n = 142$ synapses; 100 ms, 0.8 ± 0.1 , $n = 142$ synapses; knockdown: no stimulation, 1.2 ± 0.1 , $n = 137$ synapses; 100 ms after stimulation, 1.0 ± 0.1 , $n = 149$ synapses). The fraction of docked vesicles that fuse is greatly reduced by clathrin knockdown. P-values are calculated against the unstimulated control shRNA cells. The standard error of the mean is shown in each graph. ***, **, and * indicate p-value of < 0.001 , < 0.01 , < 0.05 , respectively. n.s., 'not significant'.



Extended Data Fig. 6.

Following a single stimulus, clathrin is required at endosomes to regenerate synaptic vesicles. (a) Virtual section from an electron tomogram (left) and a reconstruction (right) showing a budded synaptic endosome containing ferritin particles in the scrambled shRNA control cell. We found a total of 33 endosomes in these reconstructions. Of these 33 endosomes, none were connected to the plasma membrane or showed evidence of a tubule extending from the endosomal membrane. 17 of these 33 total endosomes were fully contained within the 200 nm tomogram. Of these 33 endosomes, 16 were ferritin-positive,


and 8 of these 16 endosomes were fully contained in the tomogram. (b) Micrographs showing ferritin-positive synaptic vesicles docked to active zone 10-20 s after stimulation. (c) Average number of ferritin particles in large endocytic vesicles, clathrin-coated vesicles, and synaptic vesicles per synaptic profile examined. At least 134 synapses were analyzed per time point. Ferritin progresses to synaptic vesicles in the control, but is trapped in large endocytic vesicles or endosomes in the clathrin knock-down. The fraction of synaptic profiles with ferritin was 27% for the control and 31% in the knockdown, suggesting that 70% of the synapses were silent. The mean number of ferritin particles found in an individual endosome, clathrin-coated vesicle, and synaptic vesicle, are 9.3 ± 1.0 , 2.0 ± 0.2 , and 1.9 ± 0.2 , respectively. The total number of ferritin particles (indicated above), declined by 40% in the control relative to the 1 s time point but not in the knockdown, either due to the fusion of the newly formed synaptic vesicles or by return of excess membrane to the surface of the synapse. The standard error of the mean is shown. (d) Distribution of ferritin-positive clathrin-coated vesicles (yellow) and synaptic vesicles (blue) relative to the active zone at defined time points after stimulation in the control cells. Numbers are binned by 50 nm. The first bin '0 nm' means vesicles are docked in active zone. Endosomes are found at 285 ± 38 nm from the active zone. Note that the data in this figure represent further analysis of the data shown in Fig. 3.



Extended Data Fig. 7.

Following high-frequency stimulation, clathrin is required at endosomes to regenerate synaptic vesicles. (a) Average number of action potentials in 10 ms bins relative to light pulses during high-frequency stimulation (10 stimuli at 20Hz, 0.5 s). Sample traces are shown above. Each light pulse triggered at least one action potential in both control and clathrin shRNA-treated cultures. (b) Average number of ferritin molecules in large endocytic vesicles, clathrin-coated vesicles, and synaptic vesicles per profile examined. Ferritin is transferred from large endocytic vesicles to synaptic vesicles in the control but is


trapped in large endocytic vesicles or endosomes in the clathrin knock-down. The number of profiles with ferritin particles after stimulation was 34% in the control and 36% in the clathrin knock-down, suggesting that 65% of the synapses were silent. On average, the number of ferritin molecules found in an individual endosome, clathrin-coated vesicle, and synaptic vesicle, are 9.2 ± 1.0 , 1.9 ± 0.3 , and 2.0 ± 0.2 . At least 142 synapses were analyzed per time point. The total number of ferritin particles (indicated above each time point), declined by 36% in the control, and by 23% in the knockdown relative to the 1 s time point. (c) Distribution of ferritin-positive clathrin-coated vesicles (yellow) and synaptic vesicles (blue) relative to the active zone at defined time points after stimulation in the control shRNA cells. Numbers are binned by 50 nm. The first bin '0 nm' means vesicles are docked at the active zone. Endosomes are found at 286 ± 43 nm from the active zone. The standard error of the mean is shown in each graph. Note that this figure is a supplemental data figure for Fig. 4 and represents the further analysis of the data from Fig. 4.



Extended Data Fig. 8.

Pitstop 2 blocks regeneration of synaptic vesicles from endosomes after high-frequency stimulation. Pitstop 2 is an inhibitor of clathrin terminal domain and blocks clathrin-

mediated endocytosis⁴⁹. (a, c) Electron micrographs showing ferritin containing vesicles in DMSO-treated (a) and Pitstop 2-treated cells (c) at different time points after stimulation. Ferritin is found in large vesicles after stimulation (middle) and in synaptic vesicles (right) in control, but it is trapped in endosomes in the Pitstop 2-treated cells. (b,d) Average increase in ferritin-positive structures per synaptic profile in DMSO-treated (b) or Pitstop 2-treated cells (d). Ferritin progressed to synaptic vesicle-like structures in the control but remained in endosomes or large endocytic vesicles in Pitstop 2-treated cells. Clathrin-coated pits on the plasma membrane were not present at any time point and were not plotted. At least 140 synapses were analyzed per time point. (e) Virtual section from an electron tomogram (left) and a reconstruction (right) showing a synaptic endosome with buds following 10 stimuli at 20 Hz. Of 32 tomograms reconstructed from 3 s time point, 25 of them showed at least one endosome in the terminal, and 7 showed budded endosomes. None of these endosomes were connected to the plasma membrane. The standard error of the mean is shown in each graph. For detailed numbers and statistical analysis, *see* Supplementary Table 1.



Extended Data Fig. 9.

Synaptic vesicles are regenerated directly from the plasma membrane in the absence of ultrafast endocytosis. Average diameter of clathrin-coated pits, clathrin-coated vesicles, and synaptic vesicles in the latrunculin-A-treated cells (a) or cells incubated at 22°C for 5 min (b). The diameter of these structures is similar suggesting a precursor-product relationship. Diameter of coated pits was determined by the full-width at the half maximum depth of the pit. For detailed numbers and statistical analysis, *see* Supplementary Table 1

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We would like to thank Dorothea Lorenz and Agnieszka Muenster-Wandowski for providing access to electron microscopy, Janet Iwasa for the original model figure, Soren Jorgensen for image processing, and Craig Garner for discussions. We thank EMBO for providing the travel funds (SW). The research was funded by the US National Institutes of Health (NS034307 EMJ), European Research Council grant (249939 SYNGLUT CR), and German

Research Council grants (Neurocure EXC 257 EMJ+CR, SFB 665 + SFB 958 CR). EMJ is an Investigator of the Howard Hughes Medical Institute and is an Alexander von Humboldt Scholar.

References

1. Heuser JE, Reese TS. Evidence for recycling of synaptic vesicle membrane during transmitter release at the frog neuromuscular junction. *J. Cell Biol.* 1973; 57:315–344. [PubMed: 4348786]
2. Maycox PR, Link E, Reetz A, Morris SA, Jahn R. Clathrin-coated vesicles in nervous tissue are involved primarily in synaptic vesicle recycling. *J. Cell Biol.* 1992; 118:1379–1388. [PubMed: 1325974]
3. Takei K, et al. Generation of coated intermediates of clathrin-mediated endocytosis on protein-free liposomes. *Cell.* 1998; 94:131–141. [PubMed: 9674434]
4. Takei K, Mundigl O, Daniell L, De Camilli P. The synaptic vesicle cycle: a single vesicle budding step involving clathrin and dynamin. *J. Cell Biol.* 1996; 133:1237–1250. [PubMed: 8682861]
5. Li C, et al. Ca²⁺-dependent and -independent activities of neural and non-neural synaptotagmins. *Nature.* 1995; 375:594–599. [PubMed: 7791877]
6. Jorgensen EM, et al. Defective recycling of synaptic vesicles in synaptotagmin mutants of *Caenorhabditis elegans*. *Nature.* 1995; 378:196–199. [PubMed: 7477324]
7. Zhang JZ, Davletov BA, Südhof TC, Anderson RGW. Synaptotagmin I is a high affinity receptor for clathrin AP-2: Implications for membrane recycling. *Cell.* 1994; 78:751–760. [PubMed: 8087843]
8. Nonet ML, et al. UNC-11, a *Caenorhabditis elegans* AP180 homologue, regulates the size and protein composition of synaptic vesicles. *Mol. Biol. Cell.* 1999; 10:2343–2360. [PubMed: 10397769]
9. Zhang B, et al. Synaptic vesicle size and number are regulated by a clathrin adaptor protein required for endocytosis. *Neuron.* 1998; 21:1465–1475. [PubMed: 9883738]
10. Watanabe S, et al. Ultrafast endocytosis at mouse hippocampal synapses. *Nature.* 2013; 504:242–247. [PubMed: 24305055]
11. Watanabe S, et al. Ultrafast endocytosis at *Caenorhabditis elegans* neuromuscular junctions. *eLife.* 2013; 2:e00723. [PubMed: 24015355]
12. Christie JM, Jahr CE. Multivesicular release at Schaffer collateral-CA1 hippocampal synapses. *J. Neurosci. Off. J. Soc. Neurosci.* 2006; 26:210–216.
13. Berndt A, et al. High-efficiency channelrhodopsins for fast neuronal stimulation at low light levels. *Proc. Natl. Acad. Sci.* 2011; 108:7595–7600. [PubMed: 21504945]
14. Rizzoli SO, et al. Evidence for early endosome-like fusion of recently endocytosed synaptic vesicles. *Traffic Cph. Den.* 2006; 7:1163–1176.
15. Royle SJ, Bright NA, Lagnado L. Clathrin is required for the function of the mitotic spindle. *Nature.* 2005; 434:1152–1157. [PubMed: 15858577]
16. Lee D, Lin B-J, Lee AK. Hippocampal Place Fields Emerge upon Single-Cell Manipulation of Excitability During Behavior. *Science.* 2012; 337:849–853. [PubMed: 22904011]
17. Sik A, Penttonen M, Ylinen A, Buzsáki G. Hippocampal CA1 interneurons: an in vivo intracellular labeling study. *J. Neurosci.* 1995; 15:6651–6665. [PubMed: 7472426]
18. Macia E, et al. Dynasore, a cell-permeable inhibitor of dynamin. *Dev. Cell.* 2006; 10:839–850. [PubMed: 16740485]
19. Park RJ, et al. Dynamin triple knockout cells reveal off target effects of commonly used dynamin inhibitors. *J. Cell Sci.* 2013; 126:5305–5312. [PubMed: 24046449]
20. Kane RE. Actin polymerization and interaction with other proteins in temperature-induced gelation of sea urchin egg extracts. *J. Cell Biol.* 1976; 71:704–714. [PubMed: 1033188]
21. Jensen V, Walaas SI, Hilfiker S, Ruiz A, Hvalby Ø. A delayed response enhancement during hippocampal presynaptic plasticity in mice. *J. Physiol.* 2007; 583:129–143. [PubMed: 17569738]
22. Hoopmann P, et al. Endosomal sorting of readily releasable synaptic vesicles. *Proc. Natl. Acad. Sci. U. S. A.* 2010; 107:19055–19060. [PubMed: 20956291]

23. Schikorski T. Readily releasable vesicles recycle at the active zone of hippocampal synapses. *Proc. Natl. Acad. Sci. U. S. A.* 2014; 111:5415–5420. [PubMed: 24706824]
24. Miller TM, Heuser JE. Endocytosis of synaptic vesicle membrane at the frog neuromuscular junction. *J. Cell Biol.* 1984; 98:685–698. [PubMed: 6607255]
25. Heuser, J.; Reese, T. Fourth Intensive Study Program Neurosci. M.I.T. press; 1979. p. 573–600.
26. Gisselsson LL, Matus A, Wieloch T. Actin redistribution underlies the sparing effect of mild hypothermia on dendritic spine morphology after in vitro ischemia. *J. Cereb. Blood Flow Metab. Off. J. Int. Soc. Cereb. Blood Flow Metab.* 2005; 25:1346–1355.
27. Hartmann-Petersen R, Walmod PS, Berezin A, Berezin V, Bock E. Individual cell motility studied by time-lapse video recording: influence of experimental conditions. *Cytometry.* 2000; 40:260–270. [PubMed: 10918277]
28. Pyott SJ, Rosenmund C. The effects of temperature on vesicular supply and release in autaptic cultures of rat and mouse hippocampal neurons. *J. Physiol.* 2002; 539:523–535. [PubMed: 11882684]
29. Kushmerick C, Renden R, von Gersdorff H. Physiological temperatures reduce the rate of vesicle pool depletion and short-term depression via an acceleration of vesicle recruitment. *J. Neurosci. Off. J. Soc. Neurosci.* 2006; 26:1366–1377.
30. Teng H, Cole JC, Roberts RL, Wilkinson RS. Endocytic active zones: hot spots for endocytosis in vertebrate neuromuscular terminals. *J. Neurosci. Off. J. Soc. Neurosci.* 1999; 19:4855–4866.
31. Teng H, Lin MY, Wilkinson RS. Macroendocytosis and endosome processing in snake motor boutons. *J. Physiol.* 2007; 582:243–262. [PubMed: 17478535]
32. Micheva KD, Smith SJ. Strong effects of subphysiological temperature on the function and plasticity of mammalian presynaptic terminals. *J. Neurosci. Off. J. Soc. Neurosci.* 2005; 25:7481–7488.
33. Renden R, von Gersdorff H. Synaptic vesicle endocytosis at a CNS nerve terminal: faster kinetics at physiological temperatures and increased endocytotic capacity during maturation. *J. Neurophysiol.* 2007; 98:3349–3359. [PubMed: 17942618]
34. Von Gersdorff H, Matthews G. Dynamics of synaptic vesicle fusion and membrane retrieval in synaptic terminals. *Nature.* 1994; 367:735–739. [PubMed: 7906397]
35. Neves G, Gomis A, Lagnado L. Calcium influx selects the fast mode of endocytosis in the synaptic terminal of retinal bipolar cells. *Proc. Natl. Acad. Sci. U. S. A.* 2001; 98:15282–15287. [PubMed: 11734626]
36. Granseth B, Odermatt B, Royle SJ, Lagnado L. Clathrin-mediated endocytosis is the dominant mechanism of vesicle retrieval at hippocampal synapses. *Neuron.* 2006; 51:773–786. [PubMed: 16982422]
37. Balaji J, Ryan TA. Single-vesicle imaging reveals that synaptic vesicle exocytosis and endocytosis are coupled by a single stochastic mode. *Proc. Natl. Acad. Sci. U. S. A.* 2007; 104:20576–20581. [PubMed: 18077369]
38. Armbruster M, Messa M, Ferguson SM, De Camilli P, Ryan TA. Dynamin phosphorylation controls optimization of endocytosis for brief action potential bursts. *eLife.* 2013; 2:e00845. [PubMed: 23908769]
39. Gad H, Löw P, Zotova E, Brodin L, Shupliakov O. Dissociation between Ca²⁺-triggered synaptic vesicle exocytosis and clathrin-mediated endocytosis at a central synapse. *Neuron.* 1998; 21:607–616. [PubMed: 9768846]
40. Wenk MR, De Camilli P. Protein-lipid interactions and phosphoinositide metabolism in membrane traffic: insights from vesicle recycling in nerve terminals. *Proc. Natl. Acad. Sci. U. S. A.* 2004; 101:8262–8269. [PubMed: 15146067]
41. Takamori S, et al. Molecular anatomy of a trafficking organelle. *Cell.* 2006; 127:831–846. [PubMed: 17110340]
42. Wucherpennig T, Wilsch-Bräuninger M, González-Gaitán M. Role of Drosophila Rab5 during endosomal trafficking at the synapse and evoked neurotransmitter release. *J. Cell Biol.* 2003; 161:609–624. [PubMed: 12743108]

43. Uytterhoeven V, Kuenen S, Kasproicz J, Miskiewicz K, Verstreken P. Loss of skywalker reveals synaptic endosomes as sorting stations for synaptic vesicle proteins. *Cell*. 2011; 145:117–132. [PubMed: 21458671]
44. Gu M, et al. AP2 hemicomplexes contribute independently to synaptic vesicle endocytosis. *eLife*. 2013; 2:e00190. [PubMed: 23482940]
45. Heerssen H, Fetter RD, Davis GW. Clathrin dependence of synaptic-vesicle formation at the *Drosophila* neuromuscular junction. *Curr. Biol. CB*. 2008; 18:401–409. [PubMed: 18356056]
46. Kasproicz J, et al. Inactivation of clathrin heavy chain inhibits synaptic recycling but allows bulk membrane uptake. *J. Cell Biol.* 2008; 182:1007–1016. [PubMed: 18762582]
47. Kononenko NL, et al. Clathrin/AP-2 mediate synaptic vesicle reformation from endosome-like vacuoles but are not essential for membrane retrieval at central synapses. *Neuron*. 2014; 82:981–988. [PubMed: 24908483]
48. Kittelmann M, et al. In vivo synaptic recovery following optogenetic hyperstimulation. *Proc. Natl. Acad. Sci.* 2013; 110:E3007–E3016. [PubMed: 23878262]

Additional references

49. Von Kleist L, et al. Role of the clathrin terminal domain in regulating coated pit dynamics revealed by small molecule inhibition. *Cell*. 2011; 146:471–484. [PubMed: 21816279]
50. Lois C, Hong EJ, Pease S, Brown EJ, Baltimore D. Germline transmission and tissue-specific expression of transgenes delivered by lentiviral vectors. *Science*. 2002; 295:868–872. [PubMed: 11786607]




Fig. 1. Synaptic vesicles are regenerated from endosomes at 34°C. Hippocampal synapses were stimulated once and frozen at the indicated times. (a) Electron micrographs showing invaginations and large endocytic vesicles (arrowheads) recovered via ultrafast endocytosis. Arrow, an endosome in the center of the bouton. (b, c) Micrographs showing single coated buds (b) and multiple coated buds (c) forming on an endosome. (d) Increase in the number of each endocytic structure per synaptic profile after a single stimulus. ~60% of synapses had endosomes in the unstimulated control; this baseline value was subtracted from endosome numbers. After stimulation two endosomes were observed in 30% of the synapses. The prevalence of large endocytic vesicles and endosomes is followed by an increase in the number of clathrin-coated vesicles. Coated pits were rarely observed on the

plasma membrane (PM). The standard error of the mean is shown in each graph. For N values, detailed numbers and statistical analysis, *see* Supplementary Table 1.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript






Fig. 2.

Ultrafast endocytosis is clathrin-independent. (a) Top, average traces for excitatory post-synaptic currents (EPSCs) in a control and a clathrin knock-down neuron from autaptic cultures. Bottom, mean amplitude of EPSCs. The amplitude is reduced by 41% in the knock-down cells. (b) An electron micrograph of a synapse frozen 100 ms after stimulation. Ultrafast endocytic pits (arrows) are observed flanking the active zone. PSD, postsynaptic density. (c) Average number of endocytic pits, and large endocytic vesicles in control and clathrin knock-down neurons with or without stimulation (control 100 ms: 0.17 ± 0.03 endocytic structures/profile; knock-down 100 ms: 0.10 ± 0.03 endocytic structures/profile, $p = 0.06$). The p-values are calculated against the matched time points in the control shRNA treated cells. *** indicates p-value of <0.001 . n.s., 'not significant'. The standard error of the mean is shown in each graph. For N values, detailed numbers and statistical analysis, see Supplementary Table 1.

**Fig. 3.**

Following a single stimulus, clathrin is required at endosomes to regenerate synaptic vesicles. (a,c) Electron micrographs showing ferritin uptake in control (a) and clathrin knock-down (c) synapses at different time points after stimulation. In control neurons, ferritin is observed in large endocytic vesicles after stimulation (middle) and in synaptic vesicles (right), but it is trapped in endosomes in the clathrin knock-down cells. (b, d) Average increase in ferritin-positive endocytic structures in all profiles. (b) In control cells, the total number of synapses containing ferritin remained at 26% at the 3, 10, and 20 s time points; suggesting that 74% of synapses were silent. The number of synaptic vesicles is higher because after an endosome is resolved it leads to ~2 vesicles that contain a clump of ferritin. The number of clathrin-coated vesicles is less than expected given that endosomes contain 5-8 synaptic vesicles worth of membrane. This discrepancy is likely because synaptic vesicles will be distributed among many thin sections. (d) In the clathrin knockdown ferritin stalled in endosomes and did not progress into synaptic vesicles. Clathrin-coated pits were not present on the plasma membrane at any time point and are therefore not plotted. Black arrows indicate ferritin-positive structures. Black arrowhead in (a) represents ferritin-positive synaptic vesicles docked in active zone. The standard error of the mean is shown in each graph. For N values, detailed numbers and statistical analysis, see Supplementary Table 1.

**Fig. 4.**

Following high-frequency stimulation, clathrin is required at endosomes to regenerate synaptic vesicles. (a, c) Electron micrographs showing ferritin uptake in control (a) and clathrin knock-down neurons (c) at different time points after stimulation. Black arrows indicate ferritin-positive structures. (b, d) Average number of ferritin-positive endocytic structures increased compared to unstimulated cells (0 ms time point) infected with control shRNA (b) or clathrin knock-down shRNA (d). Clathrin-coated pits were not present on the plasma membrane at any time points and thus not plotted. In the controls, synapses which exhibited ferritin uptake remained at 32% but the number of ferritin structures per profile increased because endosomes were typically resolved into ~2 ferritin-positive vesicles. (e) Virtual section from an electron tomogram (left) and reconstruction (right) showing a string of large endocytic vesicles trapped on the membrane by dynasore treatment following 100 stimuli at 20 Hz. Multiple large endocytic vesicles (black arrows) are attached to one another and remain connected to the plasma membrane. (f) Fraction of tomograms that contain vesicle strings attached to the plasma membrane following 1, 3, 10, 30, and 100 stimuli (20 Hz). Tomograms from 100 nm thick sections were reconstructed for each time point; only 1 vesicle string appeared in a given terminal. The number of tomograms with a


vesicle string reached 31% following 100 stimuli, suggesting that ~70% of synapses are silent in these preparations. The number of vesicles on a string increases with repetitive stimulation: 1 stimulus (no vesicle string); 3 stimuli (2 vesicles / string); 10 stimuli (2.3 vesicles /string), 30 stimuli (5.6 vesicles/ string); and 100 stimuli (4.8 vesicles/ string). However, these are likely underestimates because strings that extend into neighboring sections are not captured in these tomograms. The standard error of the mean is shown in each graph. For N values, detailed numbers and statistical analysis, *see* Supplementary Table 1.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Fig. 5.**

Clathrin regenerates synaptic vesicles from plasma membrane in the absence of ultrafast endocytosis. Electron micrographs showing a synaptic terminal 3 s after a single stimulus from cells incubated with 0.1% DMSO (a), with 10 μ M latrunculin-A (c), at 34°C (e), and at 22°C (g). (b,d,f,h) Average number of endocytic structures in profiles infected with control shRNA (b), clathrin knock-down cells (d), cells incubated at 34°C (f), and cells incubated at 22°C (h). The total number of structures per profile is plotted. Large vesicles accumulated in the center of the bouton, likely reflecting the fusion of small endocytic vesicles to endosomes. The standard error of the mean is shown in each graph. For N values, detailed numbers and statistical analysis, *see* Supplementary Table 1.