



Draft Genome Sequence of *Pseudomonas putida* CA-3, a Bacterium Capable of Styrene Degradation and Medium-Chain-Length Polyhydroxyalkanoate Synthesis

Eduardo L. Almeida,^a Lekha M. Margassery,^a Niall O'Leary,^{a,b} Alan D. W. Dobson^{a,b}

^aSchool of Microbiology, University College Cork, Cork, Ireland

^bEnvironmental Research Institute, University College Cork, Cork, Ireland

ABSTRACT *Pseudomonas putida* strain CA-3 is an industrial bioreactor isolate capable of synthesizing biodegradable polyhydroxyalkanoate polymers via the metabolism of styrene and other unrelated carbon sources. The pathways involved are subject to regulation by global cellular processes. The draft genome sequence is 6,177,154 bp long and contains 5,608 predicted coding sequences.

Styrene is a solvent used extensively in the polymer-processing industry. It is a toxic compound which is known to have numerous adverse effects on human health (1–4). As a result of this toxicity, there is considerable interest in styrene waste management solutions, including the potential for microbial bioremediation. The bacterium *Pseudomonas putida* CA-3 was isolated from an industrial bioreactor following enrichment on styrene as a sole carbon source (5). In addition to styrene degradation, the organism has also been shown to convert styrene to medium-chain-length polyhydroxyalkanoates (PHA), biodegradable polyesters with physicochemical properties suitable for a range of industrial and medical applications (6). The styrene catabolic pathway in the genus *Pseudomonas* (four major steps) has been described (7). The styrene and PHA pathways have been found to be subject to global regulatory processes in *P. putida* CA-3 (8). However, despite considerable pathway characterization to date, there is still much to be elucidated regarding the overlying cellular mechanisms, including at the genomics level, hence the importance of the draft genome of the strain presented here.

The genomic DNA of the isolate was obtained using the phenol-chloroform-isoamyl alcohol extraction method (9). The sequencing was performed by Macrogen (Seoul, South Korea) using Illumina's MiSeq paired-end technology. The sequencing generated 3,332,054 reads and 1,002,948,254 bp. The raw data had adapters trimmed using Scythe v.0.994 (see <https://github.com/vsbuffalo/scythe>) and Sickle v.1.33 programs (10). The reads were then quality filtered and trimmed using FaQCs v.1.35 for a minimum quality value (QV) score of >20, resulting in 3,321,320 reads and 916,019,954 bp, with an approximate coverage of 150-fold (11). The processed reads were assembled *de novo* using SPAdes v.3.10.1, and contigs of <500 bp were removed (12).

The quality of the assembly was assessed using QUAST 4.5, resulting in 92 contigs, 6,177,154 bp, an N_{50} of 165,779 bp, and a G+C content of 61.89% (13). The annotation was performed using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v.4.3, which predicted 5,608 coding sequences (CDSs), 6 rRNAs, and 70 tRNAs (14).

Accession number(s). This whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank under the accession no. [PIJT00000000](https://doi.org/10.1128/genomeA.01534-17). The version described in this paper is the first version, PIJT01000000.

ACKNOWLEDGMENT

This work was supported by the Brazilian National Council for Scientific and Technological Development (CNPq).

Received 8 December 2017 Accepted 15 December 2017 Published 25 January 2018

Citation Almeida EL, Margassery LM, O'Leary N, Dobson ADW. 2018. Draft genome sequence of *Pseudomonas putida* CA-3, a bacterium capable of styrene degradation and medium-chain-length polyhydroxyalkanoate synthesis. *Genome Announc* 6:e01534-17. <https://doi.org/10.1128/genomeA.01534-17>.

Copyright © 2018 Almeida et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Alan D. W. Dobson, a.dobson@ucc.ie.

REFERENCES

1. Christensen MS, Hansen J, Ramlau-Hansen CH, Toft G, Kolstad H. 2017. Cancer incidence in workers exposed to styrene in the Danish-reinforced plastics industry, 1968–2012. *Epidemiology* 28:300–310. <https://doi.org/10.1097/EDE.0000000000000608>.
2. Costa S, Ceppi M, Costa C, Silva S, Pereira C, Laffon B, Bonassi S, Teixeira JP. 2016. The cytokinesis-block micronucleus (CBMN) assay in human populations exposed to styrene: a systematic review and meta-analysis. *Mutat Res* 770:92–105. <https://doi.org/10.1016/j.mrrev.2016.06.003>.
3. McCague AB, Cox-Ganser JM, Harney JM, Alwis KU, Blount BC, Cummings KJ, Edwards N, Kreiss K. 2015. Styrene-associated health outcomes at a windblade manufacturing plant. *Am J Ind Med* 58:1150–1159. <https://doi.org/10.1002/ajim.22516>.
4. Nett RJ, Edwards NT, Ruder AM, Bertke SJ, Keumala I, Cox-Ganser J, Cummings KJ. 2017. Deaths from nonmalignant respiratory disease in styrene-exposed workers: does obliterative bronchiolitis contribute to mortality? *Ann Am Thorac Soc* 14:810–811. <https://doi.org/10.1513/AnnalsATS.201612-1026LE>.
5. O'Connor K, Buckley CM, Hartmans S, Dobson AD. 1995. Possible regulatory role for nonaromatic carbon sources in styrene degradation by *Pseudomonas putida* CA-3. *Appl Environ Microbiol* 61:544–548.
6. Ward PG, de Roo G, O'Connor KE. 2005. Accumulation of polyhydroxyalkanoate from styrene and phenylacetic acid by *Pseudomonas putida* CA-3. *Appl Environ Microbiol* 71:2046–2052. <https://doi.org/10.1128/AEM.71.4.2046-2052.2005>.
7. O'Leary ND, O'Mahony MM, Dobson AD. 2011. Regulation of phenylacetic acid uptake is σ^{54} dependent in *Pseudomonas putida* CA-3. *BMC Microbiol* 11:229. <https://doi.org/10.1186/1471-2180-11-229>.
8. Ryan WJ, O'Leary ND, O'Mahony M, Dobson ADW. 2013. GacS-dependent regulation of polyhydroxyalkanoate synthesis in *Pseudomonas putida* CA-3. *Appl Environ Microbiol* 79:1795–1802. <https://doi.org/10.1128/AEM.02962-12>.
9. Wilson K. 2001. Preparation of genomic DNA from bacteria. *Curr Protoc Mol Biol* Chapter 2:Unit 2.4. <https://doi.org/10.1002/0471142727.mb0204s56>.
10. Joshi N, Fass J. 2011. Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files (version 1.33). <https://github.com/najoshi/sickle>. Accessed 1 September 2017.
11. Lo CC, Chain PSG. 2014. Rapid evaluation and quality control of next generation sequencing data with FaQCs. *BMC Bioinformatics* 15:366. <https://doi.org/10.1186/s12859-014-0366-2>.
12. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
13. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
14. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.