



Crystal structure of 1'-(prop-2-yn-1-yl)-1,4-dihydrospiro[benzo[*d*][1,3]oxazine-2,3'-indolin]-2'-one

Y. AaminaNaaz,^a J. Kamalraja,^b G. Vimala,^a P. T. Perumal^b and A. SubbiahPandi^{a*}

^aDepartment of Physics, Presidency College (Autonomous), Chennai 600 005, India, and ^bOrganic Chemistry Division, Central Leather Research Institute, Adyar, Chennai 602 020, India. *Correspondence e-mail: aspandian59@gmail.com

Received 11 June 2015; accepted 22 June 2015

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland


In the title compound, C₁₈H₁₄N₂O₂, the six-membered oxazine ring adopts a half-chair conformation and its mean plane makes a dihedral angle of 83.23 (7)° with the pyrrolidine ring of the indoline ring system. In the crystal, molecules are linked via N—H···O hydrogen bonds, forming chains along [100]. The chains are linked by C—H···π interactions, forming slabs parallel to (001).

Keywords: crystal structure; spiro compounds; spirooxazines; oxazine; indoline; N—H···O hydrogen bonding.

CCDC reference: 1408024

1. Related literature

For the biological activity of spiro compounds, see: James *et al.* (1991); Kobayashi *et al.* (1991). For the use of 1,3-dipolar cycloaddition reactions in the construction of spiro compounds, see: Caramella & Grunanger (1984). For applications of spirooxazine derivatives, see: Chibisov & Görner (1999). For the synthetic method, see: Kamalraja *et al.* (2014).



2. Experimental

2.1. Crystal data

C ₁₈ H ₁₄ N ₂ O ₂	$\gamma = 74.125 (3)^\circ$
$M_r = 290.31$	$V = 703.17 (6) \text{ \AA}^3$
Triclinic, <i>P</i> $\bar{1}$	$Z = 2$
$a = 5.5571 (3) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 8.5404 (4) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 15.4542 (9) \text{ \AA}$	$T = 293 \text{ K}$
$\alpha = 85.884 (3)^\circ$	$0.21 \times 0.19 \times 0.18 \text{ mm}$
$\beta = 86.814 (3)^\circ$	

2.2. Data collection

Bruker SMART APEXII CCD diffractometer	16184 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2008)	3231 independent reflections
$T_{\min} = 0.981$, $T_{\max} = 0.984$	2350 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.031$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$	199 parameters
$wR(F^2) = 0.106$	H-atom parameters constrained
$S = 1.06$	$\Delta\rho_{\text{max}} = 0.15 \text{ e \AA}^{-3}$
3231 reflections	$\Delta\rho_{\text{min}} = -0.21 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å, °).

Cg3 and Cg4 are the centroids of rings C1–C6 and C9–C14, respectively.

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1···O2 ⁱ	0.86	2.13	2.9641 (16)	164
C4—H4···Cg4 ⁱⁱ	0.93	2.90	3.6572 (19)	140
C8—H8A···Cg4 ⁱⁱⁱ	0.97	2.86	3.6636 (17)	141
C16—H16B···Cg3 ^{iv}	0.97	2.79	3.5341 (18)	134

Symmetry codes: (i) $x + 1, y, z$; (ii) $x + 1, y - 1, z$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $x - 1, y, z$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINTE* (Bruker, 2008); data reduction: *SAINTE*; program(s) used to solve

structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97* and *PLATON*.

Acknowledgements

The authors thank Dr Babu Varghese, SAIF, IIT, Chennai, India, for the data collection.

Supporting information for this paper is available from the IUCr electronic archives (Reference: SU5155).

References

- Bruker (2008). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Caramella, P. & Grunanger, P. (1984). *1,3-Dipolar Cycloaddition Chemistry*, Vol. 1, edited by A. Padwa, pp. 291–312. New York: Wiley.
- Chibisov, A. K. & Görner, H. (1999). *J. Phys. Chem. A*, **103**, 5211–5216.
- James, D., Kunze, H. B. & Faulkner, D. (1991). *J. Nat. Prod.* **54**, 1137–1140.
- Kamalraja, J., Murugasan, P. & Perumal, P. T. (2014). *RSC Adv.* **4**, 19422–19432.
- Kobayashi, J., Tsuda, M., Agemi, K., Shigemori, H., Ishibashi, M., Sasaki, T. & Mikami, Y. (1991). *Tetrahedron*, **47**, 6617–6622.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supporting information

Acta Cryst. (2015). E71, o510–o511 [doi:10.1107/S2056989015011949]

Crystal structure of 1'-(prop-2-yn-1-yl)-1,4-dihydrospiro[benzo[*d*][1,3]oxazine-2,3'-indolin]-2'-one

Y. AaminaNaaz, J. Kamalraja, G. Vimala, P. T. Perumal and A. SubbiahPandi

S1. Synthesis and crystallization

A mixture of N-propargylisatin (1.0 mmol), and 2-aminobenzylalcohol (1.0 mmol) was refluxed in ethanol, in the presence of InCl₃ (10 mol%), for 2 h. After the reaction was complete as indicated by TLC, the reaction mixture was cooled to room temperature. The solid that formed was filtered, dried and recrystallized in ethanol or dichloromethane to obtain in good yield (89%) of the pure title product as block-like colourless crystals.

S2. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. The N- and C-bound H atoms were positioned geometrically (N—H = 0.86 Å, C—H = 0.93–0.97 Å) and allowed to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N,C})$.

S3. Structural commentary

Spiro compounds represent an important class of naturally occurring substances, which in many cases exhibit useful biological properties (Kobayashi *et al.*, 1991; James *et al.*, 1991). 1,3-dipolar cycloaddition reactions are widely used for construction of spiro-compounds (Caramella & Grunanger, 1984). It has also been reported that spiro-oxazine derivatives have real or potential applications in many fields such as protection, decoration, display, memory, switches, photography, photometry and photomechanics (Chibisov & Görner, 1999). Efforts have been made to design this industrially and biologically active hetrocyclic compounds by making or breaking carbon-carbon (C—C) and carbon-hetero atom (C—X) (Kamalraja *et al.*, 2014). This InCl₃-mediated compound have been synthesized as a part of the effort carried to develop eco-friendly potential compound by new synthetic method.

The molecular structure of the title compound is illustrated in Fig 1. The oxazine ring (O1/N1/C7/C8/C9/C14) adopts a half chair confirmation, and its mean plane makes a dihedral angle of 83.23 (7) ° with the pyrrolidine ring (O1/N1/C8/C9/C14) of the indolinone ring system. The indole ring system is essentially planar, with atoms C16 and O2 deviating from its mean plane by -0.0130 and 0.0273 Å, respectively. The dihedral angle between the benzene ring (C1—C6) of the indoline ring system and the benzene ring (C9—C14) of the mean plane of the 2,4-dihydro-1*H*-benzo[*d*][1,3]oxazine ring system is 76.94 (8) °.

In the crystal, molecules are linked *via* N—H···O hydrogen bonds (Table 1) forming chains along [100], as shown in Fig 2. The chains are linked by C—H··· π interactions forming slabs parallel to (001); see Table 1.





Figure 1

The molecular structure of the title compound, with atom labelling. Displacement ellipsoids are drawn at the 30% probability level.

**Figure 2**

The crystal packing of the title compound, viewed along the *c* axis. Hydrogen bonds are shown as dashed lines (see Table 1 for details).

1'-(prop-2-yn-1-yl)-1,4-dihydrospiro[benzo[*d*][1,3]oxazine-2,3'-indolin]-2'-one

Crystal data

$C_{18}H_{14}N_2O_2$

$M_r = 290.31$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 5.5571$ (3) Å

$b = 8.5404$ (4) Å

$c = 15.4542$ (9) Å

$\alpha = 85.884$ (3)°

$\beta = 86.814$ (3)°

$\gamma = 74.125$ (3)°

$V = 703.17$ (6) Å³

$Z = 2$

$F(000) = 304$

$D_x = 1.371$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2350 reflections

$\theta = 2.5$ – 27.6 °

$\mu = 0.09$ mm⁻¹

$T = 293$ K

Block, colourless

$0.21 \times 0.19 \times 0.18$ mm

Data collection

Bruker SMART APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω and ϕ scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2008)

$T_{\min} = 0.981$, $T_{\max} = 0.984$

16184 measured reflections

3231 independent reflections

2350 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\max} = 27.6$ °, $\theta_{\min} = 2.5$ °

$h = -7 \rightarrow 7$

$k = -11 \rightarrow 11$

$l = -20 \rightarrow 20$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.041$

$wR(F^2) = 0.106$

$S = 1.06$

3231 reflections

199 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0405P)^2 + 0.1883P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.15 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.21 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.44655 (19)	0.31187 (12)	0.35868 (6)	0.0365 (3)
N1	0.6584 (2)	0.48707 (15)	0.28492 (8)	0.0383 (3)
H1	0.7953	0.4869	0.2560	0.046*
O2	0.14706 (18)	0.52580 (13)	0.21423 (7)	0.0413 (3)
N2	0.3795 (2)	0.29626 (15)	0.15105 (8)	0.0352 (3)
C6	0.7141 (3)	0.21178 (17)	0.23854 (9)	0.0322 (3)
C9	0.3552 (3)	0.59601 (17)	0.39673 (9)	0.0321 (3)
C14	0.5492 (2)	0.61191 (17)	0.33911 (9)	0.0299 (3)
C7	0.5434 (2)	0.35940 (17)	0.27729 (9)	0.0296 (3)
C13	0.6337 (3)	0.75102 (18)	0.33677 (10)	0.0379 (3)
H13	0.7673	0.7600	0.2996	0.045*
C15	0.3291 (2)	0.40839 (17)	0.21203 (9)	0.0306 (3)
C8	0.2756 (3)	0.44185 (18)	0.40275 (10)	0.0369 (3)
H8A	0.2598	0.4071	0.4635	0.044*
H8B	0.1120	0.4635	0.3782	0.044*
C10	0.2424 (3)	0.7231 (2)	0.44848 (10)	0.0428 (4)
H10	0.1106	0.7142	0.4865	0.051*
C1	0.6078 (3)	0.17761 (17)	0.16557 (9)	0.0329 (3)
C12	0.5198 (3)	0.87512 (19)	0.38939 (11)	0.0453 (4)
H12	0.5767	0.9681	0.3877	0.054*
C17	0.3297 (3)	0.31873 (19)	-0.00454 (11)	0.0432 (4)
C16	0.2147 (3)	0.2998 (2)	0.08103 (10)	0.0442 (4)
H16A	0.0651	0.3894	0.0879	0.053*
H16B	0.1641	0.1994	0.0851	0.053*
C5	0.9381 (3)	0.11227 (19)	0.26579 (10)	0.0420 (4)
H5	1.0113	0.1349	0.3144	0.050*
C11	0.3219 (3)	0.8629 (2)	0.44470 (11)	0.0481 (4)
H11	0.2425	0.9483	0.4792	0.058*
C3	0.9441 (3)	-0.05574 (19)	0.14768 (12)	0.0501 (4)
H3	1.0233	-0.1477	0.1177	0.060*
C2	0.7189 (3)	0.04466 (19)	0.11901 (11)	0.0438 (4)
H2	0.6462	0.0227	0.0701	0.053*

C4	1.0533 (3)	-0.0227 (2)	0.21942 (12)	0.0496 (4)
H4	1.2056	-0.0915	0.2369	0.060*
C18	0.4183 (4)	0.3288 (2)	-0.07400 (13)	0.0641 (5)
H18	0.4891	0.3368	-0.1295	0.077*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0434 (6)	0.0375 (5)	0.0296 (5)	-0.0140 (5)	0.0021 (4)	0.0025 (4)
N1	0.0319 (6)	0.0435 (7)	0.0449 (7)	-0.0192 (6)	0.0108 (5)	-0.0129 (6)
O2	0.0295 (5)	0.0451 (6)	0.0435 (6)	0.0000 (5)	-0.0013 (4)	-0.0031 (5)
N2	0.0304 (6)	0.0400 (7)	0.0345 (7)	-0.0064 (5)	-0.0075 (5)	-0.0056 (5)
C6	0.0310 (7)	0.0313 (7)	0.0336 (7)	-0.0080 (6)	-0.0025 (6)	0.0011 (6)
C9	0.0294 (7)	0.0403 (8)	0.0267 (7)	-0.0093 (6)	-0.0037 (6)	-0.0011 (6)
C14	0.0263 (7)	0.0343 (7)	0.0294 (7)	-0.0078 (6)	-0.0055 (5)	-0.0013 (6)
C7	0.0286 (7)	0.0343 (7)	0.0270 (7)	-0.0106 (6)	-0.0007 (5)	-0.0003 (6)
C13	0.0370 (8)	0.0390 (8)	0.0406 (8)	-0.0154 (7)	-0.0039 (6)	-0.0001 (7)
C15	0.0260 (7)	0.0342 (7)	0.0321 (7)	-0.0102 (6)	0.0018 (6)	0.0012 (6)
C8	0.0350 (8)	0.0468 (9)	0.0310 (7)	-0.0155 (7)	0.0041 (6)	-0.0023 (6)
C10	0.0388 (9)	0.0531 (9)	0.0352 (8)	-0.0096 (7)	0.0022 (7)	-0.0084 (7)
C1	0.0317 (7)	0.0305 (7)	0.0365 (8)	-0.0085 (6)	-0.0032 (6)	0.0002 (6)
C12	0.0525 (10)	0.0362 (8)	0.0502 (10)	-0.0149 (7)	-0.0113 (8)	-0.0037 (7)
C17	0.0513 (10)	0.0395 (8)	0.0392 (9)	-0.0107 (7)	-0.0122 (7)	-0.0028 (7)
C16	0.0373 (8)	0.0585 (10)	0.0390 (9)	-0.0140 (7)	-0.0111 (7)	-0.0057 (7)
C5	0.0376 (8)	0.0429 (9)	0.0426 (9)	-0.0054 (7)	-0.0098 (7)	0.0008 (7)
C11	0.0531 (10)	0.0440 (9)	0.0449 (9)	-0.0060 (8)	-0.0048 (8)	-0.0142 (7)
C3	0.0519 (10)	0.0332 (8)	0.0597 (11)	-0.0012 (7)	0.0001 (8)	-0.0083 (8)
C2	0.0480 (9)	0.0363 (8)	0.0469 (9)	-0.0084 (7)	-0.0061 (7)	-0.0087 (7)
C4	0.0412 (9)	0.0395 (9)	0.0596 (11)	0.0032 (7)	-0.0066 (8)	0.0013 (8)
C18	0.0887 (15)	0.0581 (12)	0.0450 (11)	-0.0205 (11)	-0.0025 (10)	0.0033 (9)

Geometric parameters (Å, °)

O1—C7	1.4168 (16)	C9—C8	1.495 (2)
O1—C8	1.4347 (17)	C14—C13	1.390 (2)
N1—C14	1.3872 (17)	C7—C15	1.5525 (19)
N1—C7	1.4212 (17)	C13—C12	1.373 (2)
O2—C15	1.2150 (16)	C10—C11	1.378 (2)
N2—C15	1.3554 (18)	C1—C2	1.368 (2)
N2—C1	1.4078 (18)	C12—C11	1.377 (2)
N2—C16	1.4497 (18)	C17—C18	1.163 (2)
C6—C5	1.370 (2)	C17—C16	1.454 (2)
C6—C1	1.3861 (19)	C5—C4	1.384 (2)
C6—C7	1.4964 (19)	C3—C4	1.375 (2)
C9—C10	1.381 (2)	C3—C2	1.385 (2)
C9—C14	1.3888 (19)		
C7—O1—C8	114.81 (10)	C6—C7—C15	101.78 (11)

C14—N1—C7	119.77 (11)	C12—C13—C14	119.93 (15)
C15—N2—C1	111.34 (11)	O2—C15—N2	125.26 (13)
C15—N2—C16	123.33 (12)	O2—C15—C7	126.81 (13)
C1—N2—C16	125.32 (12)	N2—C15—C7	107.93 (11)
C5—C6—C1	120.23 (13)	O1—C8—C9	113.31 (11)
C5—C6—C7	130.43 (13)	C11—C10—C9	121.15 (15)
C1—C6—C7	109.30 (12)	C2—C1—C6	121.96 (14)
C10—C9—C14	118.87 (14)	C2—C1—N2	128.46 (13)
C10—C9—C8	121.35 (13)	C6—C1—N2	109.58 (12)
C14—C9—C8	119.77 (12)	C13—C12—C11	120.49 (15)
N1—C14—C9	119.32 (12)	C18—C17—C16	177.26 (18)
N1—C14—C13	120.65 (13)	N2—C16—C17	113.13 (13)
C9—C14—C13	120.03 (13)	C6—C5—C4	118.59 (15)
O1—C7—N1	111.37 (11)	C12—C11—C10	119.46 (15)
O1—C7—C6	108.82 (11)	C4—C3—C2	121.54 (15)
N1—C7—C6	113.52 (11)	C1—C2—C3	117.24 (15)
O1—C7—C15	108.92 (10)	C3—C4—C5	120.44 (15)
N1—C7—C15	111.96 (11)		
C7—N1—C14—C9	11.5 (2)	N1—C7—C15—N2	124.05 (12)
C7—N1—C14—C13	-169.20 (13)	C6—C7—C15—N2	2.48 (14)
C10—C9—C14—N1	-177.99 (13)	C7—O1—C8—C9	-41.18 (16)
C8—C9—C14—N1	3.0 (2)	C10—C9—C8—O1	-167.24 (13)
C10—C9—C14—C13	2.7 (2)	C14—C9—C8—O1	11.71 (19)
C8—C9—C14—C13	-176.28 (13)	C14—C9—C10—C11	-1.1 (2)
C8—O1—C7—N1	55.09 (15)	C8—C9—C10—C11	177.88 (14)
C8—O1—C7—C6	-179.03 (11)	C5—C6—C1—C2	0.6 (2)
C8—O1—C7—C15	-68.86 (14)	C7—C6—C1—C2	-177.30 (13)
C14—N1—C7—O1	-39.97 (17)	C5—C6—C1—N2	179.83 (13)
C14—N1—C7—C6	-163.21 (12)	C7—C6—C1—N2	1.91 (16)
C14—N1—C7—C15	82.24 (15)	C15—N2—C1—C2	178.94 (15)
C5—C6—C7—O1	-65.35 (19)	C16—N2—C1—C2	-0.4 (2)
C1—C6—C7—O1	112.28 (13)	C15—N2—C1—C6	-0.21 (16)
C5—C6—C7—N1	59.3 (2)	C16—N2—C1—C6	-179.52 (14)
C1—C6—C7—N1	-123.10 (13)	C14—C13—C12—C11	0.0 (2)
C5—C6—C7—C15	179.75 (15)	C15—N2—C16—C17	118.22 (16)
C1—C6—C7—C15	-2.62 (14)	C1—N2—C16—C17	-62.5 (2)
N1—C14—C13—C12	178.50 (13)	C18—C17—C16—N2	133 (4)
C9—C14—C13—C12	-2.2 (2)	C1—C6—C5—C4	-0.5 (2)
C1—N2—C15—O2	178.55 (13)	C7—C6—C5—C4	176.88 (15)
C16—N2—C15—O2	-2.1 (2)	C13—C12—C11—C10	1.6 (2)
C1—N2—C15—C7	-1.51 (15)	C9—C10—C11—C12	-1.1 (2)
C16—N2—C15—C7	177.82 (13)	C6—C1—C2—C3	0.0 (2)
O1—C7—C15—O2	67.59 (17)	N2—C1—C2—C3	-179.08 (15)
N1—C7—C15—O2	-56.01 (18)	C4—C3—C2—C1	-0.6 (3)
C6—C7—C15—O2	-177.58 (13)	C2—C3—C4—C5	0.7 (3)
O1—C7—C15—N2	-112.35 (12)	C6—C5—C4—C3	-0.1 (3)

Hydrogen-bond geometry (Å, °)

Cg3 and Cg4 are the centroids of rings C1–C6 and C9–C14, respectively.

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots O2 ⁱ	0.86	2.13	2.9641 (16)	164
C4—H4 \cdots Cg4 ⁱⁱ	0.93	2.90	3.6572 (19)	140
C8—H8 <i>A</i> \cdots Cg4 ⁱⁱⁱ	0.97	2.86	3.6636 (17)	141
C16—H16 <i>B</i> \cdots Cg3 ^{iv}	0.97	2.79	3.5341 (18)	134

Symmetry codes: (i) $x+1, y, z$; (ii) $x+1, y-1, z$; (iii) $-x+1, -y+1, -z+1$; (iv) $x-1, y, z$.