

## Linkage studies of catechol-O-methyltransferase (COMT) and dopamine-beta-hydroxylase (DBH) cDNA expression levels

Chao Xing<sup>1,2,3</sup>, Monica Torres-Caban<sup>1</sup>, Tao Wang<sup>1</sup>, Qing Lu<sup>1</sup>, Guan Xing<sup>1</sup> and Robert C Elston\*<sup>1</sup>

Address: <sup>1</sup>Department of Epidemiology and Biostatistics, Case Western Reserve University, Wolstein Research Building 2103, Cornell Drive, Cleveland, Ohio 44106, USA, <sup>2</sup>Department of Clinical Sciences, University of Texas Southwestern Medical Center, 5323 Harry Hines Boulevard, Dallas, Texas 75390-8591, USA and <sup>3</sup>McDermott Center for Human Growth and Development, University of Texas Southwestern Medical Center, 5323 Harry Hines Boulevard, Dallas, Texas 75390-8591, USA

Email: Chao Xing - chao.xing@utsouthwestern.edu; Monica Torres-Caban - monica.torres-caban@case.edu; Tao Wang - tao.wang@case.edu; Qing Lu - qing.lu@case.edu; Guan Xing - guan.xing@bms.com; Robert C Elston\* - rce@darwin.case.edu

\* Corresponding author

from Genetic Analysis Workshop 15  
St. Pete Beach, Florida, USA. 11–15 November 2006

Published: 18 December 2007

BMC Proceedings 2007, 1(Suppl 1):S95

This article is available from: <http://www.biomedcentral.com/1753-6561/1/S1/S95>

© 2007 Xing et al; licensee BioMed Central Ltd.

This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

The *COMT* and *DBH* genes are physically located at chromosomes 22q11 and 9q34, respectively, and both *COMT* and *DBH* are involved in catecholamine metabolism and are strong candidates for certain psychiatric and neurological disorders. Although the genetic determinants for both enzymes' activities have been widely studied, their genetic involvement on gene mRNA expression levels remains unclear. In this study we performed quantitative linkage analysis of *COMT* and *DBH* cDNA expression levels, identifying transcriptional regulatory regions for both genes. Multiple Haseman-Elston regression was used to detect both additive and interactive effects between two loci. We found that the master transcriptional regulatory region 20q13 had an additive effect on the *COMT* expression level. We also found that chromosome 19p13 showed both additive and interactive effects with 9q34 on *DBH* expression level. Furthermore, a potential interaction between *COMT* and *DBH* was indicated.

### Background

Catechol-O-methyltransferase (*COMT*) (MIM 116790) catalyzes the transfer of a methyl group from S-adenosyl-methionine to catecholamines [1], and dopamine-beta-hydroxylase (*DBH*) (MIM 223360) catalyzes the conversion of dopamine to norepinephrine [2]; both enzymes are involved in the catecholamine metabolism system, in which abnormalities are hypothesized to be related to the development of psychiatric and neurological disorders

such as schizophrenia and depression (for reviews, see [3,4]). There are two distinct forms of *COMT*, a membrane-bound *COMT* (MB-*COMT*) and a soluble *COMT* (S-*COMT*) found in the cell cytoplasm [5]; the former is predominantly expressed in brain neurons, whereas the latter is predominantly expressed in other tissues such as liver and kidney. *DBH* is expressed in the vesicles of central noradrenergic and adrenergic neurons, peripheral noradrenergic neurons, and adrenomedullary neuron-

secretory cells [6]. The *COMT* and *DBH* genes are physically located at chromosomes 22q11 and 9q34, respectively [7,8].

There has been intensive research on the genetic determinants of plasma *COMT* and *DBH* activities, and findings so far strongly suggest that the structural genes *COMT* and *DBH* encoding these proteins are the major quantitative trait loci for their respective plasma activities; moreover, in both genes single-nucleotide polymorphisms (SNPs) have been identified that may account for most of the variation in the enzyme activity levels. On the common functional SNP rs165688, a G→A substitution in exon 4 of *COMT* produces a valine→methionine (Val/Met) substitution at codons 158 and 108 in the MB-*COMT* and S-*COMT* transcripts, respectively. This polymorphism has been shown to have a functional effect on enzyme activity; in particular, Val is a predominant factor that determines higher *COMT* activity [9]. The SNP rs1611115 (-1021C→T) located in the 5' flanking region of the *DBH* gene accounts for 35% to 52% of the variance in plasma *DBH* activity in populations from diverse geographic origins [10].

Though the major effects of these common variants on enzyme activities have been replicated in independent samples, their effects on gene mRNA expression levels remain unclear. The effect of SNP rs165688 on *COMT* mRNA expression level has been controversial. Bray et al. [11] showed that it altered the *COMT* mRNA expression level; in contrast, several other studies failed to detect this association between *COMT* genotypes and expression level (for examples, see [12]). So far, studies on the determinants of *COMT* expression level have been limited to seeking *cis*-acting regulators, including SNP rs165688 and several other SNPs within *COMT*. To our knowledge, there is as yet no literature documenting effects of SNPs on *DBH* mRNA expression levels.

Phenotypic variation in the transcript level represents an intermediate stage between DNA sequence differences and the clinical expression of complex traits, and there have been studies suggesting a genetic contribution to the variation in gene expression level [13-15]. In the current study, we performed quantitative linkage analysis for both *COMT* and *DBH* expression phenotypes, identifying transcriptional regulatory regions for both genes. For *COMT*, we first identified *cis*-acting influences on the gene expression level, and then we performed a genome-wide linkage scan for *trans*-acting regulators across the genome conditional on the *cis*-acting effect at chromosome 22q11. For *DBH*, we similarly first identified the *cis*-acting influences on the gene expression level, and then, both conditional and unconditional on the *cis*-acting effect at chromosome 9q34, performed a linkage scan for *trans*-acting regulators

specifically on chromosome 19, based on the evidence from previous linkage studies [16-18] that there might be a second locus near chromosome 19p13 acting jointly with *DBH* at chromosome 9q34 to influence *DBH* plasma activity.

## Methods

The study subjects consisted of 14 three-generation CEPH (Centre d'Etude du Polymorphisme Humain) Utah pedigrees, including 371 full sibling pairs. 2,820 SNPs were relatively evenly genotyped across the 22 autosomes by the SNP Consortium [19]. Treating the expression level of *COMT* and *DBH* in lymphoblastoid cells of the above subjects as a quantitative trait, we applied Haseman-Elston regression model-free linkage analyses [20,21], as implemented in the SIBPAL program. In particular, only data on sib pairs were considered (thus avoiding the complication of possible intergenerational heterogeneity) and the generalized estimating equation technique [22] was utilized to allow for non-independence of sib pairs within a sibship. The basic Haseman-Elston (HE) regression can be written in the form

$$Y = \beta_0 + \beta_1 \hat{\pi}_i,$$

where  $Y$  is a function of traits for a sib pair,  $\hat{\pi}_i$  is the estimated proportion of alleles shared identically by descent (IBD) at marker locus  $i$  between members of a sib pair,  $\beta_0$  is the intercept, and  $\beta_1$ , the slope, is the locus-specific genetic variance of the trait attenuated by recombination between the trait and marker loci. To detect *trans*-acting effects at other genomic regions conditional on the *cis*-acting effect at the structural gene loci (22q11 and 9q34 for *COMT* and *DBH*, respectively), we performed the multiple HE regression

$$Y = \beta_0 + \beta_1 \hat{\pi}_0 + \lambda \hat{\pi}_1,$$

where the subscript 0 denotes a marker for the structural gene locus and the subscript 1 denotes another locus in the genome. The coefficient  $\lambda$  denotes the (attenuated) genetic variance at locus 1 and a value significantly different from zero indicates an additive effect of locus 1 with locus 0 on gene expression level. We further investigated genetic interaction between the structural gene locus and the other locus with the model

$$Y = \beta_0 + \beta_1 \hat{\pi}_0 + \beta_2 \hat{\pi}_1 + \gamma \hat{\pi}_0 \hat{\pi}_1,$$

where  $\gamma$  denotes variance (again, attenuated by recombination) due to the genetic interaction between locus 1 and locus 0, and a value significantly different from zero indicates an interactive effect of the two loci on gene expres-

**Table 1: Chromosome regions showing only additive *trans*-acting effects<sup>a</sup> with 22q11 on COMT expression level**

Genomic region	Closest marker	Physical distance (Mb)	p-Value for $\lambda$
3q26	rs1499812	173.2	$4.4 \times 10^{-3}$
4q35	rs1979255	190.7	$3.4 \times 10^{-4}$
5q34	rs1387395	163.8	$4.1 \times 10^{-3}$
6p22	rs1475149	21.8	$2.5 \times 10^{-3}$
7p21	rs988418	9.9	$3.7 \times 10^{-3}$
8q22	rs1075394	101.5	$1.8 \times 10^{-3}$
9q34	rs936249	138.2	$2.2 \times 10^{-3}$
10p11	rs1857063	29.4	$2.0 \times 10^{-7}$
18p11	rs321663	8.6	$2.1 \times 10^{-4}$
20q13	rs756529	47.4	$3.1 \times 10^{-3}$

<sup>a</sup>At a nominal level of 0.005

sion level. Single-point IBD sharing was estimated by the program GENIBD. Both of the above programs are included in the Statistical Analysis for Genetic Epidemiology (S.A.G.E.) software [23].

## Results

### Linkage of COMT expression levels

We detected a linkage signal at chromosome 22q11 ( $p = 6.5 \times 10^{-3}$ ). The closest SNP to the peak of the signal is rs738842, which is  $\sim 3$  megabases (Mb) centrometric to the *COMT* gene and within the *cis*-acting region defined by Morley et al. [15]. There were 10 regions showing only additive *trans*-acting effects (Table 1) and 7 regions showing only interactive *trans*-acting effects (Table 2) at the nominal significance level of 0.005; at 14 other regions, both effects attained a nominal significance level of 0.05 and at least one attained the level 0.005 (Table 3). These nominal significance levels were arbitrarily chosen in order to report the most notable results. Note that 1) chromosome 9q34, where *DBH* is located, showed an additive *trans*-acting effect with 22q11 on *COMT* expression level; and 2) chromosome 20q13, which was one of the two hot spots of transcriptional regulation detected by Morley et al. [15], was also detected to have an additive effect with 22q11 on the *COMT* expression level.

### Linkage of DBH expression levels

There was no linkage signal for the *DBH* expression level at either chromosome 9q34 or chromosome 19p13 when analyzing all of the 14 pedigrees together. There could be many reasons for the lack of any linkage signal at chromosome 9q34, the structural gene location (see Discussion). However, upon analyzing the pedigrees individually, it was noticed that three pedigrees (1340, 1345 and 1318), which included 70 full sibling pairs, showed a positive linkage signal at 9q34. Because our primary interest was to test whether there is a second locus near chromosome 19p13 related to *DBH*, in addition to the *DBH* gene, we selected these three pedigrees for further study and found that a linkage signal at 19p13 then also emerged, showing both additive and interactive *trans*-acting effects ( $p = 2.4 \times 10^{-2}$  and  $p = 2.6 \times 10^{-2}$ , respectively) with the locus at chromosome 9q34 on *DBH* expression level (Fig. 1).

## Discussion

Phenotypic variation in the transcript level of a gene involves genetic components; however, the transcriptional regulatory elements may or may not lie in the structural gene itself. Studies by Morley et al. [15] have suggested the existence of master regulatory regions, and we identified chromosome 20q13, nominated by them, as regulating *COMT* expression level. Note that Morley et al. [15] performed simple HE regression, while we performed multiple HE regression so as to condition on the

**Table 2: Chromosome regions showing only interactive *trans*-acting effects<sup>a</sup> with 22q11 on COMT expression level**

Genomic region	Closest marker	Physical distance (Mb)	p-Value for $\gamma$
1q32	rs871446	196.9	$1.9 \times 10^{-3}$
3p26	rs1385466	4.0	$1.2 \times 10^{-5}$
6q23	rs2040034	137.3	$3.9 \times 10^{-4}$
9q33	rs1326808	116.7	$2.1 \times 10^{-3}$
10q22	rs1907308	78.0	$8.3 \times 10^{-6}$
15q26	rs288423	96.0	$3.3 \times 10^{-3}$
18q21	rs1943985	45.5	$1.5 \times 10^{-4}$

<sup>a</sup>At a nominal level of 0.005

**Table 3: Chromosome regions showing both additive and interactive trans-acting effects<sup>a</sup> with 22q11 on COMT expression level**


Genomic region	Closest marker	Physical distance (Mb)	p-Value	
			$\lambda$	$\gamma$
1q41	rs1561189	213.5	$1.1 \times 10^{-4}$	$4.6 \times 10^{-3}$
2q35	rs1425118	216.8	$<1.0 \times 10^{-8}$	$3.5 \times 10^{-2}$
3q28	rs1559018	191.8	$2.3 \times 10^{-2}$	$2.9 \times 10^{-5}$
4q31	rs795995	141.1	$1.5 \times 10^{-3}$	$5.0 \times 10^{-7}$
5q14	rs1020720	80.5	$1.7 \times 10^{-2}$	$1.8 \times 10^{-3}$
6q16	rs2021678	102.1	$1.1 \times 10^{-3}$	$4.7 \times 10^{-2}$
7p14	rs2051936	40.8	$5.7 \times 10^{-3}$	$3.5 \times 10^{-3}$
8p23	rs433960	12.9	$4.7 \times 10^{-2}$	$2.9 \times 10^{-5}$
9p23	rs1412307	13.0	$1.3 \times 10^{-4}$	$6.6 \times 10^{-3}$
14q13	rs1983667	36.2	$1.4 \times 10^{-4}$	$1.1 \times 10^{-2}$
16p13	rs1981492	0.3	$1.0 \times 10^{-3}$	$5.9 \times 10^{-3}$
16q24	rs880275	84.8	$4.7 \times 10^{-3}$	$2.6 \times 10^{-3}$
17p12	rs433068	15.2	$4.8 \times 10^{-2}$	$4.4 \times 10^{-3}$
22q13	rs960362	47.4	$4.8 \times 10^{-4}$	$8.6 \times 10^{-3}$

<sup>a</sup>Both effects attained a nominal significance level of 0.05 and at least one reached the level 0.005.

IBD sharing at *COMT*, i.e., our analysis refined their original investigation and provided further evidence supporting chromosome 20q13 as a regulatory region. In the current data, overall there was a linkage signal at 22q11 for *COMT* expression level, but no signal at 9q34 for *DBH* expression level. Lack of an overall linkage signal at 9q34 for *DBH* could be due to there not being enough *DBH* polymorphism segregation in the sample, to there being less expression variation among individuals than between replicates within an individuals [15], from low levels of transcript abundance in view of its tissue-specific expres-

sion pattern (Dr. Joseph F. Cubells, personal communication), or to genetic heterogeneity, as also indicated in the earlier study of plasma *DBH* activity [16].

In this study we employed multiple HE regression to detect additive and interactive effects between two loci. In the three pedigrees selected for having a positive linkage signal at 9q34 for *DBH* expression level, we detected a boost of signal at chromosome 19p13, where previous studies [16-18] indicated the existence of a locus influencing *DBH* plasma activity. The multiple HE regression indi-



**Figure 1**  
Linkage signals of *DBH* expression levels at chromosomes 9q34 and 19p13.

cated both additive and interactive effects of the two loci, suggesting epistasis between them. For simplicity, and because of the limited amount of data available, we did not dissect the total genetic variance into additive and dominance components in this study; with more data the interaction could be characterized by more elaborate models [24], and even embedded in a general pedigree framework [25,26]. We also noticed that chromosome 9q34, where *DBH* is located, showed an additive *trans*-acting effect with 22q11 on *COMT* expression level. This is not totally unexpected because both enzymes are involved in catecholamine metabolism; however, we could not find in the literature any genetic study on the interaction between them, which is surprising. Interaction between *COMT* and *DBH* could be a potential research topic.

In summary, we performed quantitative linkage analysis for both *COMT* and *DBH* expression phenotypes, identifying transcriptional regulatory regions for both genes. Multiple HE regression was used to detect additive and interactive effects between two loci. The master transcriptional regulatory region 20q13 had an additive effect on the *COMT* expression level. Chromosome regions 9q34 and 19p13 acted epistatically on *DBH* expression level. Potential interaction between *COMT* and *DBH* was indicated.

### Competing interests

The author(s) declare that they have no competing interests.

### Acknowledgements

Some of the results in this paper were obtained by using the program package S.A.G.E., which is supported by a U.S. Public Health Service Resource Grant (RR03655) from the National Center for Research Resources. This work was supported in part by research grants GM-28356 from the National Institute of General Medical Sciences, Cancer Center Support Grant P30CAD43703 from the National Cancer Institute, and a Training grant (HL07567) from the National Heart, Lung and Blood Institute. CX was supported by a Merck Foundation fellowship. CX and QL were sponsored to attend GAW15 by the Endowment Sponsored Mentorship Program of Case Western Reserve University Graduate School.

This article has been published as part of *BMC Proceedings* Volume 1 Supplement 1, 2007: Genetic Analysis Workshop 15: Gene Expression Analysis and Approaches to Detecting Multiple Functional Loci. The full contents of the supplement are available online at <http://www.biomedcentral.com/1753-6561/1?issue=S1>.

### References

- Axelrod J, Tomchick R: **Enzymatic O-methylation of epinephrine and other catechols.** *J Biol Chem* 1958, **233**:702-705.
- Kaufman S, Friedman S: **Dopamine-beta-hydroxylase.** *Pharmacol Rev* 1965, **17**:71-100.
- Gogos JA, Gerber DJ: **Schizophrenia susceptibility genes: emergence of positional candidates and future directions.** *Trends Pharmacol Sci* 2006, **27**:226-233.
- Cubells JF, Zabetian CP: **Human genetics of plasma dopamine beta hydroxylase activity: application to research in psychiatry and neurology.** *Psychopharmacology (Berl)* 2004, **174**:463-476.
- Tenhunen J, Salminen M, Jalanko A, Ukkonen S, Ulmanen I: **Structure of the rat catechol-O-methyltransferase gene: separate promoters are used to produce mRNAs for soluble and membrane-bound forms of the enzyme.** *DNA Cell Biol* 1993, **12**:253-263.
- Kemper CM, O'Connor DT, Westlund KN: **Immunocytochemical localization of dopamine-beta-hydroxylase in neurons of the human brain stem.** *Neuroscience* 1987, **23**:981-989.
- Grossman MH, Emanuel BS, Budarf ML: **Chromosomal mapping of the human catechol-O-methyltransferase gene to 22q11.1-q11.2.** *Genomics* 1992, **12**:822-825.
- Craig SP, Buckle VJ, Lamouroux A, Mallet J, Craig IW: **Localization of the human dopamine beta hydroxylase (DBH) gene to chromosome 9q34.** *Cytogenet Cell Genet* 1988, **48**:48-50.
- Lotta T, Vidgren J, Tilgmann C, Ulmanen I, Melen K, Julkunen I, Taskinen J: **Kinetics of human soluble and membrane-bound catechol-O-methyltransferase: a revised mechanism and description of the thermolabile variant of the enzyme.** *Biochemistry* 1995, **34**:4202-4210.
- Zabetian CP, Anderson GM, Buxbaum SG, Elston RC, Ichinose H, Nagatsu T, Kim KS, Kim CH, Malison RT, Gelernter J, Cubells JF: **A quantitative-trait analysis of human plasma-dopamine beta-hydroxylase activity: evidence for a major functional polymorphism at the DBH locus.** *Am J Hum Genet* 2001, **68**:515-522.
- Bray NJ, Buckland PR, Williams NM, Williams HJ, Norton N, Owen MJ, O'Donovan MC: **A haplotype implicated in schizophrenia susceptibility is associated with reduced COMT expression in human brain.** *Am J Hum Genet* 2003, **73**:152-161.
- Chen J, Lipska BK, Halim N, Ma QD, Matsumoto M, Melhem S, Kolachana BS, Hyde TM, Herman MM, Apud J, Egan MF, Kleinman JE, Weinberger DR: **Functional analysis of genetic variation in catechol-O-methyltransferase (COMT): effects on mRNA, protein, and enzyme activity in postmortem human brain.** *Am J Hum Genet* 2004, **75**:807-821.
- Yan H, Yuan W, Velculescu VE, Vogelstein B, Kinzler KW: **Allelic variation in human gene expression.** *Science* 2002, **297**:1143.
- Cheung VG, Conlin LK, Weber TM, Arcaro M, Jen KY, Morley M, Spielman RS: **Natural variation in human gene expression assessed in lymphoblastoid cells.** *Nat Genet* 2003, **33**:422-425.
- Morley M, Molony CM, Weber TM, Devlin JL, Ewens KG, Spielman RS, Cheung VG: **Genetic analysis of genome-wide variation in human gene expression.** *Nature* 2004, **430**:743-747.
- Wilson AF, Elston RC, Siervogel RM, Tran LD: **Linkage of a gene regulating dopamine-beta-hydroxylase activity and the ABO blood group locus.** *Am J Hum Genet* 1988, **42**:160-166.
- Wilson AF, Elston RC, Sellers TA, Bailey-Wilson JE, Gersting JM, Deen DK, Sorant AJM, Tran LD, Amos CI, Siervogel RM: **Stepwise oligogenic segregation and linkage analysis illustrated with dopamine-beta-hydroxylase activity.** *Am J Med Genet* 1990, **35**:425-432.
- Province M: **Sequential methods of analysis for genome scans.** *Adv Genet* 2001, **42**:499-514.
- Matisse TC, Sachidanandam R, Clark AG, Kruglyak L, Wijsman E, Kakol J, Buyske S, Chui B, Cohen P, de Toma C, Ehm M, Glanowski S, He C, Heil J, Markianos K, McMullen I, Pericak-Vance MA, Silbergleit A, Stein L, Wagner M, Wilson AF, Winick JD, Winn-Deen ES, Yamashiro CT, Cann HM, Lai E, Holden AL: **A 3.9-centimorgan-resolution human single-nucleotide polymorphism linkage map and screening set.** *Am J Hum Genet* 2003, **73**:271-284.
- Haseman JK, Elston RC: **The investigation of linkage between a quantitative trait and a marker locus.** *Behav Genet* 1972, **2**:3-19.
- Shete S, Jacobs KB, Elston RC: **Adding further power to the Haseman and Elston method for detecting linkage in larger sibships: weighting sums and differences.** *Hum Hered* 2003, **55**:79-85.
- Liang K-Y, Zeger SL: **Longitudinal data analysis using generalized linear models.** *Biometrika* 1986, **73**:13-22.
- S.A.G.E. **Statistical analysis for genetic epidemiology. Version 5.2** [<http://darwin.case.edu/sage>]
- Tiwari HK, Elston RC: **Linkage of multilocus components of variance to polymorphic markers.** *Ann Hum Genet* 1997, **61**:253-261.
- Wang T, Elston RC: **Two-level Haseman-Elston regression for general pedigree data analysis.** *Genet Epidemiol* 2005, **29**:12-22.
- Almasy L, Blangero J: **Multipoint quantitative-trait linkage analysis in general pedigrees.** *Am J Hum Genet* 1998, **62**:1198-1211.