

Oral presentation

Open Access

Disseminated and sustained HIV-infection in CD34+ cord blood cell transplanted Rag2^{-/-}gc^{-/-} mice

Stefan Baenziger¹, Roxane Tussiwand², Erika Schlaepfer¹, Luca Mazzucchelli³, Mathias Heikenwalder¹, Michael O Kurrer¹, Silvia Behnke¹, Joachim Frey⁴, Annette Oxenius⁵, Helen Joller¹, Adriano Aguzzi¹, Markus G Manz² and Roberto F Speck^{*1}

Address: ¹University Hospital of Zurich, Zurich, Switzerland, ²Institute for Research in Biomedicine, Bellinzona, Switzerland, ³Institute for Pathology, Locarno, Switzerland, ⁴University of Berne, Berne, Switzerland and ⁵Swiss Federal Institute of Technology, Zurich, Switzerland

* Corresponding author

from 2006 International Meeting of The Institute of Human Virology
Baltimore, USA. 17–21 November, 2006

Published: 21 December 2006

Retrovirology 2006, **3**(Suppl 1):S31 doi:10.1186/1742-4690-3-S1-S31

© 2006 Baenziger et al; licensee BioMed Central Ltd.

Background

We evaluated a novel mouse model that is based on the transplantation of human cord blood CD34+ cells into immunodeficient Rag2^{-/-}gc^{-/-} newborn mice for studying HIV.

Results

Reconstituted mice (CD45+ cell engraftment: 29 ± 18%) were permissive to HIV with up to 2 × 10⁶ copies/ml 2–6 weeks after infection. Thereafter, viremia stabilized at lower levels for up to 4 months. A marked CD4+ cell depletion occurred in all mice infected with X4- strains simultaneously with the initial rise of plasma HIV RNA while this was not the case in R5-infected mice. Spleens and lymph nodes of mice infected with either R5- or X4- strains contained p24+ cells. In thymi, however, p24+ cells were detected rather exclusively following infection with X4- strain, consistent with the expression of CXCR4 but not CCR5 on human CD4+ thymocytes. Similarly as in humans, HIV-infected macrophages were only occasionally found.

Conclusion

Rag2^{-/-}gc^{-/-} mice transplanted with human CD34+ cells develop long-term, high-titer, and lymphoid organ disseminated infection irrespective of co-receptor selectivity

of HIV strain, closely resembling HIV infection in man. In particular, by using HIV strains with distinct co-receptor selectivity, we clearly illustrate the higher cytopathic potential of X4- strains as compared to R5- strains. This straightforward to generate and cost-effective in vivo model should be valuable to study virus-induced pathology, and to rapidly evaluate new approaches aiming to prevent or treat HIV infection.