

A Mobile PTS2 Receptor for Peroxisomal Protein Import in *Pichia pastoris*

Ype Elgersma,[‡] Minetta Elgersma-Hooisma,[‡] Thibaut Wenzel,[‡] J. Michael McCaffery,*
Marilyn G. Farquhar,* and Suresh Subramani[‡]

*Division of Cellular and Molecular Medicine and [‡]Department of Biology, University of California at San Diego, La Jolla, California 92093-0322

Abstract. Using a new screening procedure for the isolation of peroxisomal import mutants in *Pichia pastoris*, we have isolated a mutant (*pex7*) that is specifically disturbed in the peroxisomal import of proteins containing a peroxisomal targeting signal type II (PTS2). Like its *Saccharomyces cerevisiae* homologue, PpPex7p interacted with the PTS2 in the two-hybrid system, suggesting that Pex7p functions as a receptor. The *pex7Δ* mutant was not impaired for growth on methanol, indicating that there are no PTS2-containing enzymes involved in peroxisomal methanol metabolism. In contrast, *pex7Δ* cells failed to grow on oleate, but growth on oleate could be partially restored by expressing thiolase (a PTS2-containing enzyme) fused to the PTS1.

Because the subcellular location and mechanism of action of this protein are controversial, we used various methods to demonstrate that Pex7p is both cytosolic and intraperoxisomal. This suggests that Pex7p functions as a mobile receptor, shuttling PTS2-containing proteins from the cytosol to the peroxisomes. In addition, we used PpPex7p as a model protein to understand the effect of the Pex7p mutations found in human patients with rhizomelic chondrodysplasia punctata. The corresponding PpPex7p mutant proteins were stably expressed in *P. pastoris*, but they failed to complement the *pex7Δ* mutant and were impaired in binding to the PTS2 sequence.

THE sorting of proteins to distinct subcellular compartments is achieved by the coordinated action of organelle-specific targeting signals and receptors. Studies on protein targeting across biological membranes have uncovered two general paradigms for the action of targeting signal receptors. In certain cases, such as the SecA/B-dependent secretion of proteins across the bacterial inner membrane and the signal recognition particle (SRP)¹-dependent insertion of proteins across ER membranes or transport across the nuclear pore, the targeting

signal is recognized by a mobile receptor that typically recognizes the signal in the cytosol and then shuttles the cargo to the target organelle (for review see Schatz and Dobberstein, 1996; Görlich and Mattaj, 1996). These mobile receptors cycle either between the cytosol and the organelle membrane, as seen for SecA/B and SRP, or they cycle into and out of the organelle, as observed for importin and transportin. In other instances, such as mitochondrial (Schatz and Dobberstein, 1996) and chloroplast import (Schnell, 1995), the receptors are located on the target membrane (membrane-associated receptors), where they bind the targeting signal but their mobility is believed to be limited to the plane of the membrane.

Peroxisomal matrix proteins are synthesized on free polyribosomes and are posttranslationally imported into the peroxisome (for review see Lazarow and Fujiki, 1985). Sorting to peroxisomes requires the presence of a peroxisomal targeting signal (PTS). Most peroxisomal matrix proteins contain a PTS1 sequence that consists of the COOH-terminal tripeptide SKL or other variants (Gould et al., 1987, 1989).

A second peroxisomal targeting signal (PTS2) was first identified in the NH₂ terminus of rat peroxisomal thiolase (Osumi et al., 1991; Swinkels et al., 1991) and has been identified in several other proteins since (for review see Elgersma and Tabak, 1996). Alignments and site-directed mutagenesis of these proteins suggest a consensus PTS2

Y. Elgersma and M. Elgersma-Hooisma contributed equally to this work.

Address all correspondence to S. Subramani, Department of Biology, University of California, San Diego, Room 3230, Bonner Hall, 9500 Gilman Drive, La Jolla, CA 92093-0322. Tel.: (619) 534-2327. Fax: (619) 534-0053. E-mail: ssubramani@ucsd.edu

Y. Elgersma's and M. Elgersma-Hooisma's present addresses are Cold Spring Harbor Laboratory, 1 Bung Town Road, Cold Spring Harbor, NY 11724.

T. Wenzel's present address is Gist Brocades, Food Specialty Division, Delft, The Netherlands.

1. *Abbreviations used in this paper:* Aco1p, acyl CoA oxidase; AD, activation domain; BLE, bleomycin resistance protein; DB, DNA-binding domain; FOX3, 3-ketoacyl CoA thiolase; GFP, green fluorescent protein; Pex, peroxin; *pex*, peroxisomal biogenesis/import mutants; Pp, *Pichia pastoris*; PTS, peroxisomal targeting signal; RCDP, rhizomelic chondrodysplasia punctata; Sc, *Saccharomyces cerevisiae*; SRP, signal recognition particle; TPR, tetratricopeptide repeat.

sequence of R/K-L/V/I-X₅-H/Q-L/A (Gietl et al., 1994; Glover et al., 1994a; Tsukamoto et al., 1994).

Additional evidence for at least two pathways for peroxisomal protein import was provided by cloning of the PTS1 and the PTS2 receptors. The PTS1 receptor, Pex5p, has seven tetratricopeptide repeats (TPRs) in the COOH-terminal half of the protein, but surprisingly, it lacks a hydrophobic domain that could anchor the protein in the peroxisomal membrane (McCullum et al., 1993; Van der Leij et al., 1993; Dodt et al., 1995; Fransen et al., 1995; Nuttley et al., 1995; Szilard et al., 1995; Van der Klei et al., 1995; Wiemer et al., 1995). There has been much debate about the localization of Pex5p, but current evidence suggests that in most organisms, at least a fraction of this protein is cytosolic (Dodt et al., 1995; Van der Klei et al., 1995; Wiemer et al., 1995; Elgersma et al., 1996a; Gould et al., 1996), while the rest is peroxisome associated. This led to the proposal that Pex5p functions as a mobile receptor, shuttling between the cytosol and the peroxisome. This model is further supported by the characterization of the SH3 domain-containing peroxisomal membrane protein, Pex13p, and another protein, Pex14p, which serve as docking proteins for Pex5p (Elgersma et al., 1996a; Erdmann and Blobel, 1996; Gould et al., 1996; Albertini et al., 1997). Furthermore, kinetic evidence for Pex5p as a mobile receptor has been uncovered recently (Dodt and Gould, 1996).

The *Saccharomyces cerevisiae* (*Sc*) *pex7* mutant is characterized by the mislocalization of thiolase, whereas import of other proteins analyzed is unaffected. A phenotype analogous to that of the *pex7* mutant has also been described for a human patient cell line (Motley et al., 1994; Slawewski et al., 1995). Cloning of the yeast, and more recently the human, *PEX7* genes led to the identification of the PTS2 receptor (Pex7p; Marzioch et al., 1994; Zhang and Lazarow, 1995; Braverman et al., 1997; Motley et al., 1997; Purdue et al., 1997). Pex7p belongs to the WD-40 repeat (β -transducin) family, which is characterized by the presence of a 43-amino acid repeat domain. Like TPRs, the WD repeats are probably involved in multiple protein-protein interactions (Komachi et al., 1994). Interestingly, a functional relationship between proteins belonging to the TPR and the β -transducin (WD-40) families has been described (for review see Goebel and Yanagida, 1991; Van der Voorn and Ploegh, 1992). Although it is not yet clear whether Pex5p and Pex7p interact directly, evidence exists that they share a common import machinery. In humans, it has been shown that a certain cell line that is defective in the *PEX5* gene is not only deficient in the import of PTS1-containing proteins, but is also deficient in the import of PTS2-containing proteins (Dodt et al., 1995; Wiemer et al., 1995). Moreover, a peroxin (Pex) has been identified (ScPex14p) that is required for both import pathways and which interacts with both Pex5p and Pex7p (Albertini et al., 1997).

The location of Pex7p is not clear. An HA-tagged Pex7p (Pex7p-HA3) was found to be entirely intraperoxisomal in *S. cerevisiae*, whereas a Myc-tagged (overexpressed) ScPex7p (Myc-Pex7p) was predominantly cytosolic (Marzioch et al., 1994; Zhang and Lazarow, 1995). Human Myc-Pex7p expressed from the strong cytomegalovirus (CMV) promoter was found predominantly in the cytosol, and no peroxisome-associated Pex7p was detected (Braverman et

al., 1997). Because antibodies suitable for studying endogenous Pex7p are lacking, it is not clear which of these tagged or overexpressed proteins reflects its real location.

The model for the action of Pex7p as a receptor has remained elusive because of uncertainties regarding the precise subcellular location of this protein. The cytosolic localization of epitope-tagged yeast or human Pex7p has been cited as evidence for a mobile receptor, shuttling between the cytosol and the peroxisome (Marzioch et al., 1994; Rehling et al., 1996; Braverman et al., 1997). In contrast, the presence of an intraperoxisomal receptor that acts from within the organelle to pull proteins in would be novel in comparison with the existing paradigms for targeting signal receptors described earlier (Zhang and Lazarow, 1995, 1996).

The finding that the PTS2 import pathway may not be induced in methanol-grown *Hansenula polymorpha* (Faber et al., 1994) suggests that this pathway may not be required for growth on methanol. This may explain why despite elaborate screening procedures, mutants deficient in the PTS2 import pathway have not yet been described in methylotrophic yeasts, since the primary screening was performed by selecting mutants that were unable to grow on methanol (Cregg et al., 1990; Gould et al., 1992; Liu et al., 1992). To obtain more insight into the mechanism of import of PTS2-containing proteins in *Pichia pastoris*, we have used a new screening procedure that is more selective for the PTS2 import pathway. We have isolated the *P. pastoris* (*Pp*) *PEX7* gene, studied the function and localization of endogenous PpPex7p, and examined the use of the PTS2-dependent import pathway upon growth of *P. pastoris* on different carbon sources. Furthermore, because recent studies have identified several mutations in human Pex7p that are responsible for the disease rhizomelic chondrodysplasia punctata (RCDP; Braverman et al., 1997; Motley et al., 1997; Purdue et al., 1997), we used the corresponding mutations in PpPex7p to provide insights regarding why the mutant proteins were nonfunctional.

Materials and Methods

Strains and Media

P. pastoris strains used in this study were as follows: PPY12 (*arg4, his4*; parental wild-type strain); *pex7.1* (*his4, pex7, ARG4::pTW84* (PTS2-GFP (S65T)/*GAPDH*; PTS2-BLE/*GAPDH*)); *pex7* Δ (*his4, pex7::PpARG4*); and *fox3* Δ (*his4, fox3::PpARG4*). *S. cerevisiae* strains used in this study were as follows: BJ1991 (*MAT* α , *leu2, trp1, ura3-251, prb1-1122, pep4-3*) (wild-type strain); *pex7* (*pex7, ura3-251, leu2, his3*); and L40 (*MAT* α , *his3* Δ 200, *trp1-901, leu2-3,112, ade2, LYS2::(lexAop)₄-HIS3, URA3::(lexAop)₈-lacZ* (for two-hybrid assays). *Escherichia coli* strains used in this study were as follows: DH5 α (*recA, hsdR, supE, endA, gyrA96, thi-1, relA1, lacZ*; for cloning procedures), and SG13009 (*nal^r, str^r, rif^r, lac⁻, ara⁻, gal⁻, ml⁻, F⁻, recA⁻*; for protein expression).

Minimal methanol contained the following: 0.25% (NH₄)₂SO₄, 0.02% MgSO₄, 0.39% NaH₂PO₄, 0.05% yeast extract, 0.1% (vol/vol) Vishniac minerals mix, 0.1% vitamins, 20 mg/ml amino acids as needed, and 0.5% methanol as carbon source. Minimal oleate medium contained 0.2% oleate and 0.02% Tween 40 as carbon source. Minimal oleate plates were made according to Elgersma et al. (1993). Other media used in this study were described by Wiemer et al. (1996).

Isolation of *pex* Mutants

A culture of *P. pastoris* PPY12+pTW84 (PTS2-GFP(S65T)/*GAPDH*,

PTS2-BLE/*GAPDH*) cells was grown for 15 h on rich glucose medium (YPD), diluted 12-fold in 600 ml YPD, and grown for an additional 8 h. Cells were pelleted and washed twice with water and once with 100 mM sodium citrate, pH 5.5. The cells were then resuspended in 150 ml 100 mM Na-citrate, pH 5.5, and 25-ml batches were treated with increasing concentrations (0–2%) *N*-methyl-*N*-nitro-nitrosoguanidine (NTG) for 30 min. Mutagenized cells were washed three times with 100 mM sodium citrate, pH 5.5, resuspended in 20 ml YPD, and recovered for 2 h. Small aliquots were used to determine the survival rate, and the remainder was frozen in glycerol at –80°C. Cells with an NTG survival of 4% were used to inoculate 50 ml YPD and were grown for 24 h. These cells were pelleted by centrifugation at 2,500 g, washed with water, and transferred to 300 ml rich oleate medium (YPO) and induced for 15 h. These cells were collected, resuspended in 250 ml YPO, aliquoted in five portions of 50 ml, and induced in oleate medium for an additional 2 h. Increasing amounts of phleomycin were added to these cultures (to concentrations of 0–100 µg/ml), and the cells were incubated for 3 h at 30°C. The phleomycin-treated cells were washed twice with water and resuspended in YPD to recover for 2 h at 30°C. Cells were diluted and plated on YPD to determine the survival rate. Colonies obtained from the batch showing a 5% survival rate were tested for growth on minimal oleate and methanol medium, and they were analyzed for PTS2–green fluorescent protein (GFP) import by fluorescence microscopy.

Cloning of the *PpPEX7* and *PpPAT1* Genes

A genomic library was transformed into *pex7.1* cells. Three different plasmids (pM1, pM2, and pM3), which restored growth on oleate, were isolated. Subclone analysis revealed that pM1 contained ~1 kb of the 5' end of a gene (*PpPAT1*) that is able to suppress the *pex7.1* phenotype. pM2 and pM3 had an overlapping fragment of 2.8 kb. This 2.8-kb insert contained the *PpPEX7* gene, which was sequenced on both strands. The *PpPEX7* sequence data have been submitted to the GenBank database under accession number AF021797.

Construction of Plasmids Expressing *PpPex7p*

A 2.6-kb genomic fragment containing the *PEX7* gene was cloned in a three-fragment ligation, as a *S*all–*E*coRI and *E*coRI–*B*amHI fragment in a *S*all/*B*amHI–digested pUC18, resulting in pM19. The *PEX7* gene knockout was made by cloning the *PpARG4* gene as a blunt fragment in the blunted *N*siI sites (86 bp upstream of the ATG and 28 bp downstream of the stop codon of *PEX7*) of pM19. The polylinker of pUC18 was modified by ligating adaptors P43 and P44 (see Table I) in the *E*coRI and *B*amHI sites so that these sites were destroyed and the *B*glII, *K*pnlI, *E*coRV, *N*siI, and *X*baI sites were introduced (pM24). The *PEX7* gene was amplified by PCR using primers P33 and P34, thereby introducing a *K*pnlI site immedi-

ately upstream of the ATG, and was cloned as a *K*pnlI–*N*siI fragment in pM24. To eliminate PCR errors, the *B*amHI–*N*siI fragment was replaced by the *B*amHI–*N*siI fragment of the genomic clone, and the remainder of the *PEX7* gene was sequenced. The *PEX7* gene on this plasmid (pM27) was used for further manipulations.

For NH tagging, a linker encoding the NH tag (primers P26/P28) was ligated in the *E*coRI site of pUC18, resulting in pEL210. *PEX7* was cloned as a *K*pnlI–*N*siI fragment from pM27 in the *K*pnlI/*P*stI sites of pEL210, resulting in pM22. The NH-tagged *PEX7* was cloned downstream of the *ACO1* promoter in the *B*glII/*E*coRI-digested pTW72 (a *p*HILD2-based plasmid in which the alcohol oxidase promoter was replaced by the *ACO1* promoter), using the *B*glII/*H*indIII fragment from pM22 and a *E*coRI/*H*indIII adaptor (P49/P50), resulting in plasmid pM39. The nontagged gene was cloned downstream of the *ACO1* promoter by replacing the *B*glIII/*A*vrII fragment from pM39 (encoding *NH-PEX7*) by the *B*glIII/*X*baI fragment from pM27 (encoding *PEX7*), resulting in pM70. The *ACO1* promoter was replaced by the endogenous *PEX7* promoter by replacing the *N*deI/*B*amHI fragment from pM39, containing the *ACO1* promoter, with the genomic *X*hoI/*B*amHI fragment and the *N*deI/*X*hoI adapter P122/P123, resulting in pM78.

To introduce the C347ter mutation, we ligated linker P124/P125 in an *E*coRI/*S*allI-digested pM27, resulting in pM75, and sequenced the inserted linker. To introduce the G249V and A248R point mutations in *PEX7*, we designed primers that introduced the desired mutation and a unique restriction site at the location of the mutation. The 5' part of the gene was amplified by PCR with primers P32 (internal primer just upstream of the *B*amHI site) and P128 (G249V) or P126 (A248R). The 3' part of the gene was amplified by PCR with primers P35 (a primer just downstream of *PEX7*) and P129 (G249V) or P127 (A248R). The PCR fragments corresponding to the 5' segments of *PEX7* were digested with *B*amHI and *P*stI (G249V) or *B*amHI and *X*baI (A248R), and the PCR fragments corresponding to the 3' segments of *PEX7* were digested with *P*stI and *E*coRI (G249V) or *X*baI and *E*coRI (A248R). These fragments were ligated in a three-fragment ligation into the *B*amHI/*E*coRI-digested pM27, resulting in pM76 (G249V) and pM77 (A248R). The *B*amHI/*E*coRI inserts were sequenced. The mutated *PEX7* genes were cloned as *B*amHI/*S*allI fragments, downstream of the *ACO1* promoter in a *B*amHI/*X*hoI-digested pM70, resulting in pM85 (C347ter), pM86 (G249V), and pM87 (A248R). The mutated *PEX7* genes were cloned as *B*amHI/*S*allI fragments downstream of the *PEX7* promoter in a *B*amHI/*X*hoI-digested pM78, resulting in pM82 (C347ter), pM83 (A248R), and pM84 (G249V).

PpFox3-SKL was amplified by PCR using primers based on the sequence of the *PpFOX3* gene so that a *B*amHI site was introduced directly upstream of the ATG, and an SKL sequence and an *X*baI site were added at the 3' end of the gene. The PCR product was digested with *B*amHI and *X*baI and was cloned downstream of the *ACO1* promoter between the *B*glIII/*S*peI sites of pM39.

For the yeast two-hybrid analysis, plasmids were generated that expressed appropriate protein fusions to the DNA binding (DB) domain of LexA, and the activation domain (AD) of VP16. The LexA(DB)–*ScFox3p* (DB–*ScFox3p*) fusion protein was made by cloning *ScFOX3* as a *B*amHI/*X*baI fragment in pBTM116B, resulting in pM35. pBTM116 (Bartel et al., 1993) was modified by introducing a polylinker in all three open reading frames, resulting in pBTM116A, B, and C. The DB–*ScPTS2* fusion protein was made by digesting pM35 with *N*coI/*S*allI, followed by a filling-in reaction using Klenow polymerase and religation, thereby deleting *ScFOX3* downstream of the *PTS2* (pM52). The DB–*ScFox3p*Δ*PTS2* fusion protein was made by digesting pM35 with *Not*I/*N*coI, followed by a filling-in reaction using Klenow polymerase and religation, thereby deleting the *PTS2* signal of *ScFOX3* (pM51). The AD–*PpPex7p* fusion protein was made by cloning *PpPEX7* as a *B*glIII/*S*allI fragment from pM27 in the *B*amHI/*S*allI-digested pVP16C, resulting in pM34. pVP16 (Hollenberg et al., 1995) was modified by introducing a polylinker in all three open reading frames, resulting in pVP16A, B, and C. The RCDP mutations were introduced in the two-hybrid vector as *B*amHI/*S*allI fragments into a *B*amHI/*S*allI-digested pM34, resulting in pM79 (*Pex7p*(G249V)), pM80 (*Pex7p*(A248R)), and pM81 (*Pex7p*(C347ter)).

Raising Antibodies against *Pex7p*

To raise antibodies against *Pex7p*, we cloned the DNA encoding amino acids 1–143 of *Pex7p* as a *K*pnlI/*H*paI fragment from pM27 into pQE30 (QIAGEN Inc., Chatsworth, CA), which was first digested with *H*indIII, blunted with Klenow polymerase, and subsequently digested with *K*pnlI. The (His)₆-tagged protein was expressed in *E. coli* SG13009 and purified

Table I. Oligonucleotides Used in This Study

Primer	Sequence
P26	5'-AATTCGAGCTCATGCAAGACCTTCCAGGAAATGACA-ACAGCACAGCAGGTGGTACCC-3'
P28	5'-AATTGGGTACCACCTGCTGTGCTGTTGTCAATTCCT-GGAAGGTCTTGCATGAGGTCG-3'
P32	5'-CTGTTTTGTTACAGCATCATGGG-3'
P33	5'-AAGCTTGGAGCCACTATCGACTACGCG-3'
P34	5'-GGTACCATGTTAAGTTCCAAACAAACG-3'
P35	5'-GGCAGGGATGGTACTTAATTTCCC-3'
P43	5'-AATTAGATCTGGTACCGATATCATGCATTCTAGA-3'
P44	5'-GATCTCTAGAATGCATGATATCGGTACCAGATCT-3'
P49	5'-AGCTTACTAGTCTCATTAGGGCCCTCGAG-3'
P50	5'-AATTCTCGAGGGCCCTAGGACTAGTA-3'
P122	5'-TATGTTGCCTTGGTGGAGGG-3'
P123	5'-TCGACCCTCCAAGCAAGGCAACA-3'
P124	5'-AATTCGTAATGGGTTGAG-3'
P125	5'-TCGACTCAACCCATTACG-3'
P126	5'-GATCTCTAGAGGTGTTGACAAGCTCGCAAAGGTC-3'
P127	5'-GATCTCTAGAAGCAACTACAAAAGGTCGATAACG-3'
P128	5'-GCAGTCTGCAGTAGTTGACAAGCTCGCAAAGGTC-TGG-3'
P129	5'-GATCTACTGCAGAAGCAACTACAAAAGGTCGAT-AACG-3'

under denaturing conditions on Ni²⁺-NTA beads according to the manufacturer's manual (QIAGEN Inc.). The protein was further purified by SDS-PAGE, visualized with 0.25 M KCl, 1 mM DTT, and subsequently excised and eluted from the gel in elution buffer (50 mM Tris, pH 8.0, 0.1% SDS, 0.1 mM EDTA, 5 mM DTT, 0.15 M NaCl). This purified protein was used to immunize rabbits.

The antibody was purified using a *pex7Δ* depletion column and, subsequently, a Pex7p affinity column. The depletion column was made by coupling overnight ~20 mg of proteins from a cell-free *pex7Δ* lysate to 0.6 g of cyanogen bromide-activated Sepharose 4B (Sigma Immunochemicals, St. Louis, MO) in coupling buffer (0.1 M NaHCO₃, 0.5 M NaCl). After blocking the column for 2 h with blocking buffer (1 M ethanolamine, 0.1 M NaHCO₃, 0.5 M NaCl, pH 8.0), the column was washed extensively, alternating between coupling buffer and low pH wash buffer (20 mM sodium acetate, 0.5% [vol/vol] acetic acid, 0.5 M NaCl, pH 3.5) and subsequently with PBS, followed by 0.2 M glycine, 1 mM EGTA, pH 2.4, followed by PBS. After overnight incubation with 10 ml of serum, the serum was drained off and collected together with the first 3 ml of PBS wash. This purified serum was used for further purification on an affinity column. This was done essentially in the same manner as the depletion column, except that 2 mg of purified Pex7p was bound to the column, and after binding the serum to the column, it was washed extensively with PBS, low pH wash buffer (0.3% [vol/vol] acetic acid, 50 mM sodium acetate, 1 M NaCl, pH 4.5), high pH wash (0.1 M NaHCO₃, 1 M NaCl, pH 9.0), and again with PBS. The antibodies were eluted with 15 ml of 0.2 M glycine, 1 mM EGTA, pH 2.4, and immediately neutralized to pH 7.0 with 1 M Tris base. Finally, the antibodies were concentrated to ~1 ml using a Centrprep 30 (Amicon, Beverly, MA). The antibody was used at a 1:10,000 dilution for Western blotting.

Subcellular Fractionation and Nycodenz Gradients

Subcellular fractionations of oleate-grown cells were essentially performed as described by Van der Leij et al. (1992), in the presence of protease inhibitors (1 mM PMSF, 0.2 mg/ml NaF, and 2 μg/ml of chymostatin, leupeptin, antipain, and pepstatin). 6 ml of a 1,000 g postnuclear supernatant was layered on a continuous 16–35% Nycodenz gradient (30 ml), with a 4-ml cushion of 42% Nycodenz dissolved in 5 mM MES, pH 6.0, 1 mM EDTA, 1 mM KCl, and 8.5% sucrose. The sealed tubes were centrifuged for 2.5 h in a vertical rotor at 29,000 g at 4°C. Afterwards, fractions were collected and analyzed by SDS-PAGE and Western blotting.

Microscopy Analysis

Fluorescence microscopy for GFP analysis was done as described by Monosov et al. (1996). Fluorescence microscopy of semithin sections and immuno-EM analysis of ultrathin sections was done as follows: *P. pastoris* cells were grown to log phase and fixed in either periodate-lysine-paraformaldehyde (75 mM phosphate buffer containing 2% formaldehyde, 70 mM lysine, and 10 mM sodium periodate, pH 6.2) for 6 h, or in 4% paraformaldehyde contained in 100 mM phosphate buffer, pH 7.4, for 18–24 h at 4°C. The cells were washed, pelleted, and embedded in 1% ultralow temperature gelling agarose, and were trimmed into 1-mm³ blocks. The cells were cryoprotected in 2.3 M sucrose containing 20% polyvinyl pyrrolidone for 1 h, mounted on aluminum cryopins, and frozen in liquid nitrogen. Ultrathin cryosections were then cut on a Reichert Ultracut E microtome (Leica Inc., Heerbrugg, Switzerland) equipped with an FC4 cryostage, and sections were collected onto 300 mesh, Formvar/carbon-coated nickel grids and immunolabeled as follows. Grids were washed with several drops of PBS containing 5% FCS and 0.01 M glycine, pH 7.4. The grids were blocked in 10% FCS and incubated for 16 h with αNH antibodies in PBS containing ~10 μg/ml specific antibody. After washing the grids in PBS, they were incubated for 2 h with a 50× diluted goat anti-rabbit 5-nm gold conjugate (Amersham Life Sciences, Arlington Heights, IL). The grids were washed several times with PBS, subsequently with water, and then the grids were embedded in a mixture containing 2.7% polyvinyl alcohol (mol wt = 10,000), 0.2% methyl cellulose (400 cP), and 0.2% uranyl acetate. The sections were observed and photographed on a 1200EX II TEM microscope (Jeol Ltd., Tokyo, Japan).

Miscellaneous

Digitonin titrations were essentially performed as described by Zhang and Lazarow (1995), with modifications as described in Elgersma et al. (1996a).

For TCA lysates, cells (50 ml of culture OD₆₀₀ = 0.7) were pelleted by

centrifugation, washed, and collected in a 2-ml Eppendorf tube. After the addition of 500 μl of 10% TCA and ~200 μl of glass beads, the cells were broken by vigorous vortexing for 30 min at 4°C. Cell debris was pelleted by centrifugation for 30 s at 6,000 rpm, and the supernatant was further centrifuged for 20 min at 12,000 rpm at 4°C in a microcentrifuge. The pellet was washed once with acetone, dried, and resuspended in Laemmli sample buffer for SDS-PAGE.

Results

Isolation of PTS2 Import-deficient Mutants

The negative screening procedure for isolating *P. pastoris* *pex* mutants that has been used thus far (Gould et al., 1992; Liu et al., 1992) has not resulted in the isolation of mutants that are specifically disturbed in the PTS2 import pathway. Therefore, we developed a positive selection procedure that would enrich for such mutants. In *S. cerevisiae*, such a scheme has been developed, which is based on the ability of peroxisomes to import the bleomycin resistance protein (BLE) fused to the PTS1 (Elgersma et al., 1993). In wild-type cells, BLE-PTS1 is imported into peroxisomes, thereby sequestering the BLE protein from its toxic ligand, phleomycin. Peroxisomal import mutants (*pex*) are unable to import BLE-PTS1 into their peroxisomes, which allows the protein to bind phleomycin, thereby preventing its toxicity. We modified this procedure by fusing the BLE protein to the PTS2. Since a PTS2 sequence of an endogenous *P. pastoris* protein had not been identified, we tested the PTS2 sequence of *S. cerevisiae* thiolase (Fox3p) in *P. pastoris*. We therefore made a fusion of the first 17 amino acids of thiolase (containing the PTS2 signal) to GFP (PTS2-GFP). When expressed in oleate-grown *P. pastoris*, we observed a punctate GFP expression pattern, suggesting that PTS2-GFP was imported into the peroxisomes and that the ScPTS2 is functional in *P. pastoris* (Fig. 1 E). We also observed some cytosolic labeling, however, suggesting that PTS2-GFP import was not as efficient as GFP-SKL import (Fig. 1, E versus C).

A yeast strain (PPY12 + pTW84) that expressed PTS2-GFP was made, and the BLE fused to the PTS2 sequence (PTS2-BLE) under the control of the glyceraldehyde 3-phosphate dehydrogenase promoter (*GAPDH*) from a plasmid integrated in the genome. These cells were mutagenized, induced on oleate, and treated with phleomycin. We expected that PTS2 import mutants would be able to grow on methanol because none of the known enzymes involved in methanol metabolism contain a PTS2 import signal. Conversely, because thiolase is required for growth on oleate, we expected that a PTS2 import-deficient mutant would be unable to grow on oleate. Therefore, colonies that had survived the phleomycin selection were screened for their ability to grow on methanol but not on oleate. Two mutants (*fox3.1* and *pex7.1*) clearly fulfilled this criterion (shown for *pex7Δ* in a growth curve in Fig. 2, A and B). Inspection of PTS2-GFP import revealed, however, that one of these mutants (*fox3.1*) was unaffected in PTS2 protein import. Cloning of the complementing gene revealed that this mutant was deficient in the thiolase gene itself (Koller, A., and S. Subramani, unpublished results).

pex7.1 Cells Are Deficient in PTS2 Import Only

The second mutant (*pex7.1*) was not only specifically dis-




Figure 1. Subcellular location of GFP-PTS1 and PTS2-GFP in wild-type and *pex7.1* cells. Wild-type or *pex7.1* cells expressing either GFP-SKL or PTS2-GFP under the control of the *GAPDH* and *ACO1* promoters, respectively, were grown on YPO for 15 h and washed twice in water. (A and B) Pictures obtained using Nomarski optics of (A) wild-type (PPY12) cells and (B) *pex7.1* cells expressing GFP-SKL. (C and D) Fluorescence microscopy of (C) wild-type (D) and *pex7.1* cells expressing GFP-SKL. (E and F) Fluorescence microscopy of (E) wild-type cells and (F) *pex7.1* cells expressing PTS2-GFP. (G) Fluorescence microscopy of *pex7.1* cells with integrated NH-Pex7p/*ACO1* expressing PTS2-GFP.

turbed in its growth on oleate (Fig. 2, A and B), but was also unable to import PTS2-GFP (Fig. 1 F), whereas import of GFP-SKL was normal (Fig. 1 D). This suggested a general deficiency of *pex7.1* cells in the import of PTS2-containing proteins. The *pex7.1* mutant was therefore further characterized by biochemical subcellular fractionation. In wild-type cells, we observed an equal distribution of thiolase between the organellar pellet fraction and the cytosolic supernatant fraction. As expected, thiolase was exclusively present (27,000 g) in the cytosolic supernatant




Figure 2. Characterization of the *pex7* phenotype. Growth on (A) methanol- and (B) oleate-containing medium. Wild-type cells, *pex7Δ* cells, and *pex7Δ* cells complemented with the *PEX7* gene were pregrown in YPD medium and inoculated at (A) $OD_{600} = 0.1$ in minimal methanol medium (SM) or (B) at $OD_{600} = 0.025$ in minimal oleate medium. (C) Subcellular distribution of peroxisomal enzymes in oleate-induced wild-type and *pex7.1* cells. After subcellular fractionation, equivalent volumes of the 1,000 g postnuclear supernatant (homogenate [H]), 27,000 g pellet (P), and 27,000 g supernatant (S) were analyzed by SDS-PAGE and Western blotting using antibodies raised to the proteins as indicated. (D) Complementation of *pex7.1* on oleate by the pM1 and pM2 plasmid. *pex7.1* cells and *pex7.1* cells transformed with pM1 or pM2 were pregrown in (selective) minimal glucose medium (SD) and inoculated at $OD_{600} = 0.1$ in minimal oleate medium.

fraction of *pex7.1*, providing additional evidence that the import of PTS2-containing proteins was disturbed (Fig. 2 C). Targeting of the membrane protein Pex3p was normal, as was the import of acyl CoA oxidase (Aco1p), because these proteins were predominantly present (27,000 g) in the organellar pellet fraction. Furthermore, the import of presumed PTS1-containing proteins such as catalase (Cta1p) and trifunctional enzyme (Tfe1p) was unaffected, suggesting that the import deficiency of *pex7.1* is specific for PTS2-containing proteins only.

Cloning of *pex7.1* Complementing Genes

The *pex7.1* mutant was complemented for growth on oleate medium using a genomic library. Two plasmids (pM1 and pM2) with nonoverlapping inserts restored growth on oleate, although growth was better when using plasmid pM2 (Fig. 2 D). Further analysis of plasmid pM1 revealed that complementation of the oleate-minus phenotype was caused by a truncated protein (amino acids 1–342) with highest homology to ScPat1p, a peroxisomal ABC trans-

porter (Hetteima et al., 1996). Interestingly, it has been reported that the peroxisomal ABC transporter Pmp70 can suppress the peroxisomal protein import deficiency of a mammalian *pex2*-deficient cell line (Gartner et al., 1994). Therefore, we tested whether plasmid pM1 was able to restore the PTS2-GFP import in the *pex7.1* mutant. Microscopy analysis did not reveal any restoration of import of PTS2-GFP in these cells, suggesting that this protein did not act as a suppressor of the PTS2 import deficiency (data not shown). It has been recently demonstrated that ScPat1p and ScPat2p are involved in the transport of (activated) long-chain fatty acids across the peroxisomal membrane (Hetteima et al., 1996). We therefore speculate that the truncated PpPat1p somehow disturbed the normal functioning of the Pat1p/Pat2p complex, thereby allowing leakage of the partially completed β oxidation product (3-oxoacyl-CoA) into the cytosol, allowing cytosolic thiolase to complete the β oxidation in this *pex7.1* mutant.

The second clone (pM2) was not only able to restore growth on oleate of the *pex7.1* mutant, but it also corrected the import deficiency of PTS2-GFP (data not shown). Further analysis revealed that the complementing gene encoded a protein of 376 amino acids with a calculated molecular mass of 42.4 kD. The protein contains six WD-40 repeats and has 43% sequence identity with ScPex7p (Fig. 3). Evidence that we cloned a true orthologue of ScPex7p was obtained by expressing the gene in *S. cerevisiae*. It was able to complement the growth defect of *S. cerevisiae pex7* (Fig. 4 A); hence, the gene is designated *PpPEX7*.

Pex7p Interacts with the PTS2

Pex7p from *S. cerevisiae* has been shown to interact with the PTS2 sequence in multiple ways (Rehling et al., 1996; Zhang and Lazarow, 1996). Because PpPex7p complements the *Scpex7* mutant, we expected PpPex7p to interact with the PTS2 sequence. This was tested using the yeast two-hybrid system (Fields and Song, 1989). Therefore, we fused Pex7p to the LexA activation domain (AD-PpPex7p) and Sc-thiolase to the LexA DNA-binding domain (DB-ScFox3p). These plasmids were coexpressed in *S. cerevisiae* L40 cells with either an empty control plasmid or with each other. High induction of the *HIS3* gene and β -galactosidase was obtained when AD-PpPex7p and DB-ScFox3p were coexpressed, whereas no induction was observed when AD-PpPex7p or DB-ScFox3p was coexpressed with a control plasmid (shown for *HIS3* expression in Fig. 4 B). This indicates that indeed Pex7p interacts with Fox3p in vivo. To test whether this interaction was dependent upon the presence of the PTS2, we made a construct from which the 17 NH₂-terminal amino acids of thiolase were removed (DB-ScFox3p Δ PTS2). No induction of the *HIS3* gene and β -galactosidase was observed when DB-ScFox3p Δ PTS2 was coexpressed with AD-PpPex7p, indicating that the interaction of PpPex7p with thiolase was PTS2 dependent. To test whether the PTS2 alone would be sufficient to bind to PpPex7p, we fused the first 17 amino acids of ScFox3p to the LexA DNA-binding domain (DB-ScPTS2). When DB-ScPTS2 was coexpressed with AD-PpPex7p, we found again a strong induction of the *HIS3* gene and β -galactosidase (shown for *HIS3* expression in Fig. 4 B). These results




Figure 3. *PpPEX7* encodes a conserved WD-40 repeat protein. Alignment of PpPex7p with Pex7p from *S. cerevisiae* (*Sc*), *Kluyveromyces lactis* (*Kl*), and human (*Hs*). The WD-40 domains are overlined according to the consensus of Neer et al. (1994). The mutations found in RCDP patients are indicated.

indicate that the ScPTS2 sequence alone is sufficient for interaction with PpPex7p in vivo, supporting a receptor-like function for this protein.

Fox3p-SKL Is Efficiently Imported into Peroxisomes of pex7 Δ Cells

To study the function of Pex7p, we made a *PEX7* deletion mutant (*pex7 Δ*) in which the entire open reading frame was replaced by the *PpARG4* gene. Subcellular fractionation of this mutant showed the same phenotype as was observed for the *pex7.1* mutant: a specific import deficiency of the PTS2-containing enzyme thiolase (shown later in Figs. 5 B and 6, B and D).

Like the original *pex7.1* mutant, *pex7 Δ* did grow on methanol medium, suggesting that peroxisomal import of PTS2-containing enzymes is not necessary for growth on methanol (Fig. 2 A). Conversely, *pex7 Δ* did not grow on oleate, indicating that import of at least one PTS2-containing protein is required for growth on this medium (Fig. 2 B). We investigated whether the growth defect on oleate was caused solely by the mistargeting of thiolase. There-




Figure 4. *PpPEX7* encodes the PTS2 receptor orthologue of *ScPEX7*. (A) Complementation of *Scpex7.1* with *PpPEX7*. Wild-type *S. cerevisiae* cells (BJ1991), *Scpex7.1* cells, and *Scpex7.1* cells expressing *PpPex7p* from the *ScCTA1* promoter were grown on minimal oleate plates for 5 d. (B) Analysis of *PpPex7p* interaction with the PTS2 sequence in a two-hybrid assay by testing for *HIS3* expression. *S. cerevisiae* L40 cells were transformed with the plasmids as indicated, grown on glucose plates lacking leucine and tryptophan, spotted on plates lacking histidine, and incubated for 2 d. *AD* and *DB* denote genes cloned in frame with the LexA transcription-activation domain or LexA DNA-binding domain, respectively.

fore, a construct (Fox3p-SKL) expressing *P. pastoris* Fox3p (Koller, A., S. Subramani, unpublished results) fused to the PTS1 sequence (SKL) under the control of the *ACO1* promoter was integrated in the genome of *pex7Δ* and *fox3Δ* cells. This protein restored growth on oleate plates to nearly wild-type levels when expressed in *Ppfox3Δ*, indicating that Fox3p-SKL is active and imported into peroxisomes (Fig. 5 A). Expression of Fox3p-SKL in *pex7Δ* only partially restored the growth on oleate plates. This suggests that the inability to import thiolase is not the only reason why *pex7Δ* cells do not grow on oleate, but that other PTS2-containing proteins are also required for growth on this medium (Fig. 5 A).

The import efficiency of Fox3p-SKL was studied by subcellular fractionation (Fig. 5 B). We observed a considerable amount of endogenous thiolase in the cytosolic fraction in the control (untransformed wild-type) cells, whereas thiolase was exclusively cytosolic in untransformed *pex7Δ* cells. The control enzyme, Aco1p, was predominantly present in the organellar pellet fractions (Fig. 5 B). These results suggest that thiolase easily leaks out of peroxisomes during subcellular fractionation. Alternatively, the PTS2 import pathway in *P. pastoris* is not very efficient. This is supported by the observation that PTS2-GFP is also largely cytosolic and only partially peroxisomal (Fig.




Figure 5. Analysis of Fox3p-SKL function and targeting. (A) Growth of wild-type, *fox3Δ*, *fox3Δ* + Fox3p-SKL, *pex7Δ*, and *pex7Δ* + Fox3p-SKL on a minimal oleate plate for 5 d. (B) Subcellular location of Fox3p-SKL. Wild-type cells, *pex7Δ* cells, and *fox3Δ* and *pex7Δ* cells expressing Fox3p-SKL were induced on oleate for 15 h. After subcellular fractionation, equivalent volumes of the 1,000 g postnuclear supernatant (H, homogenate), 27,000-g pellet (P), and 27,000-g supernatant (S) were analyzed by SDS-PAGE and Western blotting using antibodies raised to the proteins as indicated.

1, E and G), although we cannot rule out that this localization is caused by the chimeric protein itself. Interestingly, Fox3p-SKL was fully imported into peroxisomes of *fox3Δ* cells, with no protein detectable in the cytosolic fraction. In *pex7Δ* cells expressing Fox3p-SKL, thiolase was equally distributed between the pellet and supernatant fractions (Fig. 5 B). From the small difference in molecular weight between Fox3p-SKL and endogenous Fox3p, we conclude that Fox3p-SKL is fully imported into these cells via its PTS1 sequence, whereas endogenous thiolase is almost exclusively cytosolic, as in the untransformed *pex7Δ* control cells (Fig. 5 B). These results suggest that the PTS2 import pathway in *P. pastoris* may be less efficient than the PTS1 import pathway (see Discussion).

Antibodies against *Pex7p*

The lack of useful antibodies against endogenous *Pex7p* has greatly hampered the localization analysis of this protein in *S. cerevisiae* and in human cells. We therefore wanted to raise antibodies against *PpPex7p* to analyze its localization. We expressed the full-length *Pex7p* in *E. coli* fused to a (His)₆ tag, but the expression of this protein was very poor. We were unable to increase its expression by fusing it to dihydrofolate reductase, nor could we increase the expression by fusing three different truncated versions of *Pex7p* to dihydrofolate reductase. Good expression in *E. coli* was obtained for a truncated protein corresponding to amino acids 1–143 fused to the (His)₆ epitope. This puri-

fied protein, (His)₆-Pex7p(143), was used to immunize rabbits. The antiserum that was obtained recognized several proteins in the wild-type strain and in the *pex7Δ* strains (Fig. 6 A, lanes 1 and 2). Because this could indicate that a (nonfunctional) pseudogene of *PEX7* might be present, we checked this possibility with PCR and low stringency Southern blotting, but did not find any evidence for the existence of such a gene (data not shown). We purified the antiserum with an immunodepletion column, using a lysate obtained from *pex7Δ* cells. The flow-through of this column was used to purify the antiserum on an affinity column using the coupled (His)₆-Pex7p(143) protein. The eluted antibody recognized two prominent bands in lysates of wild-type cells (Fig. 6, A, lane 4). The signal obtained from the lower molecular weight (cytosolic) protein was present at the same intensity in *pex7Δ* cells (Fig. 6, lane 3) and could be eliminated by using more stringent washing conditions (see figures below). The higher molecular weight band was largely reduced in *pex7Δ* cells, indicating that this signal is from Pex7p (Fig. 6 A, lane 3 versus lane 4). This could be confirmed by overexpressing Pex7p from the acyl-CoA oxidase promoter (Fig. 6, lane 5). A 25-fold dilution of this lysate (compared to a wild-type and *pex7Δ* lysate) gave only one signal at the expected molecular weight (Fig. 6 A, lane 6). Because there is still a weak signal present at this molecular weight in a *pex7Δ* lysate (from a mitochondrial protein, see below), in our subsequent analyses, we included the *pex7Δ* control and separated the peroxisomes from mitochondria where necessary.

PpPex7p Is Distributed between the Peroxisomes and the Cytosol

The intracellular location of Pex7p was determined by subcellular fractionation. Wild-type and *pex7Δ* cells were fractionated, and the organellar pellet and the cytosolic supernatant were analyzed. Aco1p, Pex3p, and thiolase (Fox3p) were present in the organellar (27,000 g) pellet fraction of wild-type cells, although varying portions of some of these proteins were also present in the cytosolic (27,000 g) supernatant fraction, possibly because of leakage (Fig. 6 B). A similar distribution was observed for *pex7Δ* cells, with the exception of thiolase, which was exclusively present in the cytosolic fraction, reflecting the PTS2 import deficiency of *pex7Δ* (Fig. 6 B). Pex7p was present in both the organellar pellet (varying from 10 to 30%) and the cytosol fraction of wild-type cells, suggesting that about a quarter of the protein is associated with peroxisomes. In Fig. 6 B, the mitochondrial protein that sometimes cross-reacts with the anti-Pex7p antibodies (see Fig. 6, A and D) would not affect our conclusion that Pex7p is partially found in the cytosol because (a) we did not observe the cross-reacting band in this particular experiment (see H and P fractions of control *pex7Δ* cells), and (b) the cross-reacting mitochondrial band was never found in the supernatant fraction during subcellular fractionation. The protein band found in the cytosolic fraction can therefore be only Pex7p.

To determine whether the pelletable Pex7p was indeed associated with peroxisomes, we used a postnuclear supernatant for further analysis on a Nycodenz gradient. A por-




Figure 6. Location of Pex7p in a subcellular fractionation. (A) Characterization of the α Pex7p antibody in a Western blot. TCA lysates were obtained from cells induced on oleate for 15 h. (Lanes 1 and 2) Antibody before purification and (lanes 3–6) antibody after purification on a depletion and affinity column. (Lanes 1 and 3) *pex7Δ* cells, (lanes 2 and 4) wild-type cells, (lane 5) *pex7Δ* cells expressing Pex7p/*ACO1*, and (lane 6) 25-fold diluted lysate (relative to other samples) of *pex7Δ* cells expressing Pex7p/*ACO1*. (B) Location of Pex7p in a subcellular fractionation. Oleate-induced wild-type and *pex7Δ* cells were fractionated, and equivalent volumes of the 1,000-g postnuclear supernatant (H, homogenate), 27,000-g pellet (P), and 27,000-g supernatant (S) were analyzed by SDS-PAGE and Western blotting using antibodies raised to the proteins as indicated. Arrow, Pex7p. (C and D) Location of Pex7p in a Nycodenz gradient. (C) Oleate-induced wild-type and (D) *pex7Δ* cells were fractionated, and the 1,000-g postnuclear supernatants were analyzed on a Nycodenz gradient. Fractions were subjected to SDS-PAGE, blotted, and probed with the antibodies listed. Fox3p and Aco1p are markers for peroxisomes, and a flavoprotein subunit of succinate dehydrogenase (Sdh1p) is a marker for the mitochondria. Fraction 1, the bottom fraction; fraction 20, the top fraction.

tion of Pex7p comigrated with the peroxisomal marker enzymes Aco1p and Fox3p, suggesting that at least some Pex7p is associated with peroxisomes (Fig. 6 C). We also found significant amounts of α Pex7p-cross-reacting mate-

rial at the top of the gradient (fractions 14–20), representing mitochondrial and cytosolic proteins. In these gradients, the mitochondrial protein (Fig. 6 D, fractions 14–18) that cross-reacted with the antibody against Pex7p was never found at the top of the gradient in fractions 19 and 20, where cytosolic proteins would be found. The presence of Pex7p in fractions 19 and 20 of the gradient from wild-type cells (Fig. 6 D) shows that some Pex7p is indeed cytosolic. Because of the cross-reacting protein present in the mitochondrial fractions (see fractions 14–18 in the *pex7Δ* control gradient, Fig. 6 D), however, it is impossible to estimate the exact amount of Pex7p present in the cytosol in this experiment.

To study the localization of Pex7p by an independent biochemical method, we selectively permeabilized spheroplasts of wild-type cells, *pex7Δ* and *pex3Δ* (a mutant disturbed in the import of PTS1- and PTS2-containing proteins; Wiemer et al., 1996) by digitonin, and we followed the subsequent leakage of Pex7p and control proteins from the cells. In this procedure, cytosolic enzymes are released at low concentrations of digitonin, whereas peroxisomal matrix enzymes are released at much higher concentrations of digitonin (Zhang and Lazarow, 1995). As can be seen in Fig. 7, the cytosolic enzyme glucose-6 phosphate dehydrogenase (G6PDH) was completely released at low concentrations (50 μg/ml) of digitonin. This was also observed for catalase in *pex3Δ*, as well as thiolase in the *pex7Δ* and *pex3Δ* strain, which reflects the peroxisomal import deficiency of these mutants. In contrast, thiolase and catalase were released at significantly higher concentrations of digitonin in wild-type cells, starting at 150 μg/ml digitonin, and release was not complete until 300–400 μg/ml digitonin was added. The membrane protein Pex3p could be completely released at only very high concentrations of digitonin (>600 μg/ml; Fig. 7). Using the αPex7p antibody, we observed release of the mitochondrial αPex7p-cross-reacting protein in *pex7Δ* cells at digitonin concentrations of ≥400 μg/ml. In wild-type cells, a substantial amount of Pex7p was already released before release of peroxisomal matrix proteins took place (before 150 μg/ml digitonin). In contrast to the cytosolic markers, however, release was not complete until 300–400 μg/ml digitonin, which is similar to what was observed for the release of Cta1p and Fox3p. In *pex3Δ* cells, Pex7p was completely released at 50 μg/ml. These results confirm that Pex7p is distributed between the peroxisome and the cytosol. Furthermore, since complete release of Pex7p is observed at digitonin concentrations <400 μg/ml, these re-

sults also suggest that some Pex7p is present in the lumen of peroxisomes.

Overexpressed PpPex7p Is Primarily Located in the Cytosol

If Pex7p functions solely as an intraperoxisomal receptor, it probably has a PTS that directs it efficiently to the peroxisomal matrix. Some experiments in *S. cerevisiae* suggest that this is indeed the case (Zhang and Lazarow, 1996). To test whether PpPex7p can direct a reporter protein to the peroxisome, we fused PpPex7p to the peroxisomal malate dehydrogenase of *S. cerevisiae*, which lacks its PTS1 (NH-Mdh3pΔPTS1). It has previously been shown that NH-Mdh3pΔPTS1 is cytosolic and that it can not complement the growth defect on oleate of *S. cerevisiae mdh3Δ* cells (Elgersma et al., 1996b). When PpPex7p was fused to NH-Mdh3pΔPTS1 (NH-Mdh3p-PpPex7p) and expressed in *S. cerevisiae mdh3Δ* cells under the control of the *ScCTA1* promoter, we observed some complementation of the *mdh3Δ* mutant (data not shown). This suggests that PpPex7p does indeed have a signal that can direct it to the peroxisomal matrix.

To study the effect of overexpressing and/or epitope-tagging Pex7p, the expression level of Pex7p was increased by expressing it under the control of the *ACO1* promoter. Western blot analysis revealed that this resulted in an ~25-fold higher expression level (see Fig. 6 A). The overexpressed protein fully complemented the growth deficiency of *pex7Δ* cells (Fig. 8 A). Full complementation was also observed when we epitope tagged Pex7p by fusing it to the NH tag (Elgersma et al., 1996b) and expressed it under the control of the *ACO1* promoter (NH-Pex7p/*ACO1*) (Fig. 8 A). Neither tagging nor overexpression affected its ability to restore the peroxisomal import of thiolase in *pex7Δ* cells, as shown by subcellular fractionation (Fig. 8 B). Interestingly, we found on average >90% of overexpressed Pex7p in the cytosol (Fig. 8 B). Comparison of the absolute amount present in the organellar pellet revealed that the amount of pelletable Pex7p was only increased 5–10-fold (Fig. 8 C) compared to untransformed cells. These experiments suggest that targeting of PpPex7p to the peroxisomal matrix is not as efficient as the PTS1 or PTS2 import pathway, and that an unknown component is limiting the import of Pex7p.

The localization of overexpressed NH-Pex7p was also addressed by fluorescence microscopy of semithin sections of yeast cells labeled with the NH antibody. This antibody




Figure 7. Location of Pex7p in digitonin-permeabilized, wild-type, *pex3Δ* and *pex7Δ* cells. Oleate-induced cells were spheroplasted, aliquoted in identical portions, and incubated for 10 min at 30°C with increasing amounts of digitonin dissolved in isotonic buffer. After quickly cooling the samples on ice, the spheroplasts were pelleted, and the supernatants were used for Western blotting with the antibodies as indicated. *Triton*, the samples have been incubated with 0.2% Triton X-100 in hypotonic buffer to get complete release of all proteins. Digitonin concentrations are given in micrograms per milliliter.




Figure 8. Analysis of overexpressed Pex7p. (A) Complementation of *pex7* on oleate medium. *pex7*Δ cells, *pex7*Δ cells expressing Pex7p/*ACO1* or Pex7p/*PEX7*, and *pex7.1* cells expressing NH-Pex7p/*ACO1* were pregrown in YPD medium and inoculated at $OD_{600} = 0.025$ in minimal oleate medium. (B) Subcellular locations of overexpressed Pex7p and NH-Pex7p. Oleate-induced *pex7*Δ cells expressing NH-Pex7p/*ACO1* or Pex7p/*ACO1* were fractionated, and equivalent volumes of the 1,000-g postnuclear supernatant (H, homogenate), 27,000-g pellet (P), and 27,000-g supernatant (S) were analyzed by SDS-PAGE and Western blotting using antibodies raised to the proteins as indicated. (C) Amounts of Pex7p in crude organellar pellets obtained from wild-type cells and *pex7*Δ cells expressing NH-Pex7p/*ACO1* or Pex7p/*ACO1*. Oleate-induced cells were fractionated, and equivalent volumes of the

27,000-g pellets were analyzed by SDS-PAGE and Western blotting. Cta1p is shown as a control that similar amounts of crude organellar pellet were layered from each fractionation.

and epitope tag have previously been shown to be very suitable for immuno-EM analysis (Elgersma et al., 1996b). Analysis of control cells expressing NH-tagged malate dehydrogenase of *S. cerevisiae* (NH-Mdh3p) under the control of the *ACO1* promoter showed a punctate labeling, indicating efficient import of NH-Mdh3p into peroxisomes (Fig. 9, A and B). In contrast, cells expressing NH-Pex7p showed only limited punctate staining and a high cytosolic staining (Fig. 9, C and D). This supports the biochemical data that a large amount of overexpressed Pex7p is cytosolic. To verify the nature of the punctate staining, we used the cells expressing NH-Mdh3p or NH-Pex7p for immuno-EM analysis. This revealed that a portion of NH-Pex7p is indeed present in the peroxisomal matrix (Fig. 10, B–D). These results confirm the predominantly cytosolic and partially intraperoxisomal location of overexpressed NH-Pex7p.

Human PEX7 Mutations Affect the Ability of PpPex7p to Bind PTS2

The human *PEX7* gene has recently been cloned and shown to be responsible for RCDP (Braverman et al., 1997; Motley et al., 1997; Purdue et al., 1997). The most frequently occurring mutations were the L292ter and A218V mutations. Five patients were genetic compounds of the L292ter as the first allele and a G217R mutation as the second allele (Braverman et al., 1997). Expression studies showed that the L292ter and A218V mutations im-




Figure 9. Immunofluorescence microscopy analysis of cells expressing NH-Mdh3p/*ACO1* or NH-Pex7p/*ACO1*. Semithin sections of cells expressing NH-Mdh3p/*ACO1* or NH-Pex7p/*ACO1* were labeled with the α NH antibody and analyzed by immunofluorescence microscopy. (A and B) Pictures obtained using Nomarski and fluorescence optics, respectively, of cells expressing NH-Mdh3p/*ACO1*. (C and D) Pictures obtained using Nomarski and fluorescence optics, respectively, of cells expressing NH-Pex7p/*ACO1*. Bar, 5 μ m.

paired Pex7p function, but the functional consequence of the G217R mutation was undetermined. To test whether the mutated proteins were nonfunctional because of instability, mistargeting, or failure to bind the PTS2 sequence, we made the *PpPEX7* equivalents of the RCDP G217R, A218V, and L292ter mutations (A248R, G249V, and C347ter, respectively, as indicated in Fig. 3). The mutated PpPex7p proteins were expressed under the control of the *PEX7* and *ACO1* promoters, and they were tested for the ability to correct the growth deficiency of *pex7*Δ on oleate. As shown in Fig. 11 A, all mutated proteins were unable to restore the growth defect on oleate. When (over)expressed under the control of the *ACO1* promoter, we observed some oleate usage by Pex7p(G249V), as judged by the formation of a halo on an oleate plate (data not shown). This correlates with the finding that patients with this corresponding allele have a phenotype milder than in classical RCDP (Braverman et al., 1997).

We made lysates of *pex7*Δ cells expressing the mutated Pex7p. As shown in Fig. 11 B, all proteins were expressed stably, suggesting that the deficiency is not caused by protein instability. We then tested whether the mutated proteins were able to bind the PTS2 sequence, using the two-hybrid system as described above. Interestingly, all mutated proteins had lost the ability to bind the PTS2 sequence (Fig. 11 C). This suggests that the mutations lead to impaired PTS2 binding.

To test whether the mutations affected the peroxisomal targeting of Pex7p, we expressed the mutated proteins in *pex7*Δ cells and compared their targeting to that observed for wild-type Pex7p by Nycodenz gradient analysis. We




Figure 10. Immuno-EM analysis of cells expressing NH-Mdh3p/ACO1 or NH-Pex7p/ACO1. Ultrathin sections of cells expressing NH-Mdh3p/ACO1 or NH-Pex7p/ACO1 were labeled with the α NH antibody and by immuno-gold, and they were analyzed by EM. (A) Cells expressing NH-Mdh3p/ACO1. (B–D) Cells expressing NH-Pex7p/ACO1. C is an enlargement of the indicated area in B. Bar, 0.1 μ m.

did not observe significant differences between the distribution of wild-type or mutated proteins (data not shown). This indicates that peroxisomal targeting of the mutated Pex7p proteins is not affected. Moreover, this suggests that binding of the PTS2 sequence is not a prerequisite for targeting Pex7p to the peroxisome.

Discussion

PpPex7p Is a Receptor for PTS2 Protein Import

Using a new screening procedure devised to yield mutants specifically compromised in the PTS2 import pathway, we have obtained a *P. pastoris pex7* mutant, cloned the *PpPEX7* gene, and shown in several ways that PpPex7p is the homologue of ScPex7p, the PTS2 receptor for protein import (Rehling et al., 1996, Zhang and Lazarow, 1996): (a) PpPex7p has high sequence identity with ScPex7p (43%), (b) *PpPEX7* complements the *ScPex7* mutant, (c) *Pppex7 Δ* is specifically disturbed in PTS2 protein import, and (d) PpPex7p interacts with the PTS2 sequence in the two-hybrid system. The observation that *Pppex7 Δ* is unaffected in its growth on methanol medium suggests that there are no PTS2-containing enzymes or peroxins required for growth on this medium. Furthermore, it ex-

plains why screening procedures for *pex* mutants, based on their inability to grow on methanol, did not result in the isolation of this mutant.

PpPex7p Functions Probably as a Mobile Receptor

The role of Pex7p in PTS2 import has been a matter of debate. Zhang and Lazarow (1995, 1996) concluded that ScPex7p is an intraperoxisomal receptor and functions by pulling PTS2 proteins inside, analogous to the role of mitochondrial HSP70. Although they found significant amounts of a triple-tagged Pex7p (Pex7p-HA3) in the cytosol upon fractionation, they observed in digitonin experiments a strikingly similar pattern of latency of Pex7p-HA3 compared to peroxisomal matrix proteins. Therefore, they concluded that the cytosolic pool found in fractionation studies was probably caused by leakage of Pex7p-HA3. In contrast, others proposed that Pex7p functions as a mobile cytosolic receptor because most of the (overexpressed and tagged) Pex7p protein was found in the cytosol by biochemical fractionation (Marzioch et al., 1994; Rehling et al., 1996) or by immunofluorescence (Braverman et al., 1997).

We raised antibodies against the endogenous Pex7p and studied its localization. Subcellular fractionation and digitonin experiments suggest that Pex7p is largely cytosolic




Figure 11. Analysis of the RCDP mutations introduced in PpPex7p. (A) Complementation of *pex7Δ* on oleate medium. *pex7Δ* cells and *pex7Δ* cells expressing Pex7p, Pex7p(C347ter), Pex7p(G249V), or Pex7p(A248R) under the control of the *PEX7* promoter were pregrown in YPD medium and inoculated at $OD_{600} = 0.025$ in minimal oleate medium. (B) The mutated Pex7p proteins are expressed stably. Western blot of TCA lysates obtained from oleate-induced *pex7Δ* cells expressing (lane 1) Pex7p, (lane 2) Pex7p(C347ter), (lane 3) Pex7p(G249V), or (lane 4) Pex7p(A248R) under the control of the *ACO1* promoter.

The mitochondrial F1 β subunit of F1 ATPase is shown as a control that is similar to the amounts of protein that were layered from each lysate. (C) Mutated PpPex7p fails to bind to ScFox3p in a two-hybrid analysis. *S. cerevisiae* L40 cells were transformed with the plasmids as indicated, grown on glucose plates lacking leucine and tryptophan, spotted on plates lacking histidine, and incubated for 2 d. ScFox3p was cloned in frame with the LexA DNA-binding domain. Pex7p and mutated Pex7p were cloned in frame with the LexA transcription-activation domain.

but that some of the protein is peroxisomal. Evidence that the peroxisome-associated protein is present in the matrix was deduced from digitonin experiments and immuno-EM using NH-tagged Pex7p, which showed clear peroxisomal matrix labeling and no obvious membrane labeling. Interestingly, a 25-fold overexpression of Pex7p results in a modest 5–10-fold increase in peroxisome-associated Pex7p and in a large increase of cytosolic Pex7p. This explains why others found an almost exclusively cytosolic localization for Pex7p when they studied the localization of overexpressed Pex7p (Marzioch et al., 1994; Rehling et al., 1996; Braverman et al., 1997).

Our analyses indicate that most of Pex7p is cytosolic (varying from 70 to 90%), suggesting a role in the cytosol where it binds to PTS2-containing proteins and delivers them at the peroxisomal membrane. The identification of Pex14p as a putative peroxisomal docking protein for Pex7p supports this model (Albertini et al., 1997). A similar model has also been proposed for the PTS1 receptor Pex5p, which binds to the peroxisomal membrane protein Pex13p (Elgersma et al., 1996a; Erdmann and Blobel, 1996; Gould et al., 1996).

In addition to the cytosolic pool of Pex7p, we found also a significant fraction of Pex7p in the peroxisomal matrix (varying from 10 to 30%). Therefore, it is possible that Pex7p is also involved in the subsequent translocation step. In that case, we expect that Pex7p has to get recycled to the cytosol because thiolase expression is at least 25-fold higher than Pex7p expression. The observation that the amount of intraperoxisomal Pex7p did not increase proportionally upon overexpression suggests that the Pex7p

import signal is not very efficient. Therefore, it is conceivable that intraperoxisomal Pex7p does not have a role at all, but that some Pex7p enters the peroxisome coincidentally and remains in the peroxisomal matrix without any function. Further kinetic experiments are required to address this point.

Targeting of Pex7p to the Peroxisomal Matrix

The ability of Pex7p to enter the peroxisomes, albeit at low efficiency, appears to be real. Some amounts of the wild-type and overexpressed Pex7p were intraperoxisomal, as judged by digitonin permeabilization and immuno-EM (Figs. 7 and 10). Furthermore, NH-Mdh3p-PpPex7p was able to complement an *mdh3Δ* strain of *S. cerevisiae* (data not shown). The mechanism involved in this targeting is unknown. Because protein unfolding is not essential for the import of proteins into the peroxisomal matrix, Pex7p might enter the matrix solely by association with PTS2-containing proteins. This is unlikely, however, because the A248R, G249V, and C347ter mutants cannot interact with the PTS2 sequence and show the same subcellular location as wild-type Pex7p. Moreover, within experimental error, we did not observe a different distribution of Pex7p when we deleted the *FOX3* gene or when we raised its expression twofold by overexpressing it from the *ACO1* promoter. An alternative possibility is that Pex7p has a new targeting signal or that it enters the peroxisome in association with some other protein. Such a targeting or protein association signal might reside in the first 56 amino acids of ScPex7p, as suggested by the experiments of Zhang and Lazarow (1996).

Yeast Model System for Human Peroxisome Biogenesis Defects

The cloning of yeast *PEX* genes has tremendously facilitated the characterization of their human orthologues (for review see Subramani, 1997). For instance, the human *PEX7* gene was cloned by screening databases using the yeast *PEX7* gene (Braverman et al., 1997; Motley et al., 1997; Purdue et al., 1997). Our studies show a new role for the yeast model system: the rapid analysis of the effect of the mutations found in the *PEX* genes of patients. We introduced the mutations found in the RCDP patients at the corresponding positions in the *PpPEX7* gene and found that this resulted in impaired growth on oleate. Furthermore, we found that the mutated Pex7p proteins in *P. pastoris* are stably expressed and not disturbed in peroxisomal targeting, but that they are unable to bind to the PTS2 sequence. The application of a similar strategy to other human *PEX* genes involved in disease is likely to provide useful insights regarding the effect of mutations.

Fox3p-SKL Only Partially Restores Growth on Oleate When Expressed in *pex7Δ* Cells

Pppex7Δ did not grow on oleate plates, but this deficiency could be partially suppressed by expressing Fox3p-SKL in these cells. This indicates that the lack of thiolase import is possibly the main, but certainly not the only, reason that *pex7Δ* cells do not grow on oleate. Hence, we expect that other PTS2-containing proteins are also required for ole-

ate metabolism. Surprisingly, it appeared that Fox3p-SKL was either imported or retained in peroxisomes much more efficiently than endogenous thiolase. Moreover, it should be noted that we did not observe significant import of endogenous thiolase in *pex7Δ* cells expressing Fox3p-SKL. This indicates that the formation and import of heterodimers of Fox3p-SKL with endogenous thiolase did not take place, in contrast to what has been observed for *S. cerevisiae* cells expressing thiolase with and without a PTS2 (Glover et al., 1994b). This may suggest that the import of Fox3p-SKL via the PTS1 pathway is much faster than the import of endogenous thiolase via the PTS2 pathway, and that import of Fox3p-SKL takes place before the folding and oligomerization of Fox3p-SKL is completed. Further kinetic experiments, as conducted by Ruigrok et al. (1996), are necessary to validate this hypothesis.

We are grateful to Dr. A. Koller for providing us the *PpFOX3* gene sequence, to Dr. H.F. Tabak for providing us thiolase antibodies and *S. cerevisiae* strains and plasmids, Dr. Peter van der Sluijs for providing the NH antibody, and Drs. P. Bartel and S. Fields for providing the two-hybrid plasmids. We thank Dr. K.-N. Faber for sharing his preliminary characterization of the *pex7* mutant with us.

The Immunocytochemistry Core Facility, University of California, San Diego, was supported by grant CA 58689 to M.G. Farquhar. This work was supported by a National Institutes of Health grant (DK41737) to S. Subramani and by a fellowship from the Netherlands Organization for Scientific Research to Y. Elgersma and by fellowships from the European Molecular Biology Organization to Y. Elgersma and T. Wenzel.

Received for publication 17 October 1997 and in revised form 18 December 1997.

References

Albertini, M., P. Rehling, R. Erdmann, W. Girzalsky, J.A.K.W. Kiel, M. Veenhuis, and W.H. Kunau. 1997. Pex14p, a peroxisomal membrane protein binding both receptors of the two PTS-dependent import pathways. *Cell*. 89:83–92.

Bartel, P., C.T. Chien, R. Sternglanz, and S. Fields. 1993. Using the two-hybrid system to detect protein-protein interactions. In *Cellular Interactions in Development: A Practical Approach*. D.A. Hartley, editor. IRL Press, Oxford. 153–179.

Braverman, N., G. Steel, C. Obie, A. Moser, H. Moser, S.J. Gould, and D. Valle. 1997. Human *PEX7* encodes the peroxisomal PTS2 receptor and is responsible for rhizomelic chondrodysplasia punctata. *Nat. Genet.* 15:369–376.

Cregg, J.M., I.J. Van der Klei, G.J. Sulter, M. Veenhuis, and W. Harder. 1990. Peroxisome-deficient mutants of *Hansenula polymorpha*. *Yeast*. 6:87–97.

Dotd, G., and S.J. Gould. 1996. Multiple *PEX* genes are required for proper subcellular distribution and stability of Pex5p, the PTS1 receptor: evidence that PTS1 protein import is mediated by a cycling receptor. *J. Cell Biol.* 135:1763–1774.

Dotd, G., N. Braverman, C. Wong, A. Moser, H.W. Moser, P. Watkins, D. Valle, and S.J. Gould. 1995. Mutations in the PTS1 receptor gene, *PXR1*, define complementation group 2 of the peroxisome biogenesis disorders. *Nat. Genet.* 9:115–125.

Elgersma, Y., and H.F. Tabak. 1996. Proteins involved in peroxisome biogenesis and functioning. *Biochim. Biophys. Acta*. 1286:269–283.

Elgersma, Y., M. Van den Berg, H.F. Tabak, and B. Distel. 1993. An efficient positive selection procedure for the isolation of peroxisomal import and peroxisome assembly mutants of *Saccharomyces cerevisiae*. *Genetics*. 135:731–740.

Elgersma, Y., L. Kwast, A. Klein, T. Voorn-Brouwer, M. Van den Berg, B. Metzger, T. America, H.F. Tabak, and B. Distel. 1996a. The SH3 domain of the *Saccharomyces cerevisiae* peroxisomal membrane protein Pex13p functions as a docking site for Pex5p, a mobile receptor for the import of PTS1-containing proteins. *J. Cell Biol.* 135:97–109.

Elgersma, Y., A. Vos, M. Van den Berg, C.W.T. Van Roermund, P. Van der Sluijs, B. Distel, and H.F. Tabak. 1996b. Analysis of the carboxyl-terminal peroxisomal targeting signal 1 in a homologous context in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 271:26375–26382.

Erdmann, R., and G. Blobel. 1996. Identification of Pex13p, a peroxisomal membrane receptor for the PTS1 recognition factor. *J. Cell Biol.* 135:111–121.

Faber, K.N., P. Haima, C. Gietl, W. Harder, G. Ab, and M. Veenhuis. 1994. The methylotrophic yeast *Hansenula polymorpha* contains an inducible import pathway for peroxisomal matrix proteins with an N-terminal targeting signal (PTS2 proteins). *Proc. Natl. Acad. Sci. USA*. 91:12985–12989.

Fields, S., and O. Song. 1989. A novel genetic system to detect protein-protein interactions. *Nature*. 340:245–246.

Fransen, M., C. Brees, E. Baumgart, J.C.T. Vanhooren, M. Baes, G.P. Mannaerts, and P.P. Van Veldhoven. 1995. Identification and characterization of the putative human peroxisomal C-terminal targeting signal import receptor. *J. Biol. Chem.* 270:7731–7736.

Gartner, J., C. Obie, P. Watkins, and D. Valle. 1994. Restoration of peroxisome biogenesis in a peroxisome-deficient mammalian cell line by expression of either the 35 kDa or the 70 kDa peroxisomal membrane proteins. *J. Inherited. Metab. Dis.* 17:327–329.

Gietl, C., K.N. Faber, I.J. van der Klei, and M. Veenhuis. 1994. Mutational analysis of the N-terminal topogenic signal of watermelon glyoxysomal malate dehydrogenase using the heterologous host *Hansenula polymorpha*. *Proc. Natl. Acad. Sci. USA*. 91:3151–3155.

Glover, J.R., D.W. Andrews, and R.A. Rachubinski. 1994a. *Saccharomyces cerevisiae* peroxisomal thiolase is imported as a dimer. *Proc. Natl. Acad. Sci. USA*. 91:10541–10545.

Glover, J.R., D.W. Andrews, S. Subramani, and R.A. Rachubinski. 1994b. Mutagenesis of the amino targeting signal of *Saccharomyces cerevisiae* 3-ketoacyl-CoA thiolase reveals conserved amino acids required for import into peroxisomes *in vivo*. *J. Biol. Chem.* 269:7558–7563.

Goebel, M., and M. Yanagida. 1991. The TPR snap helix: a novel protein repeat motif from mitosis to transcription. *Trends Biol. Sci.* 16:173–177.

Görlich, D., and I.W. Mattaj. 1996. Nucleocytoplasmic transport. *Science*. 271:1513–1518.

Gould, S.J., G.A. Keller, and S. Subramani. 1987. Identification of a peroxisomal targeting signal at the carboxy terminus of firefly luciferase. *J. Cell Biol.* 105:2923–2931.

Gould, S.J., G.A. Keller, N. Hosken, J. Wilkinson, and S. Subramani. 1989. A conserved tripeptide sorts proteins to peroxisomes. *J. Cell Biol.* 108:1657–1664.

Gould, S.J., D. McCollum, A.P. Spong, J.A. Heyman, and S. Subramani. 1992. Development of the yeast *Pichia pastoris* as a model organism for a genetic and molecular analysis of peroxisome assembly. *Yeast*. 8:613–628.

Gould, S.J., J.E. Kalish, J.C. Morrell, J. Bjorkman, A.J. Urquhart, and D.I. Crane. 1996. Pex13p is an SH3 protein of the peroxisome membrane and a docking factor for the predominantly cytoplasmic PTS1 receptor. *J. Cell Biol.* 135:85–95.

Hettema, E.H., C.W.T. Van Roermund, B. Distel, M. Van den Berg, C. Vilela, C. Rodrigues-Pousada, R.J.A. Wanders, and H.F. Tabak. 1996. The ABC transporter proteins Pat1 and Pat2 are required for import of long-chain fatty acids into peroxisomes of *Saccharomyces cerevisiae*. *EMBO J.* 15:3813–3822.

Hollenberg, S.M., R. Sternglanz, P.F. Cheng, and H. Weintraub. 1995. Identification of a new family of tissue specific basic helix-loop-helix proteins with a two-hybrid system. *Mol. Cell Biol.* 15:3813–3822.

Komachi, K., M.J. Redd, and A.D. Johnson. 1994. The WD repeats of Tup1 interact with the homeo domain protein a2. *Genes Dev.* 8:2857–2867.

Lazarow, P.B., and Y. Fujiki. 1985. Biogenesis of peroxisomes. *Annu. Rev. Cell Biol.* 1:489–530.

Liu, H., X. Tan, M. Veenhuis, D. McCollum, and J.M. Cregg. 1992. An efficient screen for peroxisome-deficient mutants of *Pichia pastoris*. *J. Bacteriol.* 174:4943–4951.

Marzioch, M., R. Erdmann, M. Veenhuis, and W.H. Kunau. 1994. *PAS7* encodes a novel yeast member of the WD-40 protein family essential for import of 3-oxoacyl-CoA thiolase, a PTS2-containing protein, into peroxisomes. *EMBO J.* 13:4908–4918.

McCollum, D., E. Monosov, and S. Subramani. 1993. The *pas8* mutant of *Pichia pastoris* exhibits the peroxisomal protein import deficiencies of Zellweger syndrome cells. The PAS8 protein binds to the COOH-terminal tripeptide peroxisomal targeting signal and is a member of the TPR protein family. *J. Cell Biol.* 121:761–774.

Monosov, E.Z., T.J. Wenzel, G.H. Luers, J.A. Heyman, and S. Subramani. 1996. Labeling of peroxisomes with green fluorescent protein in living *P. pastoris* cells. *J. Histochem. Cytochem.* 44:581–589.

Motley, A., E. Hettema, B. Distel, and H. Tabak. 1994. Differential protein import deficiencies in human peroxisome assembly disorders. *J. Cell Biol.* 125:755–767.

Motley, A.M., E.H. Hettema, E.M. Hogenhout, P. Brites, A.L.M.A. Ten Asbroek, F.A. Wijburg, F. Baas, H.S. Heijmans, H.F. Tabak, R.J.A. Wanders, et al. 1997. Rhizomelic chondrodysplasia punctata is a peroxisomal protein targeting disease caused by a non-functional PTS receptor. *Nat. Genet.* 15:377–380.

Neer, E.J., C.J. Schmidt, R. Nambudripad, and T.F. Smith. 1994. The ancient regulatory-protein family of WD-repeat proteins. *Nature*. 371:297–300.

Nuttley, W.M., R.K. Szilard, J.J. Smith, M. Veenhuis, and R.A. Rachubinski. 1995. The *PAH2* gene is required for peroxisome assembly in the methylotrophic yeast *Hansenula polymorpha* and encodes a member of the tetratricopeptide repeat family of proteins. *Gene (Amst.)*. 160:33–39.

Osumi, T., T. Tsukamoto, S. Hata, S. Yokota, S. Miura, Y. Fujiki, M. Hijikata, S. Miyazawa, and T. Hashimoto. 1991. Amino-terminal presequence of the precursor of peroxisomal 3-ketoacyl-CoA thiolase is a cleavable signal peptide for peroxisomal targeting. *Biochem. Biophys. Res. Commun.* 181:947–954.

Purdue, P.E., J.W. Zhang, M. Skoneczny, and P.B. Lazarow. 1997. Rhizomelic chondrodysplasia punctata is caused by deficiency of human *PEX7*, a homo-

- logue of the yeast PTS2 receptor. *Nat. Genet.* 15:381–384.
- Rehling, P., M. Marzioch, F. Niesen, E. Wittke, M. Veenhuis, and W.H. Kunau. 1996. The import receptor for the peroxisomal targeting signal 2 (PTS2) in *Saccharomyces cerevisiae* is encoded by the *PAS7* gene. *EMBO J.* 15:2901–2913.
- Ruigrok, C.C.M., H.F. Tabak, and I. Braakman. 1996. Import and oligomerization of peroxisomal proteins. *Mol. Biol. Cell.* 7:495.
- Schatz, G., and B. Dobberstein. 1996. Common principles of protein translocation across membranes. *Science.* 271:1519–1526.
- Schnell, D.J. 1995. Shedding light on the chloroplast protein import machinery. *Cell.* 83:521–524.
- Slawewski, M.L., G. Dodt, S. Steinberg, A.B. Moser, H.W. Moser, and S.J. Gould. 1995. Identification of three distinct peroxisomal protein import defects in patients with peroxisome biogenesis disorders. *J. Cell Sci.* 108:1817–1829.
- Subramani, S. 1997. *PEX* genes on the rise. *Nat. Genet.* 15:331–333.
- Swinkels, B.W., S.J. Gould, A.G. Bodnar, R.A. Rachubinski, and S. Subramani. 1991. A novel, cleavable peroxisomal targeting signal at the amino-terminus of the rat 3-ketoacyl-CoA thiolase. *EMBO J.* 10:3255–3262.
- Szilard, R.K., V.I. Titorenko, M. Veenhuis, and R.A. Rachubinski. 1995. Pay32p of the yeast *Yarrowia lipolytica* is an intraperoxisomal component of the matrix protein translocation machinery. *J. Cell Biol.* 131:1453–1469.
- Tsukamoto, T., S. Hata, S. Yokota, S. Miura, Y. Fujiki, M. Hijikata, T. Hashimoto, and T. Osumi. 1994. Characterization of the signal peptide at the amino terminus of the rat peroxisomal 3-ketoacyl-CoA thiolase precursor. *J. Biol. Chem.* 269:6001–6010.
- Van der Klei, I., R.E. Hilbrands, G.J. Swaving, H.R. Waterham, E.g., Vrieling, V.I. Titorenko, J.M. Cregg, W. Harder, and M. Veenhuis. 1995. The *Hansenula polymorpha* *PER3* gene is essential for the import of PTS1 proteins into the peroxisomal matrix. *J. Biol. Chem.* 270:17229–17236.
- Van der Leij, I., M. Van den Berg, R. Boot, M. Franse, B. Distel, and H.F. Tabak. 1992. Isolation of peroxisome assembly mutants from *Saccharomyces cerevisiae* with different morphologies using a novel positive selection procedure. *J. Cell Biol.* 119:153–162.
- Van der Leij, I., M.M. Franse, Y. Elgersma, B. Distel, and H.F. Tabak. 1993. PAS10 is a tetratricopeptide-repeat protein that is essential for the import of most matrix proteins into peroxisomes of *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA.* 90:11782–11786.
- Van der Voorn, L., and H.L. Ploegh. 1992. The WD-40 repeat. *FEBS Lett.* 307:131–134.
- Wiemer, E., W.M. Nuttley, B.L. Bertolaet, X. Li, U. Francke, M.J. Wheelock, U.K. Anné, K.R. Johnson, and S. Subramani. 1995. The human PTS1 receptor restores peroxisomal protein import deficiency in cells from patients with fatal peroxisomal disorders. *J. Cell Biol.* 130:51–65.
- Wiemer, E.A.C., G.H. Lüers, K.N. Faber, T. Wenzel, M. Veenhuis, and S. Subramani. 1996. Isolation and characterization of Pas2p, a peroxisomal membrane protein essential for peroxisome biogenesis in the methylotrophic yeast *Pichia pastoris*. *J. Biol. Chem.* 271:18973–18980.
- Zhang, J.W., and P.B. Lazarow. 1995. PEB1 (PAS7) in *Saccharomyces cerevisiae* encodes a hydrophilic, intra-peroxisomal protein that is a member of the WD repeat family and is essential for the import of thiolase into peroxisomes. *J. Cell Biol.* 129:65–80.
- Zhang, J.W., and P.B. Lazarow. 1996. Peb1p (Pas7p) is an intraperoxisomal receptor for the NH₂-terminal, type 2, peroxisomal targeting sequence of thiolase: Peb1p itself is targeted to peroxisomes by an NH₂-terminal peptide. *J. Cell Biol.* 132:325–334.