

# SCIENTIFIC REPORTS

OPEN

## Shortcuts to adiabatic passage for fast generation of Greenberger-Horne-Zeilinger states by transitionless quantum driving

Received: 19 June 2015

Accepted: 29 September 2015

Published: 28 October 2015

Ye-Hong Chen<sup>1</sup>, Yan Xia<sup>1</sup>, Jie Song<sup>2</sup> & Qing-Qin Chen<sup>3</sup>

Berry's approach on "transitionless quantum driving" shows how to set a Hamiltonian which drives the dynamics of a system along instantaneous eigenstates of a reference Hamiltonian to reproduce the same final result of an adiabatic process in a shorter time. In this paper, motivated by transitionless quantum driving, we construct shortcuts to adiabatic passage in a three-atom system to create the Greenberger-Horne-Zeilinger states with the help of quantum Zeno dynamics and of non-resonant lasers. The influence of various decoherence processes is discussed by numerical simulation and the result proves that the scheme is fast and robust against decoherence and operational imperfection.

"Shortcuts to adiabatic passage (STAP)"<sup>1,2</sup> which are a set of techniques to speed up a slow quantum adiabatic process usually through a non-adiabatic route, have attracted a great deal of attention in recent years. They can overcome the harmful effects caused by decoherence, noise or losses because of a long operation time. Quantum science also greatly desires fast and robust theoretical methods since high repetition rates contribute to the achievement of better signal-to-noise ratios and better accuracy. Therefore, in the last several years, STAP have been applied in a wide range of systems in theory and experiment<sup>3–23</sup>. Various reliable, fast and robust methods and schemes have been proposed to implement quantum information processing (QIP), such as fast population transfer<sup>5–7</sup>, fast entanglement generation<sup>6,8</sup>, fast implementation of quantum phase gates<sup>9</sup>.

To construct shortcuts to speed up adiabatic processes effectively, two methods which are in fact strongly related, and even potentially equivalent to each other<sup>24</sup>: are invariant-based inverse engineering based on Lewis-Riesenfeld invariant<sup>10,25</sup> and Berry's approach named "transitionless quantum driving" (TQD)<sup>26–29</sup>. Whereas, each of the two methods also has its own characteristics, for example, using Lewis-Riesenfeld invariants to construct shortcuts usually does not have to break down the form of the original Hamiltonian  $H_0(t)$ , so that the possibility of designing a Hamiltonian  $H(t)$  very difficult or impossible to implement in practice is avoided<sup>5,12</sup>. However, the invariants always have fixed forms which lead to that shortcut methods based on Lewis-Riesenfeld invariants might be limited or even hopeless in some cases to construct shortcuts to implement QIP rapidly<sup>5</sup>. For example, in the paper<sup>8</sup> proposed by Chen *et al.*, they had no choice but to make one of the atoms to be a control qubit or use auxiliary levels for the atoms to generate entangled states.

There is still plenty to do to make wide applications of STAP for fast QIP in some experimental systems, for example, the cavity quantum electronic dynamics (QED) systems. It is worth noting that, TQD provides a very effective method to construct the "counter-diabatic driving" (CDD) Hamiltonian  $H(t)$  which accurately drives the instantaneous eigenstates of  $H_0(t)$ . Nevertheless, it is almost always found that the designed CDD Hamiltonian is hard to be directly implemented in practice<sup>2,30–34</sup>, especially in

<sup>1</sup>Department of Physics, Fuzhou University, Fuzhou 350002, China. <sup>2</sup>Department of Physics, Harbin Institute of Technology, Harbin 150001, China. <sup>3</sup>Zhicheng College, Fuzhou University, Fuzhou 350002, China. Correspondence and requests for materials should be addressed to Y.X. (email: xia-208@163.com) or J.S. (email: jsong@hit.edu.cn)

multiparticle systems. Examples of ways to overcome this problem may be found in refs 34–38. Also, in a large detuning limit, Lu *et al.*<sup>6</sup> have found a simplified effective Hamiltonian equivalent to  $H(t)$ . This idea inspires us that finding an alternative physically feasible (APF) Hamiltonian which is effectively equivalent to  $H(t)$ . However, the approximation in ref. 6 is too complex to be generalized to  $N$ -qubit entanglement cases. It is known to all that, entanglement of more qubits shows more nonclassical effects and is more useful for quantum applications. For example, one of the two kinds of three-qubit entangled states named the Greenberger-Horne-Zeilinger (GHZ) states provide a possibility for testing quantum mechanics against local hidden theory without using Bell's inequality<sup>39,40</sup>. Therefore, great interest has arisen regarding the significant role of the GHZ states in the foundations of quantum mechanics measurement theory and quantum communication. In view of that we wonder if it is possible to use TQD to construct shortcuts for one-step generation of multi-qubit entanglement, i.e., the three-atom GHZ states, without abandoning any of the atoms or using auxiliary levels.

In this scenario, motivated by refs 5–8, we use TQD to construct STAP to generate the three-atom GHZ states effectively and rapidly in one step. It would be a promising idea of applying STAP to realize multi-qubit entanglement generation in cavity QED systems. Different from ref. 6, we use the quantum Zeno dynamics<sup>41,42</sup> to simplify the system first and then under the large detuning condition, we obtain the effective Hamiltonian which is equivalent to the corresponding CDD Hamiltonian to speed up the evolution process. Therefore, the adiabatic process for a multi-qubit system is speeded up, and the STAP is easy to be achieved in experiment. Comparing with ref. 8, we use TQD in this paper so that the laser pulses are not strongly limited and we do not need to use auxiliary levels or multi-step operations to generate the three-atom GHZ states. Moreover, we find that any quantum system whose Hamiltonian is possible to be simplified into the form in eq. (15), the corresponding APF Hamiltonian can be built and the STAP can be constructed with the same approach presented in this paper. The above advantages mean the present scheme is much more useful in dealing with the fast and noise-resistant generation of multi-qubit entanglement or even other QIP.

### Basic theories

**Transitionless quantum driving.** Consider an arbitrary time-dependent Hamiltonian  $H_0(t)$ , with instantaneous eigenstates and energies given by

$$H_0(t)|\varphi_n(t)\rangle = \zeta_n(t)|\varphi_n(t)\rangle. \quad (1)$$

When this system satisfies the adiabatic condition,  $H_0(t)$  will drive the system into

$$|\psi(t)\rangle = e^{i\vartheta_n(t)}|\varphi_n(t)\rangle, \quad (2)$$

where

$$\vartheta_n(t) = -\frac{1}{\hbar} \int_0^t dt' \zeta_n(t') + i \int_0^t dt' \langle \varphi_n(t') | \partial_{t'} \varphi_n(t') \rangle. \quad (3)$$

To find the Hamiltonian  $H(t)$  that drives the eigenstates  $\{|\varphi_n(t)\rangle\}$ , we define a unitary operator

$$U = \sum_n e^{i\vartheta_n(t)} |\varphi_n(t)\rangle \langle \varphi_n(0)|, \quad (4)$$

which obeys

$$i\hbar \partial_t U = H(t) U \Rightarrow H(t) = i\hbar (\partial_t U) U^\dagger. \quad (5)$$

Then the Hamiltonian  $H(t)$  is obtained

$$H(t) = H_0(t) + H_1(t), \quad H_1(t) = i\hbar \sum_n (|\partial_t \varphi_n\rangle \langle \varphi_n| - \langle \varphi_n | \partial_t \varphi_n \rangle |\varphi_n\rangle \langle \varphi_n|). \quad (6)$$


The simplest choice is  $\zeta_n = 0$ , for which the bare states  $|\varphi_n(t)\rangle$ , with no phase factors, are driven by<sup>26</sup>

$$H(t) = i\hbar \sum_n |\partial_t \varphi_n\rangle \langle \varphi_n|, \quad (7)$$

reflecting

$$i\hbar \partial_t |\varphi_n\rangle = i\hbar \sum_m |\partial_t \varphi_m\rangle \langle \varphi_m | \varphi_n \rangle. \quad (8)$$

**Quantum Zeno dynamics.** The quantum Zeno dynamics was named by Facchi and Pascazio in 2002<sup>42</sup>. It is derived from the quantum Zeno effect which describes a phenomenon that the system can actually evolve away from its initial state while it still remains in the so-called Zeno subspace determined



**Figure 1.** The cavity-atom combined system and the atomic level configuration for the original Hamiltonian.

by the measurement when frequently projected onto a multidimensional subspace. According to von Neumann's projection postulate, the quantum Zeno dynamics can be achieved via continuous coupling between the system and an external system instead of discontinuous measurements<sup>42</sup>. In general, we assume that a dynamical evolution process is governed by the Hamiltonian

$$H_Z = H_{obs} + KH_{meas}, \quad (9)$$

where  $H_{obs}$  is the Hamiltonian of the quantum system investigated,  $K$  is a coupling constant, and  $H_{meas}$  is viewed as an additional interaction Hamiltonian performing the measurement. In the "infinitely strong measurement" limit  $K \rightarrow \infty$ <sup>41,42</sup>, The Hamiltonian for the whole system is nearly equivalent to

$$H_{Zeno} = \sum_n (P_n H_{obs} P_n + \varepsilon_n P_n), \quad (10)$$

with  $P_n$  being the  $n$ th orthogonal projection onto the invariant Zeno subspace  $\forall P_n$  and the eigenspace of  $KH_{meas}$  belonging to the eigenvalue  $\varepsilon_n$ , i.e.,  $KH_{meas}P_n = \varepsilon_n P_n$ .

### Model

We consider three  $\Lambda$ -type atoms are trapped in a bimodal-mode cavity as shown in Fig. 1. Atoms 1, 2, and 3 have three sets of ground states  $\{|f\rangle_1, |g_l\rangle_1\}$ ,  $\{|g_l\rangle_2, |g_r\rangle_2\}$ , and  $\{|f\rangle_3, |g_r\rangle_3\}$ , respectively, and each of them has an excited state  $|e\rangle$ . The atomic transition  $|f\rangle \leftrightarrow |e\rangle$  is driven resonantly through classical laser field with time-dependent Rabi frequency  $\Omega(t)$ , transition  $|g_l\rangle \leftrightarrow |e\rangle$  is coupled resonantly to the left-circularly polarized mode of the cavity with coupling  $\lambda_l$ , and transition  $|g_r\rangle \leftrightarrow |e\rangle$  is coupled resonantly to the right-circularly polarized mode of the cavity with coupling  $\lambda_r$ . Under the rotating-wave approximation (RWA), the interaction Hamiltonian for this system reads ( $\hbar = 1$ ):

$$\begin{aligned} H_I &= H_{AL} + H_{AC}, \\ H_{AL} &= \Omega_1(t)|e\rangle_1\langle f| + e^{i\beta}\Omega_3(t)|e\rangle_3\langle f| + H.c., \\ H_{AC} &= \sum_{k=1,2} \lambda_l a_l |e\rangle_k \langle g_l| + \sum_{j=2,3} \lambda_r a_r |e\rangle_j \langle g_r| + H.c., \end{aligned} \quad (11)$$

where  $H_{AL}$  denotes the coupling between the atoms and the laser pulses, and  $H_{AC}$  denotes the coupling between the atoms and the cavities,  $a_l$  and  $a_r$  are the left- and right-circularly annihilation operators of the cavity modes, and  $\beta$  means the two Rabi frequencies are  $\beta$ -dephased from each other. If we assume the initial state is  $|f, g_l, g_r\rangle_{1,2,3} |0,0\rangle_c$ , the system will evolve within a single-excitation subspace with basis states

$$\begin{aligned}
 |\psi_1\rangle &= |f, g_l, g_r\rangle_{1,2,3} |0,0\rangle_c, \\
 |\psi_2\rangle &= |e, g_l, g_r\rangle_{1,2,3} |0,0\rangle_c, \\
 |\psi_3\rangle &= |g_l, g_l, g_r\rangle_{1,2,3} |1,0\rangle_c, \\
 |\psi_4\rangle &= |g_l, e, g_r\rangle_{1,2,3} |0,0\rangle_c, \\
 |\psi_5\rangle &= |g_l, g_r, g_r\rangle_{1,2,3} |0,1\rangle_c, \\
 |\psi_6\rangle &= |g_l, g_r, e\rangle_{1,2,3} |0,0\rangle_c, \\
 |\psi_7\rangle &= |g_l, g_r, f\rangle_{1,2,3} |0,0\rangle_c.
 \end{aligned} \tag{12}$$

In light of quantum Zeno dynamics, we rewrite the Hamiltonian  $H_I$  in eq. (11) as  $H_{re}$  through the relation  $H_{re} = \sum_n P_n H_{AL} P_n + \varepsilon_n P_n (H_{AL} \rightarrow H_{obs} \text{ and } H_{AC} \rightarrow KH_{meas})$ , where

$$P_n = \sum_m |m\rangle \langle m|, |m\rangle \in \{|\psi_1\rangle, |\psi_7\rangle, |\phi_0\rangle, |\phi_1\rangle, |\phi_2\rangle, |\phi_3\rangle, |\phi_4\rangle\}. \tag{13}$$

Here  $|\phi_0\rangle, |\phi_1\rangle, |\phi_2\rangle, |\phi_3\rangle$ , and  $|\phi_4\rangle$  are the eigenvectors of  $H_{AC}$  corresponding eigenvalues  $\varepsilon_0 = 0$ ,  $\varepsilon_1 = \lambda$ ,  $\varepsilon_2 = -\lambda$ ,  $\varepsilon_3 = \sqrt{3}\lambda$ , and  $\varepsilon_4 = -\sqrt{3}\lambda$ , respectively. And we obtain (we set  $\lambda_l = \lambda_r = \lambda$ )

$$\begin{aligned}
 H_{re} &= \sum_{k=0}^4 \varepsilon_k |\phi_k\rangle \langle \phi_k| + H_{AL}^r, \\
 H_{AL}^r &= \frac{1}{\sqrt{3}} [|\phi_0\rangle + \frac{1}{2}(|\phi_3\rangle + |\phi_4\rangle)] (\Omega_1 \langle \psi_1| + e^{i\beta} \Omega_3 \langle \psi_7|) \\
 &\quad + \frac{1}{2}(|\phi_1\rangle + |\phi_2\rangle) (-\Omega_1 \langle \psi_1| + e^{i\beta} \Omega_3 \langle \psi_7|) + H.c..
 \end{aligned} \tag{14}$$

Through performing the unitary transformation  $U_Z = e^{-i \sum \varepsilon_k |\phi_k\rangle \langle \phi_k| t}$  and neglecting the terms with high oscillating frequency by setting the condition  $\Omega_1/\sqrt{3}, \Omega_3/\sqrt{3} \ll \lambda$  (the Zeno condition), we obtain an effective Hamiltonian

$$H_{eff} = \frac{1}{\sqrt{3}} |\phi_0\rangle (\Omega_1(t) \langle \psi_1| + e^{i\beta} \Omega_3(t) \langle \psi_7|) + H.c., \tag{15}$$


which can be seen as a simple three-level system with an excited state  $|\phi_0\rangle$  and two ground states  $|\psi_1\rangle$  and  $|\psi_7\rangle$ . For this effective Hamiltonian, its eigenstates are easily obtained

$$\begin{aligned}
 |n_0(t)\rangle &= \begin{pmatrix} \cos \theta(t) \\ 0 \\ -e^{i\beta} \sin \theta(t) \end{pmatrix}, \\
 |n_{\pm}(t)\rangle &= \frac{1}{\sqrt{2}} \begin{pmatrix} \sin \theta(t) \\ \pm 1 \\ e^{i\beta} \cos \theta(t) \end{pmatrix},
 \end{aligned} \tag{16}$$

corresponding eigenvalues  $\eta_0 = 0$ ,  $\eta_{\pm} = \pm \Omega/\sqrt{3}$ , respectively, where  $\tan \theta = \Omega_1/\Omega_3$  and  $\Omega = \sqrt{\Omega_1^2 + \Omega_3^2}$ . When the adiabatic condition  $|\langle n_0 | \partial_t n_{\pm} \rangle| \ll |\eta_{\pm}|$  is fulfilled, the initial state  $|\psi_1\rangle = |n_0(0)\rangle$  will follow  $|n_0(t)\rangle$  closely, and when  $\theta(t) = \pi/4$  and  $\beta = l\pi/2$  ( $l = 0, \pm 1, \pm 2, \dots$ ), we obtain the GHZ states:  $|\psi(t_f)\rangle = |GHZ\rangle = (|\psi_1\rangle - e^{i\beta} |\psi_7\rangle)/\sqrt{2}$ . When  $\beta = \pi$ , it shows the most common form:  $|\psi(t_f)\rangle = (|\psi_1\rangle + |\psi_7\rangle)/\sqrt{2}$ . However, this process will take quite a long time to obtain the target state, which is undesirable.

### Using TQD to construct shortcuts to adiabatic passage

The instantaneous eigenstates  $|n_k\rangle$  ( $k = 0, \pm$ ) for the effective Hamiltonian  $H_{eff}(t)$  above do not satisfy the Schrödinger equation  $i\partial_t |n_k\rangle = H_{eff}(t) |n_k\rangle$ . According to Berry's general transitionless tracking algorithm<sup>26</sup>, from  $H_{eff}(t)$ , one can reverse engineer  $H(t)$  which is related to the original Hamiltonian  $H_{eff}(t)$  but drives the eigenstates exactly. From refs 6,11,12, we learn the simplest Hamiltonian  $H(t)$  is derived in the form



**Figure 2.** The atomic level configuration for the APF Hamiltonian.

$$H(t) = i \sum_{k=0,\pm} |\partial_t n_k(t)\rangle \langle n_k(t)|. \quad (17)$$

Substituting eq. (16) in eq. (17), we obtain

$$H(t) = i\partial_t e^{i\beta} |\psi_7\rangle \langle \psi_1| + H.c., \quad (18)$$

where  $\partial_t = [\dot{\Omega}_1(t)\Omega_3(t) - \dot{\Omega}_3(t)\Omega_1(t)]/\Omega^2$ . Similar to ref. 6, for this three-atom system in a real experiment, the Hamiltonian  $H(t)$  is hard or even impossible to be implemented in practice. We should find an APF Hamiltonian whose effect is equivalent to  $H(t)$ . The model used for the APF Hamiltonian is similar to that in Fig. 1 with three atoms trapped in a cavity, and the atomic level configuration is shown in Fig. 2: the transition  $|f\rangle \leftrightarrow |e\rangle$  is non-resonantly driven by classical field with time-dependent Rabi frequency  $\tilde{\Omega}$  and detuning  $\Delta$ , the transition  $|g_l\rangle \leftrightarrow |e\rangle$  is coupled non-resonantly to the cavity with coupling  $\lambda_l$  ( $\lambda_r$ ) and detuning  $\Delta$ . The rotating-frame Hamiltonian reads

$$\begin{aligned} H'_I &= H'_{AL} + H'_{AC} + H_e, \\ H'_{AL} &= \tilde{\Omega}_1(t)|e\rangle_1 \langle f| + e^{i\beta'} \tilde{\Omega}_3(t)|e\rangle_3 \langle f| + H.c., \\ H'_{AC} &= \sum_{m=1,2} \lambda_l a_l |e\rangle_m \langle g_l| + \sum_{n=2,3} \lambda_r a_r |e\rangle_n \langle g_r| + H.c., \\ H_e &= \sum_{k=1}^3 \Delta |e\rangle_k \langle e|, \end{aligned} \quad (19)$$

where  $\beta'$  is the phase difference between  $\tilde{\Omega}_1$  and  $\tilde{\Omega}_3$ . Then similar to the approximation for the Hamiltonian from eq. (11) to eq. (15), we also obtain an effective Hamiltonian for the present non-resonant system<sup>43</sup>

$$H'_{eff} = [\frac{1}{\sqrt{3}}|\phi_0\rangle \langle \tilde{\Omega}_1(t)\langle \psi_1| + e^{i\beta'} \tilde{\Omega}_3(t)\langle \psi_7| + H.c.] + \Delta|\phi_0\rangle \langle \phi_0|. \quad (20)$$


By adiabatically eliminating the state  $|\phi_0\rangle$  under the condition  $\Delta \gg \tilde{\Omega}_1/\sqrt{3}, \tilde{\Omega}_3/\sqrt{3}$ , we obtain the final effective Hamiltonian

$$\begin{aligned} H_{fe} &= -\frac{\tilde{\Omega}_1^2}{3\Delta} |\psi_1\rangle \langle \psi_1| - \frac{\tilde{\Omega}_3^2}{3\Delta} |\psi_7\rangle \langle \psi_7| \\ &\quad - \frac{e^{i\beta'} \tilde{\Omega}_1 \tilde{\Omega}_3}{3\Delta} |\psi_7\rangle \langle \psi_1| - \frac{e^{-i\beta'} \tilde{\Omega}_1 \tilde{\Omega}_3}{3\Delta} |\psi_1\rangle \langle \psi_7|. \end{aligned} \quad (21)$$

Choosing  $\tilde{\Omega}_1 = \tilde{\Omega}_3 = \tilde{\Omega}(t)$ , the first two terms of eq. (21) can be removed, and the Hamiltonian becomes

$$\tilde{H}_{eff} = e^{i\beta'} \Omega_x(t) |\psi_7\rangle \langle \psi_1| + e^{-i\beta'} \Omega_x(t) |\psi_1\rangle \langle \psi_7|, \quad (22)$$

where  $\Omega_x(t) = -\tilde{\Omega}^2/(3\Delta)$ . This effective Hamiltonian is equivalent to the CDD Hamiltonian  $H(t)$  in eq. (18) when



**Figure 3.** Dependence on  $t/t_f$  of  $\Omega_1/\Omega_0$  and  $\Omega_3/\Omega_0$ .

$$e^{i\beta'}\Omega_x = ie^{i\beta}\dot{\theta}. \quad (23)$$

Hence, the Rabi frequencies for the APF Hamiltonian are designed

$$\begin{aligned} \tilde{\Omega}_1 &= \tilde{\Omega}_3 = \sqrt{-3\Delta\dot{\theta}}, \\ \beta' - \beta &= \frac{\pi}{2} + 2l\pi, \end{aligned} \quad (24)$$

where  $l = 0, \pm 1, \pm 2, \dots$ .

### Fast and noise-resistant generation of the three-atom GHZ states with STAP

We will show that the creation of a three-atom GHZ state governed by  $H_I'$  is much faster than that governed by  $H_I$ . To satisfy the boundary condition of the fractional stimulated Raman adiabatic passage (STIRAP),


$$\lim_{t \rightarrow -\infty} \frac{\Omega_1(t)}{\Omega_3(t)} = 0, \quad \lim_{t \rightarrow +\infty} \frac{\Omega_3(t)}{\Omega_1(t)} = \tan \alpha, \quad (25)$$

the Rabi frequencies  $\Omega_1(t)$  and  $\Omega_3(t)$  in the original Hamiltonian  $H_I(t)$  are chosen as


$$\begin{aligned} \Omega_1(t) &= \sin \alpha \Omega_0 \exp \left[ \frac{-(t - t_0 - t_f/2)^2}{t_c^2} \right], \\ \Omega_3(t) &= \Omega_0 \exp \left[ \frac{-(t + t_0 - t_f/2)^2}{t_c^2} \right] \\ &\quad + \cos \alpha \Omega_0 \exp \left[ \frac{-(t - t_0 - t_f/2)^2}{t_c^2} \right], \end{aligned} \quad (26)$$

where  $\Omega_0$  is the pulse amplitude,  $t_f$  is the operation time, and  $t_0, t_c$  are some related parameters. In order to create a three-atom GHZ state, the final state  $|\psi(t_f)\rangle$  should be  $|\psi(t_f)\rangle = \frac{1}{\sqrt{2}}(|\psi_1\rangle - e^{i\beta}|\psi_7\rangle)$  according to eq. (16). Therefore, we have  $\tan \alpha = 1$ . By choosing parameters for the laser pulses suitably to fulfill the boundary condition in eq. (25), the time-dependent  $\Omega_1(t)$  and  $\Omega_3(t)$  are gotten as shown in Fig. 3 with parameters  $t_0 = 0.14t_f$  and  $t_c = 0.19t_f$ . For simplicity, we set  $\beta = 0$  in the following discussion. Fig. 4 shows the relationship between the fidelity of the generated three-atom GHZ state (governed by the APF Hamiltonian  $H_I'(t)$ ) and two parameters  $\Delta$  and  $t_f$  when  $\Omega_0 = 0.2\lambda$  satisfying the Zeno condition, where the fidelity for the three-atom GHZ state is given through  $F = |\langle GHZ | \rho(t_f) | GHZ \rangle|$  ( $\rho(t_f)$  is the density operator of the whole system when  $t = t_f$ ). We find that there is a wide range of selectable values for parameters  $\Delta$  and  $t_f$  to get a high fidelity of the three-atom GHZ state. The fidelity increases with the increasing of  $t_f$  while decreases with the increasing of  $\Delta$ . It is not hard to understand, putting eq. (26) into eq. (24) and setting  $t = t' \times t_f$ , we can find

$$\Omega'_0 \approx \sqrt{\frac{6\Delta}{t_f}}, \quad (27)$$



**Figure 4.** The fidelity  $F$  of the target state  $\frac{1}{\sqrt{2}}(|\psi_1\rangle - |\psi_7\rangle)$  versus the interaction time  $\lambda t_f$  and the detuning  $\Delta/\lambda$ .



**Figure 5.** Time evolution of the populations for the states  $|\psi_1\rangle$  and  $|\psi_7\rangle$  with  $\Omega_0 = 0.2\lambda$ ,  $t_f = 35/\lambda$  and  $\Delta = 2.2\lambda$ , (a) governed by the APF Hamiltonian  $H'_I(t)$ , (b) governed by the original Hamiltonian  $H_I(t)$ .


where  $\Omega'_0$  is the amplitude of  $\tilde{\Omega}(t')$ . That means, in order to satisfy the Zeno condition  $\tilde{\Omega} \ll \sqrt{3}\lambda$  and the large detuning condition  $\tilde{\Omega} \ll \sqrt{3}\Delta$ , the ratio  $\Delta/t_f$  should be small enough. Moreover, this relationship also explains the phenomenon in Fig. 4 that to achieve a high fidelity with a larger detuning  $\Delta$ , a longer interaction time  $t_f$  is required. Then to prove the operation time required for the creation of the three-atom GHZ state governed by  $H'_I$  is much shorter than that governed by  $H_I$ , we contrast the performances of population transfer from the initial state  $|\psi_1\rangle$  governed by the APF Hamiltonian  $H'_I$  and that governed by the original Hamiltonian  $H_I$  in Fig. 5 with  $\{t_f = 35/\lambda, \Omega_0 = 0.2\lambda, \Delta = 2.2\lambda\}$ . The time-dependent population for any state  $|\psi\rangle$  is given by the relationship  $P = |\langle\psi|\rho(t)|\psi\rangle|$ , where  $\rho(t)$  is the corresponding time-dependent density operator. The comparison of Fig. 5(a,b) shows that with this set of parameters, the APF Hamiltonian  $H'_I(t)$  can govern the evolution to achieve a near-perfect three-atom GHZ state from state  $|\psi_1\rangle$  in short interaction time while the original Hamiltonian  $H_I(t)$  can not. In fact, through solving the adiabatic condition  $|\langle n_0|\partial_t n_{\pm}\rangle| \ll |\eta_{\pm}|$ , we obtain

$$\left| \frac{\dot{\theta}}{\sqrt{2}} \right| \ll \left| \frac{\Omega}{\sqrt{3}} \right| \Rightarrow \frac{f(t)}{t_f} \ll \Omega, \quad (28)$$


where  $f(t)$  is a wave function whose amplitude is irrelevant to  $t_f$ . The result shows when  $\Omega_0$  is a constant, the longer the operation time  $t_f$  is, the better the adiabatic condition is satisfied. This is proved in Fig. 6. Figure 6 reveals the relationship between  $G(t_f)$  and  $\lambda t_f$ , where  $G(t_f) = \left. \frac{\sqrt{3}|\dot{\theta}|}{\sqrt{2}\Omega} \right|_{t=0.5t_f}$ . From this figure, we

discover that even with  $\Omega_0 = 0.5\lambda$  which does not meet the Zeno condition, the operation time required for the three-atom GHZ state generation in an adiabatic system is longer than  $100/\lambda$  (when  $t_f = 100/\lambda$ ,  $G(t_f) \approx 0.08$ ). We also plot the fidelities of the evolved states governed by  $H'_I(t)$  and  $H_I(t)$  (in different cases) in Fig. 7, with respect to the target three-atom GHZ state. Shown in the figure, even with a large laser intensity, say,  $\Omega_0 = 0.5\lambda$ , the interaction time required for creation of the three-atom GHZ state via adiabatic passage is still much longer than that via STAP. Generally speaking, the adiabatic condition is satisfied much better with a relatively larger laser intensity, while, the system would be very sensitive to





**Figure 6.** The relationship between  $G(t_f)$  and  $\lambda t_f$  for testing the adiabatic condition.



**Figure 7.** The comparison between the fidelities of the three-atom GHZ state governed by the APF Hamiltonian  $H'_I(t)$  and the original Hamiltonian  $H_I(t)$ .

the decoherence caused by the cavity decay with a relatively large laser intensity. This will be proved in the following.

Once the dissipation is considered, the evolution of the system can be modeled by a master equation in Lindblad form,


$$\dot{\rho} = i[\rho, H] + \sum_k [L_k \rho L_k^\dagger - \frac{1}{2}(L_k^\dagger L_k \rho + \rho L_k^\dagger L_k)], \quad (29)$$

where  $L_k$ 's are the Lindblad operators. For both the resonant and non-resonant systems, there are eight Lindblad operators governing the dissipation:


$$\begin{aligned} L_1^\kappa &= \sqrt{\kappa_l} a_l, & L_2^\kappa &= \sqrt{\kappa_r} a_r, \\ L_3^\gamma &= \sqrt{\gamma_1} |f\rangle_1 \langle e|, & L_4^\gamma &= \sqrt{\gamma_2} |g_l\rangle_1 \langle e|, \\ L_5^\gamma &= \sqrt{\gamma_3} |g_l\rangle_2 \langle e|, & L_6^\gamma &= \sqrt{\gamma_4} |g_r\rangle_2 \langle e|, \\ L_7^\gamma &= \sqrt{\gamma_5} |f\rangle_3 \langle e|, & L_8^\gamma &= \sqrt{\gamma_6} |g_r\rangle_3 \langle e|, \end{aligned} \quad (30)$$

where  $\kappa_l$  and  $\kappa_r$  are the decays of the cavity modes, and  $\gamma_n$  ( $n = 1, 2, \dots, 6$ ) are the spontaneous emissions of atoms. For simplicity, we assume  $\kappa_l = \kappa_r = \kappa$ , and  $\gamma_n = \gamma/2$ . Figure 8(a) shows the fidelity of the three-atom GHZ state governed by the APF Hamiltonian  $H'_I$  versus these two noise resources with  $\Omega_0 = 0.2\lambda$ ,  $\Delta = 2.2\lambda$ , and  $t_f = 35/\lambda$ . It turns out that the present shortcut scheme with this set of parameters is much more sensitive to the cavity decays than the spontaneous emissions. Ref. 5 contributes to understanding this phenomenon, in fact, with this set of parameters, the Zeno condition for the non-resonant system is not ideally fulfilled because shortening the time implies an energy cost<sup>12,24</sup> (in this system, the energy cost denotes requiring relative-large laser intensities). Known from ref. 5, destroying the Zeno condition slightly is also helpful to achieve the target state in a much shorter interaction time. However, if the Zeno condition has not been satisfied very well, the intermediate states including the cavity-excited states would be populated during the evolution, which causes that the system is sensitive to the cavity decays. However, we can find in Fig. 8(b) which shows fidelity of the three-atom GHZ





**Figure 8.** (a) Dependences on  $\kappa/\lambda$  and  $\gamma/\lambda$  of the fidelity of the three-atom GHZ state governed by the APF Hamiltonian  $H'_I(t)$  when  $t_f = 35/\lambda$ ,  $\Delta = 2.2\lambda$  and  $\Omega_0 = 0.2\lambda$ . (b) Dependences on  $\kappa/\lambda$  and  $\gamma/\lambda$  of the fidelity of the three-atom GHZ state governed by the original Hamiltonian  $H_I(t)$  when  $t_f = 100/\lambda$  and  $\Omega_0 = 0.5\lambda$ .



**Figure 9.** The fidelity  $F$  of the GHZ state versus the variations of (a)  $T$  and  $\Omega'_0$ . (b)  $\lambda$  and  $\Delta$ .

state governed by original Hamiltonian  $H_I$  with  $\Omega_0 = 0.5\lambda$  and  $t_f = 100/\lambda$  in the presence of decoherence, with large laser intensities, the adiabatic scheme is also sensitive to the cavity decays as we mentioned above. The comparison of these two figures drops a result that the present shortcut scheme is almost the same with the adiabatic one in restraining the decoherence.

The robustness against operational imperfection is also a main factor for the feasibility of the scheme because most of the parameters are hard to accurately achieve in experiment. Therefore, we define  $\delta x = x' - x$  as the deviation of any parameter  $x$ , where  $x'$  is the actual value and  $x$  is the ideal value. Then in Fig. 9(a) we plot the fidelity of the GHZ state versus the variations in total operation time  $T$  ( $T = 1.2t_f$ ) and laser amplitude  $\Omega'_0$ , and in Fig 9(b) we plot the fidelity of the GHZ state versus the variations in coupling  $\lambda$  and detuning  $\Delta$ . As shown in the figures, the scheme is robust against all of these variations. Any deviation  $\delta x/x = 10\%$  ( $x \in \{T, \Omega'_0, \lambda, \Delta\}$ ) causes a reduction less than 3% in the fidelity.

In a real experiment, the cesium atoms which have been cooled and trapped in a small optical cavity in the strong-coupling regime<sup>44,45</sup> can be used in this scheme. We take the hyperfine states of  $6^2P_{1/2}$  as the excited states and the hyperfine states of  $6^2S_{1/2}$  as the ground states. With a set of cavity QED parameters  $\lambda = 750 \times 2\pi$  MHz,  $\kappa = 3.5 \times 2\pi$  MHz, and  $\gamma = 2.62 \times 2\pi$  MHz in strong-coupling regime<sup>46–48</sup>, the fidelity of the three-atom GHZ state in this paper is 98.24%. Thus, the scheme is robust and might be promising within the limits of current technology.

## Conclusion

We have presented a promising method to construct shortcuts to adiabatic passage (STAP) for a three-atom system to generate GHZ states in the cavity QED system. Through using quantum Zeno dynamics and “transitionless quantum driving”, we are free to simplify a complicated Hamiltonian and choose the laser pulses to construct shortcuts in multi-qubit system to implement the fast quantum information processing. Numerical simulation demonstrates that the scheme is fast and robust against the decoherence caused by both atomic spontaneous emission, photon leakage and operational imperfection. The deficiency is that the present scheme might be sensitive to the cavity decays because of some inevitable factors. Compared with the previous shortcut methods, this method obviously works better at entanglement generation in multi-qubit systems. In fact, any quantum system whose Hamiltonian is possible to be simplified into the form in eq. (15), the shortcut can be constructed with the same method presented in this paper. For example, similar to refs 49,50 for the generation of the multiparticle GHZ states in an atom-fiber-cavity combined system, we can shorten the operation time using the same method in the following steps: (1) We consider the Hamiltonian  $H_{AL}$  (the Hamiltonian describing the interaction between atoms and lasers) as  $H_{obs}$  in eq. (9), and the rest of the total Hamiltonian (the interaction between atoms and cavities, and the interaction between cavities and fibers) as  $KH_{meas}$  in eq. (9). Then, choosing the “dark Zeno subspace”, that means  $\varepsilon_n = 0$  in eq. (10), we can obtain an effective Hamiltonian named the Zeno Hamiltonian  $H_{Zeno} = \sum_n P_n H_{obs} P_n$ . (2) For this effective Hamiltonian, by using TQD, we construct the CDD Hamiltonian  $H(t)$  that speeds up the adiabatic process. (3) Similar to the GHZ state generation, we find out the corresponding non-resonant system (the APF Hamiltonian) whose effective Hamiltonian  $\tilde{H}_{eff}(t)$  has the form in eq. (22). (4) Making  $\tilde{H}_{eff}(t) = H(t)$ , the parameters for the APF Hamiltonian are determined and the shortcut is constructed. Then the APF Hamiltonian would govern the system to achieve the same final result as the adiabatic process governed by the original Hamiltonian with a much shorter operation time. Similar idea can be generalized to generate other multi-qubit quantum entangled states, for example, Bell states,  $W$  states, singlet states, and so on. This might lead to a useful step toward realizing fast and noise-resistant quantum information processing for multi-qubit systems in current technology.

## References

- Chen, X., Lizuain, I., Ruschhaupt, A., Guéry-Odelin, D. & Muga, J. G. Shortcut to adiabatic passage in two- and three-level atoms. *Phys. Rev. Lett.* **105**, 123003 (2010).
- Torrontegui, E. *et al.* Chapter 2 - Shortcuts to Adiabaticity. *Adv. Atom. Mol. Opt. Phys.* **62**, 117–169 (2013).
- del Campo, A. Shortcuts to adiabaticity by counterdiabatic driving. *Phys. Rev. Lett.* **111**, 100502 (2013).
- Masuda, S. & Nakamura, K. Acceleration of adiabatic quantum dynamics in electromagnetic fields. *Phys. Rev. A* **84**, 043434 (2011).
- Chen, Y. H., Xia, Y., Chen, Q. Q. & Song, J. Efficient shortcuts to adiabatic passage for fast population transfer in multiparticle systems. *Phys. Rev. A* **89**, 033856 (2014).
- Lu, M., Xia, Y., Shen, L. T., Song, J. & An, N. B. Shortcuts to adiabatic passage for population transfer and maximum entanglement creation between two atoms in a cavity. *Phys. Rev. A* **89**, 012326 (2014).
- Lu, M., Xia, Y., Shen, L. T. & Song, J. An effective shortcut to adiabatic passage for fast quantum state transfer in a cavity quantum electronic dynamics system. *Laser Phys.* **24**, 105201 (2014).
- Chen, Y. H., Xia, Y., Chen, Q. Q. & Song, J. Shortcuts to adiabatic passage for multiparticles in distant cavities: applications to fast and noise-resistant quantum population transfer, entangled states' preparation and transition. *Laser Phys. Lett.* **11**, 115201 (2014).
- Chen, Y. H., Xia, Y., Chen, Q. Q. & Song, J. Fast and noise-resistant implementation of quantum phase gates and creation of quantum entangled states. *Phys. Rev. A* **91**, 012325 (2015).
- Muga, J. G., Chen, X., Ruschhaupt, A. & Guéry-Odelin, D. Frictionless dynamics of Bose-Einstein condensates under fast trap variations. *J. Phys. B* **42**, 241001 (2009).
- Chen, X. *et al.* Fast optimal frictionless atom cooling in harmonic traps: Shortcut to adiabaticity. *Phys. Rev. Lett.* **104**, 063002 (2010).
- Chen, X. & Muga, J. G. Transient energy excitation in shortcuts to adiabaticity for the time-dependent harmonic oscillator. *Phys. Rev. A* **82**, 053403 (2010).
- Schaff, J. F., Capuzzi, P., Labeyrie, G. & Vignolo, P. Shortcuts to adiabaticity for trapped ultracold gases. *New J. Phys.* **13**, 113017 (2011).
- Torrontegui, E. *et al.* Fast atomic transport without vibrational heating. *Phys. Rev. A* **83**, 013415 (2011).
- Chen, X., Torrontegui, E., Stefanatos, D., Li, J. S. & Muga, J. G. Optimal trajectories for efficient atomic transport without final excitation. *Phys. Rev. A* **84**, 043415 (2011).
- Torrontegui, E. *et al.* Fast transport of Bose-Einstein condensates. *New J. Phys.* **14**, 013031 (2012).
- Li, Y., Wu, L. A. & Wang, Z. D. Fast ground-state cooling of mechanical resonators with time-dependent optical cavities. *Phys. Rev. A* **83**, 043804 (2011).
- del Campo, A. Frictionless quantum quenches in ultracold gases: A quantum-dynamical microscope. *Phys. Rev. A* **84**, 031606(R) (2011); Fast frictionless dynamics as a toolbox for low-dimensional Bose-Einstein condensates. *Eur. Phys. Lett.* **96**, 60005 (2011).
- Ruschhaupt, A., Chen, X., Alonso, D. & Muga, J. G. Optimally robust shortcuts to population inversion in two-level quantum systems. *New J. Phys.* **14**, 093040 (2012).
- Schaff, J. F., Song, X. L., Vignolo, P. & Labeyrie, G. Fast optimal transition between two equilibrium states. *Phys. Rev. A* **82**, 033430 (2010).
- Schaff, J. F., Song, X. L., Capuzzi, P., Vignolo, P. & Labeyrie, G. Shortcut to adiabaticity for an interacting Bose-Einstein condensate. *Eur. Phys. Lett.* **93**, 23001 (2011).
- Walther, A. *et al.* Number-theoretic nature of communication in quantum spin systems. *Phys. Rev. Lett.* **109**, 050502 (2012).
- Tseng, S. Y. & Chen, X. Engineering of fast mode conversion in multimode waveguides. *Opt. Lett.* **37**, 5118–5120 (2012).
- Chen, X., Torrontegui, E. & Muga, J. G. Lewis-Riesenfeld invariants and transitionless quantum driving. *Phys. Rev. A* **83**, 062116 (2011).

25. Lewis, H. R. & Riesenfeld, W. B. An exact quantum theory of the time-dependent harmonic oscillator and of a charged particle in a time-dependent electromagnetic field. *J. Math. Phys.* **10**, 1458–1473 (1969).
26. Berry, M. V. Transitionless quantum driving. *J. Phys. A* **42**, 365303 (2009).
27. Bason, M. G. *et al.* High-fidelity quantum driving. *Nat. Phys.* **8**, 147–152 (2012).
28. Demiralp, M. & Rice, S. A. Adiabatic population transfer with control fields. *J. Phys. Chem. A* **107**, 9937–9945 (2003).
29. Demiralp, M. & Rice, S. A. On the consistency, extremal, and global properties of counterdiabatic fields. *J. Chem. Phys.* **129**, 154111 (2008).
30. del Campo, A., Rams, M. M. & Zurek, W. H. Assisted Finite-Rate adiabatic passage across a quantum critical point: Exact solution for the quantum ising model. *Phys. Rev. Lett.* **109**, 115703 (2012).
31. Takahashi, K. Transitionless quantum driving for spin systems. *Phys. Rev. E* **87**, 062117 (2013).
32. Takahashi, K. How fast and robust is the quantum adiabatic passage. *J. Phys. A* **46**, 315304 (2013).
33. Muga, J. G., Chen, X., Ibáñez, S., Lizuain, I. & Ruschhaupt, A. Transitionless quantum drivings for the harmonic oscillator. *J. Phys. B* **43**, 085509 (2010).
34. Ibáñez, S., Chen, X., Torrontegui, E., Muga, J. G. & Ruschhaupt, A. Multiple schrödinger pictures and dynamics in shortcuts to adiabaticity. *Phys. Rev. Lett.* **109**, 100403 (2012).
35. Martínez-Garaot, S., Torrontegui, E., Chen, X. & Muga, J. G. Shortcuts to adiabaticity in three-level systems using Lie transforms. *Phys. Rev. A* **89**, 053408 (2014).
36. Torrontegui, E., Martínez-Garaot, S. & Muga, J. G. Hamiltonian engineering via invariants and dynamical algebra. *Phys. Rev. A* **89**, 043408 (2014).
37. Opatrny, T. & Mølmer, K. Partial suppression of nonadiabatic transitions. *New J. Phys.* **16**, 015025 (2014).
38. Saberi, H., Opatrny, T., Mølmer, K. & del Campo, A. Adiabatic tracking of quantum many-body dynamics. *Phys. Rev. A* **90**, 060301(R) (2014).
39. Greenberger, D. M., Horne, M. A., Shimony, A. & Zeilinger, A. Bells theorem without inequalities. *Am. J. Phys.* **58**, 1131–1143 (1990).
40. Zheng, S. B. One-step synthesis of multiatom Greenberger-Horne-Zeilinger states. *Phys. Rev. Lett.* **87**, 230404 (2001).
41. Kwiat, P., Weinfurter, H., Herzog, T., Zeilinger, A. & Kasevich, M. A. Interaction-free measurement. *Phys. Rev. Lett.* **74**, 4763–4766 (1995).
42. Facchi, P. & Pascazio, S. Quantum Zeno subspaces. *Phys. Rev. Lett.* **89**, 080401 (2002).
43. Chen, Y. H., Xia, Y. & Song, J. Deterministic generation of singlet states for N-atoms in coupled cavities via quantum Zeno dynamics. *Quantum inf. Process.* **13**, 1857–1877 (2014).
44. Ye, J., Vernooij, D. W. & Kimble, H. J. Trapping of single atoms in cavity QED. *Phys. Rev. Lett.* **83**, 4987–4990 (1999).
45. McKeever, J. *et al.* State-insensitive cooling and trapping of single atoms in an optical cavity. *Phys. Rev. Lett.* **90**, 133602 (2003).
46. Spillane, S. M. *et al.* Ultrahigh-Q toroidal microresonators for cavity quantum electrodynamics. *Phys. Rev. A* **71**, 013817 (2005).
47. Hartmann, M. J., Brandão, F. G. S. L. & Plenio, M. B. Strongly interacting polaritons in coupled arrays of cavities. *Nat. Phys.* **2**, 849–855 (2006).
48. Brennecke, F. *et al.* Cavity QED with a Bose-Einstein condensate. *Nature* **450**, 268–271 (2007).
49. Li, W. A. & Wei, L. F. Controllable entanglement preparations between atoms in spatially-separated cavities via quantum Zeno dynamics. *Opt. Express* **20**, 13440–13450 (2012).
50. Hao, S. Y., Xia, Y., Song, J. & An, N. B. One-step generation of multiatom Greenberger-Horne-Zeilinger states in separate cavities via adiabatic passage. *J. Opt. Soc. Am. B* **30**, 468–474 (2013).

## Acknowledgement

This work was supported by the National Natural Science Foundation of China under Grants No. 11575045 and No. 11374054, the Foundation of Ministry of Education of China under Grant No. 212085, and the Major State Basic Research Development Program of China under Grant No. 2012CB921601.

## Author Contributions

Y.H.C. and Y.X. came up with the initial idea for the work and performed the simulations for the model. J.S. performed the calculations for the model. Y.H.C. and Q.Q.C. performed all the data analysis and the initial draft of the manuscript. All authors participated in the writing and revising of the text.

## Additional Information

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Chen, Y.-H. *et al.* Shortcuts to adiabatic passage for fast generation of Greenberger-Horne-Zeilinger states by transitionless quantum driving. *Sci. Rep.* **5**, 15616; doi: 10.1038/srep15616 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>