

Research Article

Integrative Analyses of Genes Associated with Fulminant Type 1 Diabetes

Xiaofeng Ye ^{1,2}, Tianshu Zeng^{1,2}, Wen Kong ^{1,2} and Lu-lu Chen ^{1,2}

¹Department of Endocrinology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430022, China

²Hubei Provincial Clinical Research Center for Diabetes and Metabolic Disorders, Wuhan 430022, China

Correspondence should be addressed to Wen Kong; wenly-kong@163.com and Lu-lu Chen; cheria_chen@126.com

Received 23 April 2020; Revised 25 August 2020; Accepted 15 September 2020; Published 6 October 2020

Academic Editor: Lihua Duan

Copyright © 2020 Xiaofeng Ye et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Objective. Fulminant type 1 diabetes (FT1D) is a type of type 1 diabetes, which is characterized by rapid onset of disease and severe metabolic disorders. We intend to screen for crucial genes and potential molecular mechanisms in FT1D in this study. **Method.** We downloaded GSE44314, which includes six healthy controls and five patients with FT1D, from the GEO database. Identification of differentially expressed genes (DEGs) was performed by NetworkAnalyst. The Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analyses of DEGs were screened by an online tool—Database for Annotation, Visualization, and Integration Discovery (DAVID). Protein-protein interaction (PPI) network and hub genes among DEGs were analyzed by NetworkAnalyst. And we also use NetworkAnalyst to find out the microRNAs (miRNAs) and transcription factors (TFs) which regulate the expression of DEGs. **Result.** We identified 130 DEGs (60 upregulated and 70 downregulated DEGs) between healthy controls and FT1D patients. GO analysis results revealed that DEGs were mostly enriched in generation of precursor metabolites and energy, neurohypophyseal hormone activity, and mitochondrial inner membrane. KEGG pathway analysis demonstrated that DEGs were mostly involved in nonalcoholic fatty liver disease. Results indicated that NCOA1, SRF, ERBB3, EST1, TOP1, UBE2S, INO80, COX7C, ITGAV, and COX6C were the top hub genes in the PPI network. Furthermore, we recognized that LDLR, POTEM, IFNAR2, BAZ2A, and SRF were the top hub genes in the miRNA-target gene network, and SRF, TSPAN4, CD59, ETS1, and SLC25A25 were the top hub genes in the TF-target gene network. **Conclusion.** Our study pinpoints key genes and pathways associated with FT1D by a sequence of bioinformatics analysis on DEGs. These identified genes and pathways provide more detailed molecular mechanisms of FT1D and may provide novel therapeutic targets.

1. Introduction

Fulminant type 1 diabetes (FT1D) is a novel type of type 1 diabetes (T1DM) raised by Imagawa et al. in 2000 [1], which is featured by abrupt disease onset, no C-peptide secretion, negative islet-related autoantibodies, and elevated pancreatic enzymes. At first, FT1D was identified as idiopathic T1DM because patients with FT1D lack autoimmune markers such as protein tyrosine phosphatase antibody or glutamic acid decarboxylase autoantibody. Over the past 20 years, the understanding of FT1D has increased. And a sequence of studies indicated that the immunity has a role in the occurrence and development of FT1D, which convinced that FT1D is possibly an autoimmune disease [2–4].

There are studies that reported that genetic and environmental factors take part in the initiation and progression of FT1D. Numbers of studies indicated that CTLA-4, HLA-B, and HLA DR-DQ are related with FT1D [5–7]. Many studies advocate that in FT1D, immune response against viral infection in islets caused the β cell destruction [8–10]. Numerous virus infections were covered in FT1D patients, including coxsackievirus, enterovirus, and human cytomegalovirus [11–13]. Genes such as lymphocyte cytosolic protein 1, melanoma differentiation-associated protein 5, DEAD box helicase 5, and C-X-C motif chemokine 10, which take part in the virus infection, have been proved to be associated with the pathogenesis of FT1D [3, 11, 14]. To further reveal the mechanism of FT1D, a microarray data




FIGURE 1: Volcano plot of differentially expressed genes. Genes with a significant change of more than 1.5-fold were selected.

numbered GSE44314 was deposited by Nakata et al., and it has reported that NKG2E-CD94 were significantly reduced in FT1D, indicating that the reduced expression of NK activating receptor gene and low proportion of NK cells are probably involved in the progression of FT1D [15]. However, there are no studies that had reported the possible regulatory mechanisms of transcription factors (TFs) and microRNAs (miRNAs) related to the development of FT1D.

In our study, we reanalyzed the dataset of GSE44314 by the method of bioinformatics, which includes screening differentially expressed genes (DEGs), functional enrichment analysis, protein-protein interaction (PPI) analysis, and the regulatory TFs/miRNAs related to DEG prediction. Through these analyses, we expect to determine novel insights for the knowledge of FT1D and provide more detailed molecular mechanisms underlying the development of FT1D.

2. Materials and Methods

2.1. Microarray Data. We downloaded the gene expression profile data of GSE44314 from the Gene Expression Omnibus (GEO) database in the National Center for Biotechnology Information (NCBI, <https://www.ncbi.nlm.nih.gov/geo/>). The microarray data was based on the platform of GPL6480 (Agilent-014850 Whole Human Genome Microarray 4x44K G4112F). The datasets available in this analysis were uploaded by Nakata et al. [15], which include 11 samples, containing 6 healthy controls and 5 patients with FT1D.

2.2. Identification of Differentially Expressed Genes. NetworkAnalyst [16, 17] (<https://www.networkanalyst.ca>), a website for integrative statistical and visualizing tool, was used to determine the DEGs between healthy controls and FT1D

patients. The cutoff of the P value was adjusted to 0.05, and $|\log \text{ fold change} | (|\log \text{ FC} |) > 0.585$ for the DEG discrimination, using the false discovery rate (FDR) found on the Benjamini-Hochberg program and moderated t -test based on the Limma algorithm.

2.3. Functional and Pathway Enrichment Analysis. We used an online tool named DAVID [18] (<https://david.ncifcrf.gov/>) in conducting the Gene Ontology (GO) term [19] and Kyoto Encyclopedia of Genes and Genomes (KEGG) [20] pathway enrichment analyses of DEGs, with the thresholds of count ≥ 2 and P value < 0.05 .

2.4. Protein-Protein Interaction (PPI) Network Analysis and Hub Gene Searching. Based on the analyzed DEGs, NetworkAnalyst [21] was used to perform the PPI Network identification with a hypergeometric algorithm, and $P < 0.05$ was identified as having statistically significant differences. Besides, we used NetworkAnalyst to recognize the most significant modules of hub genes using the “module explorer tool,” found on the random walk-dependent Walktrap algorithm.

2.5. Prediction of Target Gene-MicroRNA Network. The gene expression was affected by microRNAs in a disease condition through posttranscriptional control. In the present study, the online tool NetworkAnalyst [17] was used to search the miRNAs associated with DEGs, which integrates microRNA databases miRTarBase (<http://mirtarbase.mbc.nctu.edu.tw/php/download.php>) [22] and TarBase (<http://diana.imis.athena-innovation.gr/DianaTools/index.php?r=tarbase/index>) [23].

2.6. Prediction of Target Gene-Transcription Factor Network. The gene expression was influenced by TFs in a disease




FIGURE 2: Heat map of differentially expressed genes. The abscissa represents different samples, and the ordinate represents different genes. The red boxes indicate upregulated genes, and the green boxes indicate downregulated genes.

condition by transcriptional control. In our study, NetworkAnalyst [17] was used for recognizing the TFs associated with DEGs, which combines TF database JASPAR (<http://jaspar.genereg.net/>) [24].

3. Results

3.1. Identification of Differentially Expressed Genes in Fulminant Type 1 Diabetes. We identified 130 DEGs in FT1D patients compared to healthy controls in total, including 60 upregulated genes and 70 downregulated genes (Supplementary Table 1). We draw a volcano plot of the DEGs (Figure 1) and a hierarchical clustering heat map of

DEGs (Figure 2). It turned out that these DEGs were well distinguished between the FT1D group and the healthy control group. NK2 homeobox 3 (NKX2-3) and Ring finger protein 182 (RNF182) were, respectively, identified as the most significantly upregulated and downregulated genes in FT1D patients.

3.2. Functional Enrichment Analysis. We recognized 21 Gene Ontology terms (Table 1) and 5 KEGG pathways (Table 2) when analyzed with DAVID. The DEGs were mainly focused on the generation of precursor metabolites and energy, hydrogen ion transmembrane transport, and mitochondrial electron transport, cytochrome c to oxygen by biological

TABLE 1: The results of Gene Ontology (GO) of DEGs ranked by *P* value.

Term	Count	<i>P</i> value	Genes
GO-BPs			
Generation of precursor metabolites and energy	4	0.004	AVP, UQCR11, COX7C, COX6C
Hydrogen ion transmembrane transport	4	0.006	NDUFA4, UQCR11, COX7C, COX6C
Mitochondrial electron transport, cytochrome c to oxygen	3	0.006	NDUFA4, COX7C, COX6C
Extrinsic apoptotic signaling pathway in the absence of ligand	3	0.017	MOAP1, ERBB3, ITGAV
Positive regulation of female receptivity	2	0.018	NCOA1, OXT
Positive regulation of gene expression	6	0.02	AMH, ATF4, AVP, LDLR, ERBB3, GPER1
Maternal aggressive behavior	2	0.024	AVP, OXT
Hyperosmotic salinity response	2	0.029	AVP, OXT
Cellular response to lipopolysaccharide	4	0.03	TNFRSF1B, ADAMTS13, PAF1, CACTIN
Social behavior	3	0.033	AVP, OXT, DLG4
Positive regulation of apoptotic process	6	0.034	MOAP1, ATF4, NCOA1, ARHGEF6, GPER1, PDCD1
Male mating behavior	2	0.035	NCOA1, OXT
Positive regulation of uterine smooth muscle contraction	2	0.041	OXT, GPER1
Drinking behavior	2	0.041	HTR1B, OXT
Positive regulation of cytosolic calcium ion concentration	4	0.046	AVP, OXT, DLG4, GPER1
GO-MFs			
Neurohypophyseal hormone activity	2	0.011	AVP, OXT
Cytochrome c oxidase activity	3	0.013	NDUFA4, COX7C, COX6C
Neuregulin binding	2	0.028	ERBB3, ITGAV
GO-CCs			
Mitochondrial inner membrane	8	0.014	NDUFA4, UQCR11, SLC25A25, COX7C, ROMO1, MRPL30, NDUFB1, COX6C
Extracellular space	14	0.046	INA, AVP, CXCL5, ERBB3, ADAMTS13, OXT, FETUB, AMH, IFNAR2, C1QTNF5, CLEC3B, CD59, SEMA4D, PRSS33
Cell junction	7	0.05	CNIH2, OTOF, PRRT1, DLG4, PAF1, GPER1, GPR142

TABLE 2: The results of Kyoto Encyclopedia of Genes and Genomes (KEGG) of DEGs ranked by *P* value.

Term	Count	<i>P</i> value	Genes
Nonalcoholic fatty liver disease (NAFLD)	6	0.0017	NDUFA4, ATF4, UQCR11, COX7C, NDUFB1, COX6C
Huntington's disease	6	0.0048	NDUFA4, UQCR11, DLG4, COX7C, NDUFB1, COX6C
Oxidative phosphorylation	5	0.0071	NDUFA4, UQCR11, COX7C, NDUFB1, COX6C
Parkinson's disease	5	0.0089	NDUFA4, UQCR11, COX7C, NDUFB1, COX6C
Alzheimer's disease	5	0.0158	NDUFA4, UQCR11, COX7C, NDUFB1, COX6C

process (BP) analysis. For the cellular component (CC) group, mitochondrial inner membrane, extracellular space, and cell junction were the enriched terms. Molecular function (MF) analysis showed that the DEGs were remarkably focused on neurohypophyseal hormone activity, cytochrome c oxidase activity, and neuregulin binding. Moreover, the KEGG pathway analysis indicated that the DEGs were significantly involved in nonalcoholic fatty liver disease, Huntington's disease, Alzheimer's disease, and Parkinson's disease as well as oxidative phosphorylation.

3.3. PPI Network and Hub Gene Identification. There were 363 nodes and 409 edges in the PPI network (Figure 3). In this PPI network, sixteen genes with degrees > 10 were found

as key genes (Table 3). The node size is influenced by the fold change between FT1D patients and healthy controls, and the red or orange color nodes indicate that they have a higher score. The core of the whole PPI network was the most key genes in this cluster, including NCOA1, SRF, ERBB3, ETS1, TOP1, UBE2S, INO80, COX7C, ITGAV, COX6C, ATF4, PAF1, YARS, TTI1, UBC, EEF1B2, and AHSA1. Thence, the seventeen genes were recognized as the hub genes.

3.4. miRNA-DEG and TF-DEG Regulating Network Analysis. The miRNAs and TFs for DEGs are displayed in Figures 4 and 5, respectively. The top five targeted genes regulated by miRNA are shown in Supplementary Table 2. It turned out that 167 miRNAs regulate LDLR, 124 miRNAs regulate




FIGURE 3: Protein-protein interaction network of the differentially expressed genes. Red and orange nodes stand for hub genes.

TABLE 3: Sixteen genes with degrees < 10 in the protein-protein interaction network of the differentially expressed genes.

Gene	Regulation	Degree	Betweenness	Expression
ETS1	Up	26	15103.54	1.145
AHSA1	Up	11	3565	0.82
TOP1	Up	23	10312.37	0.764
NCOA1	Up	34	9967.16	0.752
PAF1	Up	18	5908.12	0.732
SRF	Up	31	22498.06	0.647
YARS	Up	15	4237.33	0.644
INO80	Up	22	7030.5	0.606
ITGAV	Down	20	9973.24	-0.603
ATF4	Down	18	11878.52	-0.705
COX6C	Down	20	3314.17	-0.759
COX7C	Down	22	4037.83	-0.801
EEF1B2	Down	11	11109	-0.817
UBE2S	Down	23	7532.83	-0.858
TTI1	Down	14	11460.5	-1.226
ERBB3	Down	29	19037.55	-1.422

POTEM, 109 miRNAs regulate IFNAR2, 107 miRNAs regulate BAZ2A, and 92 miRNAs regulate SRF. The top five targeted genes regulated by TFs are shown in Supplementary

Table 3. It turned out that 25 TFs regulate SRF, 18 TFs regulate TSPAN4, 16 TFs regulate CD59, 16 TFs regulate ETS1, and 15 TFs regulate SLC25A25.

4. Discussion

FT1D is a disease with a state of insulin dependency due to the rapid destruction of almost all pancreatic β cells, which causes the radical onset of ketoacidosis in a few days after the appearance of hyperglycemic symptoms [25–27]. It has been reported that most of the patients with FT1D are found in East Asia, but recently, Western countries also reported this disease [8, 28, 29]. FT1D makes up about 20% of abrupt-onset T1DM cases in Japan [8]. It is important to understand the molecular mechanisms of FT1D. We downloaded and analyzed a dataset (GSE44314) that contains five FT1D patients and six healthy controls from the GEO database. We identified 130 DEGs in total, including 60 upregulated DEGs and 70 downregulated DEGs. Among the 130 DEGs, we noticed that programmed cell death-1 (PD-1) was downregulated in FT1D patients. PD-1 is a critical member of the B7-CD28 family and is one of the important costimulatory molecules [30]. PD-1 can regulate the T cell response and keep maintaining peripheral tolerance by delivering critical inhibitory signals [30]. Inhibiting the PD-1 pathway would bring about excessive T cell proliferation, failure of tolerance, and autoimmune activation [31].




FIGURE 4: Target gene-miRNA regulatory network. Red and orange nodes stand for differentially expressed genes; blue diamonds stand for miRNA.

Therefore, PD-1 has gained popularity in the treatment of several advanced cancers [32, 33]. Studies have proved that treatment with PD-1 inhibitors can cause FT1D [34–36]. And the termination of anti-PD1 antibody therapy may preserve inherent insulin secretion capacity in “anti-PD1 antibody-induced” FT1D [37]. It seems that PD-1 should be upregulated in FT1D, which is totally opposite to our result. Various researchers have identified that cellular immunity, especially T cell, played a crucial role in β cell destruction in FT1D [38–40]. However, a Japanese study that compares PD-1 expression in peripheral CD4⁺ T cells between type 1A diabetes (classical type 1 diabetes), FT1D, and healthy controls found that there is no difference between FT1D and healthy controls in PD-1 expression and that there is lower PD-1 expression in CD4⁺ T cells in patients with type 1A diabetes [41]. Different studies have different conclusions in PD-1 expression in FT1D, which need further studies to confer this question and explore how PD-1 take part in the occurrence and progression of FT1D. Among the increased DEGs, NK2 homeobox 3 (NKX2-3) is the most upregulated gene in FT1D, and an animal study has indicated that NKX2-3 is related to T1DM [42], but further study is needed to figure out how NKX2-3 acts in FT1D.

In the current study, the most significant GO BP term for DEGs is generation of precursor metabolites. UQCR11, COX7C, and COX6C are the new biomarkers for the progression of FT1D. The most significant GO MF term for DEGs is neurohypophyseal hormone activity. Arginine vasopressin (AVP) and oxytocin are associated with type 2 diabetes but are new biomarkers for the progression of FT1D. The most significant GO CC term for DEGs is mitochondrial inner membrane. NDUFA4, SLC25A25, ROMO1, MRPL30, and NDUFB1 are novel biomarkers for the development of FT1D. Nonalcoholic fatty liver disease is the most significant KEGG pathway for DEGs. Activation of activating transcription factor 4 (ATF4) contributes to diabetic hepatotoxicity by ER stress [43]. Besides, ATF4 is a transcription factor implicated in β cell survival and susceptibility to stress [44]. ATF4 is a new biomarker for the progression of FT1D. Parkinson’s disease, Alzheimer’s disease, and Huntington’s disease also are significant KEGG pathways for DEGs. Diabetes mellitus (DM) adversely affects multiple organ systems, including the brain [45]. These evidences suggest that FT1D may also lead to neurodegenerative diseases and adversely affect cognition. Discs large MAGUK scaffold protein 4 (DLG4) is related to neurological disorders and type 2 diabetes [46–48]; DLG4 is a new biomarker for the progression of FT1D.




FIGURE 5: Transcription factor-target DEG regulatory network. Orange and yellow nodes stand for differentially expressed genes; blue diamonds stand for transcription factors.

In the present study, NCOA1, SRF, ERBB3, ETS1, TOP1, UBE2S, INO80, COX7C, ITGAV, and COX6C were recognized as top 10 hub genes in the PPI network. A genome-wide meta-analysis study confirmed that nuclear receptor coactivator 1 (NCOA1) is a T1DM susceptibility gene [49]. An animal study suggests that serum response factor (SRF) is decreased in diabetic nephropathy compared to healthy controls [50]. Many studies confirmed that ERBB3 was the most important T1DM association locus in the non-HLA gene [51–53]. ETS proto-oncogene 1 (ETS1) was found associated with T1DM in the NOD mouse and then confirmed in human population [54–56]. Tissues derived from the T1DM animals show that DNA topoisomerase I (TOP1) activity and enzyme protein level decreased, whereas the enzyme mRNA level was not altered, which demonstrates that TOP1 activity is regulated by high glucose levels and may lead to the pathogenesis of diabetic complications [57]. Ubiquitin-conjugating enzyme E2 (UBE2S) takes part in T1DM by enhancing M2 macrophage polarization [58]. Jin et al. compared integrin subunit alpha V (ITGAV) expression between diabetic nephropathy and normal human kidney and found that ITGAV is higher in diabetic nephropathy [59]. Although there are evidences that the hub genes are contacted with T1DM, they are novel biomarkers for the development of FT1D.

LDLR, POTEM, IFNAR2, BAZ2A, and SRF were identified as top five targeted genes in the miRNA-target gene regulatory network. Low-density lipoprotein receptor (LDLR) is increased in a NOD mouse compared with a nondiabetic mouse [60]. A study in *Ins2(Akita)Ldlr^{-/-}* mice revealed that lack of LDLR will accelerate atherosclerosis in T1DM animals [61]. When lacking the r type II interferon receptor (IFNAR2), diabetes happened only in female NOD mice [62]. POTEM and BAZ2A are novel biomarkers for the development of FT1D. SRF, TSPAN4, CD59, ETS1, and SLC25A25 were identified as top five targeted genes in the TF-DEG regulatory network. Due to the lack of complement regulatory protein CD59, the development of diabetes-induced atherosclerosis in mice is accelerated [63]. Besides, CD59 is reduced in diabetic subjects compared with healthy controls [64]. Tetraspanin 4 (TSPAN4) is a new biomarker for the progression of FT1D.

We noticed that there are two bioinformatics analysis of type 1 diabetes, and there are some the same conclusions between our study and theirs [65, 66]. Fang et al. reported that programmed cell death ligand 1 (PD-L1) was upregulated in the new-onset T1DM samples [66]. This is identical with our result. PD-1/PD-L1 is a negative modulatory signaling pathway for activation of T cell. The upregulated PD-L1 and downregulated PD-1 cause the same result, which are

the inactivation of T cell and the progression of immune tolerance, which play a protective role in the pathogenesis of T1DM. Liu et al. found that HLA-DQA1 and HLA-DRB4 might be targets for the treatment of T1D, and IL8 is likely to be a new marker for the diagnosis of T1D [65]. These results indicated that T1DM is an autoimmune disease, which is in accordance with our result.

5. Conclusions

Our data provide a comprehensive bioinformatics analysis of DEGs to search molecular mechanisms related to the progression of FT1D. We found a set of useful genes for future research into the molecular mechanisms of FT1D progression, while further molecular biological experiments are needed to confirm the effect of these DEGs in the progression of FT1D.

Data Availability

The data used to support the findings of this study are included within the article.

Conflicts of Interest

The authors declare that there are no conflicts of interests associated with the manuscript.

Acknowledgments

The authors thank Nakata and his colleges for depositing the dataset of GSE44314. This study is supported by grants from the National Natural Science Foundation of China (Grant Numbers 81770843 and 81974107).

Supplementary Materials

Supplementary Table 1: the list of all differentially expressed genes. Supplementary Table 2: the top five targeted genes regulated by miRNA. Supplementary Table 3: the top five targeted genes regulated by transcription factor. (*Supplementary Materials*)

References

- [1] A. Imagawa, T. Hanafusa, J. Miyagawa, and Y. Matsuzawa, "A novel subtype of type 1 diabetes mellitus characterized by a rapid onset and an absence of diabetes-related antibodies," *The New England Journal of Medicine*, vol. 342, no. 5, pp. 301–307, 2000.
- [2] S. Luo, X. Ma, X. Li, Z. Xie, and Z. Zhou, "Fulminant type 1 diabetes: a comprehensive review of an autoimmune condition," *Diabetes/Metabolism Research and Reviews*, vol. 36, article e3317, 2020.
- [3] Y. Hosokawa, T. Hanafusa, and A. Imagawa, "Pathogenesis of fulminant type 1 diabetes: genes, viruses and the immune mechanism, and usefulness of patient-derived induced pluripotent stem cells for future research," *Journal of Diabetes Investigation*, vol. 10, no. 5, pp. 1158–1164, 2019.
- [4] L. Liu, L. Zeng, D. Sang, Z. Lu, and J. Shen, "Recent findings on fulminant type 1 diabetes," *Diabetes/Metabolism Research and Reviews*, vol. 34, no. 1, 2018.
- [5] A. Imagawa, T. Hanafusa, Y. Uchigata et al., "Different contribution of class II HLA in fulminant and typical autoimmune type 1 diabetes mellitus," *Diabetologia*, vol. 48, no. 2, pp. 294–300, 2005.
- [6] E. Kawasaki, A. Imagawa, H. Makino et al., "Differences in the contribution of the CTLA4 gene to susceptibility to fulminant and type 1A diabetes in Japanese patients," *Diabetes Care*, vol. 31, no. 8, pp. 1608–1610, 2008.
- [7] Y. Kawabata, on behalf of the Committee on Type 1 Diabetes, J. D. Society et al., "Differential association of HLA with three subtypes of type 1 diabetes: fulminant, slowly progressive and acute-onset," *Diabetologia*, vol. 52, no. 12, pp. 2513–2521, 2009.
- [8] A. Imagawa, T. Hanafusa, Y. Uchigata et al., "Fulminant type 1 diabetes: a nationwide survey in Japan," *Diabetes Care*, vol. 26, no. 8, pp. 2345–2352, 2003.
- [9] A. Imagawa, T. Hanafusa, H. Makino, J. I. Miyagawa, and P. Juto, "High titres of IgA antibodies to enterovirus in fulminant type-1 diabetes," *Diabetologia*, vol. 48, no. 2, pp. 290–293, 2005.
- [10] S. Shibasaki, A. Imagawa, S. Tauriainen et al., "Expression of toll-like receptors in the pancreas of recent-onset fulminant type 1 diabetes," *Endocrine Journal*, vol. 57, no. 3, pp. 211–219, 2010.
- [11] S. Tanaka, Y. Nishida, K. Aida et al., "Enterovirus infection, CXC chemokine ligand 10 (CXCL10), and CXCR3 circuit: a mechanism of accelerated beta-cell failure in fulminant type 1 diabetes," *Diabetes*, vol. 58, no. 10, pp. 2285–2291, 2009.
- [12] N. Ohara, M. Kaneko, T. Nishibori et al., "Fulminant type 1 diabetes mellitus associated with Coxsackie virus type A2 infection: a case report and literature review," *Internal Medicine*, vol. 55, no. 6, pp. 643–646, 2016.
- [13] S. Yoneda, A. Imagawa, K. Fukui et al., "A histological study of fulminant type 1 diabetes mellitus related to human cytomegalovirus reactivation," *The Journal of Clinical Endocrinology and Metabolism*, vol. 102, no. 7, pp. 2394–2400, 2017.
- [14] K. Aida, Y. Nishida, S. Tanaka et al., "RIG-I- and MDA5-initiated innate immunity linked with adaptive immunity accelerates beta-cell death in fulminant type 1 diabetes," *Diabetes*, vol. 60, no. 3, pp. 884–889, 2011.
- [15] S. Nakata, A. Imagawa, Y. Miyata et al., "Low gene expression levels of activating receptors of natural killer cells (NKG2E and CD94) in patients with fulminant type 1 diabetes," *Immunology Letters*, vol. 156, no. 1-2, pp. 149–155, 2013.
- [16] J. Xia, C. D. Fjell, M. L. Mayer, O. M. Pena, D. S. Wishart, and R. E. Hancock, "INMEX—a web-based tool for integrative meta-analysis of expression data," *Nucleic Acids Research*, vol. 41, no. W1, pp. W63–W70, 2013.
- [17] J. Xia, E. E. Gill, and R. E. Hancock, "NetworkAnalyst for statistical, visual and network-based meta-analysis of gene expression data," *Nature Protocols*, vol. 10, no. 6, pp. 823–844, 2015.
- [18] W. H. Da, B. T. Sherman, and R. A. Lempicki, "Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources," *Nature Protocols*, vol. 4, no. 1, pp. 44–57, 2009.

- [19] M. Ashburner, C. A. Ball, J. A. Blake et al., "Gene ontology: tool for the unification of biology," *Nature Genetics*, vol. 25, no. 1, pp. 25–29, 2000.
- [20] M. Kanehisa and S. Goto, "KEGG: Kyoto Encyclopedia of Genes and Genomes," *Nucleic Acids Research*, vol. 28, no. 1, pp. 27–30, 2000.
- [21] J. Xia, M. J. Benner, and R. E. Hancock, "NetworkAnalyst—integrative approaches for protein-protein interaction network analysis and visual exploration," *Nucleic Acids Research*, vol. 42, no. W1, pp. W167–W174, 2014.
- [22] C. H. Chou, S. Shrestha, C. D. Yang et al., "miRTarBase update 2018: a resource for experimentally validated microRNA-target interactions," *Nucleic Acids Research*, vol. 46, no. D1, pp. D296–D302, 2018.
- [23] I. S. Vlachos, M. D. Paraskevopoulou, D. Karagkouni et al., "DIANA-TarBase v7.0: indexing more than half a million experimentally supported miRNA:mRNA interactions," *Nucleic Acids Research*, vol. 43, no. D1, pp. D153–D159, 2015.
- [24] A. Khan, O. Fornes, A. Stigliani et al., "JASPAR 2018: update of the open-access database of transcription factor binding profiles and its web framework," *Nucleic Acids Research*, vol. 46, no. D1, p. D1284, 2018.
- [25] T. Hanafusa and A. Imagawa, "Fulminant type 1 diabetes: a novel clinical entity requiring special attention by all medical practitioners," *Nature Clinical Practice, Endocrinology & Metabolism*, vol. 3, no. 1, pp. 36–45, 2007.
- [26] A. Imagawa and T. Hanafusa, "Fulminant type 1 diabetes—an important subtype in East Asia," *Diabetes/Metabolism Research and Reviews*, vol. 27, no. 8, pp. 959–964, 2011.
- [27] A. Imagawa, T. Hanafusa, T. Awata et al., "Report of the Committee of the Japan Diabetes Society on the research of fulminant and acute-onset type 1 diabetes mellitus: new diagnostic criteria of fulminant type 1 diabetes mellitus (2012)," *Journal of Diabetes Investigation*, vol. 3, no. 6, pp. 536–539, 2012.
- [28] T. S. Jung, S. I. Chung, M. A. Kim et al., "A Korean patient with fulminant autoantibody-negative type 1 diabetes," *Diabetes Care*, vol. 27, no. 12, pp. 3023–3024, 2004.
- [29] C. Moreau, D. Drui, G. Arnault-Ouary, B. Charbonnel, L. Chaillous, and B. Cariou, "Fulminant type 1 diabetes in Caucasians: a report of three cases," *Diabetes & Metabolism*, vol. 34, no. 5, pp. 529–532, 2008.
- [30] M. E. Keir, Y. E. Latchman, G. J. Freeman, and A. H. Sharpe, "Programmed death-1 (PD-1):PD-ligand 1 interactions inhibit TCR-mediated positive selection of thymocytes," *Journal of Immunology*, vol. 175, no. 11, pp. 7372–7379, 2005.
- [31] E. Lazar-Molnar, B. Chen, K. A. Sweeney et al., "Programmed death-1 (PD-1)-deficient mice are extraordinarily sensitive to tuberculosis," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 107, no. 30, pp. 13402–13407, 2010.
- [32] K. C. Ohaegbulam, A. Assal, E. Lazar-Molnar, Y. Yao, and X. Zang, "Human cancer immunotherapy with antibodies to the PD-1 and PD-L1 pathway," *Trends in Molecular Medicine*, vol. 21, no. 1, pp. 24–33, 2015.
- [33] A. V. Balar and J. S. Weber, "PD-1 and PD-L1 antibodies in cancer: current status and future directions," *Cancer Immunology, Immunotherapy: CII*, vol. 66, no. 5, pp. 551–564, 2017.
- [34] L. Marchand, A. Thivolet, S. Dalle et al., "Diabetes mellitus induced by PD-1 and PD-L1 inhibitors: description of pancreatic endocrine and exocrine phenotype," *Acta Diabetologica*, vol. 56, no. 4, pp. 441–448, 2019.
- [35] M. Okamoto, M. Okamoto, K. Gotoh et al., "Fulminant type 1 diabetes mellitus with anti-programmed cell death-1 therapy," *Journal of Diabetes Investigation*, vol. 7, no. 6, pp. 915–918, 2016.
- [36] M. Araujo, D. Ligeiro, L. Costa et al., "A case of fulminant type 1 diabetes following anti-PD1 immunotherapy in a genetically susceptible patient," *Immunotherapy*, vol. 9, no. 7, pp. 531–535, 2017.
- [37] G. Sakai, D. Saito, R. Nakajima et al., "Intrinsic insulin secretion capacity might be preserved by discontinuing anti-programmed cell death protein 1 antibody treatment in 'anti-programmed cell death protein 1 antibody-induced' fulminant type 1 diabetes," *Journal of Diabetes Investigation*, vol. 9, no. 2, pp. 448–449, 2018.
- [38] A. Shimada, J. Morimoto, K. Kodama et al., "T-cell-mediated autoimmunity may be involved in fulminant type 1 diabetes," *Diabetes Care*, vol. 25, no. 3, pp. 635–636, 2002.
- [39] A. Shimada, Y. Oikawa, T. Shigihara, T. Senda, and K. Kodama, "A case of fulminant type 1 diabetes with strong evidence of autoimmunity," *Diabetes Care*, vol. 25, no. 8, pp. 1482–1483, 2002.
- [40] K. Aoki, M. Taniyama, C. Nagayama, Y. Oikawa, and A. Shimada, "T cell immunity to glutamic acid decarboxylase in fulminant type 1 diabetes without significant elevation of serum amylase," *Annals of the New York Academy of Sciences*, vol. 1079, no. 1, pp. 181–185, 2006.
- [41] R. Fujisawa, F. Haseda, C. Tsutsumi et al., "Low programmed cell death-1 (PD-1) expression in peripheral CD4(+) T cells in Japanese patients with autoimmune type 1 diabetes," *Clinical and Experimental Immunology*, vol. 180, no. 3, pp. 452–457, 2015.
- [42] H. Weiss, A. Bleich, H. J. Hedrich et al., "Genetic analysis of the LEW.IAR1-iddm rat: an animal model for spontaneous diabetes mellitus," *Mammalian Genome*, vol. 16, no. 6, pp. 432–441, 2005.
- [43] V. K. Pandey, A. Mathur, M. F. Khan, and P. Kakkar, "Activation of PERK-eIF2 α -ATF4 pathway contributes to diabetic hepatotoxicity: attenuation of ER stress by Morin," *Cellular Signalling*, vol. 59, pp. 41–52, 2019.
- [44] C. A. Juliana, J. Yang, C. E. Cannon, A. L. Good, M. W. Haemmerle, and D. A. Stoffers, "A PDX1-ATF transcriptional complex governs β cell survival during stress," *Molecular Metabolism*, vol. 17, pp. 39–48, 2018.
- [45] J. Chen, L. Liang, L. Zhan et al., "ZiBuPiYin recipe protects db/db mice from diabetes-associated cognitive decline through improving multiple pathological changes," *PLoS One*, vol. 9, no. 3, article e91680, 2014.
- [46] G. H. Kim, E. C. Park, S. H. Yun et al., "Proteomic and bioinformatic analysis of membrane proteome in type 2 diabetic mouse liver," *Proteomics*, vol. 13, no. 7, pp. 1164–1179, 2013.
- [47] Y. Matsunaga, T. Negishi, A. Hatakeyama, Y. Kawagoe, E. Sawano, and T. Tashiro, "Impairment of synaptic development in the hippocampus of diabetic Goto-Kakizaki rats," *International Journal of Developmental Neuroscience*, vol. 53, no. 1, pp. 58–67, 2016.
- [48] Z. Tucsek, M. Noa Valcarcel-Ares, S. Tarantini et al., "Hypertension-induced synapse loss and impairment in synaptic plasticity in the mouse hippocampus mimics the aging phenotype: implications for the pathogenesis of vascular cognitive impairment," *GeroScience*, vol. 39, no. 4, pp. 385–406, 2017.

- [49] J. P. Bradfield, H. Q. Qu, K. Wang et al., "A genome-wide meta-analysis of six type 1 diabetes cohorts identifies multiple associated loci," *PLoS Genetics*, vol. 7, no. 9, article e1002293, 2011.
- [50] S. Kostic, B. Williams, S. Ksouri et al., "Changes in snail and SRF expression in the kidneys of diabetic rats during ageing," *Acta Histochemica*, vol. 122, no. 1, p. 151460, 2020.
- [51] C. Sun, H. Wei, X. Chen et al., "ERBB3-rs2292239 as primary type 1 diabetes association locus among non-HLA genes in Chinese," *Meta Gene*, vol. 9, pp. 120–123, 2016.
- [52] B. I. Frohnert, M. Laimighofer, J. Krumsiek et al., "Prediction of type 1 diabetes using a genetic risk model in the Diabetes Autoimmunity Study in the Young," *Pediatric Diabetes*, vol. 19, no. 2, pp. 277–283, 2018.
- [53] D. Wang and G. Pan, "The association between rs2292239 Polymorphism in ERBB3 Gene and type 1 diabetes: a meta-analysis," *BioMed Research International*, vol. 2019, Article ID 7689642, 7 pages, 2019.
- [54] M. Prochazka, D. V. Serreze, S. M. Worthen, and E. H. Leiter, "Genetic control of diabetogenesis in NOD/Lt mice. Development and analysis of congenic stocks," *Diabetes*, vol. 38, no. 11, pp. 1446–1455, 1989.
- [55] J. M. Aparicio, A. Wakisaka, A. Takada, N. Matsuura, and T. Yoshiki, "Non-HLA genetic factors and insulin dependent diabetes mellitus in the Japanese: TCRA, TCRB and TCRG, INS, THY1, CD3D and ETS1," *Disease Markers*, vol. 8, no. 5, 294 pages, 1990.
- [56] J. M. Aparicio, "HLA and non-HLA genetic factors in Japanese IDDM, [Hokkaido igaku zasshi]," *The Hokkaido Journal of Medical Science*, vol. 66, no. 6, pp. 780–793, 1991.
- [57] I. Levi, Y. Segev, and E. Priel, "Type 1 diabetes affects topoisomerase I activity and GlcNAcylation in rat organs: kidney, liver and pancreas," *Glycobiology*, vol. 22, no. 5, pp. 704–713, 2012.
- [58] F. Wang, F. Sun, J. Luo et al., "Loss of ubiquitin-conjugating enzyme E2 (Ubc9) in macrophages exacerbates multiple low-dose streptozotocin-induced diabetes by attenuating M2 macrophage polarization," *Cell Death & Disease*, vol. 10, no. 12, p. 892, 2019.
- [59] D. K. Jin, A. J. Fish, E. A. Wayner et al., "Distribution of integrin subunits in human diabetic kidneys," *Journal of the American Society of Nephrology: JASN*, vol. 7, no. 12, pp. 2636–2645, 1996.
- [60] L. M. Aaron-Brooks, T. Sasaki, R. E. Vickman et al., "Hyperglycemia and T cell infiltration are associated with stromal and epithelial prostatic hyperplasia in the nonobese diabetic mouse," *The Prostate*, vol. 79, no. 9, pp. 980–993, 2019.
- [61] C. Zhou, B. Pridgen, N. King, J. Xu, and J. L. Breslow, "Hyperglycemic Ins2AkitaLdlr(-)/(-) mice show severely elevated lipid levels and increased atherosclerosis: a model of type 1 diabetic macrovascular disease," *Journal of Lipid Research*, vol. 52, no. 8, pp. 1483–1493, 2011.
- [62] J. A. Carrero, N. D. Benschoff, K. Nalley, and E. R. Unanue, "Type I and II interferon receptors differentially regulate type 1 diabetes susceptibility in male versus female NOD mice," *Diabetes*, vol. 67, no. 9, pp. 1830–1835, 2018.
- [63] F. Liu, R. Sahoo, X. Ge et al., "Deficiency of the complement regulatory protein CD59 accelerates the development of diabetes-induced atherosclerosis in mice," *Journal of Diabetes and its Complications*, vol. 31, no. 2, pp. 311–317, 2017.
- [64] A. Soggiu, C. Piras, L. Bonizzi, H. A. Hussein, S. Pisanu, and P. Roncada, "A discovery-phase urine proteomics investigation in type 1 diabetes," *Acta Diabetologica*, vol. 49, no. 6, pp. 453–464, 2012.
- [65] H. Liu, R. Xu, X. Liu, R. Sun, and Q. Wang, "Bioinformatics analysis of gene expression in peripheral blood mononuclear cells from children with type 1 diabetes in 3 periods," *Experimental and Clinical Endocrinology & Diabetes*, vol. 122, no. 8, pp. 477–483, 2014.
- [66] C. Fang, Y. Huang, Y. Pei et al., "Genome-wide gene expression profiling reveals that CD274 is up-regulated new-onset type 1 diabetes mellitus," *Acta Diabetologica*, vol. 54, no. 8, pp. 757–767, 2017.