



Special issue in honor of Prof. George C. Papageorgiou

Exposed red leaves display adaptive adjustments in chlorophyll and photosystem ratios compatible with the shade imposed by anthocyanin accumulation

K. ZELIOU, A. KYZERIDOU, and Y. PETROPOULOU⁺

Laboratory of Plant Physiology, Department of Biology, University of Patras, Patras GR-26504, Greece

Abstract

Foliar anthocyanins shape a peculiar shade in a red leaf's interior leading to uneven energy distribution between the two photosystems. Accordingly, a readjustment of PSII/PSI stoichiometry could restore excitation balance. To test this hypothesis, 77 K fluorescence emission spectra of thylakoids from green and red leaves of seven species with different pigment profiles were compared. The ratio of F_{686}/F_{736} served as an indication of the PSII/PSI functional ratio. To avoid possible species-dependent differences in the measured parameters, plants showing intra-individual, intra-species, or intra-leaf variation in the expression of the anthocyanic character were used. Red leaves or red leaf areas displayed higher PSII/PSI ratio, irrespectively of species and anthocyanin accumulation pattern. PSII/PSI ratio declined in parallel with anthocyanin decrease. In five species, red leaves displayed also a lower Chl *a/b* ratio. We conclude that red leaves growing in full sunlight develop adaptive adjustments in their chlorophyll and photosystem ratios, compatible with the shade-acclimation syndrome.

Keywords: fluorescence emission spectra; foliar anthocyanins; photosystem ratio; shade-acclimation syndrome.

Introduction

Some plants accumulate anthocyanins in their leaves at levels high enough to mask the green chlorophyll (Chl) color. The major leaf anthocyanin is cyanidin-3-glucoside (Harborne 1976), absorbing strongly in the green/yellow, lesser in the blue, and negligibly in the red part of the spectrum (Neill and Gould 2000, Hughes and Smith 2007, Gould *et al.* 2018, Landi *et al.* 2021). Accordingly, anthocyanic leaves appear red to dark brown, depending on the contents of co-existing chlorophylls. Foliar anthocyanins may be found in epidermises (upper and/

or lower), mesophyll (palisade and/or spongy cells), trichomes, and they reside exclusively in the vacuole (Hrazdina *et al.* 1978, Lee and Collins 2001, Kytridis and Manetas 2006). Although in some species the anthocyanic trait is permanent, the transient leaf redness is more common and may be induced by developmental or environmental agents. In the first case, young or senescing leaves appear red while being green at the mature stage. In the latter, green fully expanded leaves turn temporarily red as a response to UV-B radiation, nutrient deficiency, wounding, pathogen attack, or low temperatures accompanied by high light (Gould *et al.* 2002a, Steyn *et al.*

Highlights

- Red leaves display a higher PSII/PSI ratio, compared to their green counterparts
- PSII/PSI ratio declines in parallel with the decrease of anthocyanin content
- Red leaves displayed also a lower Chl *a/b* ratio

Received 6 October 2021

Accepted 1 December 2021

Published online 13 January 2022

⁺Corresponding author

e-mail: petropo@upatras.gr

Abbreviations: Anths – anthocyanins; Car – total carotenoids; Chl – chlorophyll; FR – far-red light; R – red light.

Acknowledgments: This paper is dedicated to the memory of the late Dr. George C. Papageorgiou for his important contribution to photosynthesis research and we are honored to participate in this special issue.

Conflict of interest: The authors declare that they have no conflict of interest.

2002, Close and Beadle 2003, Gould 2004, Manetas 2006, Hughes 2011).

Since anthocyanins absorb visible radiation, their accumulation in leaves represents a photosynthetic cost because of the competition with Chls and carotenoids (Car) for photon capture (Merzlyak *et al.* 2008). Assuming that this cost should be counterbalanced by some advantages of leaf redness, numerous hypotheses for an adaptive significance of the anthocyanin accumulation have been proposed (for reviews see Chalker-Scott 1999, Hoch *et al.* 2001, Steyn *et al.* 2002, Close and Beadle 2003, Gould 2004, Lev-Yadun *et al.* 2004, Manetas 2006, Archetti *et al.* 2009, Hughes 2011). Regardless of the ascribed function(s), however, anthocyanins modify quantitatively as well as qualitatively the internal light environment of a leaf. Pietrini and Massacci (1998), using an indirect estimation, calculated that the chloroplasts of an anthocyanic leaf may enjoy as much as 40% lower photosynthetically active radiation than its anthocyanin-less counterpart. In addition, it has been repeatedly shown that red leaves display an array of morphological and physiological characteristics, partly compatible with the shade acclimation syndrome (Gould *et al.* 2002a, Manetas *et al.* 2003, Hughes and Smith 2007, Hughes *et al.* 2007, Kyparissis *et al.* 2007, Kytridis *et al.* 2008, Zeliou *et al.* 2009).

It is well established in the literature that almost all photosynthetic parameters of leaves and chloroplasts, especially pigment contents and their ratios, are strongly affected by both the quantity and quality of light they receive. Compared to the exposed sites, under a dense plant or forest canopy prevails diffuse light of low intensity, greatly depleted in red and blue while enriched in green and far-red photons. Thus, shade leaves possess significantly lower Chl *a/b* values than the corresponding sun leaves, since they invest primarily in higher amounts of light-harvesting chlorophyll proteins and lower number of reaction center pigment proteins (Anderson *et al.* 1995, Lichtenthaler and Babani 2004, Lichtenthaler *et al.* 2007, 2013). The internal light microenvironment shaped by an anthocyanin screen, however, is not neutral or canopy-like since, in contrast to natural shade, the green/yellow part of the penetrated light is strongly affected (Karabourniotis *et al.* 1999, Gould *et al.* 2002b, 2018). Hence, in the interior of an anthocyanic leaf, the light is FR-enriched (as in natural shade) but green/yellow depleted. This spectral selectivity may be superimposed on the already lower internal photon fluence rates and further modify a red leaf's photosynthetic physiology. According to Kyparissis *et al.* (2007), this hypothesis could explain some deviations from the classical shade acclimation syndrome observed in the permanently red leaves of *Prunus cerasifera*.

Based on the above, we argued that the selective attenuation mainly of green/yellow and, to a lesser degree, blue light by anthocyanins represents a potential loss for PSII, which preferentially absorbs these spectral bands (Glick *et al.* 1985, Pfannschmidt 2005, Hogewoning *et al.* 2012). Accordingly, an adjustment in photosystem stoichiometry could be inferred (Chow *et al.* 1990), to compensate for the uneven spectral distribution of photons

within a red leaf. Towards that aim, 77 K fluorescence spectra of dilute thylakoid preparations obtained from green and red leaves of various species with different pigment profiles were compared. The corresponding F_{686}/F_{736} ratio from each leaf type was used as a relative indication of the PSII/PSI functional analogy (Krause and Weis 1991, Papageorgiou and Govindjee 2004, Lamb *et al.* 2018). The main criteria for species selection were (1) the avoidance of possible species-dependent differences in the measured parameters and any confounding effect from the different irradiation history of the selected leaves and (2) the examination of species with different profiles of anthocyanin accumulation (tissue localization and/or inductive agent). Thus, we included plants displaying intra-individual, intra-species, or intra-leaf variation in the expression of the anthocyanic trait, accumulating anthocyanins permanently or transiently in the epidermis(es) or the mesophyll.

Materials and methods

Plant material and sampling: To meet the requisites mentioned in the 'Introduction', the following seven species with transient or permanent anthocyanin accumulation in their young or mature leaves were used as experimental material:

Ricinus communis L.: perennial shrub cultivated as ornamental in the Patras University Campus. During spring and autumn produces red young leaves, which become gradually green upon maturation. Two individuals were used.

Photinia × fraseri Dress.: perennial shrub with young red and mature green leaves. As with *R. communis*, new leaf production occurs during spring and autumn under Mediterranean conditions. Eight individuals were purchased from a local nursery and cultivated in plastic pots in a sun-lit site of Patras University Campus.

Rosa sp.: two varieties possessing red or green young leaves were selected, while the mature leaves of both are green. New leaves burst during spring. Six plants (three from each variety) were purchased from a local nursery and cultivated in plastic pots in a sun-lit site of Patras University Campus.

Quercus coccifera L.: Mediterranean evergreen sclerophyll shrub, growing in wild in the neighborhood of Patras University Campus. This species displays intra-species variation in the expression of leaf redness, *i.e.*, some individuals possess dark red young leaves while other neighboring plants (occupying the same habitat) possess green young ones (Manetas *et al.* 2003). New leaves burst during the spring (early April to mid-May) at the top of older branches. The mature leaves of both phenotypes are green. From previous observations at the same field site, it was confirmed that the same plants produce red or green young leaves every year (*i.e.*, there is stability in the expression of leaf redness among individuals). Thus, 12 tagged individuals (6 per phenotype) were used.

Cistus creticus L.: evergreen malacophyllous Mediterranean shrub growing also in wild in the vicinity of Patras University Campus. Certain individuals of this

species turn red at mid-winter and resume their green color in late spring (after the shedding of old leaves) while neighboring individuals under the same environmental conditions remain invariably green (Kytridis and Manetas 2006, Kytridis *et al.* 2008, Zeliou *et al.* 2009). As with *Q. coccifera*, there is a constancy in the expression of the anthocyanic trait among individuals. Throughout the remaining seasons (from late spring to mid-winter), both phenotypes have green leaves. New leaves (green) burst in mid-April. Ten tagged plants (five per phenotype) were used.

Prunus cerasifera Ehrh. (var. *atropurpurea*) grafted at the base of *Prunus domestica* L. rootstock: the two species form small two-stemmed trees, with a permanently green (*P. domestica*) and a permanently red-leafed (*P. cerasifera*) stem. Both stems are winter deciduous. Two plants grown as ornamentals in the Patras University Campus were used.

Coleus blumei Benth, a tropical understory variegated species, possessing chlorophyllous anthocyanin-rich ('red areas') and anthocyanin-less ('green areas') spots in the same leaf. Twelve individuals of about the same age cultivated in plastic pots in a shaded site of Patras University Campus were used.

All plants, growing side by side under the same environmental conditions, were equally exposed to direct solar radiation except *C. blumei*, which received $\sim 200 \mu\text{mol}(\text{photon}) \text{ m}^{-2} \text{ s}^{-1}$ (maximum mid-day PAR intensity on plant apex). During the whole experimental period, the wild-growing plants received only natural precipitation while the cultivated plants were well watered.

Sampling was performed always on clear days and each measuring date south-facing (*i.e.*, fully exposed) green and red leaves from each species were compared. In *R. communis* and *P. × fraseri*, young and mature leaves co-occur on the same individual, because of the extended duration of leaf development. Thus, two age classes of young leaves (referred to as red A and red B, respectively) were compared with mature green. Leaf selection was based on their color and position to the branch, numbered from the apex. In the case of *C. creticus*, measurements were performed in mature leaves of the two phenotypes at frequent intervals, during both the 'green' (*i.e.*, when the leaves of both phenotypes are green) and the 'red'

(*i.e.*, when leaves of some individuals turn red) period of the year. Sampling started before the appearance of redness (in early December) and ended in late spring when the red phenotype reverted to green.

On each measuring date, an adequate number of green and/or red leaves from each individual of a given species were harvested early in the morning, put in air-tight plastic bags with moistened filter paper, and transferred immediately to the laboratory for further experimentation. Discs (2–4) of known diameter were punched from each leaf to be used for pigment extraction. The remaining tissue of red or green leaves (without the main vein) was pooled, cut in small segments, and put in a mortar with a small volume of liquid N₂ to proceed for thylakoid isolation. In the case of *C. blumei*, an adequate number of discs from the corresponding 'red' and 'green' areas of each mature leaf was cut and pooled separately.

The location of anthocyanins was assessed by microscopic examination of free-hand leaf cross-sections from each species in conjunction with previous observations (Burger and Edwards 1996, Manetas *et al.* 2003, Kytridis and Manetas 2006, Kyparissis *et al.* 2007). Anthocyanin localization and the comparisons performed within each species are presented in the text table below.

Thylakoid isolation: All the procedures took place at 4°C. Thylakoids were prepared using the method of Šíffel *et al.* (2000) with some modifications (Yiotis *et al.* 2009). Briefly, the leaf segments (or discs) were grinded in the 'isolation medium' (0.4 M sorbitol, 5 mM MgCl₂, 10 mM KCl, 1 mM MnCl₂, 1 mM sodium ascorbate, 0.5–1% bovine serum albumin, 50 mM Tricine/KOH, pH 7.6). After two sequential centrifugations (at 1,500 × *g* for 2 min and 8,000 × *g* for 10 min) the pellet was resuspended in 'washing medium' (0.33 M sorbitol, 5 mM MgCl₂, 10 mM KCl, 1 mM MnCl₂, 0.05% bovine serum albumin, 50 mM Tricine/KOH, pH 7.6). Following two consecutive resuspensions–centrifugations at 8,000 × *g* for 10 min, the final pellet was resuspended in 2 ml of 'washing medium' and served as the thylakoid preparation. For each fluorescence measurement, a small aliquot (200–400 μl) of the thylakoid sample was used after appropriate dilution.

Species tested	Anthocyanin location	Leaf comparison
Intra-individual variation		
<i>Ricinus communis</i>	upper epidermis	young red vs. mature green
<i>Photinia × fraseri</i>	whole mesophyll	
Intra-species variation		
<i>Rosa</i> sp.	both epidermises	young red vs. young green
<i>Quercus coccifera</i>	mesophyll layers (subepidermal)	
<i>Cistus creticus</i>	palisade mesophyll	mature red vs. mature green
<i>Prunus cerasifera</i> × <i>Prunus domestica</i>	whole mesophyll	
Intra-leaf variation		
<i>Coleus blumei</i>	upper epidermis	red vs. green areas of the same leaf

Low-temperature fluorescence spectra: 77 K fluorescence emission spectra of dilute thylakoid preparations were recorded using a spectrofluorimeter (*F-2500*, Hitachi, High Technologies Corporation, Japan), equipped with a liquid-nitrogen sample device and a red-sensitive photomultiplier, as previously described (Yiotis *et al.* 2009). The calculated F_{686}/F_{736} ratio served as a relative indication of the PSII/PSI ratio (Krause and Weis 1991). Care was taken to keep the Chl concentration of the probed sample at low contents (2–4 $\mu\text{g ml}^{-1}$, as judged from preliminary trials), to avoid self-absorption of Chl fluorescence (Weis 1985, Lamb *et al.* 2018). Total Chls of the thylakoid samples eluted in pure methanol were estimated by the equations of Lichtenthaler and Wellburn (1983), using a double-beam spectrophotometer (Shimadzu *UV-160A*). To prevent condensation, a water vapor-free airstream was supplied in the Dewar cuvette of the spectrofluorimeter. Excitation was set at 490 nm (slit width 10 nm) and emission was recorded between 670–800 nm (slit width 2.5 nm). Every thylakoid sample was measured three times. Data were exported in the form of a smoothed curve and datasheet with the *Hitachi FL Solution* software. Raw fluorescence spectra were smoothed with the Savitsky-Golay method and analyzed by a multi-peak fitting process. Three Gaussian bands at 685, 695, and 735 nm were used for spectra fitting.

Pigment determination: The discs sampled before thylakoid isolation were frozen in a mortar by adding a small volume of liquid nitrogen and pigments extracted in dim light with pure methanol in the presence of a small amount of CaCO_3 . The extract was centrifuged at $5,000 \times g$ for 10 min at 4°C and the supernatant was used for pigment determination. Chls were measured spectrophotometrically, using a Shimadzu *UV-160A* double-beam spectrophotometer (Shimadzu Deutschland GmbH, Duisburg, Germany), and concentration was estimated according to the equations of Lichtenthaler and Wellburn (1983). For anthocyanin determination, the methanolic leaf extracts were acidified with 1% HCl and absorbance was scanned from 400–700 nm. The peak anthocyanin absorbance (528–532 nm, depending on species) was corrected for pheophytin absorbance at this wavelength (Lindoo and Caldwell 1978) and transformed to actual concentrations by using the mean molar absorption coefficient according to Murray and Hackett (1991).

Statistics: Significance of differences in the measured parameters between green and red leaves of each species was assessed with *Student's t-test* (SPSS v. 20.0 statistical package, IBM-SPSS Statistics, Armonk, NY, USA). In the case of *C. creticus*, statistically significant differences between green and red leaves were assessed separately for the ‘green’ and the ‘red’ period.

Results

Fig. 1 (left panels) presents 77 K fluorescence spectra recorded from dilute thylakoid preparations of red and

green leaves of the seven species tested. To facilitate comparison, within each species, the fluorescence values of red leaves were normalized to those of the corresponding green leaf at $\lambda = 736$ nm. In all cases, the 77 K spectra showed a typical profile with two main bands peaking at 686 and 736 nm related to the core antenna of PSII and LHCI-PSI, respectively. The third band at 695 nm, arising from the CP47 PSII core antenna, appears as a shoulder. From the corresponding fluorescence values of each leaf type and species, the F_{686}/F_{736} ratio (indicative of the relative functional analogy of the two photosystems, PSII/PSI) was calculated (Fig. 1, right panels).

As it is shown, in all species red leaves displayed higher PSII fluorescence and a higher F_{686}/F_{736} ratio compared to their green counterparts, irrespectively of their anthocyanin accumulation pattern. More specifically, in *R. communis* and *P. × fraseri* (Fig. 1A,B), where leaves of the different developmental stages were compared, PSII/PSI ratio declined in parallel to the gradual loss of anthocyanins. A higher F_{686}/F_{736} ratio of red leaves was also confirmed with the next four species (Fig. 1C–F), where leaf morphs of about the same physiological age, either young (*Rosa* sp., *Q. coccifera*) or mature (*C. creticus*, *P. cerasifera* × *P. domestica*), were compared. The same applies for *C. blumei* (Fig. 1G), where the comparisons concern adjacent red and green chlorophyllous spots of the same mature leaf. In the case of *C. creticus* (Fig. 1E), the emission spectra corresponded to thylakoids isolated from the two phenotypes during the ‘red’ period of the year. During the ‘green’ period (when the leaves of both phenotypes are green), there was no difference in fluorescence emission.

Area-based anthocyanin and total Chl concentrations along with Chl *a/b* ratios of red and green leaves from all species tested are given in Fig. 2. As expected from their colors, in most of the cases, green leaves possess negligible (less than 2%) or very low (5–9% in *C. creticus* and *C. blumei*, respectively) anthocyanin concentrations compared to the corresponding reds. The first two species displayed a common pigment profile, *i.e.*, the gradual loss of anthocyanins upon maturation was accompanied by an increase in Chl concentration (as expected), without any difference in the Chl *a/b* ratio. Thus, in *R. communis*, red leaves had *ca* 45% (red A) and 64% (red B) of the Chl content of the mature green ones, while their anthocyanins were ~120- and 85-fold higher, respectively (Fig. 2A). In *P. × fraseri*, a delayed-greening species, total Chls increased more sharply with age, with young leaves containing 18% (red A) and 35% (red B) of the corresponding mature ones. Concerning anthocyanins, though they didn't differ between the two young age classes, they were ~55-fold higher in red compared to green leaves (Fig. 2B).

In the remainder species, compared to green, red leaves displayed always a slightly, yet significantly, lower (7–17%, depending on species) Chl *a/b* ratio. Concerning total Chls, no difference was observed in three species (*Rosa* sp., *Q. coccifera*, *C. blumei*; Fig. 2C,D,G, respectively) while in *C. creticus* and *P. cerasifera* × *P. domestica*, red leaves had ~30% lower Chl values (Fig. 2E,F). We have to




Fig. 1. 77 K fluorescence emission spectra recorded from isolated thylakoids of red (*dotted line*) and green (*solid line*) leaves of the indicated species. Fluorescence values of red leaves were normalized to those of the corresponding green ones at $\lambda = 736$ nm. Bars represent means \pm SE of the F_{686}/F_{736} ratio from red (*white bar*) and green (*grey bar*) leaves, calculated from the corresponding emission spectra. *Different letters* above bars indicate statistically significant differences ($p < 0.05$) between red and green leaves within each species. In *Ricinus communis* and *Photinia* \times *fraseri*, two age classes of young leaves, red A (*dashed line, hatched bar*) and red B (*dotted line, white bar*) respectively, were used. In the case of *Cistus creticus*, the emission spectra and the corresponding F_{686}/F_{736} ratios concern the ‘red measuring period’. n – number of independent preparations.

note that in the case of *C. creticus*, the above differences were observed only during the ‘red’ period of the year and were almost abolished when both phenotypes were green.

In *C. creticus*, anthocyanin concentration and the F_{686}/F_{736} ratio was monitored for a period of six months, encompassing both the ‘green’ and the ‘red’ period of the year (Fig. 3). Measurements started before the commencement of anthocyanin accumulation in the red phenotype (first week of December), continued during the whole ‘red period’, and ended in late May, when both phenotypes were green again. As it is shown, there is no difference in the F_{686}/F_{736} ratio (Fig. 3B) at the green period of the year (early to mid-December and May), when anthocyanin content was almost zero in both leaf

types. Leaf redness appeared in the red phenotype in late December and anthocyanin concentration increased steeply within few days and more gradually thereafter (Fig. 3A). Anthocyanin content was maintained at high levels until mid-March, while in green leaves, it remained constantly low. During the same period a concurrent rise of the F_{686}/F_{736} ratio was observed in both phenotypes, which, however, was more pronounced in red leaves. In particular, the F_{686}/F_{736} ratio was gradually doubled in green leaves until mid-February and declined thereafter to reach the initial value in early April. In red leaves, the F_{686}/F_{736} ratio increased more sharply, *i.e.*, ~ 3.3 -fold in late January, and remained high until mid-March, as long as anthocyanin concentrations were at maximum. The values obtained in late-May correspond to new leaves (both green).




Fig. 2. Area-based total chlorophyll and anthocyanin concentrations and Chl *a/b* ratio from red (white bar) and green (grey bar) leaves or areas of the same species. In *Ricinus communis* and *Photinia × fraseri*, young leaves are presented by hatched bar (red A) and white bar (red B), respectively. Data are means \pm SE of the indicated (*n*) independent extractions. Within each species, different letters indicate statistically significant differences ($p < 0.05$) in the measured parameters between red and green leaves. In *Cistus creticus*, values refer to the 'red measuring period'.

Discussion

To test whether the presence of an anthocyanin screen leads to a readjustment of photosystem stoichiometry, compatible to the shade acclimation hypothesis, we took advantage of the intra-individual, intra-species, or intra-leaf variation in the expression of the anthocyanic trait of the selected plants (see the text table in 'Materials and methods'). In the first four species, transient leaf redness is developmentally determined (young red leaves which become green upon maturation), while in *C. creticus* is induced in mature leaves of certain individuals (red phenotype) by the combination of low temperatures and high light intensity during winter. The last two species display permanent anthocyanin accumulation either homogeneously in the whole leaf area (*P. cerasifera*) or in patchiness (*C. blumei*).

In *P. × fraseri* and *R. communis*, the compared leaves (young red vs. mature green) co-occur at the same individual, giving the opportunity to bypass possible species- and/or site-dependent effects in our measurements. Yet, unavoidably, the interference of the different developmental stage could not be excluded. To overcome possible age effects, in the next four species, leaves of the same physiological age (either young red vs. young green or mature red vs. mature green) were compared. Leaves were harvested from neighboring different individuals/phenotypes of the same species growing under apparently similar environmental conditions to eliminate site effects of water and nutrient availability. Although species-dependent differences could not be excluded in the case of *P. cerasifera × P. domestica*, the compared mature green and red leaves co-occur on the same chimeric tree, possessing a permanently red (*P. cerasifera*) and a




Fig. 3. Monitoring of anthocyanin contents (A) and F_{686}/F_{736} ratio (B) in mature leaves from green (grey bars) and red (open bars) phenotypes of *Cistus creticus* sampled on the indicated dates. On each measuring date, data are means from two independent extractions. Anthocyanin contents in green leaves were stable throughout the experimental period (average during red period: 1.32 ± 0.26). Leaves senesce and fall from mid-April to mid-May in parallel with new leaf emergence. The May sampling concerns new green leaves for both phenotypes. Differences between the two phenotypes are significant only during the 'red' period.

permanently green (*P. domestica*) stem. Hence, both leaf types experience the same environmental limitations. Finally, *C. blumei* seems to fulfil all the requisites, since anthocyanin-rich and anthocyanin-less chlorophyllous regions from the same leaf were compared.

It is evident from our results that, compared to green, red leaves display higher emission intensity at 686 nm and, accordingly, higher F_{686}/F_{736} values, indicative of an enhanced PSII/PSI ratio (Fig. 1). In all cases, the 77 K fluorescence spectra of isolated thylakoids showed a typical profile with two main peaks at 686 and 736 nm, corresponding to the emissions of PSII core complex and LHCI-PSI, respectively. A shoulder at 695 nm was also observed, ascribable to CP47 emission. Since the peak at 686 nm was better defined in our normalized spectra, we used the fluorescence amplitude at this wavelength to estimate the PSII/PSI ratio (Andrizhiyevskaya *et al.* 2005, Tang *et al.* 2005, Velitchkova and Popova 2005, Lamb *et al.* 2018).

The significantly higher PSII/PSI ratio of red leaves was observed in all tested species (except of *Q. coccifera*, where the same was observed as a trend), regardless of the inductive agent of anthocyanin accumulation (ontogenetic or environmental) or their histological localization (epidermis or mesophyll). The latter agrees with previous

findings suggesting that the absorption capacity of anthocyanins for green light is largely independent of their histological localization (Gould *et al.* 2002a, 2018), although in a comparative study with red and green leaves of *Quintinia serrata*, it was proposed that the tissue location of anthocyanins is important for the absorption profile of the leaf (Gould *et al.* 2002b). Moreover, as it was shown in *C. creticus*, where the differences between the two phenotypes were monitored during both the 'red' and the 'green' period, the increase of PSII/PSI ratio followed the pattern of anthocyanin accumulation (Fig. 3). The PSII/PSI ratio increased in both phenotypes during winter, indicating that photosystem stoichiometry was affected by several environmental factors, apart from incident light, such as low temperature (Walters 2005). Yet, the increase was more pronounced in the red leaves and the differences were abolished when the red phenotype reverted to green. Accordingly, we may consider the above-mentioned increase as a response to the unbalanced light quality penetrating into a red leaf to regulate the excitation energy afforded in the two photosystems, analogous to the corresponding shade acclimation (Melis 1984, Chow *et al.* 1990, Anderson *et al.* 1995).

The light prevailing under natural shade (*e.g.*, under a dense plant or forest canopy), apart from being of significantly lower intensity, it is relatively enriched in green and far-red while depleted in blue/red photons (Anderson *et al.* 1995, Lichtenthaler *et al.* 2013). Although, the light available under natural shade conditions tends to induce opposite effects in PSII/PSI ratio, *i.e.*, a decrease due to low intensity and an increase due to uneven spectral distribution, the resulting adjustment of photosystem stoichiometry is considered rather an acclimation to spectral quality than intensity (Melis 1991, Anderson *et al.* 1995, Murchie and Horton 1998). When FR-enriched light reaches chloroplasts, PSI receives much more excitation energy compared to PSII, thus PSI operation is favored. Conversely, incident light enriched in blue/red photons preferentially excites PSII while PSI excitation is limiting for photosynthesis. As it has been repeatedly shown with plants grown under illumination of different quality, the enrichment of light in FR photons (*i.e.*, PSI light) leads to an increase in the relative PSII/PSI ratio, to optimize light utilization by the two photosystems (Melis 1984, Glick *et al.* 1985, Chow *et al.* 1990, Melis 1991, Walters 2005). Accordingly, in true shade leaves, the PSII/PSI ratio is increased. The shade imposed by anthocyanin accumulation in a red leaf, however, apart from similarities to that of natural (*i.e.*, canopy-like) conditions, displays also characteristic differences. Foliar anthocyanins, being almost always red (Harborne 1976), absorb strongly in the green/yellow and less in the blue wavebands, while they are transparent in far-red band. Concerning their absorbance ability in the red portion of the spectrum, it is considered negligible or at least very low (Karabourniotis *et al.* 1999, Gould *et al.* 2002b, Steyn *et al.* 2002, Kyparissis *et al.* 2007). As a result, while natural shade is green- and FR-enriched, the mesophyll of a red leaf is exposed to light severely depleted in the green/yellow portion of the visible spectrum. In addition, anthocyanins

may also display an appreciable tail over the red band (600–630 nm), especially when they are conjugated with other (colorless) flavonoids, reducing even more the R/FR ratio compared to natural shade (Gould *et al.* 2018, Landi *et al.* 2021). Green and yellow light is better absorbed by PSII, whereas FR light is preferentially absorbed by PSI (Glick *et al.* 1985, Pfannschmidt 2005).

Several reports in the literature support the assumption that anthocyanin accumulation induces some morphological and physiological adjustments in red leaves, resembling, at least partly, with those displayed by the true shade (green) ones. Thus, compared to their green counterparts, red leaves tend to be thinner with lower LMA values, reduced Chl *a/b* and Car/Chls ratios, lower net photosynthetic rate, and lower or similar area based total Chls (Burger and Edwards 1996, Manetas *et al.* 2003, Hughes and Smith 2007, Hughes *et al.* 2007, Kyparissis *et al.* 2007, Kytridis *et al.* 2008, Zeliou *et al.* 2009). In many cases, however, some traits of red leaves are not compatible to the shade-acclimation syndrome. According to Kyparissis *et al.* (2007), these deviations could be explained by the selective absorption of anthocyanins, especially in the green/yellow part of the spectrum, leading to thinner rather than shade leaves, in which the missing layers had the more shade characteristics.

As already mentioned in ‘Introduction’, a typical ‘shade’ character of leaves growing under a dense canopy is the lower Chl *a/b* ratio, which reflects a higher investment in light harvesting (containing both Chl *a* and *b*) relative to core complexes (containing only Chl *a*) and/or a higher PSII/PSI ratio (Anderson 1986, Murchie and Horton 1998, Lichtenthaler and Babani 2004, Hughes *et al.* 2007, Lichtenthaler *et al.* 2007). In a comparative study with anthocyanic and non-anthocyanic species (Hughes *et al.* 2007), Chl *a/b* ratio increased gradually with leaf age in the developing red leaves (in parallel to the decline of anthocyanin content), while in the corresponding green ones, it remained relatively constant at much higher values during development. Accordingly, the reduced Chl *a/b* ratio in reds was attributed to the shade imposed by anthocyanins (Hughes *et al.* 2007). In the present study, red leaves/areas in five of our tested species displayed a lower Chl *a/b* ratio compared to their green counterparts (Fig. 2C–G), regardless of their developmental stage (young or mature). Similarly, the lower Chl *a/b* ratio in reds, ascribed to the shade imposed by anthocyanins, was reported in several previous investigations (Manetas *et al.* 2003, Hughes and Smith 2007, Hughes *et al.* 2007, Kyparissis *et al.* 2007, Zeliou *et al.* 2009). In *R. communis* and *P. × fraseri*, however, no differences were observed between the leaf types, despite the much higher anthocyanin content of red leaves (Fig. 2A,B). A possible explanation would be that chloroplasts in the mature leaves are self-shaded by the much higher Chl concentration (combined with the increased thickness), leading to similar Chl *a/b* values in green and red leaves. As it has been shown, changes of the chlorophyll content affect the proportion of ‘shaded’ chloroplasts within a leaf, which results in changes of the Chl *a/b* ratio (Walters 2005). Yet, in *P. cerasifera* × *P. domestica* and *C. creticus*,

where Chl concentration was also higher in green leaves, the above argument is not confirmed, unless if the steeper gradient of Chl content between red and green leaves of *R. communis* and *P. × fraseri* contributes more effectively to the modulation of Chl *a/b* ratio. In any case, in all tested species of the present study, red leaves displayed a higher PSII/PSI ratio, irrespective of the anthocyanin accumulation pattern and the difference in Chl *a/b* ratio. According to Murchie and Horton (1998), a change in Chl *a/b* is not necessarily associated to changes in PSII/PSI ratio.

References

- Anderson J.M.: Photoregulation of the composition, function, and structure of thylakoid membranes. – *Annu. Rev. Plant Phys.* **37**: 93-136, 1986.
- Anderson J.M., Chow W.S., Park Y.-I.: The grand design of photosynthesis: Acclimation of the photosynthetic apparatus to environmental cues. – *Photosynth. Res.* **46**: 129-139, 1995.
- Andrizhiyevskaya E.G., Chojnicka A., Bautista J.A. *et al.*: Origin of the F685 and F695 fluorescence in Photosystem II. – *Photosynth. Res.* **84**: 173-180, 2005.
- Archetti M., Döring T.F., Hagen S.B. *et al.*: Unravelling the evolution of autumn colours: an interdisciplinary approach. – *Trends Ecol. Evol.* **24**: 166-173, 2009.
- Burger J., Edwards G.E.: Photosynthetic efficiency and photodamage by UV and visible radiation in red versus green leaf coleus varieties. – *Plant Cell Physiol.* **37**: 395-399, 1996.
- Chalker-Scott L.: Environmental significance of anthocyanins in plant stress responses. – *Photochem. Photobiol.* **70**: 1-9, 1999.
- Chow W.S., Melis A., Anderson J.M.: Adjustments of photosystem stoichiometry in chloroplasts improve the quantum efficiency of photosynthesis. – *P. Natl. Acad. Sci. USA* **87**: 7502-7506, 1990.
- Close D.C., Beadle C.L.: The ecophysiology of foliar anthocyanins. – *Bot. Rev.* **69**: 149-161, 2003.
- Glick R.E., McCauley S.W., Melis A.: Effect of light quality on chloroplast-membrane organization and function in pea. – *Planta* **164**: 487-494, 1985.
- Gould K.S.: Nature’s Swiss army knife: the diverse protective roles of anthocyanins in leaves. – *J. Biomed. Biotechnol.* **2004**: 314-320, 2004.
- Gould K.S., Jay-Allemand C., Logan B.A. *et al.*: When are foliar anthocyanins useful to plants? Re-evaluation of the photoprotection hypothesis using *Arabidopsis thaliana* mutants that differ in anthocyanin accumulation. – *Environ. Exp. Bot.* **154**: 11-22, 2018.
- Gould K.S., Neill S., Vogelmann T.C.: A unified explanation for anthocyanins in leaves? – *Adv. Bot. Res.* **37**: 167-192, 2002a.
- Gould K.S., Vogelmann T.C., Han T., Clearwater M.J.: Profiles of photosynthesis within red and green leaves of *Quintinia serrata* A. Cunn. – *Physiol. Plantarum* **116**: 127-133, 2002b.
- Harborne J.B.: The anthocyanin pigments. – In: Harborne J.B. (ed.): *Comparative Biochemistry of the Flavonoids*. Pp. 1-36. Academic Press, London 1976.
- Hoch W.A., Zeldin E.L., McCowan B.H.: Physiological significance of anthocyanins during autumnal leaf senescence. – *Tree Physiol.* **21**: 1-8, 2001.
- Hogewoning S.W., Wientjes E., Douwstra P. *et al.*: Photosynthetic quantum yield dynamics: from photosystems to leaves. – *Plant Cell* **24**: 1921-1935, 2012.
- Hrazdina G., Wagner G.J., Siegelman H.W.: Subcellular localization of enzymes of anthocyanin biosynthesis in protochloroplasts. – *Phytochemistry* **17**: 53-56, 1978.

- Hughes N.M.: Winter leaf reddening in 'evergreen' species. – *New Phytol.* **190**: 573-581, 2011.
- Hughes N.M., Morley C.B., Smith W.K.: Coordination of anthocyanin decline and photosynthetic maturation in juvenile leaves of three deciduous tree species. – *New Phytol.* **175**: 675-685, 2007.
- Hughes N.M., Smith W.K.: Attenuation of incident light in *Galax urceolata* (Diapensiaceae): concerted influence of adaxial and abaxial anthocyanic layers on photoprotection. – *Am. J. Bot.* **94**: 784-790, 2007.
- Karabourniotis G., Bornman J.F., Liakoura V.: Different leaf surface characteristics of three grape cultivars affect leaf optical properties as measured with fibre optics: possible implication in stress tolerance. – *Aust. J. Plant Physiol.* **26**: 47-53, 1999.
- Krause G.H., Weis E.: Chlorophyll fluorescence and photosynthesis: the basics. – *Annu. Rev. Plant Phys.* **42**: 313-49, 1991.
- Kyparissis A., Grammatikopoulos G., Manetas Y.: Leaf morphological and physiological adjustments to the spectrally selective shade imposed by anthocyanins in *Prunus cerasifera*. – *Tree Physiol.* **27**: 849-857, 2007.
- Kytridis V.-P., Karageorgou P., Levizou E., Manetas Y.: Intraspecific variation in transient accumulation of leaf anthocyanins in *Cistus creticus* during winter: evidence that anthocyanins may compensate for an inherent photosynthetic and photoprotective inferiority of the red-leaf phenotype. – *J. Plant Physiol.* **165**: 952-959, 2008.
- Kytridis V.-P., Manetas Y.: Mesophyll versus epidermal anthocyanins as potential *in vivo* antioxidants: evidence linking the putative antioxidant role to the proximity of oxy-radical source. – *J. Exp. Bot.* **57**: 2203-2210, 2006.
- Lamb J.J., Røkke G., Hohmann-Marriott M.F.: Chlorophyll fluorescence emission spectroscopy of oxygenic organisms at 77 K. – *Photosynthetica* **56**: 105-124, 2018.
- Landi M., Agati G., Fini A. *et al.*: Unveiling the shade nature of cyanic leaves: A view from the "blue absorbing side" of anthocyanins. – *Plant Cell Environ.* **44**: 1119-1129, 2021.
- Lee D.W., Collins T.M.: Phylogenetic and ontogenetic influences on the distribution of anthocyanins and betacyanins in leaves of tropical plants. – *Int. J. Plant Sci.* **162**: 1141-1153, 2001.
- Lev-Yadun S., Dafni A., Flaishman M.A. *et al.*: Plant coloration undermines herbivorous insect camouflage. – *BioEssays* **26**: 1126-1130, 2004.
- Lichtenthaler H.K., Ač A., Marek M.V. *et al.*: Differences in pigment composition, photosynthetic rates and chlorophyll fluorescence images of sun and shade leaves of four tree species. – *Plant Physiol. Bioch.* **45**: 577-588, 2007.
- Lichtenthaler H.K., Babani F., Navrátil M., Buschmann C.: Chlorophyll fluorescence kinetics, photosynthetic activity, and pigment composition of blue-shade and half-shade leaves as compared to sun and shade leaves of different trees. – *Photosynth. Res.* **117**: 355-366, 2013.
- Lichtenthaler H.K., Babani F.: Light adaptation and senescence of the photosynthetic apparatus: changes in pigment composition, chlorophyll fluorescence parameters and photosynthetic activity during light adaptation and senescence of leaves. – In: Papageorgiou G.C., Govindjee (ed.): *Chlorophyll a Fluorescence: A Signature of Photosynthesis. Advances in Photosynthesis and Respiration.* Pp. 713-736. Springer, Dordrecht 2004.
- Lichtenthaler H.K., Wellburn A.R.: Determinations of total carotenoids and chlorophylls *a* and *b* of leaf extracts in different solvents. – *Biochem. Soc. T.* **11**: 591-592, 1983.
- Lindoo S.J., Caldwell M.M.: Ultraviolet-B radiation-induced inhibition of leaf expansion and promotion of anthocyanin production – lack of involvement of low irradiance phytochrome system. – *Plant Physiol.* **61**: 278-282, 1978.
- Manetas Y.: Why some leaves are anthocyanic and why most anthocyanic leaves are red? – *Flora* **201**: 163-177, 2006.
- Manetas Y., Petropoulou Y., Psaras G.K., Drinia A.: Exposed red (anthocyanic) leaves of *Quercus coccifera* display shade characteristics. – *Funct. Plant Biol.* **30**: 265-270, 2003.
- Melis A.: Light regulation of photosynthetic membrane structure, organization and function. – *J. Cell. Biochem.* **24**: 271-285, 1984.
- Melis A.: Dynamics of photosynthetic membrane composition and function. – *BBA-Bioenergetics* **1058**: 87-106, 1991.
- Merzlyak M.N., Chivkunova O.B., Solovchenko A.E., Naqvi K.R.: Light absorption by anthocyanins in juvenile, stressed, and senescing leaves. – *J. Exp. Bot.* **59**: 3903-3911, 2008.
- Murchie E.H., Horton P.: Contrasting patterns of photosynthetic acclimation to the light environment are dependent on the differential expression of the responses to altered irradiance and spectral quality. – *Plant Cell Environ.* **21**: 139-148, 1998.
- Murray J.R., Hackett W.P.: Dihydroflavonol reductase activity in relation to differential anthocyanin accumulation in juvenile and mature phase *Hedera helix* L. – *Plant Physiol.* **97**: 343-351, 1991.
- Neill S., Gould K.S.: Optical properties of leaves in relation to anthocyanin concentration and distribution. – *Can. J. Bot.* **77**: 1777-1782, 2000.
- Papageorgiou G.C., Govindjee: *Chlorophyll a Fluorescence: A Signature of Photosynthesis. Advances in Photosynthesis and Respiration.* Pp. 818. Springer, Dordrecht 2004.
- Pfannschmidt T.: Acclimation to varying light qualities: toward the functional relationship of state transitions and adjustment of photosystem stoichiometry. – *J. Phycol.* **41**: 723-725, 2005.
- Pietrini F., Massacci A.: Leaf anthocyanin content changes in *Zea mays* L. grown at low temperature: significance for the relationship between the quantum yield of PSII and the apparent quantum yield of CO₂ assimilation. – *Photosynth. Res.* **58**: 213-219, 1998.
- Šiffl P., Hunalová I., Roháček K.: Light-induced quenching of chlorophyll fluorescence at 77 K in leaves, chloroplasts and Photosystem II particles. – *Photosynth. Res.* **65**: 219-229, 2000.
- Steyn W.J., Wand S.J.E., Holcroft D.M., Jacobs G.: Anthocyanins in vegetative tissues: a proposed unified function in photoprotection. – *New Phytol.* **155**: 349-361, 2002.
- Tang Y., Wen X., Lu C.: Differential changes in degradation of chlorophyll-protein complexes of photosystem I and photosystem II during flag leaf senescence of rice. – *Plant Physiol. Bioch.* **43**: 193-201, 2005.
- Velitchkova M., Popova A.: High light-induced changes of 77 K fluorescence emission of pea thylakoid membranes with altered membrane fluidity. – *Bioelectrochemistry* **67**: 81-90, 2005.
- Walters R.G.: Towards an understanding of photosynthetic acclimation. – *J. Exp. Bot.* **56**: 435-447, 2005.
- Weis E.: Chlorophyll fluorescence at 77 K in intact leaves: characterization of a technique to eliminate artifacts related to self-absorption. – *Photosynth. Res.* **6**: 73-86, 1985.
- Yiotis C., Petropoulou Y., Manetas Y.: Evidence for light-independent and steeply decreasing PSII efficiency along twig depth in four tree species. – *Photosynthetica* **47**: 223-231, 2009.
- Zeliou K., Manetas Y., Petropoulou Y.: Transient winter leaf reddening in *Cistus creticus* characterizes weak (stress-sensitive) individuals, yet anthocyanins cannot alleviate the adverse effects on photosynthesis. – *J. Exp. Bot.* **60**: 3031-3042, 2009.