a Open Access Full Text Article

Mean Platelet Volume and Its Predictive Role in Chronic Heart Failure: Technical Observations [Letter]

John L Frater

Department of Pathology and Immunology, Washington University, St. Louis, Missouri, USA

Correspondence: John L Frater, Department of Pathology and Immunology, Washington University School of Medicine, 660 South Euclid Avenue, Box 8118, St. Louis, Missouri, 63110-1093, USA, Tel +1 314362-1553, Email jfrater@wustl.edu

Dear editor

Andrei et al have recently published "Mean Platelet Volume: A Possible Predictor for Patients with Decompensated Chronic Heart Failure" in International Journal of General Medicine.¹ I congratulate the authors for their paper, which adds to the expanding literature on complete blood cell count (CBC)-derived biomarkers such as the mean platelet volume (MPV) and their potential roles in the care of patients with cardiac disease. I would like to add some technical comments on the MPV that would be of interest to the readership of International Journal of General Medicine who are interested in using the MPV in the clinical setting.

First, it is noteworthy that the MPV is particularly sensitive to preanalytical variables that may result in erroneous values. These include prolonged time between phlebotomy and analysis, anticoagulant type, storage conditions (including ambient temperature), and transport conditions (such as pneumatic tube transport).² These factors may be particularly difficult to control, particularly in outpatient and emergency settings where specimens are collected throughout the workday and couriered to a central laboratory for analysis. I note that Andrei et al state that their specimens were processed by the hospital laboratory on the same day as phlebotomy.¹ I would add that it is considered best practice to process and analyze blood specimens within 1 hour of phlebotomy, since delays as brief as one hour have been noted to result in bias in the result of several CBC-based analytes, including platelet parameters such as the MPV.³

A second issue relates to the degree to which studies that rely on MPV data are generalizable. I note that the authors mentioned the analyzer and manufacturer used in this study (Beckman Coulter ACT 5 diff AL, Beckman Coulter, Miami, FL, USA). Since there are well-described issues with harmonization of the MPV among analyzers, even those that use a similar methodology,⁴ it is important to report this information in the interest of complete transparency. Although an internationally recognized standard for the MPV has not yet been established, the International Council for Standardization in Hematology (ICSH) has recently evaluated a commercially available fixed porcine platelet preparation for this purpose,⁵ and widespread use of this or similar product may improve the reproducibility of the MPV across different instrumentation platforms.

A third problem, which is a common issue in studies that evaluate the clinical use of the MPV, relates to the narrow dynamic range of the MPV.⁵ For example, in the study of Andrei et al the difference of the mean value of the MPV between the control and study populations was <1 fL (8.74 fL vs 9.08 fL) In view of the current technological limitations and issues with preanalytical and analytical variables, it would be difficult to meaningfully interpret individual patient results, particularly in individuals with borderline MPV values.

In closing, I again thank Andrei et al for their paper, which adds to the growing literature on clinical implementation of the MPV. Although at this time it may be premature to recommend the MPV for the use described in the paper, in a setting with stringent control of preanalytical variables and improved instrument development processes, the MPV and other CBC-derived data may be of increased use in the clinical setting.

cc 0 (so 2022 Frater. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms.php and by no incorporate the Creative Commons Attribution — Non Commercial (unported, v3.0) License (http://creativecommons.org/licenses/by-nc/3.0/). By accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (https://www.dovepress.com/terms.php).

Disclosure

The author reports no conflicts of interest in this communication.

References

- Andrei CL, Catana A, Sinescu CJ, et al. Mean platelet volume: a possible predictor for patients with decompensated chronic heart failure. Int J Gen Med. 2022;15:4131–4140. doi:10.2147/IJGM.S362257
- 2. Frater JL. Preanalytical and analytical factors in the mean platelet volume: a potential cause of heterogeneity in studies of erectile dysfunction. Int J Impot Res. 2022. doi:10.1038/s41443-022-00577-1
- Kayadibi H, Acar IA, Cam S. Stability of complete blood count parameters depends on the storage temperature, storage time, transport position and selected stability criterion. Scand J Clin Lab Invest. 2020;80(6):470–478. doi:10.1080/00365513.2020.1783570
- 4. Lippi G, Pavesi F, Pipitone S. Evaluation of mean platelet volume with four hematological analyzers: harmonization is still an unresolved issue. *Blood Coagul Fibrinolysis*. 2015;26(2):235–237. doi:10.1097/MBC.0000000000220
- 5. Harrison P, Price J, Didembourg M, et al. Feasibility of a mean platelet volume standard: an international council for standardization in hematology (ICSH) inter-laboratory study. *Platelets*;2022. 1–9. doi:10.1080/09537104.2022.2060956

Dove Medical Press encourages responsible, free and frank academic debate. The content of the International Journal of General Medicine 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the International Journal of General Medicine editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

International Journal of General Medicine

Dovepress

Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/international-journal-of-general-medicine-journal

https://doi.org/10.2147/IJGM.\$382622

6356 🛐 🏏 in 🕨 DovePress