

Abstract

Cite this article: (2021). State of the Art. *European Psychiatry* 64(S1), S2–S3.

State of the Art

The clinical characterisation of the patient with primary psychosis aimed at personalisation of management

SOA0003

The clinical characterisation of the patient with primary psychosis aimed at personalisation of management

M. Maj

Department of Psychiatry, University of Campania L. Vanvitelli, Naples, Naples, Italy
doi: 10.1192/j.eurpsy.2021.34

The view that the management of the patient with primary psychosis should be personalized is currently endorsed by the vast majority of clinicians, but this personalization is lacking or inadequate in most clinical contexts. We describe systematically the salient domains that should be considered in the characterization of the individual patient with primary psychosis aimed at personalization of management. These include positive and negative symptom dimensions, other psychopathological components, onset and course, neurocognition and social cognition, neurodevelopmental indicators; social functioning, quality of life and unmet needs; clinical staging, antecedent and concomitant psychiatric conditions, physical comorbidities, family history, history of obstetric complications, early and recent environmental exposures, protective factors and resilience, and internalized stigma. For each domain, simple assessment instruments are identified that could be considered for use in clinical practice and included in standardized decision tools. A management of primary psychosis is encouraged which takes into account all the available treatment modalities whose efficacy is supported by research evidence, selects and modulates them in the individual patient on the basis of the clinical characterization, addresses the patient's needs in terms of employment, housing, self-care, social relationships and education, and offers a focus on identity, meaning and resilience.

Disclosure: No significant relationships.

Targets and outcomes of psychotherapies for mental disorders

SOA0004

Targets and outcomes of psychotherapies for mental disorders

P. Cuijpers

Department of Clinical, Neuro and Developmental Psychology, Vrije Universiteit Amsterdam, Amsterdam, Netherlands
doi: 10.1192/j.eurpsy.2021.35

Because the causal pathways to mental disorders are largely unknown, it is also difficult to decide what the targets and outcomes of psychotherapies should be. In this presentation I will give an overview of the main types of targets and outcomes of therapies, as well as a brief overview of some of the main results of research on these types. The most important outcomes are symptom reduction, personal targets and outcomes from the patient's perspective, improvement of quality of life, intermediate outcomes depending on the theoretical framework of the therapist, negative outcomes, and economic outcomes. By far the most research has focused on symptom reduction. In this presentation I will focus on psychotherapies in

© The Author(s), 2021. Published by Cambridge University Press on behalf of the European Psychiatric Association. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



general, but will illustrate the findings with research on psychotherapies for depression, because these have been examined most extensively. Furthermore, the perspective of different stakeholders will be presented: Patients, relatives, therapists, employers, health care providers and society at large each have

their own perspectives on targets and outcomes. The perspective of patients should have more priority in research and standardization of measures across trials is much needed.

Disclosure: No significant relationships.