

# Effects of 17 $\beta$ -Estradiol and Estrogen Receptor Antagonists on the Proliferation of Gastric Cancer Cell Lines

Myung-Jin Kim, Sung-Il Cho, Kun-Ok Lee<sup>1</sup>, Hyung-Joon Han, Tae-Jin Song, and Seong-Heum Park

Department of Surgery, Korea University College of Medicine, Seoul, <sup>1</sup>HBP Research Laboratory,  
Department of Surgery, Korea University Ansan Hospital, Ansan, Korea

**Purpose:** The aims of this study were as follow: 1) to describe the expression status of estrogen receptor- $\alpha$  and - $\beta$  mRNAs in five gastric carcinoma cell lines; 2) to evaluate *in vitro* the effects of 17 $\beta$ -estradiol and estrogen receptor antagonists on the proliferation of the cell lines.

**Materials and Methods:** Detection of estrogen receptor- $\alpha$  and estrogen receptor- $\beta$  mRNA in five human gastric cancer cell lines (AGS, KATO III, MKN28, MKN45 and MKN74) was made by the reverse transcription-polymerase chain reaction system. To evaluate the effect of 17 $\beta$ -estradiol and estrogen receptor antagonists on the proliferation of gastric cancer cell line, the cell lines which expressed both estrogen receptors were chosen and treated with 17 $\beta$ -estradiol and estrogen receptor antagonists (methyl-piperidino-pyrazole and pyrazolo [1,5-a] pyrimidine). Cell proliferation was assessed with the methylthiazol tetrazolium test.

**Results:** Estrogen receptor- $\alpha$  and estrogen receptor- $\beta$  mRNAs were expressed in three (KATO III, MKN28 and MKN45) and all of the five gastric cancer cell lines, respectively. At higher concentrations, 17 $\beta$ -estradiol inhibited cell growth of MKN28, MKN45 and KATO III cell lines. Neither estrogen receptor- $\alpha$  nor estrogen receptor- $\beta$  antagonist blocked the anti-proliferative effect of 17 $\beta$ -estradiol.

**Conclusions:** Our results indicate that estrogen receptor- $\beta$  mRNAs are preferentially expressed in gastric cancers and also imply that hormone therapy rather than estrogen receptor blockers may be a useful strategy for the treatment of estrogen receptor- $\beta$  positive gastric cancer. Its therapeutic significance in gastric cancer are, however, limited until more evidence of the roles of estrogen receptors in the gastric cancer are accumulated.

**Key Words:** Estrogens; Receptors, estrogen; Stomach neoplasms; Cell line

## Introduction

Estrogens regulate the growth, differentiation, and functioning of diverse target tissues, both within and outside of the reproductive system.<sup>1</sup> Although the stomach is not known as a direct target organ of sex hormones, both epidemiologic and animal studies suggest that gastric cancer may be estrogen-sensitive. A male

dominance in the incidence of gastric cancer with a male to female ratio of about 2 : 1 sets up a hypothesis that estrogens may play a protective role against adenocarcinoma of the stomach.<sup>2</sup> The risk of developing gastric cancer was lower among patients with prostate cancer who had been treated with estrogen than in those without such treatment.<sup>3</sup> Studies which have shown that tamoxifen exposure has also been shown to be a risk factor for gastric cancer further support the hypothesis.<sup>4</sup> Reduced risk of gastric cancer in a nationwide cohort study of men with a diagnosis of prostate cancer who had been treated with estrogen were reported.<sup>3</sup> According to the experiment by Furukawa et al.,<sup>5</sup> the carcinogenic N-methyl-N-nitro-N-nitrosoguanidine added to drinking water induced gastric cancer in male rats but not in female, and castrated or estrogen-treated male rats had a lower incidence of gastric cancer compared

Correspondence to: Seong-Heum Park  
Department of Surgery, Korea University Ansan Hospital, 73 Incheon-ro, Seongbuk-gu, Seoul 136-705, Korea  
Tel: +82-2-920-6834, Fax: +82-2-928-1631  
E-mail: pshchw@korea.ac.kr  
Received June 1, 2013  
Revised September 13, 2013  
Accepted September 13, 2013

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

to untreated male rats.

Most of the actions of estrogens are exerted via the estrogen receptor (ER) of target cells, and 2 ER subtypes, ER- $\alpha$  and ER- $\beta$ , mediate the genomic actions of estrogens. The presence of ER- $\alpha$  and  $\beta$  in gastric adenocarcinoma has been also demonstrated.<sup>5,6</sup> The discovery of ERs has raised enormous interests in possible role of these receptors in the carcinogenesis, progression and treatment of gastric cancer.<sup>2</sup> To date, however, only limited data regarding the detection of ERs in gastric tumors and cell lines are available, and significance of ER expression and hormone manipulation in gastric cancer is not established.

The aims of this pilot study were: 1) to describe the expression status of ER- $\alpha$  and - $\beta$  mRNAs in five gastric carcinoma cell lines; 2) to evaluate *in vitro* the effects of 17 $\beta$ -estradiol and ER antagonists on the proliferation of the cell lines.

## Materials and Methods

### 1. Cell lines and chemicals

Six cell lines established from human breast cancer (MCF7) and gastric adenocarcinoma (AGS, KATO III, MKN28, MKN45 and MKN74) were obtained from the Korean Cell Line Bank Center, Seoul, Korea. MCF7 cells were used for positive control for ERs. 17 $\beta$ -estradiol was purchased from Sigma-Aldrich, Yongin, Korea. ER- $\alpha$  antagonist, methyl-piperidino-pyrazole (MPP) and ER- $\beta$  antagonist, pyrazolo [1,5-a] pyrimidine (PHTPP) were purchased from Tocris Bioscience, Bristol, UK. Solutions of 17 $\beta$ -estradiol and ER antagonists (MPP and PHTPP) were dissolved in ethanol (0.3%) and dimethylsulphoxide to 1 mM, respectively and then diluted in the medium to the final concentrations.

### 2. Cell culture

Cell lines were maintained in RPMI 1640 (GIBCO, Grand Island, NY, USA) without phenol red supplemented with 10% fetal bovine serum (FBS; HyClone Laboratories Inc., Logan, UT, USA), 1% penicillin/streptomycin and incubated at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub> in air. Cultured cells were harvested at confluence by means of trypsinization and serially subcultured at a 1 : 4 split ratio.

### 3. Assessment of cell proliferation

The proliferative response of the cell lines was estimated by the colorimetric methylthiazol tetrazolium (MTT) test (Sigma-Aldrich, St Louis, MO, USA). After incubation with a drug, cells seeded in

96-well plates were harvested and pre-incubated with MTT (0.5 mg/ml) for 4 hours at 37°C. The culture medium was discarded, and the blue formazan crystals were solubilized with acidic isopropanol (0.1 N HCl in absolute isopropanol). MTT conversion to formazan by metabolically viable cells was monitored by spectrophotometry at 570 nm.

### 4. Detection of estrogen receptors

For ER analysis of cell lines, Total RNA of cells was extracted from individual cell lines using the TRIzol reagent (Invitrogen, Camarillo, CA, USA) according to the manufacturer's guidelines. Quantification of total RNA was performed by measuring absorbance at OD 260 nm. For cDNA synthesis, a reverse transcription-polymerase chain reaction (RT-PCR) technique (Promega, Madison, WI, USA) was used. One microgram RNA were reverse-transcribed in a final volume of 20  $\mu$ l using the reverse transcriptase and then stored at -20°C. The semi-quantitative RT-PCR was performed in a final volume of 20  $\mu$ l using G-Taq DNA polymerase (Cosmo Genetech, Seoul, Korea) according to manufacturer's instructions. To control for variations in the reaction, all the polymerase chain reaction (PCR) primers were normalized against the  $\beta$ -actin expression. The primer sequences were as follows: ER- $\alpha$  5'-TCCTGATGATTGGTCTCGTCT-3' (forward) and 5'-ACATTTTCCCTGGTTCCTGTC-3' (reverse), ER- $\beta$  5'-AGTCTGGTTCGTGTGAAGGATG-3' (forward) and 5'-ACTTCTCTGTCTCCGCACAAG-3' (reverse), and  $\beta$ -actin 5'-TGTTACAGGAAGTCCCTTGC-3' (forward) and 5'-AAGCAATGCTATCACCTCCC-3' (reverse). Primers were designed and synthesized by Primer 3 (<http://simgene.com/Primer3>). PCR was performed for 30 cycles (denaturation at 95°C for 5 minutes, annealing at 60°C for 1 minute and extension at 72°C for 7 minutes). Five microliter of the reaction mixture was electrophoresed on a 2% agarose gel and visualized by ethidium bromide (MP Bio, Solon, OH, USA), applying 100-bp DNA ladder (RBC Bioscience, New Taipei, Taiwan) to estimate the band sizes.

### 5. Effects of 17 $\beta$ -estradiol and estrogen receptor antagonists on proliferation of gastric cancer cell lines

To investigate the effects of 17 $\beta$ -estradiol under estrogen-depleted culture conditions cell lines which expressed ER- $\alpha$  and ER- $\beta$  mRNAs were plated at a density of  $1 \times 10^4$  cells/100  $\mu$ l of phenol red-free RPMI 1640 containing 10% FBS in 96 well plates (SPL Life Science, Pocheon, Korea). After 24 hours of incubation, when

the cells were approximately 60% confluent, the 17 $\beta$ -estradiol (Sigma-Aldrich) dissolved in ethanol (0.3%) was added to the culture media at concentrations ranging from 5 nM to 32  $\mu$ M. After 24 hours of treatment with 17 $\beta$ -estradiol the cell proliferative response was estimated by MTT test. Each experiment was repeated three times, and all results represent the means produced by three times repeated experiments.

To evaluate the effect of the ER antagonists on the proliferation of gastric cancer cell line, the cell line which expressed both ERs was chosen and treated with the ER- $\alpha$  antagonist MPP (50  $\mu$ mol/L) and/or the ER- $\beta$  antagonist PHTPP (50  $\mu$ mol/L) 30 minutes before the application of 17 $\beta$ -estradiol (32  $\mu$ M) and incubated for 72 hours.

Each experiment was repeated three times, and all results represent the means produced by three times repeated experiments.

## 6. Statistical analysis

The significance of the differences between the control group and each experimental group was evaluated with the Wilcoxon matched pairs test.  $P < 0.05$  were considered statistically significant. IBM SPSS Statistics ver. 20.0 for Windows (IBM Co., Armonk, NY, USA) was used for the statistical analyses.


## Results

Semi-quantitative RT-PCR was performed to measure the levels of two ERs in the human gastric cancer cell lines: five cancer cell


lines (AGS, KATO III, MKN28, MKN45 and MKN74) were used for study and MCF7 cells were used for controls for positive ERs. As shown in Fig. 1A, ER- $\alpha$  mRNA was detected in the KATO III, MKN28, and MKN45, except AGS and MKN74. ER- $\alpha$ / $\beta$ -actin ratio of KATO III, MKN28 and MKN45 were 6.17, 0.86, and 0.04, respectively. In contrast, ER- $\beta$  mRNA was detected in all the five cell lines. ER- $\alpha$ / $\beta$ -actin ratio of AGS, KATO III, MKN28, MKN45 and MKN74 were 1.33, 1.74, 0.76, 2.06, and 1.93, respectively (Fig. 1B).

Fig. 2 shows the effects of increasing concentrations of 17 $\beta$ -estradiol on the proliferation of gastric cancer cell lines. The 17 $\beta$ -estradiol concentrations (from 5 nM to 32 M) did not significantly affect the cell viability in the MKN28 cells. In the MKN45 cells, the 17 $\beta$ -estradiol concentrations (from 5 nM to 4  $\mu$ M) did not significantly affect the cell viability in the MKN45 cells, whereas higher concentrations (8, 16 and 32  $\mu$ M) of 17 $\beta$ -estradiol caused a significant reduction in cell viability (-2.04%, -17.56%, -59.75%, respectively,  $P < 0.01$ ). Viable cells of KATO III decreased in a dose dependent manner, reaching the minimum value at a 17 $\beta$ -estradiol concentration of 32  $\mu$ M. 17 $\beta$ -estradiol at concentrations lower than 1  $\mu$ M did not significantly affect cell viability, whereas the higher concentrations (2, 4, 8, 16, and 32  $\mu$ M) of 17 $\beta$ -estradiol caused a significant reduction in cell viability, -14.21%, -24.13%, -26.8%, -46.93%, and -46.23%, respectively ( $P < 0.01$ ).

To better understand the role of ERs on the proliferation of gastric cancer cell line, KATO III cells were treated with the ER- $\alpha$  antagonist (MPP, 50  $\mu$ mol/L) and/or the ER- $\beta$  antagonist (PHTPP,



**Fig. 1.** Reverse transcription-polymerase chain reaction of estrogen receptor (ER)- $\alpha$  (A) and ER- $\beta$  (B) mRNAs in five human gastric cancer cell lines (AGS, KATO III, MKN28, MKN45 and MKN74). MCF7 cells were used for controls for positive ERs. ER- $\alpha$  mRNA was detected in the KATO III, MKN28 and MKN45. ER- $\alpha$ / $\beta$ -actin ratio of KATO III, MKN28 and MKN45 were 6.17, 0.86 and 0.04, respectively. In contrast, ER- $\beta$  mRNA was detected in all the five cell lines. ER- $\alpha$ / $\beta$ -actin ratio of AGS, KATO III, MKN28, MKN45 and MKN74 were 1.33, 1.74, 0.76, 2.06 and 1.93, respectively.




**Fig. 2.** Effect of 17 $\beta$ -estradiol on the proliferation of three gastric cancer cell lines (MKN28, 45 and KATO III). The plots represent the effects of increasing 17 $\beta$ -estradiol concentrations on the conversion of methylthiazol tetrazolium to formazan. All results represent the means and the 25th & 75th percentiles of three different experiments. \* $P < 0.05$  compared with control (Wilcoxon rank sum test).

50  $\mu\text{mol/L}$ ) 30 minutes before the application of 17 $\beta$ -estradiol. Neither ER- $\alpha$  nor ER- $\beta$  antagonist had effect on anti-proliferative effect of 17 $\beta$ -estradiol (Fig. 3). Proliferation rate of KATO III cells was decreased to 77.8% of control 72 hours after application of 17 $\beta$ -estradiol ( $P < 0.01$ ). The estrogenic effect was not blocked by ER- $\alpha$  antagonist, MPP, where the cell viability was reversed to 76.5% of 17 $\beta$ -estradiol treated control ( $P > 0.05$ , Wilcoxon matched pairs test). ER- $\beta$  antagonist, PHTPP, did not block antiproliferative effect of 17 $\beta$ -estradiol, with a cell viability of 80.8% of 17 $\beta$ -estradiol treated control ( $P > 0.05$ ). When treated with both MPP and PHTPP, cell proliferation was reversed to 87.5% of 17 $\beta$ -estradiol treated control ( $P > 0.05$ ).

## Discussion

Estrogens regulate the growth, differentiation, and functioning of various target tissues, and their effects are mediated by the receptors, ER- $\alpha$  and ER- $\beta$ . ER- $\alpha$  and ER- $\beta$  have distinct cellular distributions; ER- $\alpha$  is predominantly expressed in female sex organs such as the breast, uterus and ovaries, and ER- $\beta$  is far more widely distributed in the body than is ER- $\alpha$ .<sup>7,8</sup>

The oncologic importance of estrogens and ERs in cancers arising in classic target organs for estrogens such as breast and uterine endometrium have been well described.<sup>9-14</sup> Estrogens and their receptors also have an important role in the progression of cancers of gastrointestinal tract which has been believed to be non-target system for estrogens. In contrast to the cancer-promoting



**Fig. 3.** Effect of estrogen receptor (ER)- $\alpha$  and - $\beta$  antagonists on the viability of 17 $\beta$ -estradiol-treated KATO III cell line. Neither ER- $\alpha$  nor ER- $\beta$  antagonist blocked anti-proliferative effect of 17 $\beta$ -estradiol. 17b = 17 $\beta$ -estradiol; M = methyl-piperidino-pyrazole (MPP, ER- $\alpha$  antagonist); P = pyrazolo [1,5-a] pyrimidine (PHTPP, ER- $\beta$  antagonist).

role of estrogen in breast and gynecologic cancers, it functions as a suppressor for gastrointestinal cancers.<sup>15</sup> Current use of estrogen therapy is suggested to be associated with a decreased risk of colorectal cancer among peri- or postmenopausal women,<sup>13,16</sup> and in comparison with tumors with high ER- $\beta$  expression, tumors negative for ER- $\beta$  were associated with advanced cancer stages.<sup>17</sup> Most studies on esophageal adenocarcinoma are consistent in suggesting a detrimental effect and prognostic value for ER- $\beta$ .<sup>7</sup>

Contrasting results have been obtained with regard to the expression level of ERs in gastric cancer. Some studies show that only ER- $\beta$  is expressed in gastric cancer,<sup>18,19</sup> whereas others reported that the both ERs are expressed.<sup>20-22</sup> This prompted us to reevaluate ER status and hormonal modulation of cell growth in human gastric cancer.

Cultured gastric cancer cell line is a useful tool to study the effect of hormones in conditions that are independent of neural, hormonal, and blood flow factors.<sup>23</sup> Past studies performed on gastric cancer cell lines showed contrasting actions of estrogens on the cell growth. Diverse effects of estradiol on the growth of six human gastric xenografts in nude mice were found, including one case of stimulation, two of inhibition and three of unchanged condition.<sup>24</sup> Messa et al.<sup>23,25</sup> showed that 17 $\beta$ -estradiol administration exerts a growth-inhibitory effect on ER-positive cell lines (HGC-27, AGS). On the other hand, Wu et al.<sup>26</sup> found that steroid hormone administration in ER-negative human gastric carcinoma cell line NUGC-3 had a slight stimulatory effect. These studies, however, did not distinguish ER- $\beta$  from ER- $\alpha$ . The discovery of ER- $\beta$ ,<sup>12</sup> a

second ER isoform, has prompted renewed investigation into the mechanism of action of estrogenic molecules. However, the biologic significance of the ERs in gastric cancer still remains to be elucidated.

In the present study, ER- $\alpha$  and ER- $\beta$  mRNAs were expressed in three (KATO III, MKN28 and MKN45) and all of the five gastric cancer cell lines, respectively. ER- $\alpha$  mRNA was not expressed in AGS and MKN74 cell lines and the ER- $\alpha$ / $\beta$ -actin ratios were very low in MKN28 and MKN45 (0.28 and 0.04, respectively). Our results indicated that ER- $\beta$  mRNAs are preferentially expressed in gastric cancers, as Takano et al.<sup>20</sup> reported, and ER- $\beta$  plays more important role in both physiologic and oncologic processes in gastric tissues than ER- $\alpha$ .

Imbalanced ER- $\alpha$ /ER- $\beta$  expression, i.e., an increase in ER- $\alpha$ /ER- $\beta$  mRNA and protein ratios, is a common feature and could be a critical step of estrogen-dependent tumor progression, and ER- $\beta$  seems to play a key role in the mitogenic action of estrogen by providing protection against ER- $\alpha$ -induced hyperproliferation.<sup>14</sup> Our previous study indicated that The ER- $\beta$  positive group was associated with lower tumor stage, Lauren's intestinal type, negative perineural invasion, and free of recurrence, and the ER- $\beta$  positive group had a better 3-year survival compared with the negative group in survival analysis.<sup>19</sup> Xu et al.<sup>27</sup> also reported that ER- $\alpha$  expression and the absence of ER- $\beta$  expression are associated with poor survival.

In our results, however, relations between the levels of mRNA expression of ERs and effects of 17 $\beta$ -estradiol on proliferation of gastric cancer cell lines were not identified. Low concentrations of 17 $\beta$ -estradiol stimulated cell growth of MKN28 and MKN45 cell lines, even though the values were statistically not significant. A significant decrease in the cell growth rate is obtained with 17 $\beta$ -estradiol concentrations equal to or higher than 16  $\mu$ M. On the contrary, 17 $\beta$ -estradiol administration exerted a growth-inhibitory effect on the KATO III. A significant decrease in the cell growth rate is obtained with 17 $\beta$ -estradiol concentrations equal to or higher than 2  $\mu$ M, which was much lower than those required for MKN28 or MKN45. KATO III cells display a higher expression level of ER- $\alpha$  than MKN28 and MKN45. This finding could be explained by the fact that ER- $\alpha$  mRNA expression in KATO III cell line was higher than those in MKN28 and MKN45 cell lines.

Some previous studies suggested that the ER- $\alpha$  pathway may have a role in the progression of gastric cancer and 17 $\beta$ -estradiol enhances gastric cancer cell proliferation *in vitro*. Based on these suggestion, a few clinical trials using an estrogen antagonist,

tamoxifen, have been conducted for the management of patients with ER- $\alpha$  positive gastric cancer. However, the results have not been consistent. An earlier Japanese study with historical control suggested a survival benefit in tamoxifen treated patients with gastric cancer but the sample size was small and patients received chemotherapy.<sup>28</sup> A randomized clinical study by Harrison et al showed that ER- $\alpha$  positive patients survived for a significantly shorter time than untreated controls if they were treated with tamoxifen. ER- $\alpha$  negative patients had a longer survival and tamoxifen did not significantly affect prognosis.<sup>29,30</sup> On the contrary to these old reports, Kameda et al.<sup>31</sup> recently reported that 17 $\beta$ -estradiol-induced cell proliferation was suppressed by pure anti-estrogen drug, ICI 182, 780 and suggested that the development of clinically available antagonists that can sufficiently block estrogen-induced cell proliferation is promising for management of patients with aggressive diffuse-type gastric cancer. They suggested a possibility that different anti-estrogens used in the above studies may explain the discrepancies in results.

In the present study, MPP and PHTPP were used as ER- $\alpha$  and ER- $\beta$  antagonists, respectively. MPP is a highly selective ER- $\alpha$  antagonist with 200 fold greater affinity for ER- $\alpha$  versus ER- $\beta$ ,<sup>32</sup> and PHTPP is a potent ER- $\beta$  antagonist, with negligible activity with ER- $\alpha$ .<sup>33</sup> The present study, however, showed that neither ER- $\alpha$  nor ER- $\beta$  antagonist had effect on anti-proliferative effect of 17 $\beta$ -estradiol. In our previous study, immunohistochemical staining of ER- $\alpha$  showed no positivity in any of the tissue microarrays of gastric adenocarcinoma tissues, although ER- $\alpha$  was expressed in normal and gastric adenocarcinoma tissues on Western blot analysis.<sup>19</sup> The expression of a particular gene mRNA may not always correspond with the expression of the functional protein.<sup>21</sup>

Our present study imply that hormone therapy rather than ER blockers may be a useful strategy for the treatment of gastric cancer in cases of ER- $\beta$  positive gastric cancer. However, our study has some limitations. Our data do not show clear relationship between the effect of 17 $\beta$ -estradiol and ER- $\beta$ . 17 $\beta$ -estradiol showed no effect on two ER- $\beta$  positive cell line. Furthermore, ER antagonists failed to block the effect of 17 $\beta$ -estradiol in this study. The mechanism by which estrogens and their receptors inhibit the growth of cancer cell should be clarified by further studies.

## Acknowledgments

This study was supported by a Korea University grant.

## References

1. Katzenellenbogen BS. Estrogen receptors: bioactivities and interactions with cell signaling pathways. *Biol Reprod* 1996;54:287-293.
2. Chandanos E, Lagergren J. Oestrogen and the enigmatic male predominance of gastric cancer. *Eur J Cancer* 2008;44:2397-2403.
3. Lindblad M, Ye W, Rubio C, Lagergren J. Estrogen and risk of gastric cancer: a protective effect in a nationwide cohort study of patients with prostate cancer in Sweden. *Cancer Epidemiol Biomarkers Prev* 2004;13:2203-2207.
4. Curtis RE, Boice JD Jr, Shriner DA, Hankey BF, Fraumeni JF Jr. Second cancers after adjuvant tamoxifen therapy for breast cancer. *J Natl Cancer Inst* 1996;88:832-834.
5. Furukawa H, Iwanaga T, Koyama H, Taniguchi H. Effect of sex hormones on the experimental induction of cancer in rat stomach - a preliminary study. *Digestion* 1982;23:151-155.
6. Tokunaga A, Kojima N, Andoh T, Matsukura N, Yoshiyasu M, Tanaka N, et al. Hormone receptors in gastric cancer. *Eur J Cancer Clin Oncol* 1983;19:687-689.
7. Yang H, Sukocheva OA, Hussey DJ, Watson DI. Estrogen, male dominance and esophageal adenocarcinoma: is there a link? *World J Gastroenterol* 2012;18:393-400.
8. Sharpe RM. The roles of oestrogen in the male. *Trends Endocrinol Metab* 1998;9:371-377.
9. Beral V; Million Women Study Collaborators. Breast cancer and hormone-replacement therapy in the Million Women Study. *Lancet* 2003;362:419-427.
10. Rossouw JE, Anderson GL, Prentice RL, LaCroix AZ, Kooperberg C, Stefanick ML, et al. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results from the Women's Health Initiative randomized controlled trial. *JAMA* 2002;288:321-333.
11. Lacey JV Jr, Mink PJ, Lubin JH, Sherman ME, Troisi R, Hartge P, et al. Menopausal hormone replacement therapy and risk of ovarian cancer. *JAMA* 2002;288:334-341.
12. Folsom AR, Anderson JP, Ross JA. Estrogen replacement therapy and ovarian cancer. *Epidemiology* 2004;15:100-104.
13. Beral V, Bull D, Reeves G; Million Women Study Collaborators. Endometrial cancer and hormone-replacement therapy in the Million Women Study. *Lancet* 2005;365:1543-1551.
14. Bardin A, Boule N, Lazennec G, Vignon F, Pujol P. Loss of ERbeta expression as a common step in estrogen-dependent tumor progression. *Endocr Relat Cancer* 2004;11:537-551.
15. Nelson HD, Humphrey LL, Nygren P, Teutsch SM, Allan JD. Postmenopausal hormone replacement therapy: scientific review. *JAMA* 2002;288:872-881.
16. Lin KJ, Cheung WY, Lai JY, Giovannucci EL. The effect of estrogen vs. combined estrogen-progestogen therapy on the risk of colorectal cancer. *Int J Cancer* 2012;130:419-430.
17. Rudolph A, Toth C, Hoffmeister M, Roth W, Herpel E, Jansen L, et al. Expression of oestrogen receptor  $\beta$  and prognosis of colorectal cancer. *Br J Cancer* 2012;107:831-839.
18. Matsuyama S, Ohkura Y, Eguchi H, Kobayashi Y, Akagi K, Uchida K, et al. Estrogen receptor beta is expressed in human stomach adenocarcinoma. *J Cancer Res Clin Oncol* 2002;128:319-324.
19. Ryu WS, Kim JH, Jang YJ, Park SS, Um JW, Park SH, et al. Expression of estrogen receptors in gastric cancer and their clinical significance. *J Surg Oncol* 2012;106:456-461.
20. Takano N, Iizuka N, Hazama S, Yoshino S, Tangoku A, Oka M. Expression of estrogen receptor-alpha and -beta mRNAs in human gastric cancer. *Cancer Lett* 2002;176:129-135.
21. Wang M, Pan JY, Song GR, Chen HB, An LJ, Qu SX. Altered expression of estrogen receptor alpha and beta in advanced gastric adenocarcinoma: correlation with prothymosin alpha and clinicopathological parameters. *Eur J Surg Oncol* 2007;33:195-201.
22. Gan L, He J, Zhang X, Zhang YJ, Yu GZ, Chen Y, et al. Expression profile and prognostic role of sex hormone receptors in gastric cancer. *BMC Cancer* 2012;12:566.
23. Messa C, Russo F, Pricci M, Di Leo A. Epidermal growth factor and 17beta-estradiol effects on proliferation of a human gastric cancer cell line (AGS). *Scand J Gastroenterol* 2000;35:753-758.
24. Tokunaga A, Onda M, Kiyama T, Nishi K, Mizutani T, Yoshiyuki T, et al. Contrasting actions of estradiol on the growth of human gastric cancer xenografts in nude mice. *Jpn J Cancer Res* 1989;80:1153-1155.
25. Messa C, Pricci M, Linsalata M, Russo F, Di Leo A. Inhibitory effect of 17beta-estradiol on growth and the polyamine metabolism of a human gastric carcinoma cell line (HGC-27). *Scand J Gastroenterol* 1999;34:79-84.
26. Wu CW, Chang YF, Yeh TH, Chang TJ, Lui WY, P'Eng FK, et al. Steroid hormone receptors in three human gastric cancer cell lines. *Dig Dis Sci* 1994;39:2689-2694.
27. Xu CY, Guo JL, Jiang ZN, Xie SD, Shen JG, Shen JY, et al. Prognostic role of estrogen receptor alpha and estrogen

- receptor beta in gastric cancer. *Ann Surg Oncol* 2010;17:2503-2509.
28. Kitaoka H. Sex hormone dependency and endocrine therapy in diffuse carcinoma of the stomach. *Gan To Kagaku Ryoho* 1983;10:2453-2460.
  29. Harrison JD, Morris DL, Ellis IO, Jones JA, Jackson I. The effect of tamoxifen and estrogen receptor status on survival in gastric carcinoma. *Cancer* 1989;64:1007-1010.
  30. Harrison JD, Watson S, Morris DL. The effect of sex hormones and tamoxifen on the growth of human gastric and colorectal cancer cell lines. *Cancer* 1989;63:2148-2151.
  31. Kameda C, Nakamura M, Tanaka H, Yamasaki A, Kubo M, Tanaka M, et al. Oestrogen receptor-alpha contributes to the regulation of the hedgehog signalling pathway in ERalpha-positive gastric cancer. *Br J Cancer* 2010;102:738-747.
  32. Sun J, Huang YR, Harrington WR, Sheng S, Katzenellenbogen JA, Katzenellenbogen BS. Antagonists selective for estrogen receptor alpha. *Endocrinology* 2002;143:941-947.
  33. Compton DR, Sheng S, Carlson KE, Rebacz NA, Lee IY, Katzenellenbogen BS, et al. Pyrazolo[1,5-a]pyrimidines: estrogen receptor ligands possessing estrogen receptor beta antagonist activity. *J Med Chem* 2004;47:5872-5893.