Quantitative Imaging Informatics for Cancer Research

Andrey Fedorov, PhD1; Reinhard Beichel, PhD2; Jayashree Kalpathy-Cramer, PhD3; David Clunie, MBBS4; Michael Onken, Dipl-Inform5; Jörg Riesmeier, Dipl-Inform⁶; Christian Herz, MSc¹; Christian Bauer, PhD²; Andrew Beers, BA³; Jean-Christophe Fillion-Robin, MSc⁷; Andras Lasso, PhD8; Csaba Pinter, PhD8; Steve Pieper, PhD9; Marco Nolden, PhD10; Klaus Maier-Hein, PhD10; Markus D. Herrmann, MD, PhD11; Joel Saltz, MD, PhD12; Fred Prior, MD, PhD13; Fiona Fennessy, MD, PhD1; John Buatti, MD2; and Ron Kikinis, MD1

PURPOSE We summarize Quantitative Imaging Informatics for Cancer Research (QIICR: U24 CA180918), one of the first projects funded by the National Cancer Institute (NCI) Informatics Technology for Cancer Research program.

METHODS QIICR was motivated by the 3 use cases from the NCI Quantitative Imaging Network. 3D Slicer was selected as the platform for implementation of open-source quantitative imaging (QI) tools. Digital Imaging and Communications in Medicine (DICOM) was chosen for standardization of QI analysis outputs. Support of improved integration with community repositories focused on The Cancer Imaging Archive (TCIA). Priorities included improved capabilities of the standard, toolkits and tools, reference datasets, collaborations, and training and outreach.

RESULTS Fourteen new tools to support head and neck cancer, glioblastoma, and prostate cancer QI research were introduced and downloaded over 100,000 times. DICOM was amended, with over 40 correction proposals addressing QI needs. Reference implementations of the standard in a popular toolkit and standalone tools were introduced. Eight datasets exemplifying the application of the standard and tools were contributed. An open demonstration/connectathon was organized, attracting the participation of academic groups and commercial vendors. Integration of tools with TCIA was improved by implementing programmatic communication interface and by refining best practices for QI analysis results curation.

CONCLUSION Tools, capabilities of the DICOM standard, and datasets we introduced found adoption and utility within the cancer imaging community. A collaborative approach is critical to addressing challenges in imaging informatics at the national and international levels. Numerous challenges remain in establishing and maintaining the infrastructure of analysis tools and standardized datasets for the imaging community. Ideas and technology developed by the QIICR project are contributing to the NCI Imaging Data Commons currently being developed.

JCO Clin Cancer Inform 4:444-453. © 2020 by American Society of Clinical Oncology

Licensed under the Creative Commons Attribution 4.0 License (cc)



ASSOCIATED CONTENT

Data Supplement

Author affiliations and support information (if applicable) appear at the end of this article

Accepted on March 16, 2020 and published at

ascopubs.org/journal/ cci on May 11, 2020: DOI https://doi.org/10. 1200/CCI.19.00165

INTRODUCTION

Medical imaging is increasingly important in cancer applications. 1,2 Few existing imaging biomarkers are used to guide clinical decisions.3 Major efforts are underway to identify, validate, and deploy new imaging tools in the clinic. These efforts rely on the discovery of novel quantitative imaging (QI) biomarkers, which promise to support objective and reproducible characterization of disease and allow for more personalized approaches to diagnosis and therapy. One such effort is led by the National Cancer Institute (NCI) via its Quantitative Imaging Network (QIN) initiative, 4-6 with primary goals including collecting data from ongoing imaging clinical trials, developing innovative methods for data collection and analysis, and establishing consensus on QI methods.4

Practical QIN experience in striving toward those goals highlighted the importance of imaging informatics as applied to QI biomarker development. In the context of radiology, imaging informatics is defined as a subspecialty of radiology concerned with "the study of how information about medical images is exchanged within radiology and throughout the medical enterprise" (p657). Communication of health care information and integration of data from various institutions and sources is critical for development and validation of innovative QI research tools. Research tools need to be harmonized to support data interoperability, reuse, and aggregation in community repositories and enable federated learning and eventual translation of the matured tools into clinical trials. The Informatics Technology for Cancer Research (ITCR) program (https://itcr.cancer.gov/) was established by NCI to support development and



CONTEXT

Key Objective

To present the motivation, approach, results, and vision for the future development for the Quantitative Imaging Informatics for Cancer Research project, funded by the National Cancer Institute Informatics Technology for Cancer Research program.

Knowledge Generated

Free, commercially reusable, open-source tools were developed by the project to support imaging informatics tasks related to quantitative image analysis standardized communication of analysis findings. Contributions to the Digital Imaging and Communications in Medicine (DICOM) standard, collaborations, community engagement activities, and public datasets contributed by the project are summarized.

Relevance

Our contributions can be used to support additional technical and clinical validation of imaging biomarkers, and improve harmonization of clinical and research imaging data. We also set the stage and methodology for future projects to further improve capabilities, support, and adoption of the DICOM standard in imaging research, radiomics, and machine learning.

sustainment of open-source cancer informatics tools spanning all areas of cancer research.

Quantitative Imaging Informatics for Cancer Research (QIICR; U24 CA180918) was one of the first projects funded by ITCR under PAR-12-287. Motivated by challenges encountered by the QIN community, and with several investigators actively or formerly funded by the QIN program, QIICR embarked on a mission to close some of the informatics gaps that were perceived as limitations to the success of QIN.

The QIICR project developed a collection of reusable open-source components to support a range of informatics tasks for QI. In parallel, we worked on improving standardization of QI analysis outputs by refining and improving the existing standard, developing reference tools and datasets implementing the standard, and performing various outreach activities to demonstrate and promote best practices for sharing QI analysis results. We summarize the scope, collaborative approach, vision, and results of the QIICR project over the funded period; discuss the legacy of the project in the context of various ongoing activities; and outline remaining challenges and opportunities that we identified.

METHODS

The QIICR approach was defined by the three aims:

- 1. establishing open-source tools and workflows to support analysis of longitudinal and derived image data;
- 2. supporting standardized representations for communicating QI analysis results; and
- 3. improving interoperability with community image data repositories.

Three collaborating QIN groups joined the QIICR consortium and defined clinical applications, imaging modalities, analysis approaches, and generated data types. They also contributed to the development of tools and supplied representative datasets:

- 1. Quantitative magnetic resonance imaging (MRI) of glioblastoma treatment response (U01CA154601; J.K.-C. at Massachusetts General Hospital);
- 2. Quantitative positron emission tomography (PET)/computed tomography (CT) of head and neck cancer (U01CA140206; J.B. and R.B. at the University of Iowa);
- 3. Multiparametric MRI of prostate cancer (U01CA151261; F.F. at Brigham and Women's Hospital [BWH]).

We committed to share all software tools under a commercially friendly, nonrestrictive software license to facilitate reuse, adoption, and external contributions, without limiting them to the QIN community. 3D Slicer, an extensible open-source platform for imaging research, was selected as a delivery vehicle for end-user capabilities. The goal of the project was to produce open-source implementations of the application-specific analysis tools and processing workflows within 3D Slicer, while also making it feasible to use those tools outside of 3D Slicer and in applications beyond the scope of the QIICR project.

To harmonize the various outputs generated by QI research analysis tools, we used the Digital Imaging and Communications in Medicine (DICOM) standard.9 DICOM defines standard information objects to represent a broad range of acquired and image-derived data and is suitable for clinical, preclinical, and research applications. DICOM is a comprehensive and complex standard, used within the research community primarily as a mechanism for receiving images from clinical systems. Capabilities of DICOM beyond its support of clinical imaging modalities are not well known to researchers. In a typical research workflow, DICOM images obtained from clinical systems are converted into research formats supported by commonly used analysis tools. This approach allows for limited interoperability between specific research tools, but leads to difficulty reusing analysis results for other purposes. Most research formats do not communicate standardized

metadata describing the dataset, do not contain metadata that identifies and relates individual objects, and do not follow a common data model. Standard concepts for identification of the subject, dates, unique identifiers, and spatial or temporal frames of reference are missing. Translation of QI tools into clinical trials, and evaluation of their use for decision support in clinical practice, is a key priority for the QIN. Integration of research tools and the data they generate necessitates the use of an established standard such as DICOM. Our approach identified the types of analysis results that were relevant to the QIICR use cases and remediated the gaps related to use of DICOM for those results along the following directions:

- Refinements and extensions: We evaluated existing capabilities of the standard to understand how those related to the requirements of the QIN use cases, which needs could be addressed by improving documentation or implementing support in tools and toolkits, and which required enhancements to the standard.
- 2. Implementation of the relevant capabilities: To encourage and simplify adoption, we improved support of the relevant components of DICOM in existing open-source tools, developing new tools as necessary.
- 3. Publicly available reference datasets: To evaluate the capabilities of both the standard and its implementations, to support future adopters, and to demonstrate our recommended approach to deriving data curation via concrete examples, we worked on encoding relevant datasets.
- 4. Training and outreach: To lower barriers to adoption, we contributed scholarly publications, training materials, and tutorials explaining the best practices for using DICOM and the tools we developed. We prioritized collaborations with other groups (academic and commercial) developing imaging analysis tools and workstations to encourage, support, and evaluate adoption of DICOM.

For the third aim, we directed our effort to improved integration with The Cancer Imaging Archive (TCIA), ¹⁰ the recommended repository for the QIN investigators. TCIA implements procedures that ensure DICOM imaging data are free from protected health information and maintains resources to support sharing of de-identified data. ¹¹ Increasingly, TCIA is also distributing analysis results alongside the image data. We aimed to simplify the process of interaction of tool users with the repository and worked on DICOM conversion of the QI analysis results in other formats already deposited in TCIA.

RESULTS

Quantitative Image Analysis Tools

A range of batch-processing and interactive tools were developed to support QI workflows for each of the use cases. Batch-processing C++ Slicer Execution Model (SEM) plugins were added for 3D Slicer.⁸ These modules

can be used by other applications adopting SEM.^{12,13} The plugins can also be used as standalone command-line tools outside of 3D Slicer. Interactive tools were implemented as 3D Slicer Python scripted modules. Modules were disseminated using the 3D Slicer extensions infrastructure, which supports automatic testing and packaging of extensions for Windows, macOS, and Linux. The tools are summarized in Table 1 and Figure 1 and on the QIICR Web site. ¹⁴

We engaged in a number of efforts led by QIN to evaluate the developed tools using common reference datasets and to compare the results produced by the various tools available within the network (Table 1). We also conducted a number of studies designed to develop best practices in corresponding QI methodologies. ¹⁵⁻¹⁹

Standardized Representation of QI Analysis Results

To enhance the DICOM standard based on the needs of the use cases, we contributed numerous improvements to the standard (some of those are discussed in detail in Fedorov et al²⁰; see the Data Supplement for the complete list). Per the DICOM process, those updates passed the rigor of public review, revision, and approval by the DICOM standard committee.

Our DICOM implementation efforts improved support of relevant DICOM components in a variety of tools. We added new capabilities to the DICOM Toolkit (DCMTK) C++ library, 21 improving read/write support and the application programming interface (API) for DICOM Segmentation and Parametric Map objects and the DICOM TID 1500 Structured Report template. The API introduced an extra level of abstraction to simplify both reading and writing of those objects. We developed dcmgi,²² a standalone library and set of command-line converters (implementing the SEM interface) between standard and research-oriented representations of the aforementioned objects. dcmgi has subsequently been integrated into a number of independent platforms, such as MITK¹² and ePAD,²³ and has supported data conversion tasks for several other projects.^{24,25} The *Quantitative Reporting* extension of 3D Slicer²² was developed to hide (behind the interactive user interface) the peculiarities of reading and writing QI DICOM objects.

We worked with the CommonTK, ePad, and OHIF communities to create JavaScript implementations of key DICOM processing code to use in Web browsers and other JavaScript environments. An early experiment was to cross-compile DCMTK C++ code to JavaScript, using Emscripten. ^{26,27} More recently, we have been developing a pure-JavaScript suite of libraries to natively implement critical DICOM functionality, including a new version of dcmjs that manages conversion of traditional DICOM binary format to and from JavaScript objects and JavaScript object notation. ²⁸ The new dcmjs is also used as the core library for JavaScript implementation of client and server portions

Tool Name	Tool Type	Use Case	Tool Description	References
dcmqi	T	ŋ	Library and command-line tools for conversion between DICOM and research formats for QI analysis results.	Herz ²²
DSCMRIAnalysis	В	GBM	Modeling of DSC MRI data using the approach described by Boxerman et al. 52	Schmainda ⁴²
DWModeling (SlicerProstate extension)	В	GВМ, РСа	Modeling of DWI MRI data; multiple models were implemented, as discussed by Langkilde et al. 17	Langkilde ¹⁷ ; Newitt ⁴³
mpReview	_	PCa	Visualization and annotation of prostate mpMRI studies. Fixed hanging protocol supports reformat and synchronous scrolling of the selected mpMRI series, and access to a variety of segmentation tools for userguided region-of-interest segmentation.	
PkModeling	В	GBM, PCa	Modeling of DCE MRI data using Tofts model.53	Huang ⁴⁰
PETLiverUptakeMeasurement	В	HNC	Automated measurement of liver uptake in SUV normalized whole-body FDG-18 PET using approach proposed by Bauer et al. ⁵⁴	Bauer ⁵⁴
PETDICOMExtension	В	HNC	Vendor-neutral computation of PET SUV normalization following the recommendations of the RSNA QIBA.	
PET-IndiC	_	HNC	Calculation of various quantitative measures from the SUV-corrected PET image and a segmented region of interest.	Beichel ⁴¹ ; Beichel ⁵⁵
PETTumorSegmentation	_	HNC	Semiautomated segmentation of metabolically active regions in PET images.	Beichel ⁵⁶
QuantitativeReporting	_	9	Support of segmentation-based measurements with DICOM-based import and export of the results.	Herz ²²
SegmentationWizard	_	GBM	Workflow aiming to simplify the generation of volumetric segmentations of regions of interest with minimum amount of interaction.	
SlicerPathology	_	ŋ	Tools for automatic and semiautomatic segmentation of pathology images.	
SlicerOpenCV	٦	5	Wrapping of the OpenCV library to facilitate its use within 3D Slicer.	
TCIABrowser	_	9	Interactive interface to examine and download the content of public TCIA collections using TCIA API.	

NOTE. Both the quantitative image analysis tools and informatics tools are included. Tool type differentiates between the batch (B) and interactive (I) tools and libraries (L). Tools specific to the individual use cases are marked with the use case abbreviation, whereas those that have generic functionality not specific to a single application are marked with G.

Abbreviations: API, application progamming interface; DCE, dynamic contrast enhanced; DICOM, Digital Imaging and Communications in Medicine; DSC, dynamic susceptibility contrast; DWI, diffusionpositron emission tomography; QI, quantitative imaging; QIBA, Quantitative Imaging Biomarker Alliance; QIICR, Quantitative Imaging Informatics for Cancer Research; RSNA, Radiologic Society of North weighted imaging; FDG-18, [18F]fluorodeoxyglucose; GBM, glioblastoma; HNC, head and neck cancer; mpMRI, multiparametric MRI; MRI, magnetic resonance imaging; PCa, prostate cancer; PET, America; SUV, standardized uptake values; TCIA, The Cancer Imaging Archive.




FIG 1. Download statistics for the 3D Slicer extensions developed with the contribution of the Quantitative Imaging Informatics for Cancer Research (QIICR) project. Note that extensions were released at different times, with some extensions being available longer than others. Some of the extensions are dependent on others (eg, PETDICOMExtension can be installed individually, but is also downloaded every time PET-IndiC or QuantitativeReporting is installed). Statistics reported were collected on December 15, 2019, using the script available publicly at https://github.com/Slicer/SlicerDeveloperToolsForExtensions. These reported download counts do not include downloads of the binary packages directly from GitHub or downloads of Docker images from DockerHub (which were used for disseminating dcmqi packages), or downloads of the source code (due to technical limitations of being able to track such downloads).

of the DICOMweb standard.²⁹ These libraries have been used, for example, as part of the ITCR-Radiologic Society of North America (RSNA) Crowds Cure Cancer collaboration.³⁰

We published a number of datasets accompanied by peerreviewed articles demonstrating best practices for QI analysis results sharing (Table 2; Fig 2). Those supported the data-sharing activities of the collaborating QIN projects and demonstrated the capabilities of the standard and the developed tools. For example, test-retest measurements³¹ obtained from an annotated prostate multiparametric MRI dataset³² were instrumental in defining claims in the Quantitative Imaging Biomarker Alliance Diffusion-Weighted Magnetic Resonance Imaging profile³³ and the DCE (dynamic contrast enhanced) MRI Quantification under development; it was further used to evaluate repeatability of radiomics³⁴ features in the prostate.³⁵

In 2015, we initiated DICOM4QI (DICOM for Quantitative Imaging), a demonstration/connectathon project to evaluate and advance adoption of DICOM for communicating QI analysis results. DICOM4QI was an exhibit at the Quantitative Imaging Reading Room of the RSNA annual meeting. A publicly available Web site provided participation instructions and sample datasets, and summarized participation results.³⁶ Over 15 platforms were included in DICOM4QI. Notably, several platforms that initially did not

TABLE 2. TCIA Collections Contributed by QIICR or Harmonized Using the QIICR-Developed Tools

Collection	QIICR Role	Cancer Type	No.	DICOM Object Types	Reference Describing the Dataset
QIN-HEADNECK	С	HNC	156	PET, CT, RWVM, SEG, SR	Fedorov ²⁰
QIN-PROSTATE-Repeatability	С	PCa	15	MR, SEG, SR	Fedorov ³²
TCGA-GBM	T, H	GBM	262	MR, SEG	Bakas ⁵⁷
TCGA-LGG	T, H	GBM	199	MR, SEG	Bakas ⁵⁷
LIDC-IDRI	T, H	Lung	1,010	CT, SEG, SR	Armato ⁵⁸ ; Fedorov ⁵⁹
NSCLC-Radiomics	T	Lung	422	CT, RTSTRUCT, SEG	Aerts ⁶⁰
NSCLC-Radiomics-Interobserver1	T	Lung	22	CT, RTSTRUCT, SEG	Aerts ⁶⁰
Head-Neck-Radiomics-HN1	Т	HNC	137	CT, PT, RTSTRUCT, SEG	Aerts ⁶⁰

NOTE. Role of the project is indicated as C (contributed the original dataset), T (tools developed by the project were used for data harmonization by a third party), or H (project contributed harmonized representations of the analysis results submitted earlier by an independent entity).

Abbreviations: CT, computed tomography; DICOM, Digital Imaging and Communications in Medicine; GBM, glioblastoma; HNC, head and neck cancer; LGG, low-grade glioma; LIDC-IDRI, Lung Image Database Consortium–Image Database Resource Initiative; MR, magnetic resonance; NSCLC, non–small-cell lung cancer; PCa, prostate cancer; PET, positron emission tomography; PT, positron emission; QIICR, Quantitative Imaging Informatics for Cancer Research; RTSTRUCT, radiotherapy structure set; RWVM, real-world value mapping; SEG, segmentation; SR, structured reporting; TCGA The Cancer Genome Atlas; TCIA, The Cancer Imaging Archive.




FIG 2. Example demonstration of image analysis results interoperability enabled by DICOM. From bottom left corner clockwise, examples of platforms visualizing the same Digital Imaging and Communications in Medicine (DICOM) positron emission tomography (PET) segmentation dataset from the public QIN-HEADNECK collection^{20,61}: 3D Slicer⁸ (free, open-source desktop application), OHIF Viewer³⁸ (free, open-source Web viewer), Brainlab SmartBrush (commercial Food and Drug Administration–approved tumor outlining application). Results were collected as part of the DICOM4QI (DICOM for Quantitative Imaging) demonstration and connectathon organized by Quantitative Imaging Informatics for Cancer Research (QIICR) at the annual Radiologic Society of North America meeting since 2015.^{36,62} The histogram shows the significant increase in the The Cancer Imaging Archive–reported usage of the QIN-HEADNECK collection after the publication of the preprint in Nov 2015 (red arrow) and peer-reviewed paper in May 2016 (green arrow), which introduced imaging-related DICOM data (segmentations, measurements, clinical data) to accompany the imaging dataset.

use suitable DICOM objects for communicating analysis results implemented that capability and used DICOM4QI as a venue for testing interoperability with other platforms. Platforms included the open-source 3D Slicer, 8,37 MITK, 12 ePAD, 23 and OHIF Viewer, 38 and commercial products such as those from Brainlab and TeraRecon (Fig 2).

We organized a half-day tutorial on the use of DICOM for QI research at the Medical Image Computing and Computer Assisted Interventions conferences in 2017 and 2018. The tutorial included presentations from QIICR investigators and collaborators, and a hands-on component demonstrating the capabilities of the DICOM standard and tools. Materials for those tutorials (presentation slides, pointers to the datasets, Jupyter notebooks) were archived and are publicly available.³⁹

Interoperability With Data Repositories

We developed *TCIA Browser*, a 3D Slicer module that uses the TCIA API to explore, download, and visualize the content of public TCIA collections. It streamlines 3D Slicer user interaction with the repository, and makes it easier to evaluate various visualization and analysis tools available within the application.

Coordinating with the TCIA team, we converted some of the existing third-party-contributed analysis results to a standard representation. These datasets can now be accessed from the usual search interface of TCIA or directly via a digital object identifier (as opposed to being linked as compressed archives from collection wiki pages) and are interoperable with a number of popular open-source tools. Furthermore, several collections were harmonized using dcmqi without direct involvement of the QIICR team. Table 2 lists all such collections directly or indirectly supported by QIICR. Currently, dcmqi is the tool recommended by TCIA for harmonizing analysis results submitted to the archive, and 3D Slicer is listed as the only visualization and analysis desktop application integrated with TCIA via its API.

DISCUSSION

The QIICR project aimed to address major perceived needs of the QI community and the QIN, and proposed solutions for specific QIN projects. There is no clear convention on

evaluating contributions such as software tools or datasets that QIICR produced. Based on the evidence we presented, at least some of the tools we developed have been used in projects external to QIICR. The MRI modeling tools and PET segmentation tools we contributed scored favorably based on the results of the challenges organized by QIN. 40-43 Unlike most of the other tools evaluated in those challenges, QIICR-developed tools are readily available as 3D Slicer extensions; source code is public under commercially friendly, nonrestrictive, open-source license; and use of the tools does not require establishing material transfer agreements.

Our investigation of DICOM to represent QI analysis results had limited precedent in the research community. By the end of the project, we were encouraged to see increasing adoption of our ideas in both academic and commercial tools. Our contributions to the DICOM standard passed the rigor of the community review process and are now being reused in new contexts and areas of application. Through one of the QIICR collaborations, we have been investigating the use of the DICOM standard to support digital pathology applications. 44 DICOM Working Group 23 is actively exploring the use of the DICOM TID 1500 Structured Reporting template, as well as DICOM Segmentation objects and Parametric Maps to support a host of artificial intelligence (AI) use cases, which subsume but are not limited to QI applications. We collaborated with the Imaging Biomarker Standardization Initiative (IBSI), which, among other objectives, is establishing a consensus nomenclature and definitions for radiomics features.45 We worked with IBSI to augment the feature definitions with codes and added corresponding coded terms to the DICOM standard (see CP-1705 and CP-1764; Data Supplement). Furthermore, we augmented pyradiomics, 46 an open-source tool for extracting radiomics features, with the experimental features to accept input in DICOM format and store calculated features as DICOM Structured Reports, which use IBSI-defined codes. We consider this collaboration to be one example of successful translation of domain experts' consensus into the standard and implemented in tools for broader deployment and adoption.

The importance of metadata and the use of standards are becoming prominent with the increasing emphasis on quality of radiomics studies⁴⁷ and distributed learning, ^{48,49} where data cannot be shared outside the institution because of privacy, size, or regulatory reasons. Documentation of the acquisition protocol and potential confounding factors related to analysis, provenance of the individual annotations and analysis results, and use of unique identifiers to prevent cross-contamination of testing and training data are important for ensuring quality of the analysis. The use of standards is critical to scale production deployment of federated learning systems. They provide

improved consistency of data representation and harmonization of imaging with the analysis results.

Overall, we emphasize the importance of a collaborative approach for addressing challenges in imaging informatics at the national and international levels. Topics such as data standardization, interoperability, and sustainable opensource software development cannot be addressed by an individual group. Open coordinated effort involving a broad range of stakeholders, including clinical researchers, engineers, and industry collaborators, is essential. Joint development of the standard, reference implementations and datasets, and reusable components is ultimately beneficial to both academic and commercial efforts.

Most recently, the BWH team led a consortium of collaborators, including many of the QIICR investigators, who were awarded the contract to establish the NCI Imaging Data Commons (IDC),⁵⁰ a new component of the NCI Cancer Research Data Commons (CRDC). Our IDC approach is motivated by our QIN and QIICR experience, and will use DICOM to harmonize analysis results that IDC will share.

Challenges remain in establishing and maintaining the infrastructure of analysis tools and standardized imaging datasets. Sustainability of open-source tools developed in academia is often problematic, because continuous maintenance effort is required for changes in dependencies and operating systems, user support, and implementation of new features. Our use of 3D Slicer as a delivery platform partially mitigates this issue by leveraging infrastructure, community effort, and contributions from numerous other supporting projects. Development and dissemination of most of the QIICR tools would be extremely challenging without leveraging the 3D Slicer infrastructure. Sustained support for extensible open-source platforms that provide generic reusable components and a marketplace-like infrastructure for new contributions is critical to reduce redundancy, build community, and support tool dissemination and outreach.

Despite significant progress in community adoption of DICOM for QI results, there remains significant effort required to integrate it into academic and commercial tools and workflows. Although improving, support in commonly used tools is limited. Adoption of FAIR (Findable, Accessible, Interoperable, Reusable) data stewardship principles⁵¹ requires time and sustained community effort to improve the standards, tools, documentation, and outreach. We envision the CRDC IDC as a mechanism and laboratory to refine and demonstrate best practices for curating and interoperating with standardized image and image-derived data, and improve access to QI and AI analysis tools.

AFFILIATIONS

¹Department of Radiology, Brigham and Women's Hospital, Harvard Medical School, Boston, MA

²University of Iowa, Iowa City, IA

³Massachusetts General Hospital, Boston, MA

⁴PixelMed Publishing, Bangor, PA

⁵OpenConnections, Oldenburg, Germany

⁶Freelancer, Oldenburg, Germany

⁷Kitware, Clifton Park, NY

⁸Queen's University, Kingston, ON, Canada

9Isomics, Cambridge, MA

¹⁰German Cancer Research Center, Heidelberg, Germany

¹¹Department of Pathology, Massachusetts General Hospital, Harvard Medical School, Boston, MA

¹²Stony Brook University, Stony Brook, NY

¹³University of Arkansas for Medical Sciences, Little Rock, AR

CORRESPONDING AUTHOR

Andrey Fedorov, PhD, 1249 Boylston St #344, Boston, MA 02215; e-mail: andrey.fedorov@gmail.com.

SUPPORT

Supported in part by award U24 CA180918 from the National Cancer Institute (NCI; Quantitative Imaging Informatics for Cancer Research [QIICR]) and by contract 19X037Q from Leidos Biomedical Research (LBR) under Task Order HHSN26100071 from NCI (Imaging Data Commons). Collaborating Quantitative Imaging Network projects were supported by Grants No. U01 CACA154601, U01 CA140206, and U01 CA151261. SlicerPathology and SlicerOpenCV extensions were supported by a QIICR supplement and award U24 CA180924 (J.S.). Harmonization of the Lung Image Database Consortium—Image Database Resource Initiative collection was supported through

HHSN261200800001E, under which the University of Arkansas for Medical Sciences is funded by LBR subcontract 16X011 (F.P.).

AUTHOR CONTRIBUTIONS

Conception and design: Andrey Fedorov, Reinhard Beichel, Jayashree Kalpathy-Cramer, David Clunie, Michael Onken, Jörg Riesmeier, Christian Herz, Jean-Christophe Fillion-Robin, Andras Lasso, Csaba Pinter, Steve Pieper, Marco Nolden, Klaus Maier-Hein, Joel Saltz, Fiona Fennessy, Ron Kikinis

Collection and assembly of data: Andrey Fedorov, Reinhard Beichel, David Clunie, Christian Bauer, Andrew Beers, Jean-Christophe Fillion-Robin, Steve Pieper

Data analysis and interpretation: Andrey Fedorov, Reinhard Beichel, David Clunie, Andrew Beers, Steve Pieper, Markus D. Herrmann, Fred Prior, John Buatti **Manuscript writing:** All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the

subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs. org/cci/author-center.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians (Open Payments).

Andrey Fedorov

Research Funding: Siemens Healthcare (Inst)

Jayashree Kalpathy-Cramer

Research Funding: GE Healthcare (Inst)

David Clunie

Leadership: PixelMed

Consulting or Advisory Role: Algotec, BK Medical, Imago, Lunit, MDDX/

Bioclinica, Medigate

Expert Testimony: Confidential arbitration

Other Relationship: NEMA (501(c)6 trade association) MITA Division

Jean-Christophe Fillion-Robin Research Funding: Kitware (Inst)

Csaba Pinter

Employment: Pixel Medical **Leadership:** Pixel Medical

Stock and Other Ownership Interests: Pixel Medical **Honoraria:** Medical Augmented Intelligence

Steve Pieper
Employment: Isomics
Leadership: Isomics

Stock and Other Ownership Interests: Isomics
Consulting or Advisory Role: Boston Scientific (Inst)
Research Funding: Siemens Healthineers (Inst)

Patents, Royalties, Other Intellectual Property: I do joint research with Mass

General Brigham and jointly file patents (Inst)

Travel, Accommodations, Expenses: Siemens Healthineers

Klaus Maier-Hein

Research Funding: Siemens Healthineers

Joel Salta

Patents, Royalties, Other Intellectual Property: Patent pending: computational staining to generate and quantify tumor infiltrating lymphocytes using digitized diagnostic tissue images (Inst)

Fiona Fennessy

Consulting or Advisory Role: Virtualscopics

John Buatti

Patents, Royalties, Other Intellectual Property: UptoDate

No other potential conflicts of interest were reported.

ACKNOWLEDGMENT

We dedicate this work to the memory of the late Larry Clarke, PhD, who served as the Chief of the Image Technology Development Branch at the National Cancer Institute at the time the QIICR project was funded. QIICR would not have happened without Larry's energy, encouragement, vision, and support.

REFERENCES

- 1. Clarke LP, Croft BS, Nordstrom R, et al: Quantitative imaging for evaluation of response to cancer therapy. Transl Oncol 2:195-197, 2009
- 2. Shankar LK: The clinical evaluation of novel imaging methods for cancer management. Nat Rev Clin Oncol 9:738-744, 2012
- 3. O'Connor JPB, Aboagye EO, Adams JE, et al: Imaging biomarker roadmap for cancer studies. Nat Rev Clin Oncol 14:169-186, 2016

- 4. Clarke LP, Nordstrom RJ, Zhang H, et al: The Quantitative Imaging Network: NCl's historical perspective and planned goals. Transl Oncol 7:1-4, 2014
- 5. Nordstrom RJ: The Quantitative Imaging Network in precision medicine. Tomography 2:239-241, 2016
- 6. Farahani K, Tata D, Nordstrom RJ: QIN benchmarks for clinical translation of quantitative imaging tools. Tomography 5:1-6, 2019
- 7. Branstetter BF IV: Basics of imaging informatics. Part 1. Radiology 243:656-667, 2007
- Fedorov A, Beichel R, Kalpathy-Cramer J, et al: 3D Slicer as an image computing platform for the Quantitative Imaging Network. Magn Reson Imaging 30:1323-1341, 2012
- Bidgood WD Jr, Horii SC, Prior FW, et al: Understanding and using DICOM, the data interchange standard for biomedical imaging. J Am Med Inform Assoc 4:199-212, 1997
- Clark K, Vendt B, Smith K, et al: The Cancer Imaging Archive (TCIA): Maintaining and operating a public information repository. J Digit Imaging 26:1045-1057, 2013
- 11. Bennett W, Smith K, Jarosz Q, et al: Reengineering workflow for curation of DICOM datasets. J Digit Imaging 31:783-791, 2018
- 12. Nolden M, Zelzer S, Seitel A, et al: The Medical Imaging Interaction Toolkit: Challenges and advances. Int J CARS 8:607-620, 2013
- 13. Clarkson MJ, Zombori G, Thompson S, et al: The NifTK software platform for image-guided interventions: Platform overview and NiftyLink messaging. Int J CARS 10:301-316, 2015
- 14. QIICR: Tools catalog. http://qiicr.org/tools.html
- Mehrtash A, Gupta SN, Shanbhag D, et al: Bolus arrival time and its effect on tissue characterization with dynamic contrast-enhanced magnetic resonance imaging. J Med Imaging (Bellingham) 3:014503, 2016
- Glazer DI, Hassanzadeh E, Fedorov A, et al: Diffusion-weighted endorectal MR imaging at 3T for prostate cancer: Correlation with tumor cell density and percentage Gleason pattern on whole mount pathology. Abdom Radiol (NY) 42:918-925, 2017
- 17. Langkilde F, Kobus T, Fedorov A, et al: Evaluation of fitting models for prostate tissue characterization using extended-range b-factor diffusion-weighted imaging. Magn Reson Med 79:2346-2358, 2018
- 18. Peled S, Vangel M, Kikinis R, et al: Selection of fitting model and arterial input function for repeatability in dynamic contrast-enhanced prostate MRI. Acad Radiol 26:e241-e251, 2019
- 19. Jafari-Khouzani K, Paynabar K, Hajighasemi F, et al: Effect of region of interest size on the repeatability of quantitative brain imaging biomarkers. IEEE Trans Biomed Eng 66:864-872, 2019
- 20. Fedorov A, Clunie D, Ulrich E, et al: DICOM for quantitative imaging biomarker development: A standards based approach to sharing clinical data and structured PET/CT analysis results in head and neck cancer research. PeerJ 4:e2057, 2016
- 21. Eichelberg M, Riesmeier J, Wilkens T, et al: Ten years of medical imaging standardization and prototypical implementation: The DICOM standard and the OFFIS DICOM toolkit (DCMTK). Proc SPIE 5:57-68, 2004
- 22. Herz C, Fillion-Robin J-C, Onken M, et al: dcmqi: An open source library for standardized communication of quantitative image analysis results using DICOM. Cancer Res 77:e87-e90, 2017
- 23. Rubin DL, Ugur Akdogan M, Altindag C, et al: ePAD: An image annotation and analysis platform for quantitative imaging. Tomography 5:170-183, 2019
- 24. Stein T, Metzger J, Scherer J, et al: Efficient Web-based review for automatic segmentation of volumetric DICOM Images. Bildverarbeitung für die Medizin 2019. Heidelberg, Germany, Springer Fachmedien Wiesbaden, 2019
- 25. Sedghi A, Hamidi S, Mehrtash A, et al: Tesseract-medical imaging: Open-source browser-based platform for artificial intelligence deployment in medical imaging. https://doi.org/10.1117/12.2513004
- 26. Github: commontk/dcmjs. https://github.com/commontk/dcmjs
- 27. Emscripten: Emscription. http://emscripten.org
- 28. dcmjs-org: DICOM implementation in JavaScript. https://github.com/dcmjs-org
- 29. National Electrical Manufacturers Association (NEMA): DICOM. https://www.dicomstandard.org/dicomweb/
- 30. NCI ITCR Collaborators: Crowds cure cancer. http://crowds-cure.org
- 31. Fedorov A, Vangel MG, Tempany CM, et al: Multiparametric magnetic resonance imaging of the prostate: Repeatability of volume and apparent diffusion coefficient quantification. Invest Radiol 52:538-546, 2017
- 32. Fedorov A, Schwier M, Clunie D, et al: An annotated test-retest collection of prostate multiparametric MRI. Sci Data 5:180281, 2018
- 33. Quantitative Imaging Biomarker Alliance: QIBA profile: Diffusion-weighted magnetic resonance imaging (DWI). http://qibawiki.rsna.org/index.php/Perfusion,_ Diffusion_and_Flow-MRI_Biomarker_Ctte#DWI-MRI_Profile_Development
- 34. Gillies RJ, Kinahan PE, Hricak H: Radiomics: Images are more than pictures, they are data. Radiology 278:563-577, 2015
- 35. Schwier M, van Griethuysen J, Vangel MG, et al: Repeatability of multiparametric prostate MRI radiomics features. Sci Rep 9:9441, 2019
- 36. QIICR: Introduction. https://dicom4qi.readthedocs.io
- 37. Norton I, Essayed W, Zhang F, et al: SlicerDMRI: Open source diffusion MRI software for brain cancer research. Cancer Res 77:e101-e103, 2017 [Erratum: Cancer Res 78:2445, 2018]
- 38. Urban T, Ziegler E, Lewis R, et al: Lesiontracker: Extensible open-source zero-footprint web viewer for cancer imaging research and clinical trials. Cancer Res 77:e119-e122. 2017
- 39. QIICR: DICOM4MICCAI tutorial, http://giicr.org/dicom4miccai
- 40. Huang W, Li X, Chen Y, et al: Variations of dynamic contrast-enhanced magnetic resonance imaging in evaluation of breast cancer therapy response: A multicenter data analysis challenge. Transl Oncol 7:153-166, 2014
- 41. Beichel RR, Smith BJ, Bauer C, et al: Multi-site quality and variability analysis of 3D FDG PET segmentations based on phantom and clinical image data. Med Phys 44:479-496, 2017
- 42. Schmainda KM, Prah MA, Rand SD, et al: Multisite concordance of DSC-MRI analysis for brain tumors: Results of a National Cancer Institute Quantitative Imaging Network Collaborative Project. AJNR Am J Neuroradiol 39:1008-1016, 2018
- 43. Newitt DC, Malyarenko D, Chenevert TL, et al: Multisite concordance of apparent diffusion coefficient measurements across the NCI Quantitative Imaging Network. J Med Imaging (Bellingham) 5:011003, 2018
- 44. Herrmann MD, Clunie DA, Fedorov A, et al: Implementing the DICOM standard for digital pathology. J Pathol Inform 9:37, 2018
- 45. Zwanenburg A, Vallières M, Abdalah MA, et al: The Image Biomarker Standardization Initiative: Standardized quantitative radiomics for high throughput image-based phenotyping. Radiology 10.1148/radiol.2020191145 [epub ahead of print on March 10, 2020]
- 46. van Griethuysen JJM, Fedorov A, Parmar C, et al: Computational radiomics system to decode the radiographic phenotype. Cancer Res 77:e104-e107, 2017

- 47. Lambin P, Leijenaar RTH, Deist TM, et al: Radiomics: The bridge between medical imaging and personalized medicine. Nat Rev Clin Oncol 14:749-762, 2017
- Deist TM, Jochems A, van Soest J, et al: Infrastructure and distributed learning methodology for privacy-preserving multi-centric rapid learning health care: EuroCAT. Clin Transl Radiat Oncol 4:24-31, 2017
- 49. Chang K, Balachandar N, Lam C, et al: Distributed deep learning networks among institutions for medical imaging. J Am Med Inform Assoc 25:945-954, 2018
- 50. Imaging Data Commons Consortium: Imaging Data Commons. https://imagingdatacommons.github.io/
- 51. Wilkinson MD, Dumontier M, Aalbersberg IJJ, et al: The FAIR Guiding Principles for scientific data management and stewardship. Sci Data 3:160018, 2016 [Erratum Sci Data 6:6, 2019]
- 52. Boxerman JL, Schmainda KM, Weisskoff RM: Relative cerebral blood volume maps corrected for contrast agent extravasation significantly correlate with glioma tumor grade, whereas uncorrected maps do not. AJNR Am J Neuroradiol 27:859-867, 2006
- 53. Tofts PS, Kermode AG: Measurement of the blood-brain barrier permeability and leakage space using dynamic MR imaging. 1. Fundamental concepts. Magn Reson Med 17:357-367, 1991
- 54. Bauer C, Sun S, Sun W, et al: Automated measurement of uptake in cerebellum, liver, and aortic arch in full-body FDG PET/CT scans. Med Phys 39:3112-3123, 2012
- 55. Beichel RR, Ulrich EJ, Smith BJ, et al: FDG PET based prediction of response in head and neck cancer treatment: Assessment of new quantitative imaging features. PLoS One 14:e0215465, 2019
- 56. Beichel RR, Van Tol M, Ulrich EJ, et al: Semiautomated segmentation of head and neck cancers in ¹⁸F-FDG PET scans: A just-enough-interaction approach. Med Phys 43:2948-2964, 2016
- 57. Bakas S, Akbari H, Sotiras A, et al: Advancing The Cancer Genome Atlas glioma MRI collections with expert segmentation labels and radiomic features. Sci Data 4:170117, 2017
- 58. Armato SG III, McLennan G, Bidaut L, et al: The Lung Image Database Consortium (LIDC) and Image Database Resource Initiative (IDRI): A completed reference database of lung nodules on CT scans. Med Phys 38:915-931, 2011
- 59. Fedorov A, Hancock M, Clunie D, et al: Standardized representation of the LIDC annotations using DICOM. https://peerj.com/preprints/27378/
- 60. Aerts HJWL, Velazquez ER, Leijenaar RTH, et al: Decoding tumour phenotype by noninvasive imaging using a quantitative radiomics approach. Nat Commun 5: 4006, 2014 [Erratum: Nat Commun 5:4644, 2014]
- 61. Beichel RR, Ulrich EJ, Bauer C, et al: QIN-HEADNECK The Cancer Imaging Archive (TCIA). https://wiki.cancerimagingarchive.net/display/Public/QIN-HEADNECK
- 62. Fedorov A, Rubin D, Clunie D, et al: Standardized communications of quantitative image analysis results using DICOM: Establishing interoperability through outreach and community engagement. https://zerista.s3.amazonaws.com/item_files/072e/attachments/290677/original/fedorov-amiasummits2017-dicom4qi.pdf