



## Draft Genome Sequence of *Lactobacillus plantarum* CMPG5300, a Human Vaginal Isolate

## Shweta Malik,<sup>a,b</sup> Roland J. Siezen,<sup>c,d</sup> Bernadet Renckens,<sup>c</sup> Mario Vaneechoutte,<sup>e</sup> Jos Vanderleyden,<sup>a</sup> <sup>[b]</sup>Sarah Lebeer<sup>a,b</sup>

KU Leuven, Centre of Microbial and Plant Genetics, Kasteelpark Arenberg, Leuven, Belgium<sup>a</sup>; University of Antwerp, Department of Bioscience Engineering, Groenenborgerlaan, Antwerp, Belgium<sup>b</sup>; Radboud University Nijmegen Medical Centre, Centre for Molecular and Biomolecular Informatics (CMBI), Nijmegen, The Netherlands<sup>c</sup>; Microbial Bioinformatics, Ede, The Netherlands<sup>d</sup>; Ghent University, Laboratory of Bacteriology Research, Ghent, Belgium<sup>a</sup>

The draft genome of a highly auto-aggregating *Lactobacillus plantarum* strain isolated from a human vagina is reported. The peculiar phenotype also provides an adhesive and co-aggregative potential with various pathogens, which could be of significance in the vaginal niche. Detailed genome analysis could aid in identifying the adhesins of the strain.

Received 25 September 2014 Accepted 6 October 2014 Published 13 November 2014

Citation Malik S, Siezen RJ, Renckens B, Vaneechoutte M, Vanderleyden J, Lebeer S. 2014. Draft genome sequence of *Lactobacillus plantarum* CMPG5300, a human vaginal isolate. Genome Announc. 2(6):e01149-14. doi:10.1128/genomeA.01149-14.

Copyright © 2014 Malik et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 Unported license.

Address correspondence to Sarah Lebeer, sarah.lebeer@uantwerpen.be.

actobacillus plantarum is a highly versatile species found in a variety of environmental niches such as in the gastrointestinal tract, and food products including dairy, meat, and vegetable fermentations (1, 2). There has been considerable interest in the probiotic potential of this species (1, 3, 4), yet it has been restricted to the intestinal isolates. Being a less commonly found species in the vaginal niche (5, 6), there are no reports on the genomes of vaginal L. plantarum isolates yet. Here, we report the draft genome of a vaginal Lactobacillus plantarum strain CMPG5300 with a remarkably high auto-aggregative capacity and adhesive ability to vaginal epithelial cell line VK2/E6E7 (7). The strong adhesive and auto-aggregative property of the strain is related to proteinaceous structures, as it can be abolished by proteinase treatment. A gene deletion mutant of the sortase-encoding gene of the strain lost the capacity to auto-aggregate and bind to mannose-rich conjugates of yeast cells, suggesting a crucial role for sortase-dependent proteins for these characteristics (7). Further insight into the genome of the strain may help the identification of putative adhesins imparting the exceptional features to the strain. This could particularly be of interest for targeted approaches that aim at evading pathogens at the vaginal front.

The genomic DNA of the strain CMPG5300 was isolated as described before (7) and whole-genome sequencing was performed using the 454 GS FLX+ sequencing platform (Genomics Core, KU Leuven). The reads were assembled into contigs using Roche GS De Novo Assembler software (version 2.5p1). The read coverage within the contigs was  $50-100\times$ . The minimum overlap length was set to 40 nucleotides, with a minimum overlap identity of 90%. Open-reading frames were extracted using the getorf program from the EMBOSS suite v6.3.1 (8), automatically annotated using the RAST server (9), and manually curated by comparison against the published genome of *L. plantarum* WCFS1 (10) using ACT (11).

The genome assembled into 48 contigs larger than 750 nt (containing a total 3,503,628 nt) of which at least six contigs represent plasmids (size 2 to 40 kb; total ~143 kb), with a total G+C content of 44%. The genome contains 3,251 protein-encoding genes (at least 140 are plasmid encoded). The scaffolded chromosome (19 contigs) is highly syntenous to published L. plantarum chromosomes (10, 12-16) (accession no. AL935263.2, CP001617.1, CP002222.1, CP004082.1, AGRI00000000.1, CP006033.1, PRJNA203333). Large variations are found between all L. plantarum chromosomes, including CMPG5300, in gene clusters encoding prophages, exopolysaccharide biosynthesis, plantaricin biosynthesis, and sugar metabolism (17). The putative plasmids of strain CMPG5300 encode a few sortase-dependent proteins, which may be relevant for the auto-aggregating and adhesive properties of this strain (7). These include a cell-envelope associated serine proteinase (Cmpg5300\_3052) of 1,482 amino acids, which is 99% identical to plasmid-encoded proteinases of L. plantarum strains 16 and P8. Furthermore, a mucus-binding protein of 1,203 amino acids Cmpg5300.05\_29 is encoded that is 98% identical to a cell-surface protein of Lactobacillus antri, and 62% identical to the mannose-specific adhesin (Msa) of L. plantarum WCFS1.

**Nucleotide sequence accession numbers.** This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. AXZV000000000. The version described in this paper is version AXZV01000000.

## ACKNOWLEDGMENT

The genome sequencing was supported by the KU Leuven funding BOF program financing (spokesman, Jan Balzarini). S.M. was supported by Erasmus Mundus External Cooperation Window Lot 13 for her Ph.D. research. S.L. also acknowledges support of FWO Vlaanderen for her postdoc grant and a KaN grant.

## REFERENCES

 Kleerebezem M, Boekhorst J, van Kranenburg R, Molenaar D, Kuipers OP, Leer R, Tarchini R, Peters SA, Sandbrink HM, Fiers MW, Stiekema W, Lankhorst RM, Bron PA, Hoffer SM, Groot MN, Kerkhoven R, de Vries M, Ursing B, de Vos WM, Siezen RJ. 2003. Complete genome sequence of *Lactobacillus plantarum* WCFS1. Proc. Natl. Acad. Sci. U. S. A. 100:1990–1995.

- Siezen RJ, Tzeneva VA, Castioni A, Wels M, Phan HT, Rademaker JL, Starrenburg MJ, Kleerebezem M, Molenaar D, van Hylckama Vlieg JE. 2010. Phenotypic and genomic diversity of *Lactobacillus plantarum* strains isolated from various environmental niches. Environ. Microbiol. 12: 758–773. http://dx.doi.org/10.1111/j.1462-2920.2009.02119.x.
- 3. Connelly P. 2008. *Lactobacillus plantarum*—A literature review of therapeutic benefits. J. Aust. Tradit. Med. Soc. 14:79–82.
- Klarin B, Molin G, Jeppsson B, Larsson A. 2008. Use of the probiotic Lactobacillus plantarum 299 to reduce pathogenic bacteria in the oropharynx of intubated patients: a randomised controlled open pilot study. Crit. Care 12:R136. http://dx.doi.org/10.1186/cc7109.
- Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, Karlebach S, Gorle R, Russell J, Tacket CO, Brotman RM, Davis CC, Ault K, Peralta L, Forney LJ. 2011. Vaginal microbiome of reproductiveage women. Proc. Natl. Acad. Sci. U. S. A. 108(Suppl 1):4680–4687. http://dx.doi.org/10.1073/pnas.1002611107.
- Gustafsson RJ, Ahrné S, Jeppsson B, Benoni C, Olsson C, Stjernquist M, Ohlsson B. 2011. The *Lactobacillus* flora in vagina and rectum of fertile and postmenopausal healthy Swedish women. BMC Womens Health 11: 17. http://dx.doi.org/10.1186/1472-6874-11-17.
- Malik S, Petrova MI, Claes IJ, Verhoeven TL, Busschaert P, Vaneechoutte M, Lievens B, Lambrichts I, Siezen RJ, Balzarini J, Vanderleyden J, Lebeer S. 2013. The highly autoaggregative and adhesive phenotype of the vaginal *Lactobacillus plantarum* strain CMPG5300 is sortase dependent. Appl. Environ. Microbiol. 79:4576-4585. http:// dx.doi.org/10.1128/AEM.00926-13.
- Rice P, Longden I, Bleasby A. 2000. EMBOSS: the European molecular biology open software suite. Trends Genet. 16:276–277. http://dx.doi.org/ 10.1016/S0168-9525(00)02024-2.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using

Subsystems Technology. BMC Genomics 9:75. http://dx.doi.org/10.1186/ 1471-2164-9-75.

- Siezen RJ, Francke C, Renckens B, Boekhorst J, Wels M, Kleerebezem M, van Hijum SA. 2012. Complete resequencing and reannotation of the *Lactobacillus plantarum* WCFS1 genome. J. Bacteriol. 194:195–196. http:// dx.doi.org/10.1128/JB.06275-11.
- Carver TJ, Rutherford KM, Berriman M, Rajandream MA, Barrell BG, Parkhill J. 2005. ACT: the Artemis comparison tool. Bioinformatics 21: 3422–3423. http://dx.doi.org/10.1093/bioinformatics/bti553.
- Zhang ZY, Liu C, Zhu YZ, Zhong Y, Zhu YQ, Zheng HJ, Zhao GP, Wang SY, Guo XK. 2009. Complete genome sequence of *Lactobacillus plantarum* JDM1. J. Bacteriol. 191:5020–5021. http://dx.doi.org/10.1128/ JB.00587-09.
- Wang Y, Chen C, Ai L, Zhou F, Zhou Z, Wang L, Zhang H, Chen W, Guo B. 2011. Complete genome sequence of the probiotic *Lactobacillus plantarum* ST-III. J. Bacteriol. 193:313–314. http://dx.doi.org/10.1128/ JB.01159-10.
- Li X, Gu Q, Lou X, Zhang X, Song D, Shen L, Zhao Y. 2013. Complete genome sequence of the probiotic *Lactobacillus plantarum* strain ZJ316. Genome Announc. 1(2):e0009413. http://dx.doi.org/10.1128/ genomeA.00094-13.
- Axelsson L, Rud I, Naterstad K, Blom H, Renckens B, Boekhorst J, Kleerebezem M, van Hijum S, Siezen RJ. 2012. Genome sequence of the naturally plasmid-free *Lactobacillus plantarum* strain NC8 (CCUG 61730). J. Bacteriol. 194:2391–2392. http://dx.doi.org/10.1128/JB.00141 -12.
- Crowley S, Bottacini F, Mahony J, van Sinderen D. 2013. Complete genome sequence of *Lactobacillus plantarum* strain 16, a broad-spectrum antifungal-producing lactic acid bacterium. Genome Announc. 1(4): e00533-13. http://dx.doi.org/10.1128/genomeA.00533-13.
- Siezen RJ, van Hylckama Vlieg JE. 2011. Genomic diversity and versatility of *Lactobacillus plantarum*, a natural metabolic engineer. Microb. Cell Fact. 10:S3. http://dx.doi.org/10.1186/1475-2859-10-S1-S3.