

# *Psallus thomashenryi* sp. n. and *Psallus lucanicus* from Turkey (Hemiptera, Heteroptera, Miridae)

Attilio Carapezza<sup>1</sup>, Petr Kment<sup>2</sup>

**1** University of Palermo; corresponding address: via Sandro Botticelli 15, 90144 Palermo, Italy **2** Department of Entomology, National Museum, Cirkusová 1740, CZ-193 00 Prague 9 – Horní Počernice, Czech Republic

Corresponding author: Attilio Carapezza ([attilio.carapezza@unipa.it](mailto:attilio.carapezza@unipa.it)), Petr Kment ([sigara@post.cz](mailto:sigara@post.cz))

---

Academic editor: A. Wheeler | Received 9 October 2017 | Accepted 26 January 2018 | Published 15 November 2018

<http://zoobank.org/BFDD0CCA-0195-4256-BC60-B4324F00FF06>

---

**Citation:** Carapezza A, Kment P (2018) *Psallus thomashenryi* sp. n. and *Psallus lucanicus* from Turkey (Hemiptera, Heteroptera, Miridae). In: Wheeler Jr AG (Ed.) A Festschrift Recognizing Thomas J. Henry for a Lifetime of Contributions to Heteropteran Systematics. ZooKeys 796: 253–265. <https://doi.org/10.3897/zookeys.796.21536>

---

## Abstract

*Psallus* (*Psallus*) *thomashenryi* sp. n. (Hemiptera: Heteroptera: Miridae: Phylinae: Phylini: Phylina) is described from southern Anatolia, Turkey. Illustrations of the dorsal habitus and male genitalia are provided. Its habitus is similar to other uniformly orange species of the subgenus *Psallus* Fieber, 1858, particularly *P. asthenicus* Seidenstücker, 1966 from which it can be easily distinguished by the combination of extremely small size (2.3 mm in both sexes) and different morphology of the vesica. *Psallus* (*Psallus*) *lucanicus* Wagner, 1968 is recorded for the first time from Turkey. *Psallus* (*Psallus*) *aurora* (Mulsant & Rey, 1852) is removed from the list of Turkish fauna based on a reevaluation of the voucher specimen. An updated checklist of the species of *Psallus* known to occur in Turkey is provided. The relevance of Anatolia and the Syro-anatolian-transcaucasian region in the Palearctic distribution of *Psallus* is discussed. The westernmost record of another mirid, *Plagiognathus marivanensis* Linnavuori, 2010, is provided.

## Keywords

Hemiptera, Heteroptera, Miridae, new record, new species, Palearctic region, Phylinae, Phylini, *Psallus*, Turkey

## Introduction

The predominantly Palearctic genus *Psallus* Fieber, 1858, including 160 valid species-group taxa, is one of the most speciose in the subfamily Phylinae (Schuh 1995, 2013; Kerzhner and Josifov 1999; Yasunaga 2010; Matocq 2011; Duwal et al. 2012; Vinokurov and Luo 2012; Aukema et al. 2013; Li and Liu 2013; Schuh et al. 2014; Simon and Strauss 2014; Konstantinov 2016; Pagola-Carte 2017). *Psallus* is currently subdivided into eight subgenera based on similarities in habitus and male genitalia: *Apocremnus* Fieber, 1858, *Calopsallus* Yasunaga & Vinokurov, 2000, *Hyllopsallus* Wagner, 1952, *Mesopsallus* Wagner, 1970, *Psallus* Fieber, 1858, *Phylidea* Reuter, 1899, *Pityopsallus* Wagner, 1952, and *Supsallus* Linnavuori, 1993 (Kerzhner and Josifov 1999, Yasunaga and Vinokurov 2000, Yasunaga 2010). Some authors have regarded this division into subgenera and the definition of the genus itself far from satisfactory and in need of revision; the genus is generally considered polyphyletic and several proposals have been made to combine some subgenera, elevate some subgenera to genus level, and/or transfer some species to other genera (Yasunaga and Vinokurov 2000, Wyniger 2004, Mróz 2012, Aukema et al. 2013, Schuh and Menard 2013, Pluot-Sigwalt and Matocq 2017). Probably a better understanding of the systematics of *Psallus* will be reached only by dealing with it on a world basis and extending the use of female genitalia, whose relevant value in aiding the recognition of related and unrelated species was recently demonstrated by Pluot-Sigwalt and Matocq (2017).

Prior to this study, 34 species of *Psallus* were known to occur in Turkey. Our study documents two additional species; the first of them, *Psallus (Psallus) thomashenryi* sp. n., is described from Southern Anatolia, and the second, *Psallus (Psallus) lucanicus* Wagner, 1968, is recorded for the first time from Turkey. The West-Mediterranean *Psallus (Psallus) aurora* (Mulsant & Rey, 1852) is removed from the list of Turkish fauna based on a reevaluation of the voucher specimen.

## Material and methods

Images of the adults were taken using a Canon D40 camera equipped with a MP-E65 macro lens mounted on a photographic stand; stacked images were combined using Zerene Stacker. Drawings of 10 % KOH-macerated genitalia were made using a Leitz Laborlux S microscope equipped with camera lucida. Measurements were made using an eyepiece micrometer mounted on a Wild M5S binocular microscope. All measurements are in millimeters. Morphological terminology follows Schuh and Slater (1995); terminology of male genitalia follows Konstantinov (2003).

In the transcription of locality labels of types a slash (/) is used to indicate data in different rows of a single label; a double slash (//) is used to separate different labels; data on the labels are given verbatim.

All specimens mentioned in the text are deposited in the National Museum, Prague, Czech Republic (NMPC).

## Taxonomy

### *Psallus thomashenryi* sp. n.

<http://zoobank.org/EDE2EE97-C112-4896-9AC1-A805CF5A9AB4>

Figs 1–7

**Type locality.** Turkey, southern Anatolia, Mersin Province, Göksu Nehri river canyon, Evkaçiftliği, 36°27'23.6"N, 33°38'12.3"E.

**Type material. Holotype:** ♂, glued on a pointed cardboard with genitalia glued on the same cardboard with labels as follows: 36°27'23.6"N, 33°38'12.3"E / AS. TURKEY, İÇEL prov. / Evkaçiftliği, Göksu Nehri canyon / valley of drying brook, sweep / 5.v.2007, lgt. P. Kment [white printed label] // HOLOTYPUS / *PSALLUS* (*PSALLUS*) / *THOMASHENRYI* / sp. n. / det. Carapezza & Kment 2017 [red printed label]' (NMPC).

**Paratype:** ♀, glued on a pointed cardboard with labels as follows: 36°27'23.6"N, 33°38'12.3"E / AS. TURKEY, İÇEL prov. / Evkaçiftliği, Göksu Nehri canyon / valley of drying brook, sweep / 5.v.2007, lgt. P. Kment [white printed label] // PARATYPUS / *PSALLUS* (*PSALLUS*) / *THOMASHENRYI* / sp. n. / det. Carapezza & Kment 2017 [red printed label]' (NMPC).

**Description. Male. Coloration** (Fig. 1). Dorsal coloration almost uniformly orange. Head orange, vertex basally with four small reddish dots arranged in line, frons with five whitish lateral arcs; apex of clypeus whitish. Antennae pale yellowish, scape with faint basal annulation and with two preapical dark dots; labium pale yellowish, apical half of last segment darkened. Pronotum orange with traces of reddish dotting in anterior half; scutellum and hemelytra orange, cuneus basally and apically whitish; membrane pale, hyaline, veins concolorous. Thoracic sterna orange with reddish tinge, legs pale yellowish, femora with irregular orange to reddish-brown dots, more numerous on hind femora; tibial spines black, arising from small dark spots; tarsi uniformly pale.


**Structure.** Body elongate-ovoid (Fig. 1), about 2.8 times longer than basal width of pronotum. Head moderately projecting, in dorsal view 2.1 times wider than long, in frontal view 1.5 times wider than high, in lateral view 1.5 times longer than high; ocular index (ratio vertex/eye in dorsal view) 1.6. Antennae with segment II 0.8 times as long as basal width of pronotum. Labium slightly surpassing metacoxae. Hind femora elongate, 3.6 times longer than maximum width; tibial spines long, about twice longer than tibial diameter. Genital segment ventrally unkeeled; phallosome (Fig. 2) robust, with a preapical lateral ridge, apex rounded; left paramere (Fig. 4) broad, apical process straight and thin, sensory lobe short, apically rounded; right paramere (Fig. 3) elongate, apical process straight; vesica (Figs 5–7) short, C-shaped, provided with robust postbasal lateral spicule extending apically to middle of vesica, terminating in elongate, apically recurved blade, armed with rows of denticles along inner side, and three fingerlike, apically bent blades, almost equal in size, originating near subapical secondary gonopore.



**Figure 1.** Habitus of *Psallus thomashenryi* sp. n., holotype, male (2.29 mm).

*Pubescence.* Dorsum with reclining pale and semierect blackish setae; the latter few, mostly on head and lateral margins of pronotum.

*Female.* Coloration similar to males but paler. Structure and pubescence as in males, but body more ovoid, 2.8 times longer than basal width of pronotum; ocular



**Figure 2–7.** Male genitalia of *Psallus thomashenryi* sp. n.: **2** phallosome **3** left paramere **4** right paramere **5** vesica in lateral view **6–7** apex of vesica in different views. Scale bar: 0.2 mm.

index 2.2. Female genitalia could not be examined due to the imperfectly sclerotized single specimen.

**Measurements** (in mm). *Male*. Body length: 2.29; head width: 0.61; interocular distance: 0.27; pronotum width: 0.82; length of antennal segments: I – 0.13, II – 0.63, III – 0.29, IV – 0.24; length of tarsomeres: I – 0.11, II – 0.13, III – 0.15. *Female*. Body length: 2.38; head width: 0.61; interocular distance: 0.32; pronotum width: 0.89; length of antennal segments: I – 0.14, II – 0.58, III and IV missing.

**Differential diagnosis.** The dorsal coloration almost uniformly orange and the C-shaped vesica with elongate apical processes show clearly that the new species belongs to the subgenus *Psallus* s. str. Its total length, 2.3 mm in both sexes, makes it one of the smallest species in the subgenus; only a few species have a body length less than or equal to 2.5 mm, namely *P. corsicus* Puton, 1875 and *P. jeitensis* Wagner, 1963, but their coloration and male genitalia differ from those of the new species. By its habitus, *Psallus thomashenryi* is very close to the East-Mediterranean *P. asthenicus* Seidenstücker, 1966, from which, as from any other species of its genus, it can be distinguished by the char-



acteristic male genitalia, especially the unique apical blades of the vesica. In particular, *P. asthenicus* is larger (body length 2.8–3.1 mm), the postbasal lateral spicule of the vesica is membranous and its apical blades are horn-like, gradually tapering, apically pointed, and the central one is shaped like the head of a bird (see Seidenstücker 1966, figs 25a, 25b).

**Etymology.** The new species is named in honor of our colleague Thomas J. Henry on his 70th birthday in recognition of his great contribution to the advancement of heteropterology and as a token of personal friendship and gratitude. The specific epithet is a noun in the genitive case.

**Habitat.** The specimens were beaten from shrubs and trees growing around a small drying-up brook at the village margin. In the same habitat, the new species was collected with the following other species of Miridae: *Amblytylus concolor* Jakovlev, 1877, *Closterotomus annulus* (Brullé, 1832), *C. norwegicus* (Gmelin, 1790), *Globiceps (Paraglobiceps) syriacus* Wagner, 1969, *Heterocordylus (Bothrocranium) carbonellus* Seidenstücker, 1956, *Lepidargyrus syriacus* (Wagner, 1956), *Paredrocoris pectoralis* Reuter, 1878, *Phytocoris (Exophytocoris) parvulus* Reuter, 1880, and *Plagiognathus marivanensis* Linnavuori, 2010.

**Distribution.** Endemic to southern Anatolia.

## Faunistic records and corrections

### *Atomoscelis onusta* (Fieber, 1861)

= *Psallus aurora* (misidentification): Hoberlandt (1956): 54 (record).

**Material examined. TURKEY: Anatolia: Adana Province:** Toros Mts., Kozan, 8.–9. viii.1947, 1 ♂, Exp. N. Mus. ČSR lgt., L. Hoberlandt 1954 det. as *Psallus aurora* (NMPC).

**Comment.** *Psallus aurora* is a West-Mediterranean species known from France, Italy, Portugal, Spain, Algeria, Libya, Morocco and Tunisia (Kerzhner and Josifov 1999). The record from Turkey was considered doubtful by Kerzhner and Josifov (1999: 412). Our recent reexamination of the voucher specimen confirmed that it was misidentified and belongs to the widely distributed Palearctic species *Atomoscelis onusta*. We therefore exclude *P. aurora* from the list of Turkish fauna.

### *Plagiognathus (Plagiognathus) marivanensis* Linnavuori, 2010

*Plagiognathus (Plagiognathus) marivanensis* Linnavuori, 2010: 388 (original description).

**Material examined. TURKEY: Anatolia: Mersin Province:** Göksu Nehri river canyon, Evkafçiftliği, 36°27'23.6"N, 33°38'12.3"E, valley of drying brook, sweeping, 5.v.2007, 1 ♀, P. Kment lgt. (NMPC).

**Distribution.** The species was described recently from western Iran (provinces Kohgiluyeh and Boyerahmad, Kurdistan, and West Azerbaijan) by Linnavuori (2010). It was later recorded from Turkey, eastern Anatolia (provinces Elazığ and Diyarbakır; Matocq et al. 2014). Here the westernmost record of this poorly known species is presented, extending its distribution to southern Anatolia.

### *Psallus (Psallus) lucanicus* Wagner, 1968

*Psallus lucanicus* Wagner, 1968: 273 (original description).

*Psallus balcanicus* Josifov, 1969: 29 (original description). Synonymized by Carapezza (1988: 118, suspected) and Kerzhner and Josifov (1999: 416, confirmed).

**Material examined. TURKEY: Anatolia: Mersin Province:** Yeniköy env., slope above road to Gözne, 36°59'18.5"N 34°30'19"E, on *Quercus* cf. *cerris*, 6.v.2007, 2 ♂♂ 3 ♀♀, P. Kment lgt. (NMPC).

**Host plant.** It is generally collected on *Quercus cerris* (Carapezza 1988, Bryja and Kment 2002, Rabitsch 2003, Anonymus 2016, Denton 2016), but it also is known from *Q. pubescens* (Carapezza 1988) and *Q. macrolepis* (Rieger 2007).

**Distribution.** The species was described from Lucania, a region in Southern Italy whence its name is derived. It was later found in other Italian regions including Sicily (Wagner 1968, Carapezza 1988, Wyniger 2004), and in Austria (Rabitsch 2003), Bulgaria (Josifov 1969, as *P. balcanicus*; Wyniger 2004), Czech Republic (Bryja and Kment 2002, Wyniger 2004), Greece (Rieger 2007), Hungary (Kondorosy 2005), Slovakia (Günther 2000, Wyniger 2004), Slovenia (Gogala and Gogala 1986, Gogala 2006), and United Kingdom (Anonymus 2016, Denton 2016). This is the first record for Turkey and its easternmost occurrence.

## Conclusions

As a result of this study, a total of 36 species of *Psallus* are confirmed to occur in Turkey, as detailed in the following updated checklist (see Kerzhner and Josifov 1999, Lodos et al. 2003, Önder et al. 2006, Konstantinov and Namyatova 2008, Matocq and Pluot-Sigwalt 2011, Aukema et al. 2013, Matocq et al. 2014, Dursun and Fent 2017, Çerçi and Koçak 2017). An E\* indicates the species endemic for the country; non-endemic species are followed by a reference for Turkey.

## Checklist of *Psallus* from Turkey

*Psallus (Apocremnus) anatolicus* Wagner, 1963 E\* (Wagner 1963)

*Psallus (Apocremnus) anticus* (Reuter, 1876) (Hoberlandt 1956)

- Psallus (Apocremnus) betuleti* (Fallén, 1826) (Önder et al. 2006)
- Psallus (Apocremnus) skylia* Linnavuori, 1994 (Matocq et al. 2014)
- Psallus (Hyllopsallus) perrisi* (Mulsant & Rey, 1852) (Wagner 1975a, Lodos et al. 2003)
- Psallus (Hyllopsallus) variabilis* (Fallén, 1807) (Wagner 1975a)
- Psallus (Mesopsallus) ambiguus* (Fallén, 1807) (Kerzhner and Josifov 1999)
- Psallus (Phylidea) cerridis* Wagner, 1971 E\* (Wagner 1971a)
- Psallus (Phylidea) collaris* (Wagner, 1975) E\* (Wagner 1975b)
- Psallus (Phylidea) henshii* Reuter, 1888 (Seidenstücker 1962)
- Psallus (Phylidea) karakardes* Seidenstücker, 1959 E\* (Seidenstücker 1959)
- Psallus (Phylidea) nigripilis* Reuter, 1888 (Kerzhner and Josifov 1999, Matocq et al. 2014)
- Psallus (Phylidea) quercicola* (Reuter, 1904) E\* (Reuter 1904)
- Psallus (Phylidea) quercus* (Kirschbaum, 1856) (Seidenstücker 1959)
- Psallus (Phylidea) syriacus* (Reuter, 1883) (Lodos et al. 2003)
- Psallus (Pityopsallus) piceae* Reuter, 1878 (Hoberlandt 1956)
- Psallus (Pityopsallus) pinicola* Reuter, 1875 (Önder 1976)
- Psallus (Psallus) anaemicus* Seidenstücker, 1966 (Seidenstücker 1966a)
- Psallus (Psallus) apoplecticus* Seidenstücker, 1966 E\* (Seidenstücker 1966a)
- Psallus (Psallus) asthenicus* Seidenstücker, 1966 (Seidenstücker 1966a)
- Psallus (Psallus) brachycerus* Reuter, 1904 (Reuter 1904, Hoberlandt 1956)
- Psallus (Psallus) corsicus* Puton, 1875 (Konstantinov and Namyatova 2008)
- Psallus (Psallus) cruentatus* (Mulsant & Rey, 1852)
- Psallus (?Psallus) inancozgeni* Matocq & Pluot-Sigwalt, 2011 E\* (Matocq and Pluot-Sigwalt 2011)
- Psallus (Psallus) lentigo* Seidenstücker, 1972 (Seidenstücker 1972)
- Psallus (Psallus) lepidus* Fieber, 1858 (Önder 1976)
- Psallus (Psallus) milenae* Josifov, 1974 (Josifov 1974)
- Psallus (Psallus) mollis* (Mulsant & Rey, 1852) (Kerzhner and Josifov 1999)
- Psallus (Psallus) oenderi* Wagner, 1976 E\* (Wagner 1976)
- Psallus (Psallus) oleae* Wagner, 1963 E\* (Wagner 1963b)
- Psallus (Psallus) pardalis* Seidenstücker, 1966 (Seidenstücker 1966b)
- Psallus (Psallus) pseudopunctulatus* Linnavuori, 1984 (Matocq et al. 2014)
- Psallus (Psallus) rubinicterus* Seidenstücker, 1966 E\* (Seidenstücker 1966a)
- Psallus (Psallus) turcicus* Wagner, 1971 E\* (Wagner 1971b)
- Psallus (Psallus) thomashenryi* sp. n. E\*
- Psallus (Psallus) varians varians* (Herrich-Schaeffer, 1841) (Hoberlandt 1956)

The total number of 36 species is comparable to the number of species of *Psallus* occurring in other northern Mediterranean countries such as France (31) or Italy (34), but with an important difference in the percentage of endemic species. France has no endemic species and Italy has only two endemics, both restricted to Southern Italy and/or Sicily, which make 5.9 % of the total number (Wagner 1975a, Kerzhner and Josifov 1999, Schuh 2013). In Turkey, 12 of a total of 36 species are endemic, with



a percentage of 33.3 %. With the exception of *Psallus (Phylidea) quercicola* (Reuter, 1904), known also from the European part of Turkey, all Turkish endemic species of *Psallus* are known only from Anatolia (Seidenstücker 1959, 1962, 1966a,b, 1972; Linnavuori 1994; Lodos et al. 2003; Önder et al. 2006; Konstantinov and Namyatova 2008; Matocq and Pluot-Sigwalt 2011; Matocq et al. 2014; Dursun and Fent 2017). Moreover, one species, *Psallus dionysos* Simon & Strauss, 2014, is endemic to the Greek island of Lesbos, divided only by a narrow strait from the western coast of Anatolia (Simon and Strauss 2014) and another, *Psallus (Apocremnus) cyprius* Wagner, 1977, is endemic to Cyprus, not far from the southern coast of Anatolia (Linnavuori 1994). A similar high level of endemism occurs in two areas adjoining Anatolia, the Transcaucasian and the Syrian regions. In the three countries of Transcaucasia (Armenia, Azerbaijan and Georgia) 23 species of *Psallus* are known, 9 of which are endemic, with a percentage of 36 % (Zaitzeva 1968, Drapolyuk 1991, Kerzhner and Josifov 1999, Konstantinov and Namyatova 2008, Schuh 2013). In the countries of the Syrian region (Jordan, Iraq, Israel, Lebanon, Syria) 11 species of *Psallus* are known to occur, 9 of which are endemic, with a high percentage of 72.7 % (Wagner 1975a, Linnavuori 1984, Kerzhner and Josifov 1999, Carapezza 2002, Schuh 2013). In addition to the area considered above, the Palearctic region has two more centers of endemism for the genus *Psallus*: a minor one in the western Mediterranean (four endemic species in Spain, two in Italy, and one in Algeria) (Wagner 1975a, Kerzhner and Josifov 1999, Konstantinov and Namyatova 2008, Schuh 2013, Pagola-Carte 2017) and a major one in the Palearctic Far East of Asia (seven endemic species in the Far East of Russia, seven in the Korean Peninsula, seven in China, and nine in Japan) (Josifov 1983, Zheng and Li 1990, Li and Zheng 1991, Vinokurov 1998, Kerzhner and Josifov 1999, Yasunaga and Vinokurov 2000, Duwal et al. 2012, Schuh 2013, Duwal and Lee 2015). These data show the high relevance of the Syro-anatolian-transcaucasian region as center of origin of the speciation process of the genus *Psallus* for the Western Palearctic.

## Acknowledgements

The work was financially supported by the Ministry of Culture of the Czech Republic (DKRVO 2017/14, MK000023272, National Museum, Prague). Attilio Carapezza's visit to NMPC received financial support from the programme SYNTHESYS (CZ-TAF-5917).

## References

- Anonymus (2016) Species new to Britain: *Psallus lucanicus* Wagner, 1968 (Miridae). *HetNews* 23: 4.
- Aukema B, Rieger Ch, Rabitsch W (2013) Catalogue of the Heteroptera of the Palearctic Region. Vol. 6. Supplement. The Netherlands Entomological Society, Amsterdam, 629 pp.

- Bryja J, Kment P (2002) New and interesting records of plant bugs (Heteroptera: Miridae) from the Czech and Slovak Republics. *Klapalekiana* 38: 1–10.
- Carapezza A (1988) Settanta Eterotteri nuovi per la fauna siciliana (Insecta, Heteroptera). *Naturalista Siciliano, Série IV* 12: 107–126.
- Carapezza A (2002) Heteroptera of Jordan: new taxa and new records. *Naturalista Siciliano, Serie IV* 26: 35–76.
- Çerçi B, Koçak Ö (2017) Further contributions to the Heteroptera (Hemiptera) fauna of Turkey with a new synonymy. *Acta Biologica Turcica* 30: 121–127.
- Denton J (2016) Some Hemiptera new to Kent from Greenwich Park. *British Journal of Entomology and Natural History* 29: 154.
- Drapolyuk IS (1991) Novye vidy slepnyakov (Heteroptera, Miridae) iz Azerbaydzhana. (New species of capsid bugs (Heteroptera, Miridae) from Azerbaijan). *Entomologicheskoe Obozrenie* 70: 396–403. [In Russian with English summary; English translation in *Entomological Review* 71: 76–83, 1992]
- Dursun A, Fent M (2017) Type localities of Heteroptera (Insecta: Hemiptera) from Turkey. *Zootaxa* 4227(4): 451–494. <https://doi.org/10.11646/zootaxa.4227.4.1>
- Duwal RK, Lee S-H (2015) Additional descriptions of the plant bug genus *Psallus* from the Korean Peninsula (Hemiptera: Heteroptera: Miridae: Phylinae). *Zootaxa* 3926(4): 585–594. <https://doi.org/10.11646/zootaxa.3926.4.8>
- Duwal RK, Yasunaga T, Jung S-H, Lee S-H (2012) The plant bug genus *Psallus* (Heteroptera: Miridae) in the Korean Peninsula with descriptions of three new species. *European Journal of Entomology* 109: 603–632. <https://doi.org/10.14411/eje.2012.074>
- Gogala A (2006) Heteroptera of Slovenia, III: Miridae. *Annales for Istrian and Mediterranean Studies (Historia Naturalis)* 16: 77–112.
- Gogala A, Gogala M (1986) Seznam vrst stenic, ugotovljenih v Sloveniji (Insecta: Heteroptera). [True bugs of Slovenia (Insecta: Heteroptera)]. *Biološki Vestnik* 34: 21–52. [In Slovene, English summary]
- Günther H (2000) Contribution to the fauna of plant bugs (Heteroptera: Miridae) in Slovakia. *Entomological Problems* 3: 59–60.
- Hoberlandt L (1956) Results of the Zoological Scientific Expedition of the National Museum in Praha to Turkey. 18. Hemiptera IV. Terrestrial Hemiptera-Heteroptera of Turkey. *Acta Entomologica Musei Nationalis Pragae, Supplementum* 3[1955]: 1–264.
- Josifov M (1969) Einige neue Miriden aus Bulgarien (Hemiptera, Heteroptera). *Reichenbachia* 12: 29–36.
- Josifov M (1974) Eine neue *Psallus*-Art aus Bulgarien und eine neue *Orthotylus*-Art aus Kirgisien (Heteroptera, Miridae). *Reichenbachia* 15: 89–92.
- Josifov M (1983) Neue *Psallus*-Arten aus Nord-Korea (KDVR) (Heteroptera, Miridae). *Reichenbachia* 21: 197–211.
- Kerzhner IM, Josifov M (1999) Miridae Hahn, 1833. In: Aukema B, Rieger Ch (Eds) *Catalogue of the Heteroptera of the Palaearctic Region, Volume 3: Cimicomorpha II. The Netherlands Entomological Society, Amsterdam*, 1–577.
- Kondorosy E (2005) New true bug species in the Hungarian fauna (Heteroptera). *Folia Entomologica Hungarica* 66: 17–22.

- Konstantinov FV (2003) Male genitalia in Miridae (Heteroptera) and their significance for suprageneric classification of the family. Part I: general review, Isometopinae and Psallopinae. *Belgian Journal of Entomology* 5: 3–36.
- Konstantinov FV (2016) New synonymies in the plant bug family Miridae (Hemiptera: Heteroptera) from Northern China. *Zootaxa* 4205(5): 496–500. <https://doi.org/10.11646/zootaxa.4205.5.10>
- Konstantinov FV, Namyatova AA (2008) New records of Phylinae (Hemiptera: Heteroptera: Miridae) from the Palaearctic Region. *Zootaxa* 1870: 24–42.
- Li X-M, Liu G-Q (2013) A new species and newly recorded genus of the tribe Phylini (Hemiptera: Miridae: Phylinae) from China. *Entomotaxonomia* 35: 185–190.
- Li H-Y, Zheng L-Y (1991) A preliminary report on *Psallus* Fieber (Heteroptera: Miridae) from China. *Acta Scientiarum Naturalium Universitatis Nankaiensis* 1: 1–11. [In Chinese, English summary]
- Linnavuori RE (1984) New species of Hemiptera Heteroptera from Iraq and the adjacent countries. *Acta Entomologica Fennica* 44: 1–59.
- Linnavuori RE (1994) Studies on the Mediterranean Miridae fauna (Hemiptera, Heteroptera). *Biologia Gallo-Hellenica* 21: 5–34.
- Linnavuori RE (2010) Studies on the Miridae (Phylinae, addenda to Deraeocorinae and Orthotylinae) of Khuzestan and the adjacent provinces of Iran (Hemiptera: Heteroptera). *Acta Entomologica Musei Nationalis Pragae* 50: 369–414.
- Lodos N, Önder F, Pehlivan E, Atalay R, Erkin E, Karsavuran Y, Tezcan S, Aksoy S (2003) Faunistic studies on Miridae (Heteroptera) of western Black Sea, central Anatolia and Mediterranean regions of Turkey. *Ege Üniversitesi Basımevi, İzmir*, 85 pp.
- Matocq A (2011) Mise au point sur *Psallus vicinus* Reuter, 1899, une espèce peu connue [Hemiptera, Heteroptera, Miridae]. *Revue Française d'Entomologie (Nouvelle Série)* 33: 1–4.
- Matocq A, Pluot-Sigwalt D (2011) Une nouvelle espèce de *Psallus* de Turquie associée à *Fraxinus* sp. (Oleaceae) (Hemiptera, Heteroptera, Miridae, Phylinae). *Nouvelle Revue d'Entomologie (Nouvelle Série)* 27(2) [2010–2011]: 163–172.
- Matocq A, Pluot-Sigwalt D, Özgen İ (2014) Terrestrial Hemiptera (Heteroptera) collected in South-East Anatolia (Diyarbakır, Mardin and Elazığ Provinces) (Turkey): Second List. *Munis Entomology and Zoology* 9: 884–930.
- Mról E (2012) The structure of the male reproductive system of the genus *Psallus* Fieber (Hemiptera: Heteroptera: Miridae). *Polish Journal of Entomology* 81: 107–118. <https://doi.org/10.2478/v10200-011-0070-8>
- Önder F (1976) Türkiye Miridae (Hemiptera) faunası üzerinde sistematik çalışmalar. [Systematic studies on the fauna of Turkish Miridae (Hemiptera)]. *Ege Üniversitesi Ziraat Fakültesi, Entomoloji ve Ziraat Zooloji Kürsüsü, Doçentlik Tezi, İzmir*, 506 pp. [In Turkish]
- Önder F, Karsavuran Y, Tezcan S, Fent M (2006) Türkiye Heteroptera (Insecta) kataloğu. (Heteroptera (Insecta) catalogue of Turkey). *Ege Üniversitesi Ziraat Fakültesi, İzmir*, 164 pp. [In Turkish with English title and foreword]
- Pagola-Carte S (2017) *Psallus (Psallus) anashanti* n. sp. from the Basque Country, northern Iberian Peninsula (Hemiptera: Heteroptera: Miridae). *Heteropterus Revista de Entomología* 17: 9–20.

- Pluot-Sigwalt D, Matocq A (2017) Testing the diagnostic value of the dorsal sac and other structures of the roof of the female genital chamber within the plant bugs genus *Psallus* Fieber (Hemiptera, Heteroptera, Miridae, Phylinae). *Zootaxa* 4286(2): 215–227. <https://doi.org/10.11646/zootaxa.4286.2.5>
- Rabitsch W (2003) Neue und seltene Wanzen (Insecta, Heteroptera) aus Niederösterreich und Wien. Teil 3. *Linzer Biologische Beiträge* 35: 1293–1305.
- Reuter OM (1904) Capsidae novae mediterraneae descriptae. V. Species a dominis J. et U. Sahlberg in itinere a. 1903-1904 collectae. *Öfversigt af Finska Vetenskaps societetens Förhandlingar* 47(4): 1–26.
- Rieger Ch (2007) Neunachweise und Ergänzungen zur Wanzen-Fauna Griechenlands (Insecta: Heteroptera). *Mainzer Naturwissenschaftliches Archiv, Beiheft* 31: 199–207.
- Schuh RT (1995) *Plant bugs of the world* (Insecta: Heteroptera: Miridae). New York Entomological Society, New York, 1329 pp.
- Schuh RT (2013) On-line Systematic Catalog of Plant Bugs (Insecta: Heteroptera: Miridae). <http://research.amnh.org/pbi/catalog>
- Schuh RT, Menard KL (2013) A revised classification of the Phylinae (Insecta: Heteroptera: Miridae): arguments for the placement of genera. *American Museum Novitates* 3785: 1–72. <https://doi.org/10.1206/3785.2>
- Schuh RT, Slater JA (1995) *True bugs of the world* (Hemiptera: Heteroptera): classification and natural history. Cornell University Press, Ithaca and London, 338 pp.
- Schuh RT, Weirauch Ch, Menard K (2014) Resolving the identities of Phylinae (Heteroptera: Miridae) described by O. M. Reuter from Australia in 1904. *Entomologica Americana* 120(1): 4–6. <https://doi.org/10.1664/14-SN-014R.1>
- Seidenstücker G (1959) Heteroptera aus Anatolien II. *Revue de la Faculté des Sciences de l'Université d'Istanbul, Série B*, 23 [1958]: 119–129.
- Seidenstücker G (1962) Über einige Miriden aus Kleinasien mit Beschreibung von zwei neuen Halticinen (Heteroptera). *Reichenbachia* 1: 129–143.
- Seidenstücker G (1966a) Neue *Psallus*-Arten aus der Türkei (Heteroptera, Miridae). *Reichenbachia* 6: 291–302.
- Seidenstücker G (1966b) *Psallus pardalis* n. sp. (Heteroptera, Miridae). *Reichenbachia* 8: 85–88.
- Seidenstücker G (1972) *Psallus lentigo* n. sp. (Heteroptera, Miridae). *Notulae Entomologicae* 52: 57–64.
- Simon H, Strauss G (2014) *Psallus* (s. str.) *dionysos* n. sp. – eine neue Miridenart (Heteroptera: Miridae) von der Insel Lesbos (Griechenland). *Andrias* 20: 227–230.
- Vinokurov NN (1998) Asian plant bugs of the subgenus *Pityopsallus* E. Wagn., genus *Psallus* Fieb. (Heteroptera: Miridae). *Zoosystematica Rossica* 7: 285–296.
- Vinokurov NN, Luo Z-H (2012) *Psallus jungaricus* sp. n. – a new species of plant bugs from Xinjiang (Western China) (Hemiptera: Heteroptera: Miridae: Phylinae). *Zootaxa* 3394: 25–30.
- Wagner E (1963a) *Psallus anatolicus* n. spec., a new species of Miridae from Turkey (Hem. Het.). *Revue de la Faculté des Sciences de l'Université d'Istanbul, Série B* 26 [1961]: 201–203.
- Wagner E (1963b) Zur Systematik des *Psallus*-Komplexes (Hem., Het., Miridae). *Mitteilungen der Münchner Entomologischen Gesellschaft* 53: 150–163.

- Wagner E (1968) Über einige südeuropäische Miridae (Hemiptera, Heteroptera). Reichenbachia 10: 271–277.
- Wagner E (1971a) Über 2 neue *Psallus*-Arten (Hemiptera, Heteroptera, Miridae). Reichenbachia 14: 19–22.
- Wagner E (1971b) Eine neue und eine bereits bekannte *Psallus*art (Het. Miridae). Nachrichtenblatt der Bayerischen Entomologen 20: 65–71.
- Wagner E (1975a) Die Miridae Hahn, 1831 des Mittelmeerraumes und der Makaronesischen Inseln (Hemiptera, Heteroptera). Teil 3. Entomologische Abhandlungen Staatlichen Museum für Tierkunde in Dresden 40(suppl.): 1–483.
- Wagner E (1975b) Über *Asthenarius* Kerzhner, 1962 (Hemiptera, Heteroptera, Miridae). Reichenbachia 15: 233–244.
- Wagner E (1976) Neue Heteropteren aus der Türkei und dem Libanon (Hemiptera, Heteroptera). Reichenbachia 16: 135–141.
- Wyniger D (2004) Taxonomy and phylogeny of the Central European bugs genus *Psallus* (Hemiptera, Miridae) and faunistics of the terrestrial Heteroptera of Basel and surroundings (Hemiptera). PhD thesis, Philosophisch-Naturwissenschaftlichen Fakultät der Universität Basel, 108 pp.
- Yasunaga T (2010) Plant bugs of the tribe Phylini in Thailand (Heteroptera: Miridae: Phylinae), with descriptions of six new species from additional areas in tropical and subtropical Asia. Entomologica Americana 116: 50–92. <https://doi.org/10.1664/10-RA-006.1>
- Yasunaga T, Vinokurov NN (2000) The phylinae plant bug genus *Psallus* Fieber in Japan (Heteroptera: Miridae: Phylinae). Entomological Science 3: 653–668.
- Zaitzeva IF (1968) Obzor vidov poluzhestkokrylykh roda *Psallus* Fieb. (Heteroptera, Miridae) Kavkaza. (A survey of species of the genus *Psallus* Fieb. (Heteroptera Miridae) from the Caucasus.) Entomologicheskoe Obozrenie 47: 864–877. [In Russian; English translation in Entomological Review 47: 526–533]
- Zheng L-Y, Li H-Y (1990) Four new species of *Psallus* Fieb. from China (Insecta, Hemiptera, Heteroptera: Miridae). Reichenbachia 28: 15–19.