

Determination of Aflatoxin M1 Levels in Produced Pasteurized Milk in Ahvaz City by Using HPLC

Abdolazim Behfar¹, Zahra Nazari Khorasgani^{2*}, Ziyaaddin Alemzadeh¹, Mehdi Goudarzi², Rezvan Ebrahimi², Najmedin Tarhani¹

¹ Department of Food Science and Medical Hydrology, Pharmacy School, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran

² Department of Pharmacology and Toxicology, Pharmacy School, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran

ARTICLE INFO

Article type:
Original Article

Article history:
Received: 1 Feb 2011
Revised: 28 Feb 2011
Accepted: 3 Mar 2011

Keywords:
Aflatoxins
Aflatoxin M₁
Milk
Chromatography, High-Performance
Liquid Chromatography

ABSTRACT

Background: Aflatoxins are one of the most potent toxic substances that occur naturally. Nowadays extensive attention has been taken to their existence in food and environment, as there is the possibility of harm to humans following chronic exposure to extremely low levels via food chain. Aflatoxin M₁ (AFM₁) is a hepatic carcinogenic metabolite found in the milk of lactating animals fed with contaminated feed contaminated by aflatoxin B₁ (AFB₁).

Objectives: This study aimed to determine the levels of AFM₁ in produced pasteurized milk in the Ahvaz of city.

Materials and Methods: For this purpose, 100 samples of pasteurized milk from the Jamus Factory were analyzed to determine AFM₁ content by using an immunoaffinity column for clean-up and high-performance liquid chromatography (HPLC) with a C₁₈ column, a fluorescence detector (excitation 365 nm, emission 435 nm) and a mobile phase of acetonitrile-water (25:75, v/v) at a flow rate of 1 mL/min.

Results: AFM₁ was detected in all 100 samples of pasteurized milk at concentrations ranging from 0.45 to 9.760 ng/L.

Conclusions: The mean concentration of AFM₁ in the the pasteurized milk samples was 2.7 ng/L, which was below the 50 ng/L, accepted as level of for milk in Iran.

Copyright © 2012 DocS.

► Implication for health policy/practice/research/medical education:

This study aimed to increase the knowledge about aflatoxin which produces naturally in food materials.

► Please cite this paper as:

Behfar A, Nazari Khorasgani Z, Alemzadeh Z, Gudarzi M, Ebrahimi R, Tarhani N. Determination of Aflatoxin M1 Levels in Produced Pasteurized Milk in Ahvaz City by Using HPLC. *Jundishapur J Nat Pharm Prod.* 2012;7(2);80-4.

1. Background

Aflatoxins are produced by the *Aspergillus* species under suitable conditions. They are found in a wide variety of products and commodities, including cereals, peanuts, walnuts, and dried fruits (1-8). Five billion people in developing countries all over the world are at risk of

chronic exposure to aflatoxins through contaminated foods (9). One of the metabolites of AFB₁ by cytochrome P₄₅₀ enzyme system in the liver is 4-hydroxy AFB₁ (AFM₁) which is excreted into milk when lactating animals are given feed known to contain aflatoxins (3, 10). The amount of AFM₁ excreted is directly related to the level of AFB₁ in the feed. Milk and milk products are good sources of many nutrients such as proteins, calcium, vitamins, and essential fatty acids. On the other hand, contamination of milk with AFM₁ is considered as a potential risk for human health (11-13). AFM₁ was classified by the International Agency for Research on Cancer (IARC) as a group

* Corresponding author: Zahra Nazari Khorasgani, Department of Pharmacology & Toxicology, Pharmacy School, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran. Tel.: +98-6113738378, Fax: +98-6113738381, Email: znazarikh@yahoo.com
Copyright © 2012 DocS.

2B agent (possibly carcinogenic to humans). It has been experimentally shown to confer high hepatotoxic and mutagenic risk. AFM₁ is relatively stable during pasteurization, sterilization, preparation, and storage of dairy products (13). There is very little data in the literature on AFM₁ levels in the milk produced in Ahvaz, the capital city of Khouzestan province, Iran. Therefore, it is difficult to estimate the daily intake of AFM₁ from milk or other dietary sources, thus there is a need to detect and quantify AFM₁ in milk. Various methods to determine AFM₁ have been developed, including radioimmunoassay, enzyme-linked immunoassay, and high-performance liquid chromatography (HPLC).

2. Objectives

This study was carried out to evaluate AFM₁ levels in pasteurized milk produced in Ahvaz city by using HPLC.

3. Materials and Methods

3.1. Chemicals, Reagents, and Materials

AFM₁ standard was obtained from Sigma Chemical Co. in Iran. Aflatest immunoaffinity columns were purchased from VICAM Co. USA. Acetonitrile HPLC grade was purchased from Merck Co. The stock solution of AFM₁ was prepared in acetonitrile at a concentration of 0.5 µg/ml and was kept at -20°C. Working standard solutions were prepared by of stock standard solution diluting acetonitrile stock solution at concentrations ranging from 0.05 to 100 ng/ml.

3.2. Samples

In this study, 100 composite milk samples, each comprising 5 packs of pasteurized milk, were taken on site at the Jamus Factory from February 2009 to June 2009, and transferred to the Toxicology Lab of the Department of Toxicology and Pharmacology, Pharmacy School of Ahvaz

Jundishapur University of Medical Sciences. All samples were stored at -20°C until analyzed.

3.3. Apparatus

The Shimadzu 10ADvp HPLC system (Japan) was equipped with a Shimadzu RF-10AXL fluorescence detector. Shimadzu LC-10 ADvp pump u, isocratic mode, Shimadzu DGU-14A Degasser, Shimadzu SCL-10Avp System Controller, Shimadzu FCL-10ALvp flow controller, LC solution software. The column (4.6 × 150 mm), which was packed with particles of silica modified with octadecylsilyl groups (5 µm in diameter), was purchased from Capital Co., England.

3.4. Clean-up by Immunoaffinity Column Chromatography

Each sample was warmed at 37°C and centrifuged at 2000×g. The fat layer was removed completely and milk was passed through a paper filter. Then, a 50 ml portion of this prepared sample was taken into a syringe barrel attached to an Aflatest column and passed at the flow rate of 2–3 ml min⁻¹. The column was washed with 20 ml of water and discarded. The sorbent bed was dried and the AFM₁ in the samples was eluted with 4 ml acetonitrile. The solution was evaporated under nitrogen gas and the residue was dissolved in 1 ml of mobile phase.

3.5. Quantitative Analysis

The above solution (200 µl) was injected into the HPLC. Excitation and emission wavelengths were 365 nm and 435 nm, respectively. Acetonitrile–water (25:75 v/v) was used as the mobile phase at the flow rate of 1 ml/min. AFM₁ peak in the chromatogram was identified by comparing its retention time with that of the analyzed AFM₁ standard under the same conditions. The peak was quantified from the area under the curve of sample chromatogram by using the equation of calibration curve ($y = .94481x +$

Table 1. Recoveries for AFM₁ From Spiking Into the one of the Milk Samples (n=6)

Sample type	Spiking levels, ppb	Measurable levels, ppb	Recovery, %
Milk	0.1	0.094	94
	0.5	0.48	97
	1	0.96	98

Table 2. Intra-day and inter-day Precision of Method (n=6)

AFM ₁ concentration, ng/ml	Intra-day, Mean ± SD (µ V*s)	Precision, RSD, %	Inter-day, Mean ± SD (µ V*s)	Precision, RSD, %
0.05	5158.27±578.073	11.224	5005.05±578.973	11.568
0.1	10479.229±442.672	4.224	10559.90±442.672	4.192
0.5	48398.475±887.005	1.833	48769.78±887.005	1.819
1	99154.367 ±1930.621	1.947	97011.275±1930.621	1.990
5	466528.600±4320.988	0.926	324436.4±4320.988	1.332
10	948743.350±3169.465	0.334	939349.233±16424.708	1.749

875.9, $R^2 = 0.9999$). Calibration curve drawn at concentrations of 0.05, 0.1, 0.5, 1, 5, and 10 ng/ml of AFM₁.

The limits of detection and quantitation were 15.5 and 50 ng/L, respectively. Recovery was performed by the standard addition method. To do so, 18 portions (1 ml each) of 0.1, 0.5, and 1 ng/ml of standard solutions (6 repeats for each level) were transferred into 50 ml volumetric flasks and evaporated under nitrogen gas. The residues in the volumetric flasks were diluted to the mark by adding the required amount of one of the milk samples whose content of AFM₁ was being analyzed. Then, the procedures above were followed. The results are summarized in Table 1. All recoveries were more than 94%, indicating good accuracy. Intra-day and inter-day precision is shown in Table 2. All measurements were repeated 6 times. The %RSDs of intra-day and inter-day analyses were in the range of 0.334–11.224 and 1.332–11.568, respectively. These data indicate that the method has acceptable precision.

4. Results

The average recoveries and relative standard deviation of the analytical method applied for AFM₁ in milk were investigated. The results are shown in Tables 1 and 2. The highest and lowest concentrations of AFM₁ were 9.76 and 0.45 ng/L respectively (Table 3). The mean of AFM₁ concentration in samples was 2.7 ng/L (Table 3). Retention time under this condition was 9.478 ± 0.236 min (Figures 1 and 2).

5. Discussion

Since milk and dairy products are an important source of nutrition in the human diet, the presence of AFM₁ in milk and milk products has been investigated worldwide. In 1996, Galvano, F *et al.* examined for the presence of AFM₁ in 161 samples of milk, 92 samples of dry milk for infant formula, and 120 samples of yogurt obtained from supermarkets and drug stores in 4 large Italian cities by using immunoaffinity column extraction and HPLC. AFM₁ was detected in 125 (78%) of milk samples (ranging from <0.001 µg/L to 0.0235 µg/L; mean level 0.00628 µg/L), 49 (53%) of dry milk samples (ranging from <0.001 µg/L to 0.0796 µg/kg; mean level 0.0322 µg/kg), and 73 (61%) of yogurt samples (ranging from <0.001 µg/kg to 0.0321 µg/kg;

mean level 0.00906 µg/kg).

Only 4 samples of dry milk were over the legal limit established by the European Community (EC) in 1999 (14). In October–July 2000, Bognanno, M. *et al.* analyzed 240 samples of dairy ewes' milk from farms in Enna (Sicily, Italy) for AFM₁ by using HPLC equipped with a fluorescence detector. The limit of detection was 0.250 µg/L for AFM₁. All positive milk samples for AFM₁ were confirmed by LC-MS. AFM₁ was detected in 81% of milk samples, ranging from 0.002 to 0.108 µg/L. Three samples were over the permission limit (0.05 µg/L) (15). Zinedine, A. *et al.*, Jordi investigated 54 samples of pasteurized milk produced in 5 different dairies from Morocco for the presence of AFM₁ using immunoaffinity columns, liquid chromatography, fluorescence. Their results showed that 88.8% samples were contaminated with AFM₁; 7.4% were above the maximum level of 0.05 µg/L set by Moroccan and European regulations for AFM₁ in liquid milk. The incidence of AFM₁ in milk from these 5 different dairies were 100, 92.3, 90, 83.3, and 77.7% respectively, with AFM₁ levels ranging from 0.001 to 0.117 µg/L and a mean value of 0.0186 µg/L (16).

Tekinsen, K. Kaan and Eken, H. Semih analyzed 100 UHT milk and 132 Kasha cheese samples from retail outlets in 5 large cities (Istanbul, Izmir, Konya, Tekirdag, and Edirne) for AFM₁ by using ELISA. Sixty-seven percent UHT milk samples and 82.6% Kasha cheese samples contained AFM₁. The incidence of AFM₁ in the UHT milk and Kasha cheese samples ranged from 0.010 to 0.630 µg/kg and from 0.050 to 0.690 µg/kg, respectively. AFM₁ levels in 31 (31%) UHT milk samples and 36 (27.3%) Kasha cheese samples exceeded the maximum tolerable limit proposed by EC and TFC. AFM₁ levels in the samples indicate high aflatoxin levels, thereby constituting a human health risk in Turkey (17). Srivastava, V. P. *et al.* measured 54 samples of fresh full cream and skimmed skim milk, powdered milk, yogurt, and infant formula for AFM₁ by using HPLC after sample clean-up using immune affinity columns in Ku-




Figure 1. HPLC Chromatogram of 100 ng/ml AFM₁ Standard Solution




Figure 2. HPLC-FD Chromatogram of Milk Containing Aflatoxin M₁.

Table 3. Descriptive Statistics of Data of Investigated Milk Samples (ng/L)

Type sample	N	Minimum	Maximum	Mean	Std. Deviation	Std. Error of Mean
milk	100	0.45	9.76	2.7	1.878256	0.419991

wait. A total of 28% of samples were contaminated with AFM₁, with 6% above the maximum permissible limit of 0.2 µg/L. According to their results, 3 fresh cow milk samples collected from a private local producer showed the highest level of 0.21 µg/L AFM₁. There was no contamination with AFM₁ in powdered milk and infant formula (18).

In 1984, Piva, G. *et al.* tested 313 samples of imported liquid milk and 159 samples of imported cheese for AFM₁; 225 milk samples were obtained from Federal Republic of (FR) Germany and 88 from France, while 82 cheese samples were obtained from France, 34 from FR Germany, and 43 from the Netherlands. The number of positive samples was low for both German (13.8%) and for French (12.5%) milk, and the contamination levels were very low (maximum 23 ng/L). As regards the cheeses, AFM₁ was detected in 19.5, 26.5, and 53.5% French, German, and Dutch samples, respectively, but only 2 French samples exceeded 250 ng/kg (the limit set by Swiss law). In 1985, 2 surveys were carried out on 276 milk samples mostly obtained from individual farms and on 416 cheese samples obtained from all parts of the country. As regards the milk samples, 70 (25.3%) contained AFM₁, but generally at very low levels; in fact only 7 (2.5%) samples exceeded 50 ng/L. AFM₁ was found in 130 (31.3%) cheese samples, but again only 9 (2.2%) exceeded 250 ng/kg. There was no significant difference in AFM₁ levels between Italian, German, and French cheese samples, but these were significantly lower ($P < 0.01$) than in Dutch samples (19).

Sefidgar, S. A. *et al.* collected raw cow's milk samples from milk churns at 40 traditional and semi-industrial cattle farms located in Babol (Northern Iran) in the winter of 2006. In total, they analyzed 120 raw milk samples for AFM₁ contamination by ELISA. Sixty-eight out of 120 samples (56.7%) had AFM₁ levels ranging from 50 to 352.3 ng/L. Fifty-two samples (43.3%) contained AFM₁ at 4–50 ng/L. AFM₁ contamination levels were 4–352.3 ng/L with an average of 102.73 ng/L. Their results indicated that 56.7% of samples were above the limit of European community regulations (0.050 µg/L). In other words, AFM₁ contamination levels in raw milk were more than twice as high as permitted levels (20).

Mohamadi Sani, A. *et al.* evaluated AFM₁ contamination and antibiotic presence in milk samples in the Khorasan province in Iran. For 4 months (March to June 2008), 196 milk samples were collected from 7 dairies. The presence and concentration range of AFM₁ in the samples were investigated by ELISA. AFM₁ was found in 100% of the examined milk samples with an average concentration of 0.07792 µg/kg. The concentrations of AFM₁ in all samples were lower than the Iranian national standard and the FDA limit (0.5 µg/L), but 80.6% samples had an AFM₁ level

greater than the maximum limit (0.050 µg/L) accepted by the European Union and the Codex Alimentarius Commission. There was no significant difference between the mean AFM₁ concentrations in the milk samples obtained from different factories ($P > 0.05$) (21).

Heshmati, Ali *et al.* determined the levels of AFM₁ in 210 UHT milk samples obtained from supermarkets in Tehran, Iran by using ELISA. AFM₁ was found in 116 (55.2%) of 210 UHT milk samples. The levels of AFM₁ in 70 (33.3%) samples were higher than the maximum limit (0.05 µg/L) accepted by Iran and some European countries, while none of the samples exceeded the prescribed limit of US regulations. The highest mean concentration of AFM₁ was recorded at 0.087 µg/L and the lowest at 0.021 µg/L. The incidence of AFM₁ levels exceeding legal limits in UHT milk samples (33.3%) was much higher relative to some other countries. It was therefore concluded that the levels of AFM₁ in the UHT milk samples in Iran were high and seemed to pose a threat to public health (22).

The results of this study showed that all 100 investigated pasteurized milk samples were contaminated with AFM₁ at levels ranging from 0.45 to 9.7 ng/L (mean, 2.7 ng/L). Therefore, all milk samples contained AFM₁ below the maximum limit of 50 ng/L for milk in Iran. These results highlight the necessity of a survey involving a larger number of milk and milk product samples, and suggest that currently, the contamination of milk and milk products with AFM₁ does not appear to pose a serious health problem to Ahvaz city in the Khuzestan province of Iran. Nevertheless, a continuous surveillance program may be warranted to monitor the occurrence of aflatoxins in animal feeds responsible for the present limited contamination. In addition, prolonged storage of cereal and nuts in warm and humid conditions should be avoided in order to minimize the risk of aflatoxin contamination.

Financial Disclosure

None declared.

Funding/Support

None declared.

Acknowledgments

None declared.

References

1. Vasanthi S, Bhat RV. Mycotoxins in foods—occurrence, health & economic significance & food control measures. *Indian J Med Res.* 1998;108:212-24.
2. van Egmond HP, Jonker MA. Regulations and Limits for Mycotoxins in Fruits and Vegetables. In: Rivka B-G, Nachman P, editors.

- Mycotoxins in Fruits and Vegetables*. San Diego: Academic Press; 2008. p. 45-74.
3. Unusan N. Occurrence of aflatoxin M1 in UHT milk in Turkey. *Food Chem Toxicol*. 2006;**44**(11):1897-900.
 4. van Egmond HP. Mycotoxins in dairy products. *Food Chem*. 1983;**11**(4):289-307.
 5. Saleemullah, Iqbal A, Khalil IA, Shah H. Aflatoxin contents of stored and artificially inoculated cereals and nuts. *Food Chem*. 2006;**98**(4):699-703.
 6. Reiter EV, Vouk F, Böhm J, Razzazi-Fazeli E. Aflatoxins in rice - A limited survey of products marketed in Austria. *Food Control*. 2010;**21**(7):988-91.
 7. Richard E, Heutte N, Bouchart V, Garon D. Evaluation of fungal contamination and mycotoxin production in maize silage. *Animal Feed Sci Tech*. 2009;**148**(2-4):309-20.
 8. van Egmond HP, Dragacci S. Liquid chromatographic method for aflatoxin M1 in milk. *Methods Mol Biol*. 2001;**157**:59-69.
 9. Shephard GS, Sewram V. Determination of the mycotoxin fumonisin B1 in maize by reversed-phase thin-layer chromatography: a collaborative study. *Food Addit Contam*. 2004;**21**(5):505.
 10. Sassahara M, Pontes Netto D, Yanaka EK. Aflatoxin occurrence in foodstuff supplied to dairy cattle and aflatoxin M1 in raw milk in the North of Paraná state. *Food Chem Toxicol*. 2005;**43**(6):981-4.
 11. Udagawa S. [Fungal spoilage of foods and its risk assessment]. *Nippon Ishinkin Gakkai Zasshi*. 2005;**46**(1):1-5.
 12. van Egmond HP, Stavenuiter JF. [Developments in research on mycotoxins]. *Tijdschr Diergeneesk*. 1985;**110**(23):1002-7.
 13. Stoloff L, Trucksess M, Hardin N, Francis OJ, Hayes JR, Polan CE, et al. Stability of Aflatoxin M in Milk. *J Dairy Sci*. 1975;**58**(12):1789-93.
 14. Galvano F, Galofaro V, Ritieni A, Bognanno M, De Angelis A, Galvano G. Survey of the occurrence of aflatoxin M1 in dairy products marketed in Italy: second year of observation. *Food Addit Contam*. 2001;**18**(7):644-6.
 15. Bognanno M, La Fauci L, Ritieni A, Tafuri A, De Lorenzo A, Micari P, et al. Survey of the occurrence of Aflatoxin M1 in ovine milk by HPLC and its confirmation by MS. *Mol Nutr Food Res*. 2006;**50**(3):300-5.
 16. Zinedine A, Mañes J. Occurrence and legislation of mycotoxins in food and feed from Morocco. *Food Control*. 2009;**20**(4):334-44.
 17. Tekinsen KK, Eken HS. Aflatoxin M1 levels in UHT milk and kashar cheese consumed in Turkey. *Food Chem Toxicol*. 2008;**46**(10):3287-9.
 18. Srivastava VP, Bu-Abbas A, Alaa B, Al-Johar W, Al-Mufti S, Siddiqui MK. Aflatoxin M1 contamination in commercial samples of milk and dairy products in Kuwait. *Food Addit Contam*. 2001;**18**(11):993-7.
 19. Piva G, Pietri A, Galazzi L, Curto O. Aflatoxin M1 occurrence in dairy products marketed in Italy. *Food Addit Contam*. 1988;**5**(2):133-9.
 20. Sefidgar SA, Azizi G, Khosravi AR, Roudbar-Mohammadi S. Presence of Aflatoxin M1 in raw milk at cattle farms in Babol, Iran. *Pak J Biol Sci*. 2008;**11**(3):484-6.
 21. Mohamadi Sani A, Nikpooyan H, Moshiri R. Aflatoxin M1 contamination and antibiotic residue in milk in Khorasan province, Iran. *Food Chem Toxicol*. 2010;**48**(8-9):2130-2.
 22. Heshmati A, Milani JM. Contamination of UHT milk by aflatoxin M1 in Iran. *Food Control*. 2010;**21**(1):19-22.