

## RESEARCH ARTICLE

# Effect of HIV status and retinol on immunogenicity to oral cholera vaccine in adult population living in an endemic area of Lukanga Swamps, Zambia

Charlie Chaluma Luchen<sup>1,2\*</sup>, John Mwaba<sup>1,3</sup>, Harriet Ng'ombe<sup>1,3</sup>, Peter Ibukun Oluwa Alabi<sup>1</sup>, Michelo Simuyandi<sup>1</sup>, Obvious N. Chilyabanyama<sup>1</sup>, Luiza Miyanda Hatyoka<sup>1</sup>, Cynthia Mubanga<sup>1</sup>, Samuel Bosomprah<sup>1,4</sup>, Roma Chilengi<sup>1</sup>, Cleopatra Caroline Chisenga<sup>1</sup>

**1** Enteric Diseases and Vaccine Research Unit, Centre for Infectious Disease Research in Zambia (CIDRZ), Lusaka, Zambia, **2** Amsterdam UMC, University of Amsterdam, Institute for Infection and Immunity, Amsterdam, the Netherlands, **3** University of Zambia, School of Health Sciences, Lusaka, Zambia, **4** Department of Biostatistics, School of Public Health, University of Ghana, Accra, Ghana

\* [Chaluma.Luchen@cidrz.org](mailto:Chaluma.Luchen@cidrz.org)



## OPEN ACCESS

**Citation:** Luchen CC, Mwaba J, Ng'ombe H, Alabi PIO, Simuyandi M, Chilyabanyama ON, et al. (2021) Effect of HIV status and retinol on immunogenicity to oral cholera vaccine in adult population living in an endemic area of Lukanga Swamps, Zambia. *PLoS ONE* 16(12): e0260552. <https://doi.org/10.1371/journal.pone.0260552>

**Editor:** Aftab A. Ansari, Emory University School of Medicine, UNITED STATES

**Received:** June 19, 2021

**Accepted:** October 20, 2021

**Published:** December 2, 2021

**Copyright:** © 2021 Luchen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Data will be made available to any interested researchers upon request however, the data set cannot be made publicly available because it contains human research participant data. The CIDRZ Ethics and Compliance Committee is responsible for approving data requests. Therefore, the data will be available to any interested persons upon request provided they write to the Secretary of the Committee/Head of Research Operations through this email address: [info@cidrz.org](mailto:info@cidrz.org). The request

## Abstract

### Background

We set out to assess the impact of human immunodeficiency virus (HIV) and micronutrient deficiency as indicated by serum retinol levels on the immune responses to Oral Cholera Vaccine (Shanchol™) in a cohort of participants in Lukanga Swamps, Zambia. Cholera remains endemic in Zambia with vaccines being the only effective preventive measures. However, the effect of these vaccines on populations living with HIV has not been widely documented.

### Methods

HIV testing and confirmation was done using the Alere Determine™ HIV-1/2 and Uni-Gold™ kits while vibriocidal antibody assay was applied for vaccine immunogenicity. Serum retinol analysis was assessed by Shimadzu Prominence HCT-2010 High Performance Liquid Chromatography (HPLC). The primary outcome was log transformed geometric mean titre.

### Results

From 47 participants screened for HIV, 51% (24) tested positive. There was a statistically significant reduction in Ogawa geometric mean ratio (GMR) by 67% (GMR = 0.33; 95% CI: -0.15, 0.76; p-value = 0.009) attributable to HIV positivity with a non-significant reduction in Inaba GMR by about 50% due to HIV positivity. When doubling of retinol levels modelled, GMR reduction against Ogawa were non-significant but that against Inaba resulted in a significant reduction in geometric mean titer (GMT) (GMT-0.33, C.I 0.16–0.66, p-value 0.002). At 1000copies/ml viral load cut off and 350 cells/μl CD4 counts, Ogawa GMT was two times higher 11.16 (95%CI: 8.20–15.19) versus 6.06 (95%CI: 4.04–9.10) in low viremia

should include the contact information, and intended use of the data. The requested data should only be used for purposes related to the original research or study. The CIDRZ Ethics and Compliance Committee will normally review all data requests within 48–72 hrs (Monday–Friday), and provide notification if access has been granted or additional project information is needed, before access can be granted.

**Funding:** Dr C C Chisenga received a career development fellowship award. Grant number: TMA2016CDF-1550. This study was funded by The European Developing Countries Clinical Trials Partnership (EDTCP). <http://www.edctp.org/projects-2/edctp2-projects/career-development-fellowships-2016/#> The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

participants, and three times higher in above threshold CD4 count participants; 24.81 (95% CI: 18.94–32.50) versus 7.07 (95%CI: 5.22–9.58).

## Conclusion

Our results show that while Shanchol™ is immunogenic in both HIV+/- individuals, HIV + participants responded poorly. Viral load and CD4 count affected vaccine immunogenicity. More research is required for detailed understanding of this in order to appropriately inform policy and practice.

## Introduction

Cholera is an acute watery diarrheal disease caused by *Vibrio cholerae* serogroups O1 and O139. Serogroup O1 is divided into two major serotypes of Inaba and Ogawa [1]. Cholera outbreaks are common in endemic settings, which also tend to have the world's poorest and vulnerable populations [2,3]. The disease affects at least 47 countries worldwide with an estimated 2.9 million cases and about 95,000 annual deaths [4]. However, these figures only account for about 5–10% of the actual number as most of the cases go unreported [5].

Sub-Saharan Africa remains a hub of cholera outbreaks since 1970 with Zambia reporting its first outbreak in 1977 [6]. Since then, Zambia has had recurrent outbreaks occurring almost yearly [7]. According to the WHO country stratification on the incidence rate and fatality rate, Zambia is classified in the (E) region, which is a region with high child and very high adult mortality attributed to cholera [5]. The Lukanga Swamps (study site), is located in Central Province of Zambia and often records recurrent cholera outbreaks. For instance, in 2013 and 2014, the area recorded 47 and 41 cholera cases respectively while in 2016 alone, a total of 27 cases with two fatalities were recorded [8].

A review of the burden of HIV in lower-to-middle income countries (LMIC) reported an estimated 33.4 million HIV infected individuals living in LMICs [9]. As of 2018, approximately 1.2 million people were living with HIV in Zambia according to a report by UNAIDS [10]. A study carried out in Mozambique reported that people living with HIV have a high prevalence of Cholera (23%) compared to the control group (13%) [11].

Vaccination with oral cholera vaccines (OCV) is an effective way to reduce the high cholera incidence when deployed in combination with improved water, sanitation and hygiene (WASH) facilities [6].

Poor nutrition resulting in malnutrition has been known to affect the immune responses to OCV in individuals living in LMIC [12]. Nutritional status such as retinol levels, influence innate immune cells and epithelial cells associated with mucosal surfaces, thus in the presence of retinol, epithelial cells express retinaldehyde dehydrogenase which enhances antibody production by stimulated B-cells [13–15]. However, in settings of poor nutrition coupled with a high prevalence of HIV infection, it is not clear how these interact to affect OCV uptake. In this study, we evaluated the independent effects of HIV infection and serum retinol level as a derivative of Vitamin A on immunogenicity to Shanchol™ in a cholera endemic setting.

## Materials and methods

### Study setting

The Lukanga Swamps (260,000 hectares, 14° 24'S 027° 38'E), located in the Central Province of Zambia, is one of the cholera hotspots with recurrent annual cholera outbreaks before the

Shanchol™ intervention in 2016. This area has shallow swamps and several lagoons and makes up the largest permanent water body in the Kafue basin. The area has approximately 16,000 inhabitants and the major economic activity of the population is fishing.

## Study design and participants

This was an observational study nested on a cohort study aimed at profiling immunological characteristics of a population at risk of cholera before and after receiving 1<sup>st</sup> and 2<sup>nd</sup> dose of OCV (Shanchol™) intervention. The parent cohort study was part of the Zambian Ministry of Health's cholera prevention and control program, which deployed Shanchol™ and WASH interventions to the affected population in Lukanga Swamps in 2016. A total of 223 adults aged 18 to 65 years, living in the study area, and available for the duration of the study were enrolled in the study and were followed up for 4 years. Adults who had medical conditions such as hepatic disease, diarrhoea within the previous 7 days or a history of persistent diarrhoea (defined as diarrhoea that lasts for 14 days or longer), and ever having received an OCV were excluded. This parent study has been registered as a clinical trial on clinicaltrials.gov with trial # NCT04423159. Of the 223, only 47 consented to voluntary HIV testing and these are the ones included in our study.

Ethical approval was obtained from the University of Zambia Biomedical and Research Ethics Committee (UNZABREC) reference number 007-12-16. Written informed consent was obtained from all participants of the study.

## Procedures

**Vaccine administration and blood sample collection and processing.** The vaccine was stored at a temperature of 2–8°C prior to administration. The first dose of 1.5 ml OCV was given at baseline (day 0) and the second dose on day 28. Both doses were administered orally. Up to 10mls of venous blood was obtained from each participant, in vacutainers, before vaccination (day 0) and on day 28 followed by 6, 12, 24 and 30 months after the first dose, serum was separated and stored at -20°C until laboratory testing.

**Retinol quantification.** Retinol quantification was performed by using High-Performance Liquid Chromatography (HPLC) [16]. The quantification was done on the SHIMADZU Prominence HCT2010 HPLC system. Briefly, 100µl of serum sample/control was mixed with an equal volume of an ethanolic retinyl acetate mixture in a clean capped glass tube to release the retinol from its complex. The retinyl acetate served as the internal standard for the analysis. The resulting mixture was vortexed for 30 seconds. The released retinol was then recovered from the ethanol solution by two extractions of 500µl each of n-hexane mixed into a separate capped tube. The combined hexane extracts were evaporated to dryness under a gentle stream of nitrogen gas. Thereafter, the resulting residue was then dissolved in 100µl of ethanol. A volume of 60 µl of this residue solution was applied on the reverse phase HPLC column (5-µm C18, 25-cm) protected by a Supelcosil™ C-18 guard column and eluted at 1.5 ml/min, using a 98:2, methanol: water solvent, with detection at 325nm.

**HIV testing.** The blood-based HIV test kit, Alere Determine™ HIV1/2 (Alere Scarborough, Inc.) was used to screen for HIV. A second confirmatory test, the Uni-Gold™ HIV (Trinity Biotech Manufacturing Ltd) was used and if the results from both tests were discordant a tie-breaker test SD Bioline (Standard Diagnostics, Inc) was used.

**CD4+ T lymphocytes enumeration.** To enumerate percentage and absolute cell counts of CD4+ T lymphocytes, the FACS Calibur instrument was used using lysed whole blood collected in vacutainers with EDTA. 50µl of the sample was stained directly in a BD Trucount

tube. Fluorescent beads were released by the lyophilized pellet in the tube as it dissolved. During analysis, the absolute number (cells/ $\mu$ l) of positive cells in the sample was determined by comparing cellular events to bead events. The BD Multiset software was used to determine the number of positive cells per microliter of blood (absolute counts).

**HIV-1 viral load testing.** Human plasma collected in EDTA anticoagulant was used for viral load analysis. HIV-1 viral load testing was performed using the COBAS<sup>®</sup> Ampliprep/COBAS<sup>®</sup> Taqman<sup>®</sup> 48 HIV-1 Test. The specimen was prepared using the automated COBAS<sup>®</sup> AmpliPrep Instrument with amplification and detection was done using the COBAS<sup>®</sup> TaqMan<sup>®</sup> 48 Analyzer as per manufacturer's instructions.


**Cholera vibriocidal assay.** The vibriocidal responses were assessed according to previously described methods [17] with some modifications. The strains were incubated with heat-inactivated serum and exogenous guinea pig complement (Sigma Aldrich S1639-5ML) at 37°C for 1 hour, shaking (50 revs/min). Vibriocidal titers were defined as the reciprocal of the highest serum dilution resulting in a 50% reduction in optical density (595 nm) compared to controls without serum. Standard monoclonal antibody (mAb) (Boston, MA, USA, CF29.1.A2) and high titer pooled sera were used to normalize the results in case of inter-assay variations. Seroconversion was defined as a 4-fold or greater increase in vibriocidal titers after vaccination in comparison to the baseline (D0) titers with a titer of 5 assigned in cases where no vibriocidal activity was observed.

### Sample size calculation

The participants of this study were a subset of a larger cohort study that has been described above. The sample size for the large study was calculated to be 176 with 95% confidence ( $\alpha = 0.05$  (two tailed)), 80% power ( $\beta = 0.2$ ), a difference of 0.3 and conservative estimates of 0.5 variance for pre-vaccine and post-vaccine groups. Adding 20% to account for anticipated attrition yields a total required sample size of 212. Thus, for the parent study, 212 participants were enrolled. This current study excluded individuals from the parent study who did not consent to HIV testing (161) or had no baseline titers (4) (Fig 1). Therefore, a total of 47 participants were included in this study.

### Statistical analysis

The primary outcome was geometric mean titer of Vibriocidal immune responses. The titers were transformed using natural logarithm prior to statistical analyses. Participants' socio-demographic and clinical characteristics were presented as frequencies (percentages) and mean (standard deviation). Geometric mean titres and 95% confidence intervals were calculated for key background characteristics. The repeated measurements of the outcomes introduce panel structure into the dataset. Therefore, random-effects log-normal regression model (i.e. 'xtgee' gaussian-family; identity-link command in Stata) was used to estimate the independent effects of HIV status and Retinol on immune response to OCV, adjusting for age, sex, education and occupation. The participant ID represents the panel and the measurement occasions represent the time variable. For the secondary outcome of seroconversion, random-effects log-binomial model (i.e. 'xtlogit' with 'margins' in Stata) was used to estimate the effects of HIV status and Retinol on seroconversion controlling for the confounding effects of age, sex, education and occupation. In all the models, Retinol was transformed into log base 2 so that the effect is interpreted as doubling in retinol levels. Level of statistical significance was set at a 2-tailed p-value of 0.05 or less. Data analysis was performed using Stata 14.2 for Windows (StataCorp, College Station, TX, USA).



**Fig 1. Flow diagram of the participants who took part in the study.**

<https://doi.org/10.1371/journal.pone.0260552.g001>

## Results

### Characteristics of participants

A total of 47 adults were included in the analysis. Of these, 42 (89%) were females. The median age was 38 years (IQR = 31–45). More than half of the participants had an education level of grade 8–12 (56%). The major occupation engaged in by the participants was fishing (77%). Among the HIV+ participants, 15(65%) had CD+4 count of 350cells/ $\mu$ l or more while 78% had a viral load of 1000 copies/ml or less (Table 1).

Overall, the geometric mean titres (GMTs) of *V cholerae* O1 Ogawa and Inaba were [16.95 (95%CI: 13.94–20.60)] and 10.70 [(95%CI: 9.02–12.69)] respectively (Table 1). Ogawa GMT was higher among female participants [25.70 (95%CI: 12.16–54.20)] than male participants [16.10 (95%CI: 13.17–19.66)] and those 35 years old or younger [19.81 (95%CI: 13.57–28.92)] than those older than 35 years old [15.76 (95%CI: 12.55–19.78)]. Study participants who completed grades 1–7 [19.82 (95%CI: 14.03–28.01)] had higher Ogawa GMT than those who had grades 8–12 [13.75 (95%CI: 10.43–18.13)] while those who engaged in fishing [17.38 (95%CI: 13.77–21.95)] also had higher Ogawa GMTs compared to those involved in other occupation [14.74 (95%CI: 9.58–22.68)].

**Table 1. Geometric mean ratios of *V. cholerae* O1 Ogawa and *V. cholerae* O1 Inaba by background characteristics of participants.**

Characteristics	Number (% of total)	<i>V. cholerae</i> O1 Ogawa	<i>V. cholerae</i> O1 Inaba
		GMTs (95% CI)	GMTs (95% CI)
<b>Sex</b>			
Female	5 (11)	25.67 (12.16–54.20)	8.24 (5.43–12.50)
Male	42(89)	16.10 (13.17–19.66)	11.05 (9.18–13.30)
<b>Age(years)</b>			
≤35	16 (34)	19.81 (13.57–28.92)	12.60 (8.61–18.44)
>35	31 (66)	15.76 (12.55–19.78)	9.91 (8.29–11.85)
<b>Education<sup>‡</sup></b>			
Grade 1–7	16(44)	19.82 (14.03–28.01)	11.53 (8.61–15.44)
Grade 8–12	20(56)	13.75 (10.43–18.13)	11.68 (8.72–15.66)
<b>Occupation<sup>‡</sup></b>			
Fishing	34(77)	17.38 (13.77–21.95)	12.69 (10.18–15.81)
Others*	10(23)	14.74 (9.58–22.68)	7.07 (5.56–8.99)
<b>Viral loads (copies/ml)<sup>§§</sup></b>			
>1000	4 (22)	6.06 (4.04–9.10)	6.80 (4.95–9.35)
≤1000	14 (78)	11.16 (8.20–15.19)	6.76 (5.63–8.10)
<b>CD4count (cells/μl)<sup>§§</sup></b>			
<350	8(35)	7.07 (5.22–9.58)	7.33 (5.42–9.93)
≥350	15(65)	24.81 (18.94–32.50)	7.18 (5.84–8.83)
<b>Total</b>	<b>47 (100)</b>	<b>16.95 (13.94–20.60)</b>	<b>10.70 (9.02–12.69)</b>

<sup>‡</sup>Total was not equal to 47 due to missing information.

<sup>§§</sup>Total was not equal to 24 due to missing information.

Others\* refers to traders, farmers and students.

GMT refers to Geometric Mean Titre, CI (confidence interval).

<https://doi.org/10.1371/journal.pone.0260552.t001>

Ogawa GMT was about two times higher in participants with viral loads of 1000copies/ml or less [11.16 (95%CI: 8.20–15.19)] compared to those with more than 1000 copies/ml [6.06 (95%CI: 4.04–9.10)] and about three times higher in participants with CD4 counts of 350 cells/μl or more [24.81 (95%CI: 18.94–32.50)] compared to those with CD4 counts lower than 350 cells/μl [7.07 (95%CI: 5.22–9.58)] (Table 1). Inaba GMT was higher in male participants [11.05 (95%CI: 9.18–13.30)] than the females [8.24 (95%CI: 5.43–12.50)] but lower in the older ones [9.91 (95%CI: 8.29–11.85)] than the young [12.60 (95%CI: 8.61–18.54)]. Inaba GMT does not vary widely by education, viral load and CD4 count, however, participants whose primary occupation was fishing [12.69 (95%CI: 10.18–15.81)] had higher Ogawa GMTs (about two times more) compared to those involved in other occupation [7.07 (95%CI: 5.56–8.99)]. Despite the observed trends the GMT results were not statistically significant.

### Effects of HIV status and retinol on the immune response to OCV

After adjusting for the confounding effects of age, sex, occupation, and education, there was a statistically significant reduction in Ogawa GMR by 67% due to HIV positivity (GMR = 0.33; 95% CI: -0.15, 0.76; p-value = 0.009) (Table 2). Similarly, there was a reduction in Inaba GMR by about 50% due to HIV positivity, but this is likely to be due to chance as it was not statistically significant (GMR = 0.46; 95%CI: 0.19, 1.15; p-value = 0.096) (Table 2). In the HIV positive group, CD4 cell counts and viral loads did not have any significant effects on responses to OCV (S1 Table). For retinol level, there was a reduction in Ogawa GMR due to a doubling of retinol level but again this was not statistically significant (GMR = 0.86; 95%CI: 0.40, 1.86; p-



Table 2. Effects of HIV status and retinol on the immune response to OCV.

HIV Status	Number (% of total)	<i>V cholerae</i> O1 Ogawa			
		Crude GMR (95%CI)	P-value	Adjusted GMR (95% CI)*	P-value
Negative	23(49)	Ref		Ref	
Positive	24(51)	0.46 (0.23–0.93)	0.029	0.33 (0.15–0.76)	<b>0.009</b>
<b>Retinol (µmol/l)**</b>					
Mean ± SD	3.14 ± 0.64	1.10 (0.63–1.94)	0.73	0.86 (0.40–1.86)	0.698
HIV Status	Number (% of total)	<i>V cholerae</i> O1 Inaba			
		Crude GMR (95%CI)	P-value	Adjusted GMR (95% CI)*	P-value
Negative	23(49)	Ref		Ref	
Positive	24(51)	0.49 (0.27–0.92)	0.026	0.46 (0.19–1.15)	0.096
<b>Retinol (µmol/l)**</b>					
Mean ± SD	3.14 ± 0.64	0.65 (0.39–1.07)	0.07	0.33 (0.16–0.66)	<b>0.002</b>

\*Adjusted for sex age, education, & occupation

\*\*Transformed to log2, GMR: Geometric Mean Ratio, CI: Confidence Interval.

<https://doi.org/10.1371/journal.pone.0260552.t002>

value = 0.698) while for Inaba the doubling of retinol level resulted into a significant reduction in GMR (GMR = 0.33; 95%CI: 0.16, 0.66; p-value = 0.002) (Table 2).

### Kinetics of vibriocidal geometric mean titers against Ogawa and Inaba by HIV status

The GMTs against Ogawa and Inaba are shown in Fig 2 below. Against both serotypes, HIV–participants had higher GMTs than HIV + participants across all the time points (DO, D28, M6, M12, M24, M30). The only significant differences in GMTs were at D28 for Ogawa (HIV- 49.79, HIV + 10.76, p-value 0.002) and for Inaba D28, M2 and M24 were significant with GMTs of (22.31, 7.20 p = 0.02), (14.86, 6.55 p = 0.03) and (21.01, 6.89 p = 0.04) respectively.




Fig 2. Kinetics of vibriocidal geometric mean titers (95% CI) against Ogawa and Inaba by HIV status. Red dots indicate HIV + participants and blue dots HIV- participants.

<https://doi.org/10.1371/journal.pone.0260552.g002>

Table 3. Percent seroconversion by HIV status and retinol level.

HIV Status	<i>V cholerae</i> O1 Ogawa				
	Number (% of total)	Crude RR (95%CI)	P-value	Adjusted RR (95% CI)*	P-value
Negative	23(49)	Ref		Ref	
Positive	24(51)	0.88 (0.32–2.39)	0.801	0.90 (0.22–3.63)	0.877
<b>Retinol (<math>\mu\text{mol/l}</math>)**</b>					
Mean $\pm$ SD	3.14 $\pm$ 0.64	0.96 (0.85–1.09)	0.553	0.92 (0.77–1.10)	0.368
HIV Status	<i>V cholerae</i> O1 Inaba				
	Number (% of total)	Crude RR (95%CI)	P-value	Adjusted RR (95% CI)*	P-value
Negative	23(49)	Ref		Ref	
Positive	24(51)	0.37 (0.10–1.36)	0.127	0.50 (0.11–2.40)	0.375
<b>Retinol (<math>\mu\text{mol/l}</math>)**</b>					
Mean $\pm$ SD	3.14 $\pm$ 0.64	0.96 (0.85–1.09)	0.551	0.88 (0.72–1.08)	0.227

RR = Crude Risk Ratio, RR = Risk Ratio, CI = Confidence Interval, Ref = reference \*Adjusted for sex age, education, & occupation.

<https://doi.org/10.1371/journal.pone.0260552.t003>

### Effects of HIV status and retinol on seroconversion

HIV + individuals compared to the negatives have about 10% lower risk of seroconverting to O1 Ogawa after vaccination although this association is likely to have occurred by chance (RR = 0.90; 95%CI: 0.22, 3.63; p-value = 0.877). Similarly, against Inaba, HIV + respondents are about 2 times less likely to seroconvert after vaccination compared to those who are negative but this association was not statistically significant (RR = 0.50; 95%CI: 0.11, 2.40; p-value = 0.375). A doubled level of retinol reduced the risk of sero-conversion to O1 Ogawa and Inaba by 8% and 12% respectively, although both associations were not statistically significant (RR = 0.92; 95%CI: 0.77, 1.10; p-value = 0.375 & RR = 0.88; 95%CI: 0.72, 1.08; p-value = 0.227) (Table 3).

### Discussions

A majority of the participants enrolled in this study were fishermen; these were at higher risk of cholera infection due to their way of life such as practicing open defecation in the water lagoons that are later used for fetching drinking water. A study in Uganda reported Cholera as the leading cause of morbidity and mortality in Ugandan fishermen [18]. There was a difference in sex distribution among the participants with 10.6% being female and 89.4% male participants as shown in Table 1. This is not a surprising finding for the Zambian setting because fishing is mostly dominated by men in most fishing camps and areas. Despite the unequal distribution of sex, it had no confounding effects on the reported outcomes of immunogenicity and seroconversion.

A noticeable trend, although not statistically significant, pointed towards HIV negative participants having higher vibriocidal GMTs from baseline up until month 30 compared to HIV positive individuals, against both serogroups (Fig 2). This trend is in agreement with what was observed in Haitian adults [19]. It's also consistent with the reports on immune response in the HIV/AIDS population to other vaccines as this population generally has lower antibody titers as compared to healthy individuals [20]. This could be explained by susceptibility to severe disease occurring in HIV infected individuals when secretory IgA, which is supposed to agglutinate the vibrios by binding to the surface antigens, therefore, preventing them from attaching to the mucosal membranes, is depleted [11].

Also, our study observed a better response in our target population against the Ogawa than Inaba serotype. One possible explanation for this is the relative fluctuation of the two serotypes



(Ogawa and Inaba) in an endemic population between epidemics of the disease at a given time-point were one of the two serotypes being responsible for the majority of cases [21,22]. The *V. cholerae* serotypes have been reported to undergo serotype conversions in both directions in most of these endemic areas [23]. Zambia had a cholera outbreak in 2017 and this was predominantly caused by the 01 Ogawa serotype [6]. Given that CD4+ T cell priming is a critical step in vaccination as there is a direct relationship between CD4+ cells and protective immunity and long term humoral responses [24], we hypothesized that the population could have been better primed against Ogawa as compared to Inaba resulting in the observable differences in the trends of GMTs between the two serotypes.

Micronutrients such as vitamin A have been reported to play a crucial role in immune regulation [25,26]. Retinol a derivative of vitamin A has been shown to increase antibody response to T-cell-dependent and T-cell-independent antigens and restoration and maintenance of mucosal function as well as integrity [27]. While retinol supplementation has been known to reduce the severity and pathogen-induced mortality, also at the same time, retinol deficiency leads to impaired mucosal and systemic immune responses henceforth affecting uptake of orally administered vaccines [28,29]. We analyzed retinol levels before administration of Shanchol™ and the results indicated that all our participants had serum retinol levels above 0.7umol/L, a cut-off for retinol deficiency [30].

When the retinol levels were doubled, our study reveals a reduction in GMRs across both serotypes which is contrary to what has been reported on the effects of retinol supplementation on efficacies of other vaccines. For instance, a study elsewhere reported the use of vitamin A as an adjuvant in oral vaccines increasing the vaccine efficacies in settings that have reported low efficacies such as Asia, Latin America and Africa [31]. Our findings might suggest that retinol supplementation during the administration of Shanchol™ in the HIV infected population might not be of significant benefit.

Our study has shown a lack of statistical significance in differences in the seroconversion rates among the HIV- and the HIV + participants following uptake of the OCV Shanchol™. In addition, our analysis of results showed that there was no appreciable boosting in vibriocidal titers following the second dose at 28 days after the first dose in this cohort (Fig 2). This is compatible with what previous studies reported over the lack of benefit of a second dose [32,33]. A hypothesis that could explain this lack of significant impact of the second dose, is that the first dose elicits an immune response in the intestinal mucosal, thus this response possibly blocks the uptake of the second dose [33].

## Limitations

Our study had some limitations. Firstly, the assay is limited in its inability to detect re-exposure; therefore, re-exposure to cholera could have been missed if it did occur and shortly waned off. Secondly, we had several participants that did not show up at all the time-points. We underestimated how mobile fishermen can be, hence the missing data was excluded in some of the analysis, and therefore, going forward this should be put into consideration in such populations.

Thirdly, the second dose was given 28 days past the first dose, which was a deviation from the recommended dose intervals as per manufacturer's instructions. This deviation was due to logistical challenges in getting back to the vaccinated population which is in a remote setting after the recommended 14 days coupled with limited resources. It was going to be interesting if we included a randomized controlled trial sub-study of 14 days versus 28 days Shanchol™ administration so that we could have answered which interval is superior.

Lastly, this study did not look at how other immunological factors like cytokines of the HIV +/- participants and how these could have affected immunogenicity to the vaccine as these

have been reported to qualitatively contribute to different immune responses. Hence, going forward for similar studies, we recommend that such factors be put into consideration when assessing vaccine immune responses. Thus, despite us observing non-significant results in our study, we observed a trend which can guide a hypothesis to be explored with a larger sample size and also by looking at other immunological parameters as mentioned above.

## Conclusion

Our study has demonstrated that the OCV Shanchol™ can elicit an immune response in the HIV+ populations. The observed immunogenicity was dependent on the serotype, the amounts of CD4 + T cells as well as viral load, hence these parameters should be put into consideration when developing as well as deploying a vaccine for this vulnerable population. In addition, the role played by retinol in Shanchol™ immunogenicity was unclear. Also, our study suggests that supplementing with retinol may not be beneficial in OCV uptake as we observed a reduction in GMRs when the retinol amounts were doubled.

## Supporting information

**S1 Table. Effects of viral load and cd4 count on immune response to OCV.**  
(DOCX)

## Acknowledgments

We are grateful for the help rendered by staff from the Ministry of Health and the Enteric Disease Unit at the Center for Infectious Disease Research in Zambia, for assisting in the collection of data.

## Author Contributions

**Conceptualization:** Charlie Chaluma Luchen, John Mwaba, Michelo Simuyandi, Samuel Bosomprah, Roma Chilengi, Cleopatra Caroline Chisenga.

**Data curation:** Charlie Chaluma Luchen, Harriet Ng'ombe, Peter Ibukun Oluwa Alabi, Obvious N. Chilyabanyama, Luiza Miyanda Hatyoka.

**Formal analysis:** Peter Ibukun Oluwa Alabi, Obvious N. Chilyabanyama, Samuel Bosomprah.

**Funding acquisition:** Cleopatra Caroline Chisenga.

**Investigation:** Michelo Simuyandi.

**Methodology:** Charlie Chaluma Luchen, Harriet Ng'ombe, Luiza Miyanda Hatyoka, Cynthia Mubanga.

**Project administration:** Roma Chilengi.

**Supervision:** Samuel Bosomprah, Roma Chilengi, Cleopatra Caroline Chisenga.

**Visualization:** Roma Chilengi.

**Writing – original draft:** Charlie Chaluma Luchen.

**Writing – review & editing:** Charlie Chaluma Luchen, John Mwaba, Luiza Miyanda Hatyoka, Cynthia Mubanga, Roma Chilengi, Cleopatra Caroline Chisenga.

## References

1. Finkelstein RA. Cholera, *Vibrio cholerae* O1 and O139, and Other Pathogenic Vibrios. 4th ed. In: Baron S, editor. *Medical Microbiology*. 4th ed. Galveston (TX): University of Texas Medical Branch at Galveston; 1996. Available: <http://www.ncbi.nlm.nih.gov/books/NBK8407/>.
2. Legros D. Global Cholera Epidemiology: Opportunities to Reduce the Burden of Cholera by 2030. *J Infect Dis*. 2018; 218: S137–S140. <https://doi.org/10.1093/infdis/jiy486> PMID: 30184102
3. Lopez AL, Deen J, Azman AS, Luquero FJ, Kanungo S, Dutta S, et al. Immunogenicity and Protection From a Single Dose of Internationally Available Killed Oral Cholera Vaccine: A Systematic Review and Metaanalysis. *Clin Infect Dis*. 2018; 66: 1960–1971. <https://doi.org/10.1093/cid/cix1039> PMID: 29177437
4. Wierzba TF. Oral cholera vaccines and their impact on the global burden of disease. *Human Vaccines & Immunotherapeutics*. 2019; 15: 1294–1301. <https://doi.org/10.1080/21645515.2018.1504155> PMID: 30183486
5. Ali M, Nelson AR, Lopez AL, Sack DA. Updated Global Burden of Cholera in Endemic Countries. *PLoS Negl Trop Dis*. 2015; 9. <https://doi.org/10.1371/journal.pntd.0003832> PMID: 26043000
6. Sinyange N, Brunkard JM, Kapata N, Mazaba ML, Musonda KG, Hamoonga R, et al. Cholera Epidemic—Lusaka, Zambia, October 2017–May 2018. *MMWR Morb Mortal Wkly Rep*. 2018; 67: 556–559. <https://doi.org/10.15585/mmwr.mm6719a5> PMID: 29771877
7. Kapata N, Sinyange N, Mazaba ML, Musonda K, Hamoonga R, Kapina M, et al. A multisectoral emergency response approach to a cholera outbreak in Zambia: October 2017–February 2018. *Journal of Infectious Diseases*. 2018; 218: S181–S183. <https://doi.org/10.1093/infdis/jiy490> PMID: 30215738
8. Murebwa-Chirambo R, Mwanza R, Mwinuna C, Mazaba M, Mweene-Ndumba I, Mufunda J. Occurrence of cholera in Lukanga fishing camp, Kapiri-mposhi district, Zambia.: 7.
9. Gupta A, Nadkarni G, Yang W-T, Chandrasekhar A, Gupte N, Bisson GP, et al. Early Mortality in Adults Initiating Antiretroviral Therapy (ART) in Low- and Middle-Income Countries (LMIC): A Systematic Review and Meta-Analysis. *PLoS One*. 2011; 6. <https://doi.org/10.1371/journal.pone.0028691> PMID: 22220193
10. Zambia | UNAIDS. [cited 4 Jun 2020]. Available: <https://www.unaids.org/en/regionscountries/countries/zambia>.
11. Seidlein LV, Wang X-Y, Macuamule A, Mondlane C, Puri M, Hendriksen I, et al. Is HIV infection associated with an increased risk for cholera? Findings from a case–control study in Mozambique. *Tropical Medicine & International Health*. 2008; 13: 683–688. <https://doi.org/10.1111/j.1365-3156.2008.02051.x> PMID: 18331384
12. Zimmermann P, Curtis N. Factors That Influence the Immune Response to Vaccination. *Clinical Microbiology Reviews*. 2019; 32. <https://doi.org/10.1128/CMR.00084-18> PMID: 30867162
13. Penkert RR, Rowe HM, Surman SL, Sealy RE, Rosch J, Hurwitz JL. Influences of Vitamin A on Vaccine Immunogenicity and Efficacy. *Front Immunol*. 2019; 10. <https://doi.org/10.3389/fimmu.2019.01576> PMID: 31379816
14. Surman SL, Penkert RR, Jones BG, Sealy RE, Hurwitz JL. Vitamin Supplementation at the Time of Immunization with a Cold-Adapted Influenza Virus Vaccine Corrects Poor Mucosal Antibody Responses in Mice Deficient for Vitamins A and D. *Clin Vaccine Immunol*. 2016; 23: 219–227. <https://doi.org/10.1128/CVI.00739-15> PMID: 26740391
15. Rudraraju R, Jones BG, Surman SL, Sealy RE, Thomas PG, Hurwitz JL. Respiratory tract epithelial cells express retinaldehyde dehydrogenase ALDH1A and enhance IgA production by stimulated B cells in the presence of vitamin A. *PLoS ONE*. 2014; 9: e86554. <https://doi.org/10.1371/journal.pone.0086554> PMID: 24466150
16. Thurnham D, Smith E, Flora P. Concurrent liquid-chromatographic assay of retinol,  $\alpha$ -tocopherol,  $\beta$ -carotene,  $\alpha$ -carotene, lycopene, and  $\beta$ -cryptoxanthin in plasma, with tocopherol acetate as internal standard. *Clinical chemistry*. 1988; 34: 377–81. <https://doi.org/10.1093/clinchem/34.2.377> PMID: 3342512
17. Qadri F, Mohi G, Hossain J, Azim T, Khan AM, Salam MA, et al. Comparison of the vibriocidal antibody response in cholera due to *Vibrio cholerae* O139 Bengal with the response in cholera due to *Vibrio cholerae* O1. *Clinical and diagnostic laboratory immunology*. 1995; 2: 685–688. <https://doi.org/10.1128/cdli.2.6.685-688.1995> PMID: 8574829
18. Bwire G, Munier A, Ouedraogo I, Heyerdahl L, Komakech H, Kagirita A, et al. Epidemiology of cholera outbreaks and socio-economic characteristics of the communities in the fishing villages of Uganda: 2011–2015. *PLoS Neglected Tropical Diseases*. 2017; 11: e0005407. <https://doi.org/10.1371/journal.pntd.0005407> PMID: 28288154

19. Ivers LC, Charles RC, Hilaire IJ, Mayo-Smith LM, Teng JE, Jerome JG, et al. Immunogenicity of the Bivalent Oral Cholera Vaccine Shanchol in Haitian Adults With HIV Infection. *J Infect Dis.* 2015; 212: 779–783. <https://doi.org/10.1093/infdis/jiv108> PMID: 25722294
20. Fritzsche C, Bergmann L, Loebermann M, Glass A, Reisinger EC. Immune response to hepatitis A vaccine in patients with HIV. *Vaccine.* 2019; 37: 2278–2283. <https://doi.org/10.1016/j.vaccine.2019.02.064> PMID: 30890384
21. Khan AI, Chowdhury F, Harris JB, Larocque RC, Faruque ASG, Ryan ET, et al. Comparison of clinical features and immunological parameters of patients with dehydrating diarrhoea infected with Inaba or Ogawa serotypes of *Vibrio cholerae* O1. *Scand J Infect Dis.* 2010; 42: 48–56. <https://doi.org/10.3109/00365540903289688> PMID: 19883159
22. Seas C, Miranda J, Gil AI, Leon-Barua R, Patz J, Huq A, et al. New insights on the emergence of cholera in Latin America during 1991: the Peruvian experience. *The American Journal of Tropical Medicine and Hygiene.* 2000; 62: 513–517. <https://doi.org/10.4269/ajtmh.2000.62.513> PMID: 11220769
23. Garg P, Nandy RK, Chaudhury P, Chowdhury NR, De K, Ramamurthy T, et al. Emergence of *Vibrio cholerae* O1 Biotype El Tor Serotype Inaba from the Prevailing O1 Ogawa Serotype Strains in India. *J Clin Microbiol.* 2000; 38: 4249–4253. <https://doi.org/10.1128/JCM.38.11.4249-4253.2000> PMID: 11060101
24. Ciabattini A, Pettini E, Medagliani D. CD4+ T Cell Priming as Biomarker to Study Immune Response to Preventive Vaccines. *Front Immunol.* 2013; 4. <https://doi.org/10.3389/fimmu.2013.00421> PMID: 24363656
25. Albert MJ, Qadri F, Wahed MA, Ahmed T, Rahman ASMH, Ahmed F, et al. Supplementation with zinc, but not vitamin A, improves seroconversion to vibriocidal antibody in children given an oral cholera vaccine. *J Infect Dis.* 2003; 187: 909–913. <https://doi.org/10.1086/368132> PMID: 12660937
26. West KP. Vitamin A Deficiency Disorders in Children and Women. *Food Nutr Bull.* 2003; 24: S78–S90. <https://doi.org/10.1177/15648265030244S204> PMID: 17016949
27. Qadri F, Ahmed T, Wahed MA, Ahmed F, Bhuiyan NA, Rahman ASMH, et al. Suppressive effect of zinc on antibody response to cholera toxin in children given the killed, B subunit-whole cell, oral cholera vaccine. *Vaccine.* 2004; 22: 416–421. <https://doi.org/10.1016/j.vaccine.2003.07.005> PMID: 14670323
28. Sinha R, Howlader DR, Ta A, Mitra S, Das S, Koley H. Retinoic acid pre-treatment down regulates *V. cholerae* outer membrane vesicles induced acute inflammation and enhances mucosal immunity. *Vaccine.* 2017; 35: 3534–3547. <https://doi.org/10.1016/j.vaccine.2017.05.036> PMID: 28545924
29. Kozakova H, Hanson LA, Stepankova R, Kahu H, Dahlgren UI, Wiedermann U. Vitamin A deficiency leads to severe functional disturbance of the intestinal epithelium enzymes associated with diarrhoea and increased bacterial translocation in gnotobiotic rats. *Microbes and Infection.* 2003; 5: 405–411. [https://doi.org/10.1016/s1286-4579\(03\)00045-5](https://doi.org/10.1016/s1286-4579(03)00045-5) PMID: 12737996
30. Lima ABM de, Garcêz LS, Oliveira IKF, Santos MM dos, Da Paz SMRS, Paiva A de A, et al. Vitamin A deficiency and factors associated with retinol levels in public school students. *Revista de Nutrição.* 2017; 30: 605–614. <https://doi.org/10.1590/1678-98652017000500006>
31. Mwanza-Lisulo M, Kelly P. Potential for use of retinoic acid as an oral vaccine adjuvant. *Philos Trans R Soc Lond B Biol Sci.* 2015; 370. <https://doi.org/10.1098/rstb.2014.0145> PMID: 25964457
32. Taylor DN, Cárdenas V, Perez J, Puga R, Svennerholm AM. Safety, immunogenicity, and lot stability of the whole cell/recombinant B subunit (WC/rCTB) cholera vaccine in Peruvian adults and children. *The American Journal of Tropical Medicine and Hygiene.* 1999; 61: 869–873. <https://doi.org/10.4269/ajtmh.1999.61.869> PMID: 10674661
33. Kanungo S, Paisley A, Lopez AL, Bhattacharya M, Manna B, Kim DR, et al. Immune responses following one and two doses of the reformulated, bivalent, killed, whole-cell, oral cholera vaccine among adults and children in Kolkata, India: A randomized, placebo-controlled trial. *Vaccine.* 2009; 27: 6887–6893. <https://doi.org/10.1016/j.vaccine.2009.09.008> PMID: 19761838