



CASE REPORT

Open Access

Echocardiography of isolated subacute left heart tamponade in a patient with cor pulmonale and circumferential pericardial effusion

Tomaž Marš¹, Helena Mikolavčič², Barbara Salobir³, Matej Podbregar^{2*}

Abstract

Patients with advanced idiopathic pulmonary artery hypertension have often a chronic pericardial effusion. It is the result of increased transudation and impaired re-absorption due to elevated venous pressure. These patients have pre-existent symptoms and signs of chronic right heart failure. High degree of suspicion is required to detect of development of an atypical form of tamponade with isolated compression of left heart chambers as shown in present case report. Transthoracic echocardiography provides a rapid access to the correct diagnosis, a prompt relief of symptoms following the ultrasound guided pericardiocentesis and important diagnostic tool for regular follow up of patients thereafter as shown in our case report.

Background

Circumferential pericardial effusion typically results in biventricular tamponade and equalization of intracardiac and pericardial pressure during diastole. In classic subacute tamponade the rising of pericardial pressure causes a progressive collapse of right atrium and ventricle preventing venous return to the right atrium. Symptoms and signs referable to increased filling pressure and diminished cardiac output ensue. Patients presents with dyspnea, orthopnea, peripheral edema, fatigability, hepatic engorgement. The three principal features: jugular venous distention, soft or absent heart sounds and hypotension (Beck's triad), tachycardia and pulsus paradoxus are present. However, tamponade may involve the right or left heart. While isolated left ventricular tamponade can occur as a postoperative complication form localized posterior pericardial effusions or hematoma, circumferential pericardial effusions leading to left heart tamponade are rare [1].

Case presentation

We present a clinical course and echocardiographic examination in a patient with idiopathic pulmonary artery hypertension (IPAH) and chronic pericardial

effusion who developed a subacute isolated left heart tamponade.

A 57-year-old patient with IPAH and a previously known chronic pericardial effusion presented in an out-patient clinic with symptoms of dyspnea on exertion, in the last days even at rest, orthopnea and leg edema. During the past few months he was in good physical condition. He was on therapy with sildenafil, amlodipine, acenocoumarol and had a combination inhaler containing fluticasone propionate and salmeterol xinafoate. The clinical examination showed distended jugular veins, leg edema and an accentuated second heart sound. A chest radiogram showed an enlarged heart shadow. The echocardiographic examination showed a dilated right atrium and ventricle (Figure 1) with reduced ejection fraction, severe tricuspid insufficiency (Figure 2), systolic right ventricular pressure 82 mm Hg plus central venous pressure (Figure 3), the inferior vena cava larger than 2.5 cm with no respiration variability. Clinically estimated central venous pressure was approximately 20 mm Hg. Compared with the previous transthoracic echocardiographic examination, there was enlarged pericardial effusion, 3 cm behind the left ventricular posterior (Figure 4) wall with diastolic collapse of left atrium and ventricle (Additional files 1, 2 and 3).

After the normalisation of the haemostasis with fresh frozen plasma echocardiographically guided pericardiocentesis using a parasternal approach was performed.

* Correspondence: matej.podbregar@guest.arnes.si

²Clinical Department for Internal Intensive Care, University Medical Center Ljubljana, Slovenia




Figure 1 Transthoracic echocardiography from apical four chamber view shows enlarged right ventricle (RV) and right atrium (RA). There is pericardial effusion (PE) compressing left ventricle (LV) and left atrium (LA).




Figure 2 Transthoracic echocardiography from apical four chamber view shows massive tricuspid regurgitation (TR).




Figure 3 Right ventricular (RV) pressure was estimated by measuring tricuspid regurgitation (TR) velocity as stated in the Bernoulli equation: $RV\ pressure = 4 \times (tricuspid\ regurgitation\ velocity)^2 + central\ venous\ pressure$.




Figure 4 Transthoracic echocardiography from short-axis plane at the papillary muscle level shows enlarged right ventricle (RV) with paradoxical movement of intraventricular septum. Left ventricle is compressed by RV and pericardial effusion (PE).

Immediately after removal of 250 ml of yellow fluid the patient became normopnoic. There was still 1.5 cm large pericardial effusion behind the left ventricular posterior wall without left heart collapse (Additional file 4).

The pericardial fluid was a borderline transudate with slightly elevated proteins. Cytological studies for malignancy, microbiological microscopic examination and cultures (included *Mycobacterium tuberculosis*) were negative. Rheumatic markers (antinuclear antibodies, anti-double-strand DNA, anticardiolipin antibodies, lupus anticoagulants, rheumatoid factor), angiotensin-converting enzyme and tumor markers (tissue polypeptide antigen, neuron-specific enolase, alpha-fetoprotein, prostate-specific antigen, carcinoembryonic antigen) were all in normal range. The function of the thyroid gland was normal.

Discussion

Patients with cor pulmonale and circumferential pericardial effusion develop an atypical form of cardiac tamponade with isolated left heart compression. Pre-existing pulmonary arterial hypertension can modify the classic presentation. Symptoms and signs of right heart failure could already be present, so a high index of suspicion for tamponade is required in every worsening of right heart failure symptoms. When the pericardial pressure starts to increase in a patient with cor pulmonale, elevated pressure in right heart chambers prevent right atrial and ventricular compression, but while the pericardial pressure rises to the point to exceed left chambers pressure, this results first in diastolic collapse of left atrium and later on in left ventricle collapse due to a transient reversal of the transmural pressure [2,3]. Signs of impaired filling of left ventricle ensue leading to a drop in cardiac output.

The most probable mechanism of accumulation of pericardial fluid in patients with IPAH is transudation and impaired re-absorption of pericardial fluid due to elevated venous hydrostatic pressure in the setting of cor pulmonale.

In the setting of pulmonary arterial hypertension large hemodynamically significant pericardial effusions might be treated surgically and/or conservative and it is known that prognosis of patients with this complication is poor [4,5]. However, our patient has been stable throughout one year period after pericardiocentesis on his regular therapy after titration of diuretic furosemide (one tablets of 40 mg two to four times weekly) according to signs of right heart failure and measurements of NT-pro BNP (NT-proBNP before 600 ng/L, at the time of detection 1029 ng/L and one year after detection of circumferential pericardial effusion 601 ng/L). Additional regular repeated echocardiographic examinations were performed.

The right ventricle-to-right atrial pressure gradient may be difficult to estimate in the setting of severe tricuspid regurgitation, when there is a large color flow regurgitant jet. In this case, the peak velocity may not reflect the true pressure gradient.

In conclusion, patients with advanced IPAH have often a chronic pericardial effusion. It is the result of increased transudation and impaired re-absorption due to elevated venous pressure. These patients have pre-existent symptoms and signs of chronic right heart failure. High degree of suspicion is required to detect development of an atypical form of tamponade with isolated compression of left heart chambers. Transthoracic echocardiography provides a rapid access to the correct diagnosis, a prompt relief of symptoms following the ultrasound guided pericardiocentesis and important diagnostic tool for regular follow up of patients thereafter.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of written consent is available for review by the Editor-in-Chief of this journal.

Additional material

Additional file 1: Transthoracic echocardiography from apical four chamber view shows enlarged right ventricle (RV) and right atrium (RA). There is pericardial effusion (PE) compressing left ventricle (LV) and left atrium (LA).

Additional file 2: Transthoracic echocardiography from short-axis plane at the papillary muscle level shows enlarged right ventricle (RV) with paradoxical movement of intraventricular septum. Left ventricle is compressed by RV and pericardial effusion (PE).

Additional file 3: Transthoracic echocardiography from apical four chamber view shows enlarged right ventricle (RV) and tricuspid regurgitation (TR).

Additional file 4: Transthoracic echocardiography from apical four chamber view after pericardiocentesis. Right ventricle (RV), right atrium (RA), left ventricle (LV), left atrium (LA).

Author details

¹Institute of Pathophysiology, Faculty of Medicine, University of Ljubljana, Slovenia. ²Clinical Department for Internal Intensive Care, University Medical Center Ljubljana, Slovenia. ³Clinical Department for Pulmonary Disease and Allergy, University Medical Center Ljubljana, Slovenia.

Authors' contributions

TM: carried out interpretation and drafted the manuscript
HM: carried out interpretation and drafted the manuscript
BS: treated the patient, carried out interpretation and drafted the manuscript
MP: treated the patient, made acquisition of data, carried out interpretation and drafted the manuscript
All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 31 May 2010 Accepted: 14 July 2010 Published: 14 July 2010

References

1. Chuttani K, Pandian NG, Mohanty PK, Rosenfield K, Schwartz SL, Udelson JE, Simonetti J, Kusay BS, Caldeira ME: **Left ventricular diastolic collapse. An echocardiographic sign of regional cardiac tamponade.** *Circulation* 1991, **83**:1999-2006.
2. Frey MJ, Berko B, Palevsky H, Hirshfeld JW Jr, Herrmann HC: **Recognition of cardiac tamponade in the presence of severe pulmonary hypertension.** *Ann Intern Med* 1989, **111**:615-617.
3. Gollapudi RR, Yeager M, Johnson AD: **Left ventricular cardiac tamponade in the setting of cor pulmonale and circumferential pericardial effusion. Case report and review of the literature.** *Cardiol Rev* 2005, **13**:214-217.
4. Aqel RA, Aljaroudi W, Hage FG, Tallaj J, Raybum B, Nanda NC: **Left ventricular collapse secondary to pericardial effusion treated with pericardiocentesis and percutaneous pericardiectomy in severe pulmonary hypertension.** *Echocardiography* 2008, **6**:658-61.
5. Hemmes AR, Gaine SP, Wiener CM: **Poor outcomes associated with drainage of pericardial effusions in patients with pulmonary arterial hypertension.** *South Med* 2008, **5**:490-4.

doi:10.1186/1476-7120-8-27

Cite this article as: Marš *et al.*: Echocardiography of isolated subacute left heart tamponade in a patient with cor pulmonale and circumferential pericardial effusion. *Cardiovascular Ultrasound* 2010 **8**:27.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

