

ORIGINAL RESEARCH

Complete mitochondrial genome sequence of the Himalayan Griffon, *Gyps himalayensis* (Accipitriformes: Accipitridae): Sequence, structure, and phylogenetic analyses

Lichun Jiang^{1,2} | Liqing Peng² | Min Tang² | Zhangqiang You² | Min Zhang¹ | Andrea West³ | Qiping Ruan¹  | Wei Chen^{1,2}  | Juha Merilä⁴

¹Key Laboratory for Molecular Biology and Biopharmaceutics, School of Life Science and Technology, Mianyang Normal University, Mianyang, Sichuan, China

²Ecological Security and Protection Key Laboratory of Sichuan Province, Mianyang Normal University, Mianyang, Sichuan, China

³Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin University, Geelong, Victoria, Australia

⁴Ecological Genetics Research Unit, Organismal and Evolutionary Biology Research Programme, Faculty Biological & Environmental Sciences, University of Helsinki, Helsinki, Finland

Correspondence

Wei Chen, Ecological Security and Protection Key Laboratory of Sichuan Province, Mianyang Normal University, Mianyang, Sichuan 621006, China.
Email: wchen1949@163.com

Qiping Ruan, Key Laboratory for Molecular Biology and Biopharmaceutics, Mianyang Normal University, Mianyang, Sichuan 621006, China.
Email: qpruan20141230@163.com

Funding information

National Natural Science Foundation of China, Grant/Award Number: 31670392 and 31872216; Natural Sciences Foundation for Distinguished Young Scholar of Sichuan, Grant/Award Number: 2016JQ0038; Key Foundation of Sichuan Provincial Department of Education, Grant/Award Number: 18ZA0255; Technology Foundation for Selected Overseas Chinese Scholar; Ministry of Personnel of Sichuan Province; Research Project of Ecological Security and Protection Key Laboratory of Sichuan Province, Grant/Award Number: ESP1706; the Research Project of Education Office Project of Sichuan Province, Grant/Award Number: 18ZA0261; Scientific Research Foundation Projects of Mianyang Teacher's College, Grant/Award Number: QD2015A003; Scientific Research Fund of Mianyang Teacher's College, Grant/Award Number: PY-2016-A03, MYSY2017JC02 and Mnu-JY18285; Finnish Cultural Foundation; Academy of Finland, Grant/Award Number: 129662, 134728 and 218343

Abstract

This is the first study to describe the mitochondrial genome of the Himalayan Griffon, *Gyps himalayensis*, which is an Old World vulture belonging to the family Accipitridae and occurring along the Himalayas and the adjoining Tibetan Plateau. Its mitogenome is a closed circular molecule 17,381 bp in size containing 13 protein-coding genes, 22 tRNA coding genes, two rRNA-coding genes, a control region (CR), and an extra pseudo-control region (CCR) that are conserved in most Accipitridae mitogenomes. The overall base composition of the *G. himalayensis* mitogenome is 24.55% A, 29.49% T, 31.59% C, and 14.37% G, which is typical for bird mitochondrial genomes. The alignment of the Accipitridae species control regions showed high levels of genetic variation and abundant AT content. At the 5' end of the domain I region, a long continuous poly-C sequence was found. Two tandem repeats were found in the pseudo-control regions. Phylogenetic analysis with Bayesian inference and maximum likelihood based on 13 protein-coding genes indicated that the relationships at the family level were (Falconidae + (Cathartidae + (Sagittariidae + (Accipitridae + Pandionidae))). In the Accipitridae clade, *G. himalayensis* is more closely related to *Aegypius monachus* than to *Spilornis cheela*. The complete mitogenome of *G. himalayensis* provides a potentially useful resource for further exploration of the taxonomic status and phylogenetic history of *Gyps* species.

KEYWORDS

Accipitridae, genome organization, *Gyps himalayensis*, mitochondrial genome, phylogenetic status

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.

1 | INTRODUCTION

The Himalayan vulture (*Gyps himalayensis*) from the family Accipitridae has a wide distribution in Southeast Asia. In China it is found along the Himalayas and the adjoining Tibetan Plateau during the breeding season, and from Yunnan province during the winter season (BirdLife International, 2016; Lu, Ke, & Zeng, 2009). It inhabits mountainous areas at altitudes ranging from 1,200 to 6,000 m (Ferguson-Lees & Christie, 2001). The species' population trend is currently stable, but future declines are possible because vultures feed on livestock carcasses containing diclofenac, a nonsteroidal anti-inflammatory pharmaceutical known to cause reproductive failures in birds (Das, Cuthbert, Jakati, & Prakash, 2011; Oaks et al., 2004; Shultz et al., 2004; Swan et al., 2006).

There are currently eight recognized species in the genus *Gyps* including *G. bengalensis*, *G. indicus*, *G. tenuirostris*, and *G. himalayensis* from Asia, *G. africanus*, *G. coprotheres*, and *G. rueppellii* from Africa, and *G. fulvus* from Europe, Africa, and Asia (Ferguson-Lees & Christie, 2001; Pain et al., 2003). So far, the complete mitochondrial genomes of all species from this genus have not been amplified. In addition, the phylogenetic position of *Gyps* and related genus species has not been fully solved. Phylogenetic relationships in the family Accipitridae have traditionally been proven to be difficult to resolve based on morphological traits (Griffiths, 1994; Jollie, 1976). Due to its simple genomic organization, high rates of evolution and maternal inheritance mode, complete mitochondrial genomes have been proposed to show potential in resolving evolutionary history and taxonomic status (Arshad, Gonzalez, El-Sayed, Osborne & Wink, 2009; Boore, 2006; Downton, Castro, & Austin, 2002; Gibb, Kardailsky, Kimball, Braun, & Penny, 2007; Kan, Yang, et al., 2010; Ren et al., 2014). In the past 10 years, a number of studies performed using mitochondrial genome sequences to clarify avian phylogeny (Kan, Li, et al., 2010; Li, Liu, Zhou, & Gu, 2015; Li, Liu, et al., 2015; Song et al., 2015; Zhang et al., 2012).

The systematic taxonomic status among Falconiform taxa is rather complex and controversial (Jiang et al., 2015; Liu, Li, Du, & Liu, 2017). The family Accipitridae (hawks, eagles, and kites), with other four families (Pandionidae, Sagittariidae, Cathartidae, and Falconidae), was traditionally classified within the order Falconiformes (Cracraft, 1981; Haring, Kruckenhauser, Gamauf, Riesing, & Pinsker, 2001; Livezey & Zusi, 2007). However, according to more detailed morphological studies of several families, accipitrids and falconids are not closely in evolutionary relationship (Haring et al., 2001; Jollie, 1976). Based on his point of view, Falconiformes is polyphyletic, including New World vultures (Cathartidae), and the result is supported by studies of some behavioral traits and molecular data (Wink & Sauer-Gürth, 2004). Although the taxonomy of order Falconiformes is supported morphologically (Livezey & Zusi, 2001; Mayr & Clarke, 2003), recent studies have shown that they do not support close phylogenetic relationships between the Falconidae and other families included in the order Falconiformes (Burleigh, Kimball, & Braun, 2015; Hackett et al., 2008; Jarvis et al., 2014; Mahmood, Mclenachan, Gibb, & Penny, 2014). According to the viewpoint of Hackett et

al. (2008) and the South American Classification Committee, the Falconiformes is only included in the family Falconidae, and the remaining four families are placed in the order Accipitriformes. For a long time, the majority view has been to include accipitrids with the falcons in the Falconiformes, but some authorities have begun to recognize them as a separate group, Accipitriformes (Chesser et al., 2010). Additionally, a recent DNA study indicated that falcons not closely to the accipitrids, accordingly some authors propose that it should not be a part of Falconiformes (Chesser et al., 2012). Instead, it should be placed within Accipitriformes, and include most of the diurnal birds of prey: hawks, eagles, vultures, and many others (Chesser et al., 2012; Liu et al., 2017). However, its phylogenetic position of *Sagittarius serpentarius* is still confusing, and a lot of controversy about its validity is ongoing (Ferguson-Lees & Christie, 2001). Finally, some authors propose that *S. serpentarius* become an independent family, Sagittariidae (Wink & Sauer-Gürth, 2004).

In this study, the complete mitochondrial genome of *G. himalayensis* was sequenced and reported for the first time. Based on new data generated from *G. himalayensis* and obtaining complete mitogenome sequence data from GenBank, we tried to elucidate: (a) sequences, features, and structures of the mitogenomes of *G. himalayensis*, (b) taxonomic status phylogenetic relationships of *G. himalayensis* and related genus species, and (c) the phylogenetic relationships Falconidae, Accipitridae, Pandionidae, Sagittariidae, and Cathartidae.

2 | MATERIALS AND METHODS

2.1 | Sample collection and genomic DNA extraction

Blood of the Himalayan Griffon, *G. himalayensis* (Figure 1), was sampled from an adult vulture captured in the Dangxiong County (91°5'54.50"E, 30°28'50.79"N, 4,300 m.a.s.l), the Tibet Autonomous Region, China, in July 2016. The distribution of this




FIGURE 1 The Himalayan Griffon, *Gyps himalayensis*; photograph taken by Wei Chen



FIGURE 2 Species distribution map of the Himalayan Griffon, *Gyps himalayensis*

species is shown in Figure 2. Five milliliters of blood was sampled from the brachial vein using heparinized syringes and stored in liquid nitrogen in field and stored at -80°C in the laboratory. Total genomic DNA was extracted from blood tissue using Phenol: Chloroform: Isoamyl alcohol method (Sambrook & Russell, 2001). After a quality inspection, DNA was purified on 0.8% agarose gel. After measuring DNA concentration using a spectrophotometer, it was used for the PCR amplification.

2.2 | Mitochondrial DNA amplification and sequencing

The complete mitochondrial genome was amplified in 13 overlapping segments by PCR with Taq DNA Polymerase (TaKaRa), using 20 ng of genomic DNA from the sample as a template. Complete mtDNA was amplified as concatenated sequences using selectively amplified mtDNA template and 11 primer pairs from literature (Jiang, Wang, Peng, Peng, & Zou, 2014; Sorenson; , Ast, Dimcheff, Yuri, & Mindell, 1999; Zhao et al., 2012). Partial PCR primers were designed based on the conserved regions of the related species,

Aegypius monachus (NC_022957), *Buteo buteo* (NC_003128), and *Spilornis cheela* (NC_015887). The PCR amplifications were conducted in 25 μl reactions containing 1 \times PCR Buffer, 200 nM of each primer, 400 μM dNTP, 1 μl template (as per kit), and 1 U LA-Taq or Taq DNA Polymerase (TaKaRa). The PCR protocol was as follows: an initial denaturation at 95°C for 3 min, followed by 30 amplification cycles with a S1000TM thermal cycler at 94°C for 30 s, $53\text{--}65^{\circ}\text{C}$ for 30 s, and 72°C for 80–180 s, continued with a final extension at 72°C for 10 min. The PCR products were preserved at 12°C after the last cycle. Each amplicon was then purified using the Omega Cycle-Pure Kit (Omega Bio-Tek) and subjected to automated sequencing using an ABI 3730 sequencer, either directly or following sub-cloning into the pUC19 DNA vector (TaKaRa). To ensure high accuracy, each amplicon was sequenced twice independently with all the PCR primers.

2.3 | Sequence assembly, annotation, and analysis of the mitochondrial genome

The complete mitochondrial genome sequence of the Himalayan Griffon, *G. himalayensis*, was assembled using the SeqMan module

TABLE 1 Characteristics of the mitochondrial genome of *Gyps himalayensis*

Gene	Position		Sizes Nucleotide (bp)	Codon		Intergenic nucleotide ^b	Strand ^c	A + T%
	From	To		Start	Stop ^a			
tRNA-Phe	1	70	70				H	48.6
12S ribosomal RNA	71	1,055	985			-19	H	50.9
tRNA-Val	1,037	1,108	72			-8	H	55.6
16S ribosomal RNA	1,101	2,730	1,630			-20	H	53.4
tRNA-Leu	2,711	2,784	74			9	H	47.3
ND1	2,794	3,771	978	ATG	AGG	-2	H	53.2
tRNA-Ile	3,770	3,841	72			13	L	56.9
tRNA-Gln	3,855	3,925	71			-1	H	66.2
tRNA-Met	3,925	3,993	69				H	49.3
ND2	3,994	5,038	1,045	ATG	T--		H	51.9
tRNA-Trp	5,039	5,110	72			1	H	62.5
tRNA-Ala	5,112	5,180	69			2	L	56.5
tRNA-Asn	5,183	5,255	73			2	L	47.9
tRNA-Cys	5,258	5,324	67			-1	L	49.3
tRNA-Tyr	5,324	5,394	71			1	L	54.9
COXI	5,396	6,946	1,551	GTG	AGG	-9	H	52.5
tRNA-Ser	6,938	7,011	74			4	L	52.7
tRNA-Asp	7,016	7,084	69			2	H	59.4
COXII	7,087	7,770	684	ATG	TAA		H	53.2
tRNA-Lys	7,772	7,843	72			1	H	54.2
ATP8	7,844	8,008	165	ATG	TAA	-10	H	52.1
ATP6	7,999	8,682	684	ATG	TAA	-1	H	54.5
COXIII	8,682	9,464	783	ATG	TAA	1	H	52.6
tRNA-Gly	9,466	9,534	69				H	66.7
ND3	9,535	9,885	351	ATT	TAA	4	H	56.1
tRNA-Arg	9,890	9,958	69			1	H	60.9
ND4L	9,960	10,256	297	ATG	TAA	-7	H	54.2
ND4	10,250	11,626	1,377	ATG	TAA		H	51.2
tRNA-His	11,627	11,696	70			1	H	65.7
tRNA-Ser	11,698	11,762	65				H	52.7
tRNA-Leu	11,763	11,833	71				H	62.0
ND5	11,834	13,648	1,815	ATG	TAA	12	H	54.3
Cytb	13,661	14,803	1,143	ATG	TAA	2	H	52.1
tRNA-Thr	14,806	14,873	68				H	64.7
Control region	14,874	16,075	1,202			8	H	56.7
tRNA-Pro	16,084	16,153	70			22	L	62.9
ND6	16,176	16,694	519	ATT	TAG	-3	L	50.1
tRNA-Glu	16,692	16,762	71				L	62.0
Pseudo-control region	16,763	17,381	619				H	69.0

^aT--represents incomplete stop codons.

^bIntergenic bp indicates gap nucleotides (positive value) or overlapped nucleotides (negative value) between two adjacent genes.

^cH and L indicate genes transcribed on the heavy and light strands, respectively.

of the Lasergene 8.0 software (DNASTAR; Burland, 2000). Protein-coding genes (PCGs) were identified through sequence comparisons with the known sequences of other birds, using CLUSTAL W

program. Transfer RNA (tRNA) genes and their secondary structures were identified using tRNAscan-SE 1.21 (<http://lowelab.ucsc.edu/tRNAscan-SE>; Lowe & Eddy, 1997) or by their proposed secondary

structures (Kumazawa & Nishida, 1993) and anticodons. Ribosomal RNA genes (rRNAs) were identified by NCBI Internet BLAST search and comparisons. Codon usage and nucleotide composition statistics were computed using MEGA 6.06 (Tamura, Stecher, Peterson, Filipski, & Kumar, 2013). Composition skew analysis was carried out with the formulas $AT\text{-skew} = [A-T]/[A + T]$ and $GC\text{-skew} = [G-C]/[G + C]$, respectively (Perna & Kocher, 1995). The complete mitogenome of *G. himalayensis* was deposited in GenBank under the accession number KY594709.

2.4 | Phylogenetic analysis

The sequence data were initially compared using program ClustalX 1.83 (Thompson, Gibson, Plewniak, Jeanmougin, & Higgins, 1997) with default parameters. The mitogenome of *G. himalayensis* was analyzed in a phylogenetic tree using a total of 32 other order Falconiformes mitogenomes retrieved from NCBI GenBank (Table S1). The concatenated sequences of the 13 PCGs of complete mitochondrial genomes were used. On the basis of the best-fit model predicted by jModelTestv.0.1.1, the optimal nucleotide substitution model was selected using jModeltest v.0.1.1 (Posada, 2008) and the Akaike information criterion (Posada & Buckley, 2004). Maximum likelihood (ML) analysis was carried out with PhyML v.3.0 (Guindon & Gascuel, 2003). The confidence level (Felsenstein, 1985) at each

branch was evaluated by allowing four substitution rate categories and performing 1,000 bootstrap replicates. BI analysis was performed with MrBayes v.3.1.2 (Huelsenback & Ronquist, 2001). The best-fitting model of TVM+I+G (nst = 6; rates = gamma) was selected for subsequent Bayesian phylogenetic analyses. Bayesian posterior probabilities were estimated with the Markov Chain Monte Carlo sampling. The analysis was initiated with randomly generated trees and ran for 2×10^7 generations from which a total of 2×10^5 trees were sampled at the intervals of every 100 generations, and among the sampled trees, the top 25% were used as burn-in and discarded. Tracer v.1.4 (Rambaut & Drummond, 2007) was used to check that sampled values of log likelihood, plotted against generation time, reached stationarity. Finally, a majority rule consensus tree was generated and posterior probabilities were computed from the remaining postburn in trees.

3 | RESULTS AND DISCUSSION

3.1 | Genome content and organization

The complete mitogenome of *G. himalayensis* is a closed circular molecule 17,381 bp in length, similar to the other raptors (Li, Liu, et al., 2015; Li, Liu, et al., 2015; Qin, Guan, Shi, Hou, & Qin, 2013). It contains the typical set of 37 genes present in vertebrate mitogenomes, including 13 PCGs, 22 tRNA genes, two genes for ribosomal RNA subunits (12 s rRNA and 16 s rRNA), a control region (CR), and an extra pseudo-control region (CCR; Table 1, Figure 3). ND6 and eight tRNAs are transcribed from the light strand (L), however the other 12 PCGs, 14 tRNAs, two rRNAs, and two noncoding region (CR, CCR) genes are located on the heavy (H) strand. The gene arrangement pattern of the *G. himalayensis* mitochondrial genome is identical to that in genera *Aegypius*, *Buteo*, and *Falco*, all which have the remnant CCR gene order, but differ from the standard bird gene order (Barker, Benesh, Vandergon, & Lanyon, 2012; Gibb et al., 2007; Li, Huang, et al., 2015). The *G. himalayensis* mitogenome nucleotide composition is biased toward A and T (56.14%: A = 224.55%, T = 29.49%, G = 14.37% and C = 31.59%), which is similar to that seen in the other *Gyps* mitogenomes (52.71%–57.65%; Table S1; Haring et al., 2001). The A + T content of the control region was clearly higher than that of PCGs, tRNAs, and rRNAs (Table S1).

Mitochondrial genes overlapped by a total of 151 bp in 12 different locations from 1 to 70 bp, and the longest overlap (70 bp) existed between genes tRNA-Glu and pseudo-control region. The shortest overlap (1bp) existed in ATP6-COXIII, tRNA-Cys-tRNA-Tyr, and tRNA-Gln-tRNA-Met genes. In addition, some protein-coding genes shared 1–20 nucleotides in common with adjacent tRNA genes. This is a common feature of mitochondrial DNA which is very compact and economical (Cuore & Kocher, 1999). Furthermore, 17 intergenic spacers were present in the *G. himalayensis* mitogenome and involved a total of 85 bp. The longest spacer sequence was 22 nucleotides long and located between ND6 and tRNA-Pro (Table 1).




FIGURE 3 Complete mitochondrial genome organization and gene arrangement of *Gyps himalayensis*. Genes coded on the H-strand are indicated in the outer ring, while the genes coded on the L-strand are indicated in the inner ring. Genes are abbreviated as follows: ATP6 and ATP8 (subunits 6 and 8 of ATPase), COXI-COXIII (cytochrome c oxidase subunits 1–3), Cytb (cytochrome b), ND1-ND6 and ND4L (NADH dehydrogenase subunits 1–6 and 4L), 12S rRNA and 16S rRNA (ribosomal RNA of 12S and 16S), CR (control region); pCR (pseudo-coding region). One-letter amino acid abbreviations were used to label the corresponding tRNA genes

TABLE 2 AT/CG skews in the mitochondrial protein-coding genes (PCG), 2 rRNA genes, CR and the entire mitogenomes of Falconiformes birds, *Gyps himalayensis*, *Aegypius monachus*, *Falco peregrinus*, and *Micrastur gilvicollis*

Gene	AT-skew = A - T/A + T				GC-skew = G - C/G + C			
	<i>G. himalayensis</i>	<i>A. monachus</i>	<i>F. peregrinus</i>	<i>M. gilvicollis</i>	<i>G. himalayensis</i>	<i>A. monachus</i>	<i>F. peregrinus</i>	<i>M. gilvicollis</i>
ND1	0.000	0.006	0.023	0.016	-0.410	-0.417	-0.464	-0.446
ND2	0.162	0.144	0.126	0.153	-0.555	-0.549	-0.564	-0.558
ND3	-0.015	0.032	0.095	0.000	-0.455	-0.509	-0.516	-0.421
ND4	0.110	0.110	0.150	0.143	-0.545	-0.564	-0.570	-0.569
ND4L	-0.056	-0.012	0.031	0.051	-0.397	-0.467	-0.456	-0.447
ND5	0.145	0.152	0.206	0.174	-0.477	-0.508	-0.502	-0.524
ND6	-0.477	-0.496	-0.594	-0.591	0.552	0.580	0.595	0.624
COX1	0.032	0.054	0.062	0.019	-0.303	-0.309	-0.336	-0.304
COX2	0.154	0.163	0.142	0.105	-0.369	-0.375	-0.350	-0.350
COX3	0.000	0.022	0.104	0.058	-0.353	-0.367	-0.388	-0.398
ATP6	0.072	0.102	0.114	0.079	-0.569	-0.576	-0.599	-0.590
ATP8	0.186	0.209	0.283	0.196	-0.646	-0.707	-0.789	-0.775
Cytb	0.047	0.074	0.102	0.106	-0.463	-0.455	-0.458	-0.488
12S rRNA	0.206	0.202	0.244	0.195	-0.157	-0.343	-0.176	-0.147
16S rRNA	0.238	0.241	0.280	0.243	-0.170	-0.181	-0.209	-0.203
CR	-0.169	-0.182	0.027	/	-0.194	-0.141	-0.379	/
13PCG	0.054	0.066	0.087	0.067	-0.402	-0.414	-0.425	-0.422
overall	0.092	0.098	0.180	0.133	-0.375	-0.382	-0.396	-0.416

The bold values indicated significant differences between the 4 species

AT/CG skews were essentially similar in the mitochondrial protein-coding genes (PCG), 2 rRNA genes, CR and the mitogenomes of other Falconiformes birds, (e.g. *A. monachus*, *F. peregrinus*, and *Micrastur gilvicollis*), except for a few genes (ND3, ND4L and CR; Table 2). Moreover, *G. himalayensis* and *A. monachus* share negative AT-skew in ND4L and CR genes, whereas *F. peregrinus* and *M. gilvicollis* have positive AT-skew (Table 2). For ND3 and COX3 genes, *G. himalayensis* has negative AT-skew, while the other three species have positive AT-skews. Apparent positive AT-skew and negative GC-skew biases were identified for all 11 PCGs (except for ND3 and ND4L) encoded by the H-strand, whereas the reverse was detected in ND6 encoded by the L-strand, which has been observed also in previous studies (Table 2; Jiang et al., 2015; Liu et al., 2017).

3.2 | Protein-coding genes and codon usage patterns

The 13 mitochondrial PCGs of *G. himalayensis* were 11,393 bp in length, which accounted for 65.54% of the total mitogenome sequence. The PCG regions of the *G. himalayensis* mitogenome consisted of 3,797 codons in total. Most of the PCGs had ATG as start codon, except for COX1 and ND6, which had GTG and ATT start codons, respectively. This is the case in the other *Gyps* species as well (Table 1; Jiang et al., 2015; Liu et al., 2017). As in the

case of mitogenomes of other vertebrates, TAA was the most frequent stop codon in *G. himalayensis*. ND2 had incomplete stop codon consisting of T-, COX I, and ND1 ending with AGG and ending with TAG. These have been presumably added by post-transcriptional polyadenylation using poly-A tail (Ojala, Montoya, & Attardi, 1981). All other stop codons had the standard terminal codon (TAA). Among the 64 available codons, the most commonly used ones were isoleucine (L; AUU; 7.14%), phenylalanine (F; UUU; 4.99%), leucine (L; CUU; 4.49%), and leucine (L; CUA; 3.82%). The most frequently used amino acid in *G. himalayensis* was isoleucine (Table 3). In contrast, the codons ACG, CCG, UCG, CGG, AAG, GCG, and CAG were rarely used, accounting for only 1.38% of the amino acids. Moreover, just like in other vertebrates (Li, Liu, et al., 2015; Li, Huang, et al., 2015; Zhang, Nie, Wang, & Hu, 2009), a strong bias against G at the third codon position was found in all the 13 PCGs. The base composition of three codon positions of the 13 PCGs is shown in Table S2. The A + T content of the first (56.37%), the second (56.12%), and the third positions (56.41%). Hence, the third position has the highest A + T content, whereas the second position has the highest C + G content (43.88%; first position: 43.63%; third position: 43.59%). Moreover, the first positions have a positive AT- and negative GC-skews, but the second and third positions have negative AT- and GC-skews (Table S2). Among the 13 PCGs in the *G. himalayensis* mitogenome (Table 1),

TABLE 3 Codon usage in *Gyps himalayensis* mitochondrial protein-coding genes

Codon	Count	RSCU	%	Codon	Count	RSCU	%	Codon	Count	RSCU	%	Codon	Count	RSCU	%
UUU(F)	188	1.41	4.99	UCU(S)	59	1.22	1.57	UAU(Y)	59	1.13	1.57	UGU(C)	17	1.26	0.45
UUC(F)	79	0.59	2.1	UCC(S)	41	0.85	1.09	UAC(Y)	45	0.87	1.2	UGC(C)	10	0.74	0.27
UUA(L)	137	1.40	3.64	UCA(S)	137	2.83	3.64	UAA(*)	3	1.71	0.08	UGA(W)	95	1.76	2.52
UUG(L)	14	0.14	0.37	UCG(S)	8	0.17	0.21	UAG(*)	1	0.57	0.03	UGG(W)	13	0.24	0.35
CUU(L)	169	1.73	4.49	CCU(P)	20	0.40	0.53	CAU(H)	27	0.57	0.72	CGU(R)	13	0.70	0.35
CUC(L)	92	0.94	2.44	CCC(P)	54	1.08	1.43	CAC(H)	68	1.43	1.81	CGC(R)	19	1.03	0.5
CUA(L)	144	1.47	3.82	CCA(P)	117	2.34	3.11	CAA(Q)	79	1.80	2.1	CGA(R)	38	2.05	1.01
CUG(L)	31	0.32	0.82	CCG(P)	9	0.18	0.24	CAG(Q)	9	0.20	0.24	CGG(R)	4	0.22	0.11
AUU(I)	269	1.53	7.14	ACU(T)	69	0.95	1.83	AAU(N)	64	1.01	1.7	AGU(S)	21	0.43	0.56
AUC(I)	83	0.47	2.2	ACC(T)	86	1.19	2.28	AAC(N)	63	0.99	1.67	AGC(S)	24	0.50	0.64
AUA(M)	139	1.56	3.69	ACA(T)	127	1.75	3.37	AAA(K)	74	1.80	1.97	AGA(*)	1	0.57	0.03
AUG(M)	39	0.44	1.04	ACG(T)	8	0.11	0.21	AAG(K)	8	0.20	0.21	AGG(*)	2	1.14	0.05
GUU(V)	63	1.39	1.67	GCU(A)	73	0.91	1.94	GAU(D)	29	0.77	0.77	GGU(G)	45	0.79	1.2
GUC(V)	37	0.82	0.98	GCC(A)	147	1.84	3.9	GAC(D)	46	1.23	1.22	GGC(G)	69	1.21	1.83
GUA(V)	63	1.39	1.67	GCA(A)	94	1.18	2.5	GA(E)	64	1.52	1.7	GGA(G)	73	1.28	1.94
GUG(V)	18	0.40	0.48	GCG(A)	6	0.07	0.16	GAG(E)	20	0.48	0.53	GGG(G)	42	0.73	1.12




FIGURE 4 Putative tRNA secondary structures predicted from the 22 tRNA gene sequences found in the *Gyps himalayensis* mitogenome

TABLE 4 Sequence characteristics of pseudo-control regions (CCRs) in Falconiformes birds

Family	Genus	Scientific name	Accession number	CCR		Tandem repeats	Single repeat unit
				length (bp)	Types		
Accipitridae	<i>Gyps</i>	<i>Gyps himalayensis</i>	KY594709	689	2	12.8 × (11)	5'-TCTTTTTCAT
							3'-CCCTAAACAAGTAATAATAAGTAGATGAGCTATCTACAAAGC
Accipitridae	<i>Aegypius</i>	<i>Aegypius monachus</i>	KF682364	1,025	5	2.0 × (20)	5'-TAATCTTTCTTTCCACTTTT
							5'-CACTTTATTTT
Accipitridae	<i>Aegypius</i>	<i>Aegypius monachus</i>	KF682364	1,025	5	2.0 × (20)	5'-CACTTTATTTT
							5'-CAAGTCTACAAGCCCTAAACAAGTAATAATAAGCAGATGAG
Accipitridae	<i>Aegypius</i>	<i>Aegypius monachus</i>	KF682364	1,025	5	2.0 × (20)	5'-CAAGTCTACAAGCCCTAAACAAGTAATAATAAGCAGATGAG
							3'-CAAGTCTACAAGCCCTAAACAAGTAATAATAAGCAGATGAG
Accipitridae	<i>Aegypius</i>	<i>Aegypius monachus</i>	KF682364	1,025	5	2.0 × (20)	TAAACAAGTAATAATAATAAGCAGATGAGCTATCTACAAAGCCCTAAACAAGTAATAATAAGCAGATGAG
							ATAAGCAGATGAGCAAGTCTACAAGCCCTAAACAAGTAATAATAAGCAGATGAG
Accipitridae	<i>Aegypius</i>	<i>Aegypius monachus</i>	KF682364	1,025	5	2.0 × (20)	3'-CAAGTCTACAAGCCCTAAACAAGTAATAATAAGCAGATGAG
							CCCTAAACAAGTAATAATAAGCAGATGAGCTATCTACAAAGCCCTAAACAAGTAATAATAAGCAGATGAG
Accipitridae	<i>Aegypius</i>	<i>Aegypius monachus</i>	KF682364	1,025	5	2.0 × (20)	ATATATAAGCAGATGAGCAAGTCTACAAGCCCTAAACAAGTAATAATAAGCAGATGAG
							GCAAGTCTACAAGCCCTAAACAAGTAATAATAAGCAGATGAG
Accipitridae	<i>Spilornis</i>	<i>Spilornis cheela</i>	NC_015887	1532	3	7.6 × (10)	5'-TATGTATATA
							5'-TATATACATGTATATA
Accipitridae	<i>Spizaetus</i>	<i>Spizaetus alboniger</i>	AP008239	1,273	1	17.5 × (46)	3'-AACAGATGAGCAATCTACAAGCCCTAAACAAGTAATAATAAGCAGATGAG
							3'-TAACCAACAATAATAAGTAAATGATACGTTCTATAAACCGTCAATCAA
Accipitridae	<i>Accipiter</i>	<i>Accipiter virgatus</i>	KJ699124	1,092	4	3.5 × (11)	5'-CATCTTTATTA
							5'-ATAACTAGTAAACCCCGGCGCAGTGCATAATTTCAAAAATCATGAAATAT
Accipitridae	<i>Accipiter</i>	<i>Accipiter virgatus</i>	KJ699124	1,092	4	3.5 × (11)	3'-AAGCCGTGAAATAGCAACTACTATATAAAGAAATGAGCAGCTTTATAG
							5'-AGCAGTCTTATAGAAGCCCGTGAAT
Accipitridae	<i>Hieraetus</i>	<i>Hieraetus fasciatus</i>	KP329567	1799	1	18.4 × (49)	3'-CCAACAATAATAATAAGTAAATGATGCGCTAATAAAGCCGTGAGCAAGTAA
							5'-AGCAATACTACCAAGCCCTAAACAAGTAATAATAAGCAGATGAG
Accipitridae	<i>Buteo</i>	<i>Buteo buteo</i>	AF380305	1,455	1	23.2 × (48)	3'-AAACAACAATAATAATAAGTAAATGATGCGCTAATAAAGCCGTGAGCAAGTAA
							3'-AAACAACAATAATAATAAGTAAATGATGCGCTAATAAAGCCGTGAGCAAGTAA
Falconidae	<i>Falco</i>	<i>Falco peregrinus</i>	AF090338	950	2	28.9 × (27)	3'-ATAAACAACAATAATAATAAGTAAATGATGCGCTAATAAAGCCGTGAGCAAGTAA
							3'-ATAAACAACAATAATAATAAGTAAATGATGCGCTAATAAAGCCGTGAGCAAGTAA
Falconidae	<i>Falco</i>	<i>Falco tinnunculus</i>	NC_011307	796	3	36.2 × (18)	3'-ATAAACAACAATAATAATAAGTAAATGATGCGCTAATAAAGCCGTGAGCAAGTAA
							3'-ATAAACAACAATAATAATAAGTAAATGATGCGCTAATAAAGCCGTGAGCAAGTAA
Falconidae	<i>Micrastrur</i>	<i>Micrastrur gilvicolis</i>	NC_008548	1539	1	6.4 × (69)	5'-CAACAACAACAACAATAATAATAAGTAAATGATGCGCTAATAAAGCCGTGAGCAAGTAA
							3'-TAAACGAAA
Pandionidae	<i>Pandion</i>	<i>Pandion haliaetus</i>	KF961184	-	-	-	65.2 × (9)
							3'-TAAACGAAA
Sagittariidae	<i>Sagittarius</i>	<i>Sagittarius serpentarius</i>	NC_008550	-	-	-	3'-CATCACTTTTATCATGACAACAACAATAATAATAAGTAAATGATGCGCTAATAAAGCCGTGAGCAAGTAA
							ATCAAACTTTTTT

Notes: CCR = pseudo-control region; 7 (48) = 7 repeats of 48 bp (single repeat unit is 48-bp in length); 3' - = the repetitive region close to 3' region of CCR.




(a) CR and CCR in *G. Himalayensis* (KY594709)(b) CCR in *Aegyptus monachus* (KF682364) 1025 bpCCR in *Falco tinnunculus* (NC_011307) 796 bpCCR in *Spilornis cheela* (NC_015887) 1532 bpCCR in *Buteo buteo* (AF380305) 1455 bp

FIGURE 5 (a) Structure of CR and CCR in *Gyps himalayensis*. Positions of the conserved motifs in CR and the division into the three domains DI, DII and DIII are shown. (b) Structure of CCR in *Aegyptus monachus*, *Spilornis cheela*, *Buteo buteo*, and *Falco tinnunculus*. nr-CRRs are depicted as gray bars. Regions containing incomplete units are depicted as hatched bars. CCR, pseudo-control region; nr-CRR, nonrepetitive region in CCR; r-CRR, repetitive region in CCR

the ND3 has the highest (56.1%) and the ND6 the lowest A + T content (50.1%).

3.3 | Transfer and ribosomal RNA genes

In total, the typical 22 tRNA genes with conventional secondary structures were found from the mitogenome of *G. himalayensis*. The total length of mitochondrial tRNA genes was 1,548 bp and size of individual genes ranged from 65 bp (tRNA-Ser2) to 74 bp (tRNA-Leu1 and tRNA-Leu2). Most of the tRNAs could be folded into the canonical cloverleaf secondary structure, while the tRNA-Ser (S1) contains a predicted secondary structure with the T Ψ C arm and loop, but lacks the DHU arm and loop. This phenomenon is considered to be a typical feature of vertebrate mitogenomes (Lu, Lu, Li, & Jiang, 2016; Wolstenholme, 1992). Moreover, most of anticodons were identical to those observed in other *Gyps* species. The CCA 3'-terminal group was added during processing. A total of 24 unmatched base pairs were found in the tRNAs of *G. himalayensis*, and these contained G-U pairs in the 9 bp AA, 6 bp A-C, 5 bp DHU and 4 bp T Ψ C stems (Figure 4).

Stem mismatches appear to be ordinary in tRNA genes and possibly repaired through a post-transcriptional editing process (Lavrov, Brown, & Boore, 2000). Compared to other birds, most mismatched nucleotides were G-U pairs, which can form weak bonds in tRNAs and non-canonical pairs in tRNA secondary structures (Gutell, Lee, & Cannone, 2002; Li, Liu, et al., 2015; Li, Liu, et al., 2015).

The ribosomal genes (12S rRNA and 16S rRNA) were identified as to location, length, and base composition in the mitogenome of the *G. himalayensis*. The 12S rRNA gene is situated between tRNA-Phe and tRNA-Val, while the 16S rRNA gene is located between tRNA-Val and tRNA-Leu (Figure 3). The length of 12S rRNA and 16S rRNA were 985 bp and 1,630bp, respectively (Table 1). The base composition of the 12S rRNA was 20.2%T, 28.4%C, 30.7%A, and 20.7%G, with an A + T content of 50.9%, while the 16S ribosomal gene composition was 20.4%T, 27.2%C, 33.1%A, and 19.3%G, with an A + T content of 53.4%. The gene order and arrangement of two rRNA genes, including gene length, base composition, and RNA structure, were similar to those in other Accipitridae birds (Jiang et al., 2015; Li, Liu, et al., 2015; Li, Liu, et al., 2015; Song et al., 2015).




FIGURE 7 Results of phylogenetic analyses based on Bayesian inference (BI) and maximum likelihood (ML) analyses for 32 raptor taxa based on 13 PCGs sequences. *Phodilus badius* (KF961183) and *Strix leptogrammica* (KC953095) were used as outgroups. Tree topologies produced by BI and ML analyses were identical. Bayesian posterior probability and bootstrap support values from ML analyses, respectively, are shown on the nodes. The asterisks indicate new sequence generated in this study

Sorenson, & Dimcheff, 1998), 338 bp in *B. buteo* (Haring, Riesing, Pinsker, & Gamauf, 1999), 360 bp in *Haliaeetus albicilla* (Haring et al., 2001), and 472 bp in *Aquila chrysaetos* (Masuda et al., 1998). None of them had recognizable repeats from tRNA-Glu or ND6 genes, as was observed in *Amazona* parrots (Eberhard, Wright, & Bermingham, 2001). These variable tandem repeats have been identified as the main reason for the length variability of mitochondrial genome control regions, as well as for the whole mitogenome (Kim, Seong, Ho, & Ha, 1998; Zhang et al., 2016).

3.5 | Phylogenetic relationships

The nucleotide data for phylogenetic analyses included 11,364 nucleotide sites from 13 protein-encoding genes. Of these sites, 5,472 were conservative, 5,889 were variable, and 4,835 sites were parsimoniously informative. So far, based on a total of 33 other complete mitogenome sequences retrieved from NCBI GenBank (Table S1), BI and ML trees showed identical topology, and most internal nodes were well supported by posterior probabilities and bootstrap values, respectively (Figure 7). The results indicate that mitochondrial genome sequences of 33 Raptors could be divided into three branches: Accipitriformes, Cathartiformes, and Falconiformes (Figure 7). Accipitriformes and Cathartiformes are sister branches, which are then grouped with Falconiformes. Accipitriformes includes three subclades: Accipitridae,

Pandionidae, and Sagittariidae. The monophyly of Accipitridae was strongly supported (Figure 7). New World vultures (Cathartidae) have some similar characters in common with storks, and a close phylogenetic relationship with storks was proposed (Seibold & Helbig, 1995; Sibley & Monroe, 1990; Wink, 1995). However, recent molecular studies have shown that the New World vultures clearly have their affinity with many raptors and not with storks (Burleigh et al., 2015; Hackett et al., 2008; Wink & Sauer-Gürth, 2004). Our phylogeny indicated that *C. aura* (Cathartidae) was basal to the remaining Accipitriformes, which is consistent with Jiang et al.'s results (Jiang et al., 2015). According to a relatively short sequence, Wink and Sauer-Gürth (2004) placed *S. serpentarius* with storks. In our phylogenetic analysis, the *S. serpentarius* is deepest on the branch with *Pandion haliaetus* (Pandionidae) and Accipitridae (Figure 7). This result is consistent with some previous studies (Hackett et al., 2008; Lerner & Mindell, 2005; Mahmood et al., 2014). Furthermore, *P. haliaetus* is the sister to Accipitridae. The relationships among the four families of Accipitriformes are consistent with Burleigh et al.'s results (Burleigh et al., 2015; Jiang et al., 2015). In addition, the phylogenetic results indicate that there were four clades in Accipitridae. Clade 1 was (*Gys* + *Aegyptius* + *Spilornis*), Clade 2 was (*Aquila* + *Hieraaetus* + *Spizaetus*), Clade 3 was only *Accipiter*, and the Clade 4 was (*Buteo* + *Butastur*) (Figure 7).

Our results indicated that the relationships at the family level were (Falconidae + (Cathartidae + (Sagittariidae + (Accipitridae + P

andionidae))), and this inference was supported by both BI (posterior probabilities ≥ 0.89) and ML (bootstrap $\geq 85.3\%$) analyses (Figure 7). The Falconidae clade contained members from the genera *Falco*, *Phalcoboenus*, and *Micrastur*, and all the *Falco* species formed a monophyletic cluster (Figure 7). *Falco* and *Phalcoboenus* species were resolved as the sister group of the clade in genus *Micrastur* (1.00, 100%). In agreement with traditional taxonomy (Mahmood et al., 2014; Sibley, Ahlquist, & Monroe, 1988; Wang et al., 2016), *F. tinnunculus* and *F. naumanni* formed the earliest diverging lineage within the *Falco*. The Accipitridae included the genera *Gyps*, *Aegypius*, *Spilornis*, *Spizaetus*, *Aquila*, *Hieraetus*, *Butastur*, *Accipiter*, *Buteo*, and *Spizaetus*. In this clade, *S. cheela* was a taxon ancestral to *G. himalayensis* and *A. monachus*. Furthermore, the results of the phylogenetic analyses support a closer relationship between *G. himalayensis* and *A. monachus*, rather than between *G. himalayensis* and *S. cheela* (Figure 7). This is in accordance with the results of Jiang et al. (2014) and Wink (1995). *P. haliaetus* forms a sister group with the family Accipitridae clade. Clade Sagittaridae comprised of *S. serpentarius* as the sister taxon to the Accipitridae and Pandionidae (1.00, 98.5%). Also, this result was consistent with those of the previous studies (Lerner, Klaver, & Mindell, 2008; Liu et al., 2017; Song et al., 2015), but differed from those of Hackett et al. (2008) and Pacheco et al. (2011). The difference is mainly due to the different samples used and molecular markers, and their evolutionary relationships need further investigating and searching for more evidences from binding nuclear gene data and morphological characters. In the current study, all clades were well resolved, with only a few exceptions less than 80%, while Bayesian posterior probabilities were 1.00 (except for one node). Despite their fast evolutionary rates, mitogenomes contain species-specific evolutionary relationships, which can be efficiently recovered by improving taxon sampling (Rubinstein et al., 2013).

4 | CONCLUSIONS

In short, this is the first study to report and analyze the complete mitogenome of the *G. himalayensis*. The sequence structure of the *G. himalayensis* mitogenome is typical for birds and possesses high similarity to other reported Accipitridae mitogenomes. However, at the 5' end of the DI region, a long continuous poly-C sequence was discovered. Additionally, two types of tandem repeat units were also found in the pseudo-control region. The phylogenetic analyses support a closer relationship between *G. himalayensis* and *A. monachus* than that between *G. himalayensis* and *S. cheela*. Consequently, the results should provide a useful resource for future study of *Gyps* mitochondrial genomic evolution. However, the proposed evolutionary relationships among *G. himalayensis*, *A. monachus*, and *S. cheela* based on the findings that emerged in the current study should be accepted with caution due to limited taxon sampling. Many aspects of the phylogeny of the genus *Gyps* remain to be resolved and further analysis based on more molecular markers (binding nuclear gene) and a large number of taxon sampling is necessary to clarify the phylogenetic relationships among species of genus *Gyps*, *Aegypius*, and *Spilornis*.

ACKNOWLEDGMENTS

This research was supported by the National Natural Science Foundation of China (Nos. 31670392, 31872216), Natural Sciences Foundation for Distinguished Young Scholar of Sichuan (No. 2016JQ0038), Key Foundation of Sichuan Provincial Department of Education (No. 18ZA0255), Technology Foundation for Selected Overseas Chinese Scholar, Ministry of Personnel of Sichuan Province, the Research Project of Ecological Security and Protection Key Laboratory of Sichuan Province (No. ESP1706), the Research Project of Education Office Project of Sichuan Province (No. 18ZA0261), Scientific Research Foundation Projects of Mianyang Teacher's College (No. QD2015A003), the Scientific Research Fund of Mianyang Teacher's College (No. PY-2016-A03, MYSY2017JC02 and Mnu-JY18285) and the Finnish Cultural Foundation (grant to PM) and the Academy of Finland (grants 129662, 134728, and 218343 to JM). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest.

AUTHOR CONTRIBUTIONS

Jiang L and Ruan Q conceived and designed the experiments, performed the experiments, analyzed the data. Peng L, Tang M, and Zhang M performed field work and laboratory work. You Z and Zhang M analyzed the data. Jiang L, West A, Merilä J, and Chen W participated in writing the manuscript.

DATA ACCESSIBILITY

The following information was supplied regarding the deposition of DNA sequences: GenBank accession numbers: KY594709.

ORCID

Qiping Ruan  <https://orcid.org/0000-0002-0150-3000>

Wei Chen  <https://orcid.org/0000-0001-7437-3238>

REFERENCES

- Arshad, M., Gonzalez, J., El-Sayed, A. A., Osborne, T., & Wink, M. (2009). Phylogeny and phylogeography of critically endangered *Gyps* species based on nuclear and mitochondrial markers. *Journal of Ornithology*, 2, 419–430. <https://doi.org/10.1007/s10336-008-0359-x>
- Baker, A. J., & Marshall, H. D. (1997). *Mitochondrial control region sequences as tools for understanding evolution*. Avian Molecular Evolution & Systematics, 51–82. San Diego: Academic Press.
- Barker, F. K., Benesh, M. K., Vandergon, A. J., & Lanyon, S. (2012). Contrasting evolutionary dynamics and information content of the avian mitochondrial control region and ND2 gene. *PLoS ONE*, 7, e46403. <https://doi.org/10.1371/journal.pone.0046403>

- BirdLife International. (2016). *Gyps himalayensis*. The IUCN Red List of Threatened Species 2016: e.T22695215A93496722. Downloaded on 23 February 2017.
- Boore, J. L. (1999). Animal mitochondrial genomes. *Nucleic Acids Research*, 27, 1767–1780. <https://doi.org/10.1093/nar/27.8.1767>
- Boore, J. L. (2006). The use of genome-level characters for phylogenetic reconstruction. *Trends in Ecology & Evolution*, 21, 439–446. <https://doi.org/10.1016/j.tree.2006.05.009>
- Boore, J. L., Macey, J. R., & Medina, M. (2005). Sequencing and comparing whole mitochondrial genomes of animals. *Method. Enzymol.*, 395, 311–348.
- Brown, L. H., & Amadon, D. (1968). *Eagles, Hawks and Falcons of the world*. London, UK: Country Life Books.
- Burland, T. G. (2000). DNASTAR's Lasergene sequence analysis software. In S. Misener & S. A. Krawetz (Eds.), *Bioinformatics methods and protocols*. Methods in Molecular Biology™, vol 132. Totowa, NJ: Humana Press.
- Burleigh, J. G., Kimball, R. T., & Braun, E. L. (2015). Building the avian tree of life using a large-scale, sparse supermatrix. *Molecular Phylogenetics and Evolution*, 84, 53–63. <https://doi.org/10.1016/j.ympev.2014.12.003>
- Cadahía, L., Pinsker, W., Negro, J. J., Pavlicev, M., Urios, V., & Haring, E. (2009). Repeated sequence homogenization between the control and pseudo-control regions in the mitochondrial genomes of the subfamily aquilinae. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*, 312B, 171–185. <https://doi.org/10.1002/jez.b.21282>
- Chen, W., Jiang, L. C., You, Z. Q., & Chen, M. (2015). Breeding biology of the upland buzzard (*Buteo hemilasius*) on the Tibetan Plateau. *Journal of Raptor Research*, 49, 320–324.
- Chesser, R. T., Banks, R. C., Barker, F. K., Cicero, C., Dunn, J. L., Kratter, A. W., ... Winker, K. (2010). Fifty-first supplement to the American Ornithologists' union check-list of North American birds. *The Auk*, 127, 726–744.
- Chesser, R. T., Banks, R. C., Barker, F. K., Cicero, C., Dunn, J. L., Kratter, A. W., ... Winker, K. (2012). Fifty-third supplement to the American Ornithologists' union check-list of North American birds. *The Auk*, 29, 573–588.
- Cracraft, J. (1981). Toward a phylogenetic classification of the recent birds of the world (class Aves). *The Auk*, 98, 681–714.
- Curole, J. P., & Kocher, T. D. (1999). Mitogenomics: Digging deeper with complete mitochondrial genomes. *Trends in Ecology & Evolution*, 14, 394–398. [https://doi.org/10.1016/S0169-5347\(99\)01660-2](https://doi.org/10.1016/S0169-5347(99)01660-2)
- Das, D., Cuthbert, R. J., Jakati, R. D., & Prakash, V. (2011). Diclofenac is toxic to the Himalayan Vulture *Gyps himalayensis*. *Bird Conservation International*, 21, 72–75.
- Dowton, M., Castro, L. R., & Austin, A. D. (2002). Mitochondrial gene rearrangements as phylogenetic characters in the invertebrates: The examination of genome 'morphology'. *Invertebrate Systematics*, 16, 345–356. <https://doi.org/10.1071/ISO2003>
- Eberhard, J. R., Wright, T. F., & Bermingham, E. (2001). Duplication and concerted evolution of the mitochondrial control region in the parrot genus *Amazona*. *Molecular Biology and Evolution*, 18, 1330–1342. <https://doi.org/10.1093/oxfordjournals.molbev.a003917>
- Felsenstein, J. (1985). Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 39, 783–791. <https://doi.org/10.1111/j.1558-5646.1985.tb00420.x>
- Ferguson-Lees, I. J., & Christie, D. A. (2001). *Raptors of the world*. London, UK: Christopher Helm.
- Gibb, G. C., Kardailsky, O., Kimball, R. T., Braun, E. L., & Penny, D. (2007). Mitochondrial genomes and avian phylogeny: Complex characters and resolvability without explosive radiations. *Molecular Phylogenetics and Evolution*, 24, 269–280. <https://doi.org/10.1093/molbev/msl158>
- Griffiths, C. S. (1994). Monophyly of the falconiformes based on syringeal morphology. *The Auk*, 111, 787–805. <https://doi.org/10.2307/4088811>
- Guindon, S., & Gascuel, O. (2003). A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology*, 52, 696–704. <https://doi.org/10.1080/10635150390235520>
- Gutell, R. R., Lee, J. C., & Cannone, J. J. (2002). The accuracy of ribosomal RNA comparative structure models. *Current Opinion in Structural Biology*, 12, 301–310. [https://doi.org/10.1016/S0959-440X\(02\)00339-1](https://doi.org/10.1016/S0959-440X(02)00339-1)
- Hackett, S. J., Kimball, R. T., Reddy, S., Bowie, R. C., Braun, E. L., Braun, M. J., ... Yuri, T. (2008). A phylogenomic study of birds reveals their evolutionary history. *Science*, 320, 1763–1768. <https://doi.org/10.1126/science.1157704>
- Haring, E., Kruckenhauser, L., Gamauf, A., Riesing, M. J., & Pinsker, W. (2001). The complete sequence of the mitochondrial genome of *Buteo buteo* (Aves, Accipitridae) indicates an early split in the phylogeny of raptors. *Molecular Biology and Evolution*, 18, 1892–1904. <https://doi.org/10.1093/oxfordjournals.molbev.a003730>
- Haring, E., Riesing, M. J., Pinsker, W., & Gamauf, A. (1999). Evolution of a pseudo-control region in the mitochondrial genome of *Palaearctic buzzards* (genus *Buteo*). *Journal of Zoological Systematics and Evolutionary Research*, 37, 185–194. <https://doi.org/10.1111/j.1439-0469.1999.tb00982.x>
- Houston, D. C. (1983). The adaptive radiation of the Griffon Vultures. In S. R. Wilbur, & J. A. Jackson (Eds.), *Vulture biology and management* (pp. 360–363). Berkeley, CA: University of California Press.
- Huelsenback, J. P., & Ronquist, F. (2001). MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics*, 17, 754–755. <https://doi.org/10.1093/bioinformatics/17.8.754>
- Jarvis, E. D., Mirarab, S., Aberer, A. J., Li, B., Houde, P., Li, C., ... Zhang, G. (2014). Whole-genome analyses resolve early branches in the tree of life of modern birds. *Science*, 346, 1320–1331. <https://doi.org/10.1126/science.1253451>
- Jiang, L., Chen, J., Wang, P., Ren, Q., Yuan, J., Qian, C., ... Kan, X. (2015). The mitochondrial genomes of *Aquila fasciata* and *Buteo lagopus* (Aves, Accipitiformes): Sequence, structure and phylogenetic analyses. *PLoS ONE*, 10, e0141037.
- Jiang, L., Wang, G., Peng, R., Peng, Q., & Zou, F. (2014). Phylogenetic and molecular dating analysis of Taiwan blue pheasant (*Lophura swinhoii*). *Gene*, 539, 21–29. <https://doi.org/10.1016/j.gene.2014.01.067>
- Jollie, M. (1976). A contribution to the morphology and phylogeny of the Falconiformes. *Evolutionary Theory*, 1, 198–298.
- Kan, X. Z., Li, X. F., Lei, Z. P., Wang, M., Chen, L., Gao, H., & Yang, Z. Y. (2010). Complete mitochondrial genome of Cabot's tragopan, *Tragopan caboti* (Galliformes: Phasianidae). *Genetics and Molecular Research*, 9, 1204–1216. <https://doi.org/10.4238/vol9-2gmr820>
- Kan, X. Z., Yang, J. K., Li, X. F., Chen, L., Lei, Z. P., Wang, M., ... Yang, Z. Y. (2010). Phylogeny of major lineages of galliform birds (Aves: Galliformes) based on complete mitochondrial genomes. *Genetics and Molecular Research*, 9, 1625–1633. <https://doi.org/10.4238/vol9-3gmr898>
- Kim, K. S., Seong, E. L., Ho, W. J., & Ha, J. H. (1998). The complete nucleotide sequence of the domestic dog (*Canis familiaris*) mitochondrial genome. *Molecular Phylogenetics and Evolution*, 10, 210–220.
- Kumazawa, Y., & Nishida, M. (1993). Sequence evolution of mitochondrial tRNA genes and deep-branch animal phylogenetics. *Journal of Molecular Evolution*, 37, 380–398. <https://doi.org/10.1007/BF00178868>
- Lavrov, D. V., Brown, W. M., & Boore, J. L. (2000). A novel type of RNA editing occurs in the mitochondrial tRNAs of the centipede *Lithobius forficatus*. *Proceedings of the National Academy of Sciences of the United States of America*, 97, 13738–13742. <https://doi.org/10.1073/pnas.250402997>

- Lerner, H. R. L., Klaver, M. C., & Mindell, D. P. (2008). Molecular phylogenetics of the buteonine birds of prey (Accipitridae). *The Auk*, *125*, 304–315. <https://doi.org/10.1525/auk.2008.06161>
- Lerner, H. R., & Mindell, D. P. (2005). Phylogeny of eagles, Old World vultures, and other Accipitridae based on nuclear and mitochondrial DNA. *Molecular Phylogenetics and Evolution*, *37*, 327–346. <https://doi.org/10.1016/j.ympev.2005.04.010>
- Li, B., Liu, G., Zhou, L., & Gu, C. (2015). Complete mitochondrial genome of cinereous vulture *Aegypius monachus* (Falconiformes: Accipitridae). *Mitochondrial DNA*, *26*, 910–911.
- Li, X., Huang, Y., & Lei, F. (2015). Comparative mitochondrial genomics and phylogenetic relationships of the *Crossoptilon* species (Phasianidae, Galliformes). *BMC Genomics*, *16*, 42. <https://doi.org/10.1186/s12864-015-1234-9>
- Liu, G., Li, C., Du, Y., & Liu, X. (2017). The complete mitochondrial genome of a Japanese sparrow hawk (*Accipiter gularis*) and the phylogenetic relationships among some predatory birds. *Biochemical Systematics and Ecology*, *70*, 116–125. <https://doi.org/10.1016/j.bse.2016.11.007>
- Livezey, B. C., & Zusi, R. L. (2001). Higher-order phylogenetics of modern aves based on comparative anatomy. *Netherlands Journal of Zoology*, *51*, 179–205. <https://doi.org/10.1163/156854201X00260>
- Livezey, B. C., & Zusi, R. L. (2007). Higher-order phylogeny of modern birds (Theropoda, Aves: Neornithes) based on comparative anatomy. II. Analysis and discussion. *Zoological Journal of the Linnean Society*, *149*, 1–95. <https://doi.org/10.1111/j.1096-3642.2006.00293.x>
- Lowe, T., & Eddy, S. (1997). TRNAscan-SE: A program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Research*, *25*, 955–964. <https://doi.org/10.1093/nar/25.5.955>
- Lu, X., Ke, D. H., & Zeng, X. H. (2009). Status, ecology, and conservation of the Himalayan Griffon Gyps himalayensis (Aves, Accipitridae) in the Tibetan Plateau. *Ambio*, *38*, 166–173.
- Lu, J., Lu, J., Li, X. F., & Jiang, H. (2016). Complete mitochondrial genome of the saker falcon, *Falco cherrug* (Falco, Falconidae). *Mitochondrial DNA Part A*, *27*, 3226–3227.
- Mace, G., Masundire, H., & Baillie, J.E.M. (2005). Biodiversity. In R. Hassan, R. Scholes, & N. Ash (Eds.), *Ecosystems and human well-being: Current states and trends: findings of the condition and trends working group*. (pp. 77–122). The Millennium Ecosystem Assessment series. Washington, DC: Island Press.
- Mahmood, M. T., Mclenachan, P. A., Gibb, G. C., & Penny, D. (2014). Phylogenetic position of avian nocturnal and diurnal raptors. *Genome Biol. Evol.*, *6*, 326–332. <https://doi.org/10.1093/gbe/evu016>
- Masuda, R., Noro, M., Kurose, N., Nishida-Umehara, C., Takechi, H., Yamazaki, T., ... Yoshida, M. (1998). Genetic characteristics of endangered Japanese golden eagles (*Aquila chrysaetos japonica*) based on mitochondrial DNA D-loop sequences and karyotypes. *Zoo Biology*, *17*, 111–121. [https://doi.org/10.1002/\(SICI\)1098-2361\(1998\)17:2<111::AID-ZOO6>3.0.CO;2-C](https://doi.org/10.1002/(SICI)1098-2361(1998)17:2<111::AID-ZOO6>3.0.CO;2-C)
- Mayr, G., & Clarke, J. (2003). The deep divergences of neornithine birds: A phylogenetic analysis of morphological characters. *Cladistics*, *19*, 527–553. <https://doi.org/10.1111/j.1096-0031.2003.tb00387.x>
- Mendelssohn, H., & Leshem, Y. (1983). The status and conservation of vultures in Israel. In S. R. Wilbur, & J. A. Jackson (Eds.), *Vulture biology and management* (pp. 86–98). Berkeley, CA: University of California Press.
- Mindell, D. P., Sorenson, M. D., & Dimcheff, D. E. (1998). Multiple origin of mitochondrial gene order in birds. *Proceedings of the National Academy of Sciences of the United States of America*, *95*, 19693–19697.
- Oaks, J. L., Gilbert, M., Virani, M. Z., Watson, R. T., Meteyer, C. U., Rideout, B. A., ... Khan, A. A. (2004). Diclofenac residues as the cause of vulture population decline in Pakistan. *Nature*, *427*, 630–633. <https://doi.org/10.1038/nature02317>
- Ojala, D., Montoya, J., & Attardi, G. (1981). tRNA punctuation model of RNA processing in human mitochondria. *Nature*, *290*, 470–474. <https://doi.org/10.1038/290470a0>
- Pacheco, M. A., Battistuzzi, F. U., Lentino, M., Aguilar, R. F., Kumar, S., & Escalante, A. A. (2011). Evolution of modern birds revealed by mitogenomics: Timing the radiation and origin of major orders. *Molecular Biology and Evolution*, *28*, 1927. <https://doi.org/10.1093/molbev/msr014>
- Pain, D. J., Cunningham, A. A., Donald, P. F., Duckworth, J. W., Houston, D. C., Katzner, T., ... Timmins, R. (2003). Causes and effects of temporospatial declines of Gyps vulture in Asia. *Conservation Biology*, *17*, 661–671.
- Perna, N. T., & Kocher, T. D. (1995). Patterns of nucleotide composition of four fold degenerate sites of animal mitochondrial genomes. *Journal of Molecular Evolution*, *41*, 353–358.
- Posada, D. (2008). jModelTest: Phylogenetic model averaging. *Molecular Biology and Evolution*, *25*, 1253–1256. <https://doi.org/10.1093/molbev/msn083>
- Posada, D., & Buckley, T. R. (2004). Model selection and model averaging in phylogenetics: Advantages of Akaike information criterion and Bayesian approaches over likelihood ratio test. *Systematic Biology*, *53*, 793–808. <https://doi.org/10.1080/10635150490522304>
- Qin, X. M., Guan, Q. X., Shi, J. P., Hou, L. X., & Qin, P. S. (2013). Complete mitochondrial genome of the *Spilornis cheela* (Falconiformes, Accipitridae): Comparison of *S. cheela* and *Spizaetus alboniger*. *Mitochondrial DNA*, *24*, 255–256.
- Rambaut, A., & Drummond, A. J. (2007). Tracer v1.5. Retrieved from <https://the-tracer.en.softonic.com/>
- Ren, Q., Yuan, J., Ren, L., Zhang, L., Zhang, L., Jiang, L., ... Zhang, B. (2014). The complete mitochondrial genome of the yellow-browed bunting, *Emberiza chrysophrys* (Passeriformes: Emberizidae), and phylogenetic relationships within the genus *Emberiza*. *Journal of Genetics*, *93*, 699–707. <https://doi.org/10.1007/s12041-014-0428-2>
- Roques, S., Godoy, J. A., Negro, J. J., & Hiraldo, F. (2004). Organization and variation of the mitochondrial control region in two vulture species, *Gypaetus barbatus* and *Neophron percnopterus*. *Journal of Heredity*, *95*, 332–337. <https://doi.org/10.1093/jhered/esh047>
- Rubinstein, N. D., Feldstein, T., Shenkar, N., Botero-Castro, F., Griggio, F., Mastrototaro, F., ... Huchon, D. (2013). Deep sequencing of mixed total DNA without barcodes allows efficient assembly of highly plastic ascidian mitochondrial genomes. *Genome Biology and Evolution*, *5*, 1185–1199. <https://doi.org/10.1093/gbe/evt081>
- Ruokonen, M., & Kvist, L. (2002). Structure and evolution of the avian mitochondrial control region. *Molecular Phylogenetics and Evolution*, *23*, 422–432. [https://doi.org/10.1016/S1055-7903\(02\)00021-0](https://doi.org/10.1016/S1055-7903(02)00021-0)
- Saccone, C., Pesole, G., & Sbisà, E. (1991). The main regulatory region of mammalian mitochondrial DNA: structure-function model and evolutionary pattern. *Journal of Molecular Evolution*, *33*, 83–91.
- Sambrook, J., & Russell, D. (2001). *Molecular cloning: A laboratory manual* (3rd ed.). Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- Sbisà, E., Tanzariello, F., Reyes, A., Pesole, G., & Saccone, C. (1997). Mammalian mitochondrial D-loop region structural analysis: Identification of new conserved sequences and their functional and evolutionary implications. *Gene*, *205*, 125–140. [https://doi.org/10.1016/S0378-1119\(97\)00404-6](https://doi.org/10.1016/S0378-1119(97)00404-6)
- Seibold, I., & Helbig, A. J. (1995). Evolutionary history of New and Old World vultures inferred from nucleotide sequences of the mitochondrial cytochrome b gene. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *350*, 163–178.
- Shultz, S., Baral, H. S., Charman, S., Cunningham, A. A., Das, D., Ghalsasi, G. R., ... Prakash, V. (2004). Diclofenac poisoning is widespread in declining vulture populations across the Indian subcontinent. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *271*, S458–S460. <https://doi.org/10.1098/rsbl.2004.0223>
- Sibley, C. G., Ahlquist, J. E., & Monroe, B. L. (1988). A classification of the living birds of the world based on DNA-DNA hybridization studies. *The Auk*, *105*, 409–423.

- Sibley, C. G., & Monroe, B. L. (1990). *The Distribution and Taxonomy of the Birds of the World*. New Haven, CT: Yale University Press.
- Song, X., Huang, J., Yan, C., Xu, G., Zhang, X., & Yue, B. (2015). The complete mitochondrial genome of *Accipiter virgatus*, and evolutionary history of the pseudo-control regions in Falconiformes. *Biochemical Systematics and Ecology*, 58, 75–84. <https://doi.org/10.1016/j.bse.2014.10.013>
- Sorenson, M. D., Ast, J. C., Dimcheff, D. E., Yuri, T., & Mindell, D. P. (1999). Primers for a PCR-based approach to mitochondrial genome sequencing in birds and other vertebrates. *Molecular Phylogenetics and Evolution*, 12, 105–114. <https://doi.org/10.1006/mpev.1998.0602>
- Swan, G., Naidoo, V., Cuthbert, R., Green, R. E., Pain, D. J., Swarup, D., ... Wolter, K. (2006). Removing the treat of diclofenac to critically endangered Asian vultures. *PLoS Biology*, 4, e66.
- Taanman, J. W. (1999). The mitochondrial genome: Structure, transcription, translation and replication. *Biochimica Et Biophysica Acta (BBA) - Bioenergetics*, 1410, 103–123. [https://doi.org/10.1016/S0005-2728\(98\)00161-3](https://doi.org/10.1016/S0005-2728(98)00161-3)
- Tamura, K., Stecher, G., Peterson, D., Filipowski, A., & Kumar, S. (2013). MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Molecular Biology and Evolution*, 30, 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Thompson, J., Gibson, T., Plewniak, F., Jeanmougin, F., & Higgins, D. (1997). The CLUSTAL_X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research*, 25, 4876–4882. <https://doi.org/10.1093/nar/25.24.4876>
- Wang, H. W., Zhang, H. F., Ren, L., Xu, Y., Zeng, Y. J., Miao, Y. L., ... Wang, K. H. (2016). The whole mitochondrial genome of the lesser kestrel (*Falco naumanni*). *Mitochondrial DNA*, 27, 2385–2386.
- Wang, X., Sun, Y., Yuan, X. D., Tang, M. Q., Wang, L., Yu, Y. F., & Li, Q. W. (2004). Comparative study of mitochondrial tRNA gene sequence and secondary structure among fifteen predatory birds. *Acta Genetica Sinica*, 31, 411–419.
- Wink, M. (1995). Phylogeny of Old and New World vultures (Aves: Accipitridae and Cathartidae) inferred from nucleotide sequences of the mitochondrial cytochrome b gene. *Zeitschrift Für Naturforschung C*, 50, 868–882. <https://doi.org/10.1515/znc-1995-11-1220>
- Wink, M., & Sauer-Gürth, H. (2004). *Phylogenetic relationships in diurnal raptors based on nucleotide sequences of mitochondrial and nuclear marker genes* (pp. 483–498). Berlin, Germany: Raptors Worldwide WWGBP.
- Wolstenholme, D. R. (1992). Animal mitochondrial DNA: Structure and evolution. *International Review of Cytology*, 141, 173–216.
- Yamamoto, Y., Murata, K., Matsuda, H., Hosoda, T., Tamura, K., & Furuyama, J. (2000). Determination of the complete nucleotide sequence and haplotypes in the D-loop region of the mitochondrial genome in the oriental white stork, *Ciconia boyciana*. *Genes & Genetic Systems*, 75, 25–32. <https://doi.org/10.1266/ggs.75.25>
- Zhang, H., Dou, H., Yang, X., Zhao, C., Liu, G., & Zhang, J. (2016). The complete mitochondrial genome sequence of the Sparrowhawk (*Accipiter nisus*). *Mitochondrial DNA*, 27, 1648–1649.
- Zhang, J. F., Nie, L. W., Wang, Y., & Hu, L. L. (2009). The complete mitochondrial genome of the large-headed frog, *Limnonectes banaensis* (Amphibia: Anura), and a novel gene organization in the vertebrate mtdna. *Gene*, 442, 119–127. <https://doi.org/10.1016/j.gene.2009.04.018>
- Zhang, L., Wang, L., Gowda, V., Wang, M., Li, X., & Kan, X. (2012). The mitochondrial genome of the Cinnamon Bittern, *Ixobrychus cinnamomeus* (Pelecaniformes: Ardeidae): Sequence, structure and phylogenetic analysis. *Molecular Biology Reports*, 39, 8315–8326. <https://doi.org/10.1007/s11033-012-1681-1>
- Zhao, S., Ma, Y., Wang, G., Li, H., Liu, X., Yu, J., ... Zou, F. (2012). Molecular phylogeny of major lineages of the avian family Phasianidae inferred from complete mitochondrial genome sequences. *Journal of Natural History*, 46, 757–767. <https://doi.org/10.1080/00222933.2011.653588>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Jiang L, Peng L, Tang M, et al. Complete mitochondrial genome sequence of the Himalayan Griffon, *Gyps himalayensis* (Accipitriformes: Accipitridae): Sequence, structure, and phylogenetic analyses. *Ecol Evol*. 2019;9:8813–8828. <https://doi.org/10.1002/ece3.5433>