



Review

# Factors That Affect the Enlargement of Bacterial Protoplasts and Spheroplasts

Hiromi Nishida

Department of Biotechnology, Toyama Prefectural University, 5180 Kurokawa, Imizu, Toyama 939-0398, Japan; hnishida@pu-toyama.ac.jp; Tel.: +81-766-56-7500

Received: 7 August 2020; Accepted: 25 September 2020; Published: 27 September 2020



**Abstract:** Cell enlargement is essential for the microinjection of various substances into bacterial cells. The cell wall (peptidoglycan) inhibits cell enlargement. Thus, bacterial protoplasts/spheroplasts are used for enlargement because they lack cell wall. Though bacterial species that are capable of gene manipulation are limited, procedure for bacterial cell enlargement does not involve any gene manipulation technique. In order to prevent cell wall resynthesis during enlargement of protoplasts/spheroplasts, incubation media are supplemented with inhibitors of peptidoglycan biosynthesis such as penicillin. Moreover, metal ion composition in the incubation medium affects the properties of the plasma membrane. Therefore, in order to generate enlarged cells that are suitable for microinjection, metal ion composition in the medium should be considered. Experiment of bacterial protoplast or spheroplast enlargement is useful for studies on bacterial plasma membrane biosynthesis. In this paper, we have summarized the factors that influence bacterial cell enlargement.

**Keywords:** bacterial cell enlargement; microinjection; protoplasts; spheroplasts; metal ion composition; DNA replication; vacuole formation


## 1. Introduction

Bacteria usually grow asexually, inheriting their DNA by clonal production (cell division). Therefore, these clones have the same genetic information. If clonal production were the only form of inheritance in bacteria, bacteria would display limited genetic variation. However, bacteria are so diverse that more than 99% are thought to be unknown at the species level. The horizontal transfer of genetic information between heterologous bacteria greatly contributes to genetic variation [1–7].

Which genetic elements have been maintained or lost during bacterial evolution? Has the acceptance of genetic information occurred randomly? Thus far, these remain open questions in the field of molecular biology. This is because, by using gene manipulation technologies, experiments have been done in specific genes only or in a limited number of genes, making it impossible to investigate the system of bacterial genome evolution. In contrast, the genome manipulation technology that has been recently reported is based on the repetition of existing gene manipulation technologies [8–12]. Unfortunately, the number of bacterial species capable of gene manipulation, such as *Bacillus subtilis* and *Mycoplasma mycoides*, used in this kind of experiments, is still limited. In addition, these experiments are time-consuming, labor-intensive, and costly. Thus, it is difficult to perform them in regular laboratories.

For genomic sciences, changing from gene manipulation technologies to genome manipulation technology, in which a whole genome may be treated as if it were one gene, could be very useful. For example, for the introduction of *B. subtilis* genomic DNA into *Escherichia coli* cells at one time, to be able to examine how many genes are expressed from the heterologous genome in such a system. Currently, there is no way to introduce whole heterologous genomic DNA into bacterial host cells at once.

In this review, the bacterial cells lacking cell wall with an outer membrane and a plasma membrane are called spheroplasts, and those without an outer membrane are called protoplasts. Thus, protoplasts and spheroplasts are produced from Gram-positive and Gram-negative bacteria, respectively. Cell wall (peptidoglycan) biosynthesis plays a crucial role in bacterial cell shape maintenance. Bacterial cells cannot enlarge in the presence of an intact peptidoglycan sacculus. Thus, it is necessary to stop peptidoglycan biosynthesis in order to enlarge bacterial cells. On the other hand, cell wall is not essential for bacteria to survive because many bacteria can change to L-form bacteria that are capable of dividing, increasing the number of cells without cell wall. L-form bacteria have been detected and isolated from various environments and most of them are antibiotic-resistant [13–18]. The spheroplast incubation method is used to enlarge bacterial cells [19]. Bacterial protoplasts/spheroplasts are produced by lysing the cell wall with lysozyme or by penicillin [20]. In the spheroplast incubation method, protoplasts/spheroplasts are produced by lysozyme. Though bacterial protoplasts/spheroplasts cannot divide, they may enlarge under suitable culture conditions where cell wall synthesis is inhibited (Figure 1). Enlarged bacterial cells have already been used in patch clamp analyses [19,21–23], but only recently, a microinjection method has been established for these enlarged cells [24]. Noteworthy, different bacterial species have different patterns of cell enlargement.



**Figure 1.** Comparison between intact bacterial cell and enlarged protoplast.

Here, we summarize the factors that influence the enlargement of bacterial cells, based on available literature. In our laboratory, the following bacterial cells have been enlarged: *Bacillus subtilis*, *Deinococcus grandis*, *Enterobacter hormaechei*, *Enterococcus faecalis*, *Erythrobacter litoralis*, *Escherichia coli*, *Lelliottia amnigena*, and *Rhodospirillum rubrum*. The native forms of these 8 species are a few micrometers in diameter. Among them, *B. subtilis* and *E. faecalis* are Gram-positive. The other bacteria are Gram-negative. Commonality and diversity are observed in bacterial protoplast or spheroplast enlargement. We chose to work with protoplasts/spheroplasts as cells that could not divide, which differ from L-form bacteria. L-form bacteria have been detected from various environments because they can divide. In contrast, it has been exceedingly difficult to detect bacterial protoplasts/spheroplasts in

nature because they do not grow. More works are needed to elucidate functions for bacterial protoplast or spheroplast formation in nature.

## 2. Osmotic Pressure

The osmotic pressure of incubation media plays an important role in the maintenance of bacterial protoplasts/spheroplasts [25]. They do not enlarge when under an osmotic pressure higher than the suitable one [26], whereas their plasma membranes break when the osmotic pressure is below the optimum. For example, in spheroplasts of the Gram-negative radiation-resistant bacterium *Deinococcus grandis*, the plasma membrane breaks under a low osmotic pressure, whereas the outer membrane does not [26]. Thus, *D. grandis* cells seem to be maintained under low osmotic pressure; however, plasma membrane-broken cells are dead and cannot enlarge [26]. In *D. grandis* spheroplasts, the outer membrane has a higher osmotic pressure resistance than the inner (plasma) membrane. This may be related to the fact that both the cell wall and the outer membrane have an important role in cell shape maintenance [27].

We used Difco Marine Broth (5 g/L peptone, 1 g/L yeast extract, 0.1 g/L ferric citrate, 19.45 g/L NaCl, 5.9 g/L MgCl<sub>2</sub>, 3.24 g/L MgSO<sub>4</sub>, 1.8 g/L CaCl<sub>2</sub>, 0.55 g/L KCl, 0.16 g/L NaHCO<sub>3</sub>, 0.08 g/L KBr, 34 mg/L SrCl<sub>2</sub>, 22 mg/L H<sub>3</sub>BO<sub>3</sub>, 8 mg/L Na<sub>2</sub>HPO<sub>4</sub>, 4 mg/L Na<sub>2</sub>SiO<sub>3</sub>, 2.4 mg/L NaF, and 1.6 mg/L NH<sub>4</sub>NO<sub>3</sub>; BD, Franklin Lakes, NJ) containing penicillin (DMBp) as an incubation medium for cell enlargement of both marine and non-marine bacteria. We chose this medium because it has a higher osmotic pressure than other media and because most bacterial protoplasts/spheroplasts live stably under seawater osmotic conditions [24,26,28].

During cell enlargement, the lipid composition of the plasma membrane changes [28], indicating that the properties of the membrane differ between bacterial cells capable of dividing and enlarged protoplasts/spheroplasts. Bacterial protoplast or spheroplast enlargement is not accompanied by swelling due to water invasion into the cells, but it is accompanied by plasma (and outer) membrane biosynthesis [24,28]. More studies are needed to elucidate whether osmotic pressure affects lipid composition. When using metal salts or sugars as osmotic stabilizers, it should be noted that the function is often not limited to osmotic pressure adjustment (see below).

## 3. Metal Ions

Tryptone glucose yeast extract (TGY: 5 g/L tryptone, 1 g/L glucose, and 3.0 g/L yeast extract) medium is commonly used for culturing *Deinococcus*. *D. grandis* spheroplast enlargement was not observed in TGY plus penicillin medium containing sucrose as an osmotic stabilizer, but it was observed when the medium was supplemented with metal salts instead [28]. In order to investigate which metal salt plays an important role in spheroplast enlargement, we compared the cell morphology among *D. grandis* spheroplasts in incubation media with different metal salt compositions. We determined that either calcium or magnesium ions were essential for *D. grandis* spheroplast enlargement [28].

It is known that calcium and magnesium ions bind lipopolysaccharides (LPS) on the outer membrane of Gram-negative bacteria to maintain the cell surface structure [29–31]. However, *Deinococcus* has a unique lipid composition [32,33] and no LPS [34], suggesting that *D. grandis* may have another component that binds calcium/magnesium ions on its cell surface. In *E. coli*, calcium and magnesium ions play a role in lipid biosynthesis [35,36]. It suggests that calcium/magnesium ions may also play a role in lipid biosynthesis in *D. grandis*.

Interestingly, outer membrane fusion occurs when incubation is done in a medium containing calcium ions [26,37], but not when incubated in a medium containing magnesium ions. Outer membrane fusion was observed at 200 mM of calcium ions, but was not observed at 100 mM of both calcium and magnesium ions as well as at 200 mM of magnesium ions [26]. These results indicate that calcium and magnesium ions do not have the same function in the enlargement of the outer membrane of *D. grandis* [26,37].

During *D. grandis* spheroplast enlargement, the synthesis of the outer membrane is much faster than that of the inner membrane in the presence of calcium ions [26,28,37]. This strongly suggests that the outer membrane has a calcium ion-binding component associated with its expansion and fusion.

*Deinococcus* has surface (S)-layer proteins in its cell surface [38,39]. S-layer is two-dimensional protein array that are frequently found on the bacterial and archaeal surfaces [40]. *D. radiodurans* has two major S-layer proteins, Hpi and SlpA [39]. Hpi locates on outer membrane, and SlpA binds Hpi on one side and peptidoglycan on the other side [39]. Thus, we constructed an S-layer protein coding gene (*slpA*) deletion mutant of *D. grandis*. These mutant's spheroplasts enlarged as those from the wild type strain (unpublished data), suggesting that the SlpA homologous protein may not be the calcium ion-binding component that we were searching for.

*D. grandis* is rod-shaped and has a *rodZ* gene, a homolog to that of *E. coli* [41]. *E. coli* RodZ is a transmembrane protein that binds MreB, a cell shape-determining protein, inside the plasma membrane [42–45]. A *D. grandis rodZ* deletion mutant has a higher sensitivity to calcium ion than wild type strains [46]. Thus, at low calcium ion concentrations, where wild type spheroplasts cannot enlarge, *rodZ*-deletion mutants can [46]. The mechanism underlying increased sensitivity for calcium ion by the *rodZ*-deletion spheroplasts of *D. grandis* needs further investigation. However, this result indicates that calcium ion sensitivity is associated with spheroplast enlargement. Interestingly, an *mreB* deletion mutant in the Gram-positive bacterium *B. subtilis* requires magnesium ion for normal growth [47]. These findings suggest that the MreB-RodZ system for bacterial cell shape maintenance may be associated with calcium or magnesium ions.

Enlarged protoplasts of the Gram-positive lactic acid bacterium *Enterococcus faecalis* incubated in DMBp are useful for microinjection [24]. However, for the Gram-negative bacterium *Lelliottia amnigena* spheroplasts, the success rate of microinjection is very low when incubation is done in DMBp [24]. However, after cultivating these spheroplasts in media supplemented with different metal salt compositions, it was determined that the optimum metal salt composition for their enlargement was 62 mM CaCl<sub>2</sub>, 7.4 mM KCl, and 16.2 mM MgCl<sub>2</sub> [24]. This metal salt composition was more suitable for enlargement than the composition of 16.2 mM CaCl<sub>2</sub>, 7.4 mM KCl, and 62 mM MgCl<sub>2</sub> [24], which is the same concentration of CaCl<sub>2</sub>, KCl, and MgCl<sub>2</sub> in DMB. This indicates that the plasma membrane properties depend on the metal salt composition in the incubation medium.

The response to metal ions differs among different bacterial species. For example, the magnesium ion influx-related gene *corA* is upregulated in enlarged spheroplasts of the photosynthetic bacterium *Erythrobacter litoralis*, but it is downregulated in *L. amnigena* enlarged spheroplasts [48]. In contrast, the magnesium ion efflux-related gene *corD* is downregulated in *E. litoralis*, but upregulated in *L. amnigena* [48]. This suggests that *E. litoralis* enlarged spheroplasts need magnesium ions for their maintenance, but *L. amnigena* spheroplasts do not. Transcriptome and proteome analyses are useful for studies on the relationship between metal ions and cell enlargement. Based on our previous studies, it appears that there is no metal ion with common functions in all bacterial protoplasts/spheroplasts.

#### 4. Photosynthesis and Respiration

The aerobic anoxygenic photosynthetic bacterium *E. litoralis* was used to investigate how photosynthesis and respiration were involved in spheroplast enlargement. *E. litoralis* spheroplasts enlarge in DMBp under dark conditions in the presence of oxygen, indicating that their respiration mechanisms are active during cell enlargement [49,50]. In addition, they enlarge under light/dark conditions in the absence of oxygen, indicating that their photosynthesis is also active during cell enlargement [50]. *Erythrobacter* synthesizes ATP through photosynthesis under aerobic conditions, but not under anaerobic conditions [51]. Thus, spheroplasts have different photosynthetic characteristics than intact cells. They do not enlarge when cultured under anaerobic and dark conditions, indicating that either respiration or photosynthesis is essential for cell enlargement [50].

Continuous light or blue light inhibits *E. litoralis* spheroplast enlargement [49,52]. Considering that the photosynthetic apparatus is generally synthesized in the dark [53,54], the enlarged spheroplasts of

*E. litoralis* may also generate their photosynthetic apparatuses in the dark. It is known that blue light represses photosynthesis gene expression in the photosynthetic bacterium *Rhodobacter sphaeroides* [55]. Thus, blue light may also inhibit photosynthesis gene expression in *E. litoralis* spheroplasts [56]. *E. litoralis* spheroplasts do not enlarge under blue light but enlarge under green and red lights, indicating that they respond to different light signals [52]. These results support the notion that *E. litoralis* spheroplasts have an active photosynthetic apparatus.

Not only *E. litoralis* spheroplasts but also other bacterial protoplasts/spheroplasts enlarge in the presence of oxygen. On the other hand, aerobic culture conditions inhibit L-form bacterial growth [57,58]. Thus, the response to oxygen differs between bacterial protoplasts/spheroplasts and L-form cells. Although L-form bacteria also lack cell walls, their cells can divide [15,58]. Further investigation is needed to elucidate how oxygen affects cell division and cell enlargement of cell wall-lacking bacteria.

## 5. Chromosomal DNA (Replication)

Gene expression patterns differ between cells with normal division and enlarged bacterial protoplasts/spheroplasts [48,59]. In addition, both transcription and translation are essential for bacterial protoplast or spheroplast enlargement because inhibitors of transcription and translation efficiently prevent *D. grandis* spheroplast enlargement [28]; however, it is uncertain whether DNA replication is also essential for this enlargement. DNA replication occurs generally before cell division in all living organisms. Bacteria have several checkpoints along the cell division cycle [60–68]. Bacterial protoplasts/spheroplasts cannot divide but are enlarged in the presence of inhibitors of peptidoglycan synthesis [69]. To recover to their native forms, cell wall resynthesis is necessary [70,71]. Although bacterial protoplasts/spheroplasts cannot divide, DNA replication occurs during cell enlargement [19,21,49,72]. In the presence of the peptidoglycan biosynthetic inhibitor penicillin, *E. coli* forms a bulge at the site where the new cell wall is normally formed, which requires DNA replication [73]. These results indicate that DNA replication affects plasma membrane synthesis.

In the protoplasts of *E. faecalis*, the amount of DNA increases during the protoplast enlargement between 0 and 120 h of incubation in DMBp [74]. Both cell enlargement and DNA replication stop at 120 h of incubation, with no further change in cell size and DNA amount [74]. In order to investigate whether DNA replication influences bacterial protoplast or spheroplast enlargement, the DNA replication inhibitor novobiocin [75,76] was added to the incubation medium during the enlargement of *E. faecalis* protoplasts [74,77]. As a result, novobiocin inhibited not only DNA replication but also cell enlargement [74,77]. Although mitomycin C degrades chromosomal DNA of *E. faecalis* protoplasts, novobiocin does not [74]. When novobiocin was removed from the medium, *E. faecalis* protoplasts enlarged again [74,77]. Thus, DNA replication is associated with the biosynthesis of the plasma membrane. Bacterial protoplast/spheroplast enlargement can be controlled using novobiocin.


Bacterial chromosomal DNA attaches to the plasma membrane for an accurate distribution during cell division [78–83]. However, it is uncertain whether chromosomal DNA attaches to the plasma membrane of enlarged protoplasts/spheroplasts, whether this attachment is required for enlargement; and whether the DNA replication system is associated with plasma membrane biosynthesis in the cytoplasm. Therefore, further studies are needed to further understand these matters.

Interestingly, bacteria with large cell sizes have multiple copies of the genome in a single cell [84–86]. In *B. subtilis* and the large bacterium *Epulopiscium*, cell size and copy number of the genome have been shown to be positively correlated [84,87].

## 6. Vacuole Formation

In this review, “vacuole” is used for “provacuole” or “vacuole-like structure” in previous studies [19,21]. The size of *E. litoralis* enlarged spheroplasts is limited to a diameter of 6–7  $\mu\text{m}$ ; characteristically, they lack vacuoles [48,49]. In contrast, other bacterial protoplasts/spheroplasts exceed 15  $\mu\text{m}$  in diameter and have vacuoles in their cytoplasm (Figure 1) [19,21,24]. Thus, vacuole generation and growth occur in the cytoplasm after plasma membrane expansion. If these vacuoles were not

generated, cell enlargement would stop. Noteworthy, vacuole characteristics differ among bacterial species. For example, among the bacterial protoplasts/spheroplasts used in our laboratory, those from *L. amnigena* produce more vacuoles than any other, whereas those from *E. faecalis* produce the largest (Figure 2) [24]. It is known that the biosynthesis of the vacuole membrane continues during cell enlargement [24]. Thus, vacuole generation and expansion may be essential for bacterial protoplast or spheroplast enlargement over 15  $\mu\text{m}$  in diameter. In other words, plasma membrane and vacuole membrane syntheses may be interrelated.



**Figure 2.** Transmission electron microscopy images of Gram-positive *Enterococcus faecalis* and Gram-negative *Lelliottia amnigena*.

Transmission electron microscopy images show that enlarged *E. faecalis* protoplasts bear discoidal structures in the cytoplasm; structures that have not been observed in any other bacterial protoplast/spheroplast (Figure 2) [24]. These structures contain no DNA or vacuoles [24]. Thus, more studies are needed to elucidate whether they are associated with vacuoles in *E. faecalis* protoplasts.

Mature vacuoles are not connected to the plasma membrane [24]. However, the vacuolar membrane has proteins that are similar to those of the plasma membrane, such as respiratory chain proteins,  $F_0F_1$ -ATPase, and penicillin binding proteins [19,21,24]. This indicates that the vacuolar membrane components are the same or, at least, they are very similar to those of the plasma membrane. In addition, vacuoles of *E. coli* and *B. subtilis* enlarged protoplasts/spheroplasts have everted membranes, strongly suggesting that the vacuolar membrane may be generated by endocytosis of the plasma membrane [19,21].

After the DNA replication inhibitor novobiocin is introduced prior to vacuole formation during the enlargement of the *E. faecalis* protoplasts, the cell size is limited to 6  $\mu\text{m}$  in diameter and the cells lack vacuoles [74]. It indicates that plasma membrane biosynthesis and vacuole formation require DNA replication in *E. faecalis* protoplasts.

Interestingly, bacteria with large cell sizes also generate vacuoles in their cytoplasm [86,88,89]. For example, the largest bacterium known, *Thiomargarita namibiensis*, has a large vacuole [88,89] in which nitrate accumulates. Given that sulfur is present in the cytoplasm, away from nitrate, it is

thought that the *T. namibiensis* vacuole has a different function than that of vacuoles from enlarged protoplasts/spheroplasts.

## 7. Conclusions

Bacterial protoplasts/spheroplasts enlarge in the incubation medium containing an inhibitor of peptidoglycan synthesis under osmotically protective conditions.

Plasma membrane biosynthesis plays a central role in bacterial protoplast or spheroplast enlargement. Metal ions, especially calcium and magnesium ions, affect the properties of plasma membranes of bacterial protoplasts/spheroplasts such as flexibility suggesting that metal ion composition in the incubation medium may influence the final lipid composition of the plasma membrane of these cells. In order to generate bacterial protoplasts/spheroplasts suitable for microinjection, it is important to consider the metal salt composition in the incubation medium.

Plasma membrane biosynthesis (cell enlargement) requires ATP, which may be generated either during respiration or photosynthesis. Thus, it is important that respiration or photosynthesis systems are maintained in enlarged plasma membranes of bacterial protoplasts/spheroplasts. Plasma membrane expansion induces an increase in cytoplasmic volume. Thus, as protoplasts/spheroplasts become larger, ATP production in the plasma membrane alone becomes insufficient for maintaining the cytoplasm.

Besides, during plasma membrane expansion, DNA replication occurs, and vacuoles are generated and enlarged in the cytoplasm. Generation and enlargement of vacuoles in the cytoplasm repress cytoplasmic volume increase. In addition, an increase in chromosomal copy number may activate transcription in the cytoplasm. Thus, plasma membrane biosynthesis in bacterial protoplasts/spheroplasts is associated with DNA replication and vacuole formation. More work is needed to elucidate the relationships among plasma membrane biosynthesis, DNA replication, and vacuole formation at the molecular level.

**Funding:** This research received no external funding.

**Conflicts of Interest:** The author declares no conflict of interest.

## References

1. Garcia-Vallvé, S.; Romeu, A.; Palau, J. Horizontal gene transfer in bacterial and archaeal complete genomes. *Genome Res.* **2000**, *10*, 1719–1725. [[CrossRef](#)] [[PubMed](#)]
2. Koonin, E.V.; Makarova, K.S.; Aravind, L. Horizontal gene transfer in prokaryotes: Quantification and classification. *Annu. Rev. Microbiol.* **2001**, *55*, 709–742. [[CrossRef](#)] [[PubMed](#)]
3. Sun, D. Pull in and push out: Mechanisms of horizontal gene transfer in bacteria. *Front. Microbiol.* **2018**, *9*, 2154. [[CrossRef](#)] [[PubMed](#)]
4. Suzuki-Minakuchi, C.; Navare, W.W. Xenogeneic silencing and horizontal gene transfer. In *DNA Traffic in the Environment*; Nishida, H., Oshima, T., Eds.; Springer: Singapore, 2019; pp. 1–27.
5. Ito, T.; Okura, M.; Maruyama, F. Acquired and innate immunity in prokaryotes define their evolutionary story. In *DNA Traffic in the Environment*; Nishida, H., Oshima, T., Eds.; Springer: Singapore, 2019; pp. 47–75.
6. Nagata, Y.; Kato, H.; Ohtsubo, Y.; Tsuda, M. Mobile genetic elements involved in the evolution of bacteria that degrade recalcitrant xenobiotic compounds. In *DNA Traffic in the Environment*; Nishida, H., Oshima, T., Eds.; Springer: Singapore, 2019; pp. 215–244.
7. Hooton, S.P.; Millard, A.D.; Baker, M. DNA traffic in the environment and antimicrobial resistance. In *DNA Traffic in the Environment*; Nishida, H., Oshima, T., Eds.; Springer: Singapore, 2019; pp. 245–271.
8. Itaya, M.; Tsuge, K.; Koizumi, M.; Fujita, K. Combining two genomes in one cell: Stable cloning of the *Synechocystis* PCC6803 genome in the *Bacillus subtilis* 168 genome. *Proc. Natl. Acad. Sci. USA* **2005**, *102*, 15971–15976. [[CrossRef](#)]
9. Lartigue, C.; Glass, J.I.; Alperovich, N.; Pieper, R.; Parmar, P.P.; Hutchison, C.A., III; Smith, H.O.; Venter, J.C. Genome transplantation in bacteria: Changing one species to another. *Science* **2007**, *317*, 632–638. [[CrossRef](#)]

10. Gibson, D.G.; Glass, J.I.; Lartigue, C.; Noskov, V.N.; Chuang, R.-Y.; Algire, M.A.; Benders, G.A.; Montague, M.G.; Ma, L.; Moodie, M.M.; et al. Creation of a bacterial cell controlled by a chemically synthesized genome. *Science* **2010**, *329*, 52–56. [[CrossRef](#)]
11. Hutchinson, C.A., III; Chuang, R.-Y.; Noskov, V.N.; Assad-Garcia, N.; Deerinck, T.J.; Ellisman, M.H.; Gill, J.; Kannan, K.; Karas, B.J.; Ma, L.; et al. Design and synthesis of a minimal bacterial genome. *Science* **2016**, *351*, aad6253. [[CrossRef](#)]
12. Itaya, M.; Sato, M.; Hasegawa, M.; Kono, N.; Tomita, M.; Kaneko, S. Far rapid synthesis of giant DNA in the *Bacillus subtilis* genome by a conjugation transfer system. *Sci. Rep.* **2018**, *8*, 8792. [[CrossRef](#)]
13. Dienes, L.; Weinberger, H.J. The L forms of bacteria. *Bacteriol. Rev.* **1951**, *15*, 245–288. [[CrossRef](#)]
14. Allan, E.J.; Hoischen, C.; Gumpert, J. Bacterial L-forms. *Adv. Appl. Microbiol.* **2009**, *68*, 1–39.
15. Errington, J. L-form bacteria, cell walls and the origin of life. *Open Biol.* **2013**, *3*, 120143. [[CrossRef](#)]
16. Errington, J. Cell wall-deficient, L-form bacteria in the 21st century: A personal perspective. *Biochem. Soc. Trans.* **2017**, *45*, 287–295. [[CrossRef](#)]
17. Kawai, Y.; Mickiewicz, K.; Errington, J. Lysozyme counteracts  $\beta$ -lactam antibiotics by promoting the emergence of L-form bacteria. *Cell* **2018**, *172*, 1038–1049. [[CrossRef](#)]
18. Kawai, Y.; Mercier, R.; Mickiewicz, K.; Serafini, A.; de Carvalho, L.P.S.; Errington, J. Crucial role for central carbon metabolism in the bacterial L-form switch and killing by  $\beta$ -lactam antibiotics. *Nat. Microbiol.* **2019**, *4*, 1716–1726. [[CrossRef](#)] [[PubMed](#)]
19. Kuroda, T.; Okuda, N.; Saitoh, N.; Hiyama, T.; Terasaki, Y.; Anazawa, H.; Hirata, A.; Mogi, T.; Kusaka, I.; Tuchiya, T.; et al. Patch clamp studies on ion pumps of the cytoplasmic membrane of *Escherichia coli*. Formation, preparation, and utilization of giant vacuole-like structures consisting of everted cytoplasmic membrane. *J. Biol. Chem.* **1998**, *273*, 16897–16904. [[CrossRef](#)] [[PubMed](#)]
20. Lederberg, J. Bacterial protoplasts induced by penicillin. *Proc. Natl. Acad. Sci. USA* **1956**, *42*, 574–577. [[CrossRef](#)] [[PubMed](#)]
21. Nakamura, K.; Ikeda, S.; Matsuo, T.; Hirata, A.; Takehara, M.; Hiyama, T.; Kawamura, F.; Kusaka, I.; Tsuchiya, T.; Kuroda, T.; et al. Patch clamp analysis of the respiratory chain in *Bacillus subtilis*. *Biochim. Biophys. Acta* **2011**, *1808*, 1103–1107. [[CrossRef](#)] [[PubMed](#)]
22. Cox, C.D.; Bavi, N.; Martinac, B. Bacterial mechanosensors. *Annu. Rev. Physiol.* **2018**, *80*, 71–93. [[CrossRef](#)]
23. Morgan, J.L.W.; Evans, E.G.B.; Zagotta, W.N. Functional characterization and optimization of a bacterial cyclic nucleotide-gated channel. *J. Biol. Chem.* **2019**, *294*, 7503–7515. [[CrossRef](#)]
24. Takahashi, S.; Mizuma, M.; Kami, S.; Nishida, H. Species-dependent protoplast enlargement involves different types of vacuole generation in bacteria. *Sci. Rep.* **2020**, *10*, 8832. [[CrossRef](#)]
25. Tabor, C.W. Stabilization of protoplasts and spheroplasts by spermine and other polyamines. *J. Bacteriol.* **1962**, *83*, 1101–1111. [[CrossRef](#)]
26. Nishino, K.; Nishida, H. Calcium ion induces outer membrane fusion of *Deinococcus grandis* spheroplasts to generate giant spheroplasts with multiple cytoplasm. *FEMS Microbiol. Lett.* **2019**, *366*, fny282. [[CrossRef](#)] [[PubMed](#)]
27. Rojas, E.R.; Billings, G.; Odermatt, P.D.; Auer, G.K.; Zhu, L.; Miguel, A.; Chang, F.; Weibel, D.B.; Theriot, J.A.; Huang, K.C. The outer membrane is an essential load-bearing element in Gram-negative bacteria. *Nature* **2018**, *559*, 617–621. [[CrossRef](#)]
28. Nishino, K.; Morita, Y.; Takahashi, S.; Okumura, M.; Shiratani, S.; Umemura, K.; Narumi, I.; Kondo, C.; Ochiai, R.; Oshima, T.; et al. Enlargement of *Deinococcus grandis* spheroplasts requires  $Mg^{2+}$  or  $Ca^{2+}$ . *Microbiology* **2018**, *164*, 1361–1371. [[CrossRef](#)] [[PubMed](#)]
29. Kučerka, N.; Papp-Szabo, E.; Nieh, M.-P.; Harroun, T.A.; Schooling, S.R.; Pencer, J.; Nicholson, E.A.; Beveridge, T.J.; Katsaras, J. Effect of cations on the structure of bilayers formed by lipopolysaccharides isolated from *Pseudomonas aeruginosa* PAO1. *J. Phys. Chem.* **2008**, *112*, 8057–8062. [[CrossRef](#)]
30. Lam, N.H.; Ma, Z.; Ha, B.-Y. Electrostatic modification of the lipopolysaccharide layer: Competing effects of divalent cations and polycationic or polyanionic molecules. *Soft Matter* **2014**, *10*, 7528–7544. [[CrossRef](#)]
31. Clifton, L.A.; Skoda, M.W.A.; Le Brun, A.P.; Ciesielski, F.; Kuzmenko, I.; Holt, S.A.; Lakey, J.H. Effect of divalent cation removal on the structure of Gram-negative bacterial outer membrane models. *Langmuir* **2015**, *31*, 404–412. [[CrossRef](#)] [[PubMed](#)]
32. Anderson, R.; Hansen, K. Structure of a novel phosphoglycolipid from *Deinococcus radiodurans*. *J. Biol. Chem.* **1985**, *260*, 12219–12223.

33. Huang, Y.; Anderson, R. Structure of a novel glucosamine-containing phosphoglycolipid from *Deinococcus radiodurans*. *J. Biol. Chem.* **1989**, *264*, 18667–18672.
34. Raetz, C.R.H.; Whitfield, C. Lipopolysaccharide endotoxins. *Annu. Rev. Biochem.* **2002**, *71*, 635–700. [[CrossRef](#)]
35. Pieringer, R.A.; Kunnes, R.S. The biosynthesis of phosphatidic acid and lysophosphatidic acid by glyceride phosphokinase pathways in *Escherichia coli*. *J. Biol. Chem.* **1965**, *240*, 2833–2838.
36. Pieringer, R.A.; Bonner, H.; Kunnes, R.S. Biosynthesis of phosphatidic acid, lysophosphatidic acid, diglyceride, and triglyceride by fatty acyltransferase pathways in *Escherichia coli*. *J. Biol. Chem.* **1967**, *242*, 2719–2724. [[PubMed](#)]
37. Nishino, K.; Tsuchikado, R.; Nishida, H. Sugar enhances outer membrane fusion in *Deinococcus grandis* spheroplasts to generate calcium ion-dependent extra-huge cells. *FEMS Microbiol. Lett.* **2019**, *366*, fnz087. [[CrossRef](#)] [[PubMed](#)]
38. Farci, D.; Bowler, M.W.; Kirkpatrick, J.; McSweeney, S.; Tramontano, E.; Piano, D. New features of the cell wall of the radio-resistant bacterium *Deinococcus radiodurans*. *Biochim. Biophys. Acta* **2014**, *1838*, 1978–1984. [[CrossRef](#)] [[PubMed](#)]
39. Misra, C.S.; Basu, B.; Apte, S.K. Surface (S)-layer proteins of *Deinococcus radiodurans* and their utility as vehicles for surface localization of functional proteins. *Biochim. Biophys. Acta* **2015**, *1848*, 3181–3187. [[CrossRef](#)]
40. Fagan, R.P.; Fairweather, N.F. Biogenesis and functions of bacterial S-layers. *Nat. Rev. Microbiol.* **2014**, *12*, 211–222. [[CrossRef](#)] [[PubMed](#)]
41. Morita, Y.; Nishida, H. The common ancestor of *Deinococcus* species was rod-shaped. *Open Bioinform. J.* **2018**, *11*, 252–258. [[CrossRef](#)]
42. Shiomi, D.; Sakai, M.; Niki, H. Determination of bacterial rod shape by a novel cytoskeletal membrane protein. *ENBO J.* **2008**, *27*, 3081–3091. [[CrossRef](#)]
43. Bendezú, F.O.; Hale, C.A.; Bernhardt, T.G.; de Boer, P.A.J. RodZ (YfgA) is required for proper assembly of the MreB actin cytoskeleton and cell shape in *E. coli*. *EMBO J.* **2009**, *28*, 193–204. [[CrossRef](#)]
44. Alyahya, S.A.; Alexander, R.; Costa, T.; Henriques, A.O.; Emonet, T.; Jacobs-Wagner, C. RodZ, a component of the bacterial core morphogenic apparatus. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 1239–1244. [[CrossRef](#)]
45. Ago, R.; Shiomi, D. RodZ: A key-player in cell elongation and cell division in *Escherichia coli*. *AIMS Microbiol.* **2019**, *5*, 358–367. [[CrossRef](#)]
46. Morita, Y.; Okumura, M.; Narumi, I.; Nishida, H. Sensitivity of *Deinococcus grandis* rodZ deletion mutant to calcium ions results in enhances spheroplast size. *AIMS Microbiol.* **2019**, *5*, 176–185. [[CrossRef](#)] [[PubMed](#)]
47. Formstone, A.; Errington, J. A magnesium-dependent *mreB* null mutant: Implications for the role of *mreB* in *Bacillus subtilis*. *Mol. Microbiol.* **2005**, *55*, 1646–1657. [[CrossRef](#)]
48. Takahashi, S.; Takayanagi, A.; Takahashi, Y.; Oshima, T.; Nishida, H. Comparison of transcriptomes of enlarged spheroplasts of *Erythrobacter litoralis* and *Lelliottia amnigena*. *AIMS Microbiol.* **2016**, *2*, 152–189. [[CrossRef](#)]
49. Takayanagi, A.; Takahashi, S.; Nishida, H. Requirement of dark condition for enlargement of the aerobic photosynthetic marine bacterium *Erythrobacter litoralis*. *J. Gen. Appl. Microbiol.* **2016**, *62*, 14–17. [[CrossRef](#)] [[PubMed](#)]
50. Nakazawa, M.; Nishida, H. Effects of light and oxygen on the enlargement of *Erythrobacter litoralis* spheroplasts. *J. Gen. Appl. Microbiol.* **2017**, *63*, 58–61. [[CrossRef](#)] [[PubMed](#)]
51. Okamura, K.; Mitsumori, F.; Ito, O.; Takamiya, K.-I.; Nishimura, M. Photophosphorylation and oxidative phosphorylation in intact cells and chromatophores of an aerobic photosynthetic bacterium, *Erythrobacter* sp. strain OCh114. *J. Bacteriol.* **1986**, *168*, 1142–1146. [[CrossRef](#)]
52. Nishino, K.; Nishida, H. Blue light inhibits the enlargement of *Erythrobacter litoralis* spheroplasts. *J. Gen. Appl. Microbiol.* **2017**, *63*, 203–206. [[CrossRef](#)]
53. Harashima, K.; Kawazoe, K.; Yoshida, I.; Kamata, H. Light-stimulated aerobic growth of *Erythrobacter* species OCh114. *Plant Cell Physiol.* **1987**, *28*, 365–374.
54. Yurkov, V.V.; Beatty, J.T. Aerobic anoxygenic phototrophic bacteria. *Microbiol. Mol. Biol. Rev.* **1998**, *62*, 695–724. [[CrossRef](#)]
55. Masuda, S.; Bauer, C.E. AppA is a blue light photoreceptor that antirepresses photosynthesis gene expression in *Rhodobacter sphaeroides*. *Cell* **2002**, *110*, 613–623. [[CrossRef](#)]

56. Nishino, K.; Takahashi, S.; Nishida, H. Comparison of gene expression levels of *appA*, *ppsR*, and *EL368* in *Erythrobacter litoralis* spheroplasts under aerobic and anaerobic conditions, and under blue light, red light, and dark conditions. *J. Gen. Appl. Microbiol.* **2018**, *64*, 117–120. [[CrossRef](#)]
57. Kawai, Y.; Mercier, R.; Wu, L.J.; Domínguez-Cuevas, P.; Oshima, T.; Errington, J. Cell growth of wall-free L-form bacteria is limited by oxidative damage. *Curr. Biol.* **2015**, *25*, 1613–1618. [[CrossRef](#)]
58. Errington, J.; Mickiewicz, K.; Kawai, Y.; Wu, L.J. L-form bacteria, chronic diseases and the origins of life. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **2016**, *371*, 20150494. [[CrossRef](#)]
59. Takahashi, S.; Nishida, H. Comparison of gene expression among normally divided cells, elongated cells, spheroplasts at the beginning of growth, and enlarged spheroplasts at 43 h of growth in *Lelliottia amnigena*. *Gene Rep.* **2017**, *7*, 87–90. [[CrossRef](#)]
60. Autret, S.; Levine, A.; Holland, I.B.; Séror, S.J. Cell cycle checkpoints in bacteria. *Biochimie* **1997**, *79*, 549–554. [[CrossRef](#)]
61. Grainge, I. FtsK—A bacterial cell division checkpoint? *Mol. Microbiol.* **2010**, *78*, 1055–1057. [[CrossRef](#)] [[PubMed](#)]
62. Modell, J.W.; Kambara, T.K.; Perchuk, B.S.; Laub, M.T. A DNA damage-induced, SOS-independent checkpoint regulates cell division in *Caulobacter crescentus*. *PLoS Biol.* **2014**, *12*, e1001977. [[CrossRef](#)]
63. Ghusinga, K.R.; Vargas-García, C.A.; Singh, A. A mechanistic stochastic framework for regulating bacterial cell division. *Sci. Rep.* **2016**, *6*, 30229. [[CrossRef](#)]
64. Hajduk, I.V.; Rodrigues, C.D.A.; Harry, E.J. Connecting the dots of the bacterial cell cycle: Coordinating chromosome replication and segregation with cell division. *Semi. Cell Dev. Biol.* **2016**, *53*, 2–9. [[CrossRef](#)]
65. Willis, L.; Huang, K.C. Sizing up the bacterial cell cycle. *Nat. Rev. Microbiol.* **2017**, *15*, 606–620. [[CrossRef](#)] [[PubMed](#)]
66. Osella, M.; Tans, S.J.; Lagomarsino, M.C. Step by step, cell by cell: Quantification of the bacterial cell cycle. *Trends Microbiol.* **2017**, *25*, 250–256. [[CrossRef](#)] [[PubMed](#)]
67. Zhang, Z.; Miliás-Argeitis, A.; Heinemann, M. Dynamic single-cell NAD(P)H measurement reveals oscillatory metabolism throughout the *E. coli* cell division cycle. *Sci. Rep.* **2018**, *8*, 2162. [[CrossRef](#)] [[PubMed](#)]
68. Si, F.; Le Treut, G.; Sauls, J.T.; Vadia, S.; Levin, P.A.; Jun, S. Mechanistic origin of cell-size control and homeostasis in bacteria. *Curr. Biol.* **2019**, *29*, 1760–1770. [[CrossRef](#)] [[PubMed](#)]
69. Kusaka, I. Growth and division of protoplasts of *Bacillus megaterium* and inhibition of division by penicillin. *J. Bacteriol.* **1967**, *94*, 884–888. [[CrossRef](#)]
70. Ranjit, D.K.; Young, K.D. The Rcs stress response and accessory envelope proteins are required for de novo generation of cell shape in *Escherichia coli*. *J. Bacteriol.* **2013**, *195*, 2452–2462. [[CrossRef](#)]
71. Ranjit, D.K.; Jorgenson, M.A.; Young, K.D. PBP1B glycosyltransferase and transpeptidase activates play different essential roles during the de novo regeneration of rod morphology in *Escherichia coli*. *J. Bacteriol.* **2017**, *199*, e00612–16. [[CrossRef](#)]
72. Takahashi, S.; Nishida, H. Quantitative analysis of chromosomal and plasmid DNA during the growth of spheroplasts of *Escherichia coli*. *J. Gen. Appl. Microbiol.* **2015**, *61*, 262–265. [[CrossRef](#)]
73. Schwarz, U.; Asmus, A.; Frank, H. Autolytic enzymes and cell division of *Escherichia coli*. *J. Mol. Biol.* **1969**, *41*, 419–429. [[CrossRef](#)]
74. Tsuchikado, R.; Kami, S.; Takahashi, S.; Nishida, H. Novobiocin inhibits membrane synthesis and vacuole formation of *Enterococcus faecalis*. *Microb. Cell* **2020**, in press.
75. Gellert, M.; O’Dea, M.H.; Tomizawa, J. Novobiocin and coumermycin inhibit DNA supercoiling catalyzed by DNA gyrase. *Proc. Natl. Acad. Sci. USA* **1976**, *73*, 4474–4478. [[CrossRef](#)] [[PubMed](#)]
76. Sugino, A.; Higgins, N.P.; Brown, P.O.; Peebles, C.L.; Cozzarelli, N.R. Energy coupling in DNA gyrase and the mechanism of action of novobiocin. *Proc. Natl. Acad. Sci. USA* **1978**, *75*, 4838–4842. [[CrossRef](#)] [[PubMed](#)]
77. Kami, S.; Tsuchikado, R.; Nishida, H. DNA replication and cell enlargement of *Enterococcus faecalis* protoplasts. *AIMS Microbiol.* **2019**, *5*, 347–357. [[CrossRef](#)] [[PubMed](#)]
78. Marvin, D.A. Control of DNA replication by membrane. *Nature* **1968**, *219*, 485–486. [[CrossRef](#)] [[PubMed](#)]
79. Sueoka, N.; Quinn, W.G. Membrane attachment of the chromosome replication origin in *Bacillus subtilis*. *Cold Spring Harb. Symp. Quant. Biol.* **1968**, *33*, 695–705. [[CrossRef](#)]
80. Earhart, C.F.; Tremblay, G.Y.; Daniels, M.J.; Schaechter, M. DNA replication studied by a new method for the isolation of cell membrane–DNA complexes. *Cold Spring Harb. Symp. Quant. Biol.* **1968**, *33*, 707–710. [[CrossRef](#)]

81. Leibowitz, P.J.; Schaechter, M. The attachment of the bacterial chromosome to the cell membrane. *Int. Rev. Cytol.* **1975**, *41*, 1–28.
82. Garner, J.; Crooke, E. Membrane regulation of the chromosomal replication activity of *E. coli* DnaA requires a discrete site on the protein. *EMBO J.* **1996**, *15*, 3477–3485. [[CrossRef](#)]
83. Boeneman, K.; Crooke, E. Chromosomal replication and the cell membrane. *Curr. Opin. Microbiol.* **2005**, *8*, 143–148. [[CrossRef](#)]
84. Mendell, J.E.; Clements, K.D.; Choat, J.H.; Angert, E.R. Extreme polyploidy in a large bacterium. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 6730–6734. [[CrossRef](#)]
85. Levin, P.A.; Angert, E.R. Small but mighty: Cell size and bacteria. *Cold Spring Harb. Perspect. Biol.* **2015**, *7*, a019216. [[CrossRef](#)] [[PubMed](#)]
86. Ionescu, D.; Bizic-Ionescu, M.; De Maio, M.; Cypionka, H.; Grossart, H.-P. Community-like genome in single cells of the sulfur bacterium *Achromatium oxaliferum*. *Nat. Commun.* **2017**, *8*, 455. [[CrossRef](#)]
87. Sharpe, M.E.; Hauser, P.M.; Sharpe, R.G.; Errington, J. *Bacillus subtilis* cell cycle as studied by fluorescence microscopy: Constancy of cell length at initiation of DNA replication and evidence for active nucleoid partitioning. *J. Bacteriol.* **1998**, *180*, 547–555. [[CrossRef](#)]
88. Schulz, H.N.; Brinkhoff, T.; Ferdelman, T.G.; Hernández Mariné, M.; Teske, A.; Jørgensen, B.B. Dense populations of a giant sulfur bacterium in Namibian shelf sediments. *Science* **1999**, *284*, 493–495. [[CrossRef](#)] [[PubMed](#)]
89. Cunningham, J.A.; Thomas, C.-W.; Bengtson, S.; Marone, F.; Stampanoni, M.; Turner, F.R.; Bailey, J.V.; Raff, R.A.; Raff, E.C.; Donoghue, P.C.J. Experimental taphonomy of giant sulfur bacteria: Implications for the interpretation of the embryo-like Ediacaran Doushantuo fossils. *Proc. Biol. Sci.* **2012**, *279*, 1857–1864. [[PubMed](#)]



© 2020 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).