

BMJ Open Early Life Exposure in Mexico to ENvironmental Toxicants (ELEMENT) Project

Wei Perng,^{1,2,3} Marcela Tamayo-Ortiz,⁴ Lu Tang,⁵ Brisa N Sánchez,⁵ Alejandra Cantoral,⁴ John D Meeker,⁶ Dana C Dolinoy,^{2,6,7} Elizabeth F Roberts,⁸ Esperanza Angeles Martinez-Mier,⁹ Hector Lamadrid-Figueroa,¹⁰ Peter X K Song,⁵ Adrienne S Ettinger,² Robert Wright,¹¹ Manish Arora,¹¹ Lourdes Schnaas,¹² Deborah J Watkins,⁶ Jaclyn M Goodrich,⁶ Robin C Garcia,² Maritsa Solano-Gonzalez,¹³ Luis F Bautista-Arredondo,¹³ Adriana Mercado-Garcia,¹³ Howard Hu,¹⁴ Mauricio Hernandez-Avila,¹⁵ Martha Maria Tellez-Rojo,¹³ Karen E Peterson^{2,6}

To cite: Perng W, Tamayo-Ortiz M, Tang L, *et al.* Early Life Exposure in Mexico to ENvironmental Toxicants (ELEMENT) Project. *BMJ Open* 2019;**9**:e030427. doi:10.1136/bmjopen-2019-030427

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-030427>).

WP and MT-O contributed equally.

Received 14 March 2019
Revised 23 July 2019
Accepted 25 July 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Martha Maria Tellez-Rojo; mmtellez@insp.mx

ABSTRACT

Purpose The Early Life Exposure in Mexico to ENvironmental Toxicants (ELEMENT) Project is a mother–child pregnancy and birth cohort originally initiated in the mid-1990s to explore: (1) whether enhanced mobilisation of lead from maternal bone stores during pregnancy poses a risk to fetal and subsequent offspring neurodevelopment; and (2) whether maternal calcium supplementation during pregnancy and lactation can suppress bone lead mobilisation and mitigate the adverse effects of lead exposure on offspring health and development. Through utilisation of carefully archived biospecimens to measure other prenatal exposures, banking of DNA and rigorous measurement of a diverse array of outcomes, ELEMENT has since evolved into a major resource for research on early life exposures and developmental outcomes.

Participants n=1643 mother–child pairs sequentially recruited (between 1994 and 2003) during pregnancy or at delivery from maternity hospitals in Mexico City, Mexico.

Findings to date Maternal bone (eg, patella, tibia) is an endogenous source for fetal lead exposure due to mobilisation of stored lead into circulation during pregnancy and lactation, leading to increased risk of miscarriage, low birth weight and smaller head circumference, and transfer of lead into breastmilk. Daily supplementation with 1200 mg of elemental calcium during pregnancy and lactation reduces lead resorption from maternal bone and thereby, levels of circulating lead. Beyond perinatal outcomes, early life exposure to lead is associated with neurocognitive deficits, behavioural disorders, higher blood pressure and lower weight in offspring during childhood. Some of these relationships were modified by dietary factors; genetic polymorphisms specific for iron, folate and lipid metabolism; and timing of exposure. Research has also expanded to include findings published on other toxicants such as those associated with personal care products and plastics (eg, phthalates, bisphenol A), other metals (eg, mercury, manganese, cadmium), pesticides (organophosphates) and fluoride; other biomarkers (eg, toxicant levels in plasma, hair and teeth); other outcomes (eg, sexual maturation, metabolic

Strengths and limitations of this study

- In-person collection of exposure and outcome measures using research-quality standard protocols.
- Extensive biorepository of specimens from gestation through adolescence.
- Rich covariate data including physician-assessed pubertal status, newly collected high-dimensional ‘omics data and ethnographical data.
- Limited information on maternal perinatal conditions (eg, data on prepregnancy body mass index in a small subset of mothers, no information on maternal glycaemia).
- Potential non-generalisability of results to those of other race/ethnicities.

syndrome, dental caries); and identification of novel mechanisms via epigenetic and metabolomics profiling.

Future plans As the ELEMENT mothers and children age, we plan to (1) continue studying the long-term consequences of toxicant exposure during the perinatal period on adolescent and young adult outcomes as well as outcomes related to the original ELEMENT mothers, such as their metabolic and bone health during perimenopause; and (2) follow the third generation of participants (children of the children) to study intergenerational effects of in utero exposures.

Trial registration number NCT00558623.

INTRODUCTION

Why was the cohort set up?

Original cohorts

In the early 1990s, a series of epidemiological studies on the detrimental effects of childhood lead exposure on neurodevelopment^{1–4} and an emerging concern on the neurotoxic effects of prenatal lead exposure⁵ motivated HH, MH-A and Eduardo Palazuelos (EP) to

explore the two original research questions of the Early Life Exposure to ENvironmental Toxicants (ELEMENT) Project: (1) *Does enhanced mobilisation of lead from maternal bone stores during pregnancy pose a risk to fetal development and subsequent offspring neurodevelopment?*⁵; (2) *Can calcium supplementation during early-life suppress the mobilisation of lead from maternal bone and mitigate the adverse health effects of lead exposure?*⁶? These inquiries were critically important given high levels of lead exposure in several countries worldwide^{7,8} including Mexico⁹; biological evidence of enhanced bone mobilisation (and thus, mobilisation of lead stored in bone) during the in utero and post-partum lactation periods;^{10,11} and growing evidence of vulnerability during the fetal and infancy life stages to environmental exposures in relation to a range of health outcomes.^{12–14} Together, these concerns stimulated the search for cost-effective and population-level intervention strategies to mitigate the adverse effects of lead.

Mexico City was an ideal setting to address these questions for two key reasons. First, because the phase-out of leaded gasoline in Mexico occurred later than in the USA,^{9,15} levels of lead in bone were expected to be higher in women in Mexico. Second, even after the phase-out of leaded gasoline, blood lead concentrations among children in Mexico City remained three to four times higher than that of children in the USA¹⁶ due, in large part, to continued use of lead-glazed ceramics, which remains prevalent in Mexico, especially in the central and southern regions of the nation. While unfortunate from a population health perspective, the relatively high lead exposure in Mexico City provided a powerful context within which to explore the first research question, especially given that k-X-ray fluorescence (KXRF), a non-invasive technology for measuring bone lead levels, had recently become available.¹⁷ Thus, in 1993, HH, MHA and EP founded the ELEMENT Project. Additional investigators, including KEP (current principal investigator (PI) in the USA) and M-MT-R (current PI in Mexico) became involved as the project expanded.

Today, ELEMENT has become a 20+ year international multi-institutional partnership. The ELEMENT Project is based on three sequentially recruited distinct cohorts of mother–child pairs recruited from a homogenous low-income to middle-income population attending family clinics in Mexico City, most belonging to the *Instituto Mexicano del Seguro Social* (IMSS). Cohort 1 (recruited 1994–1995) was a prospective study of measures of maternal and fetal lead biomarkers, determinants of levels of lead in the biomarkers and perinatal outcomes coupled with a postnatal randomised placebo-controlled trial (RCT). The trial, which focused on lactating women, sought to examine the impact of early life lead exposure on neuro-behavioural and physical developmental outcomes and the potential modifying effect of a daily 1200 mg calcium supplementation on biomarkers of lead in maternal blood and breast milk while breast feeding.¹⁸ Cohort 2 (recruited 1997–2000) was initiated, in part, based on findings from cohort 1 and was an observational cohort

study that examined the relationship between periconceptual and gestational maternal blood, plasma and bone lead biomarkers in two different groups of women with follow-up of offspring neurocognitive outcomes. Cohort 3 (recruited 2001–2003) was an RCT with similar goals to that of cohort 1, but with the goal of assessing effects of calcium supplementation during pregnancy (table 1).

Background and sociodemographic characteristics of the three original cohorts are similar, as shown in table 2, with the exception of delivery mode wherein vaginal deliveries were of highest prevalence in cohort 2B. In analyses involving multiple cohorts, researchers evaluated for potential heterogeneity across cohorts by examining bivariate associations of key study characteristics by cohort membership, followed by covariate adjustment for cohort membership or inclusion of a random effect for cohort if appropriate.^{19,20} Additionally, development of more sophisticated statistical methods is underway to deal with heterogeneity across cohorts.^{21,22}

These cohorts comprise 1643 mother–child pairs, some of whom we have followed for over two decades. Findings from these cohorts motivated the reduced use of lead in consumer products in Mexico^{23,24} and informed US lead exposure and management guidelines.^{25–28}

Maternal informed consent and child assent were provided for all participants. The institutional review boards of the Mexico National Institute of Public Health (CI-03121993, CI-124-30052000, CI-362-16122003, CI-599-13062007, CI-599-2-07112010, CI-599-17072014, CI-599-9-15102014, CI-599-8-14102014, CI-599-3-23062014, CI-1377-15062016, CI-1594-18122018), Harvard School of Public Health (P10345-107/9910CONT) and the University of Michigan approved research protocols (#HUM00034344, HUM00155958).

Follow-up studies

ELEMENT 2008

In 2008, we pooled together mother–child pairs from the original cohorts to study interactions of perinatal lead exposure and polymorphisms of genes relevant to cholesterol metabolism and their influence on offspring neurodevelopment and behaviour at age 6–15 years.

ELEMENT 2011

Between 2011 and 2012, we re-recruited a convenience sample (based on availability of prenatal biological samples) of offspring from cohorts 2 and 3 for participation in follow-up studies concerned with: (1) whether prenatal and peripubertal exposure to endocrine disrupting chemicals (EDCs) were associated with obesity, metabolic risk and onset of sexual maturation at age 8–15 years; and (2) to examine epigenetic, metabolomics and oxidative stress as mechanisms linking EDC exposure to health outcomes.

ELEMENT 2014

In 2014, we commenced ethnographic observations of six families (the ‘Mexican Exposures Project’) to obtain a

Table 1 Original ELEMENT cohorts and follow-up history

	Study information							
	Cohort 1		Cohort 2				Cohort 3	
			2A		2B			
Recruitment dates	1994–1995		1997–1999		1999–2000		2001–2003	
Recruitment life stage	Delivery		First trimester		Delivery		First trimester	
Eligible participants	2193 mothers		2143 mothers		2191 mothers		1855 mothers	
Study design	RCT		Cohort study		Cohort study		RCT	
	Mother	Child	Mother	Child	Mother	Child	Mother	Child
N individuals	639	632	327	253	533	433	670	393
N mother–child dyads	631		252		367		393	
Follow-up visits for the original cohorts								
Pregnancy								
First trimester			x				x	
Second trimester			x				x	
Third trimester			x				x	
Delivery	x	x	x	x	x	x	x	x
Postpartum months								
1	x	x	x		x		x	
3	x	x		x		x	x	x
4			x					
5		x					x	
6				x		x		x
7	x	x	x				x	
9							x	
11							x	
12		x	x	x		x	x	x
18		x		x		x		x
24		x		x		x		x
30		x		x		x		x
36		x		x		x		x
42		x						
48		x		x		x		x
60				x		x		

Total mothers = 2169; total children = 1711.

ELEMENT, Early Life Exposure in Mexico to ENvironmental Toxicants; RCT, randomised placebo-controlled trial.

fuller picture of how chemical exposures and the changing food environment in Mexico City impact health.^{29 30} This study provided a context for mixed-methods approaches as we continue to follow participants over their life course and a platform for combining ethnographic and biomarker data about ELEMENT participants. Building off of this project, we recently began multidisciplinary home visit study called ‘Neighbourhood Environments as Socio-Techno-Bio Systems: Water Quality, Public Trust and Health in Mexico City (NESTSMX)’ with 60 ELEMENT families. The aim of the study is to combine

ethnographic and environmental health and environmental engineering to understand household water management, water trust and water quality.

ELEMENT 2015

Between 2015 and 2018, we followed-up children in ELEMENT 2011 through late adolescence and included additional similarly aged participants from the original cohorts. Goals of this study were similar in scope to ELEMENT 2011, but included new research foci: (1) characterising long-term consequences of early life exposure

Table 2 Characteristics of the original ELEMENT cohort participants

	Mean±SD or %			
	Cohort 1	Cohort 2		Cohort 3
		2A	2B	
Maternal characteristics	n=639	n=327	n=533	n=670
Age at enrolment (years)	24.6±5.1	26.9±5.3	25.2±5.1	26.4±5.5
Gestational age at enrolment (weeks)	39.1±1.6	15.4±5.9	38.9±1.3	13.1±2.1
Marital status (%)				
Married or cohabiting	91.10	91.10	93.30	88.50
Single	8.90	8.90	6.70	11.50
Maternal education (years)	9.34±3.1	10.75±3.2	10.84±3.0	10.7±2.9
Parity (including index birth) (%)				
0	0	1.50	0	0
1-2	73.60	70.90	78.40	70.40
≥3	26.40	27.50	21.60	29.60
Smoked during pregnancy (%)	4.30	4.00	9.50	2.20
Calcium treatment (% , N)	47.00	N/A	N/A	50.00
Delivery method (%)				
Vaginal	77.30	46.20	91.10	50.70
Forceps	1.70	3.50	3.40	1.10
C-section	19.70	40.00	5.00	21.30
Missing	1.30	12.80	0.40	26.90
Offspring characteristics at delivery	n=632	n=253	n=433	n=393
Sex (%)				
Male	54.50	51.90	54.10	49.40
Female	45.50	48.10	45.90	50.60
Gestational age at delivery (weeks)	39.2±1.5	38.6±1.9	39.0±1.2	38.6±1.7
Birth weight (g)	3153±474	3102±531	3130±484	3131±484

ELEMENT, Early Life Exposure in Mexico to ENvironmental Toxicants.

to fluoride; (2) exploring epigenetic and metabolomics mechanisms.

Pilot studies

In addition to the many pilot studies conducted in the past, two recent pilot studies are currently underway at the time of writing and will soon reach completion. Although we do not formally describe participants of these cohorts in this article, we provide a description of study goals and sample sizes below and in [figure 1](#).

ELEMENT 2016

Between 2016 and 2017, 100 offsprings from cohort 1 were recruited for MRI of hepatic fat fraction to identify determinants of fatty liver during young adulthood.

ELEMENT 2017

We recruited 101 mothers from all three cohorts to evaluate associations of chemical exposures during pregnancy and later adulthood with metabolic and bone

health (measured via intelligent dual X-ray absorptiometry) during the perimenopausal period.

COHORT DESCRIPTION

Who is in the cohort?

[Table 2](#) shows characteristics of participants in the original cohorts. [Table 1](#) and [figure 1](#) summarise study design, recruitment timing and number of eligible and enrolled participants, with added details on the follow-up and pilot studies in the figure. Inclusion criteria included healthy women in Mexico City with a normal pregnancy and/or delivery, intention to breast feed and no previous diagnoses of severe disease. We excluded women if they had a non-singleton birth, a history of preeclampsia or pregnancy-related hypertension, psychiatric or cardiac disease, diagnosis of gestational diabetes, a history of repeated urinary tract infections or seizure disorders requiring daily medication. The same research team followed all cohorts, using the same facilities, questionnaires and protocols.




Figure 1 Study participant flow for the original ELEMNT cohorts, follow-up studies and pilot studies. Ch, children; ELEMNT, Early life exposure in Mexico to ENvironmental toxicants; F, follow-up study; M, mothers; P, pilot study.

In the section below, we describe participant characteristics for the original three cohorts and key follow-up studies (ELEMNT 2008, 2011, and 2015).

Original cohorts

As outlined in table 1, between January 1994 and June 1995, we interviewed 2945 women admitted for labour and delivery for cohort 1. Of them, 2193 were eligible for the study and 639 (45.5%) agreed to participate. After excluding 22 who stopped breast feeding by 1 month postpartum, the remaining 617 women were randomly assigned to receive 1200 mg of elemental calcium per day (n=296) versus a placebo (n=321).

Cohort 2 comprises two different groups of women. For cohort 2A (recruitment 1997–1999), we interviewed 2215 women before pregnancy or during the first trimester. Of the 2143 eligible women, 327 were recruited. Cohort 2B includes 533 women recruited from 1999 to 2000 at delivery (2191 eligible and 2194 interviewed). Exclusion criteria for cohort 2 were similar to cohort 1, but also included missing data on cord blood lead or cord blood lead >10 µg/dL, conditions interfering with calcium metabolism, conditions associated with low birth weight, gestational age of <37 weeks, admittance to the neonatal intensive care unit, infant Apgar score of ≤6, birth weight <2000 g or serious birth defects.

Following findings from cohort 1 that calcium supplementation reduced maternal blood lead by 15%–20%¹⁸ and breast milk lead by 5%–10%,³¹ and results of cohort 2, which clarified the relationship between lead in whole blood versus plasma,^{32–34} we initiated cohort 3 to examine effects of

calcium supplementation on blood lead levels during pregnancy. For cohort 3, we recruited women between 2001 and 2003 from IMSS family clinics. Eligibility criteria included gestational age <14 weeks; no hypertensive disorders, history of type 1 or 2 diabetes or gestational diabetes; and planned residence in the Mexico City area for the next 5 years. Of the 3837 women interviewed, 1855 eligible women were eligible, 36% (n=670) of whom agreed to participate and were randomised to receive 1200 mg calcium per day (n=334) or a placebo (n=336).

Follow-up studies

ELEMNT 2008

Of the 1036 eligible children between 6 and 15 years of age from the original three cohorts, we excluded participants born <37 gestational weeks, those with birth weight <2000 g and Apgar score ≤6, admittance to the neonatal intensive care unit or a serious birth defect, arriving at a final sample size of 824 mother–child pairs.

ELEMNT 2011

Between 2011 and 2012, we recontacted 250 children and adolescents from cohorts 2 and 3. Criteria for eligibility included availability of archived maternal urine and cord-blood for toxicant assays of interest and offspring age between 8 and 15 years.

ELEMNT 2015

In 2015, we rerecruited 224 of 250 ELEMNT 2011 participants. We also recruited 330 additional offsprings who were of similar age to ELEMNT 2011 participants, giving

priority to those who had archived cord blood and urine samples and/or having been part of ELEMENT 2008. Additional data collected for 2015 included physical activity and sleep data from accelerometers; measures of fluoride exposure in food/beverages, water and plasma; and dental examinations for caries, enamel fluorosis and hypomineralisation.

How often have the cohorts been followed-up?

Depending on the specific cohort, women were interviewed two to four times during the first postpartum year. Offsprings were assessed every 6 months during their first four (cohort 1) or five (cohorts 2 and 3) years of life, with one to two follow-up visits at ages 6–15 years in ELEMENT 2008, once at ages 8–15 years in ELEMENT 2011 and twice at 10–18 years in ELEMENT 2015. [Table 1](#) and [figure 1](#) provide an overview of the timing of research visits.

WHAT HAS BEEN MEASURED?

While some of the specific measures differ across the original cohorts and follow-up studies, there are several key assessments common across studies. These common assessments include: (1) assessment of lead exposure (blood lead concentration from maternal venous, child venous and umbilical cord samples using graphite furnace atomic absorption spectroscopy until the late 2000s; then, inductively coupled plasma mass spectrometry; all in laboratories participating in the quality assurance/quality control (QA/QC) protocols of the US Centres for Disease Control and the State of Wisconsin; breast-milk lead concentration via mass spectrometry³⁵) and maternal bone lead levels using a KXRF instrument^{17 36}; (2) dietary intake via a food frequency questionnaire that has been validated for use among women living in Mexico City³⁷; (3) evaluation of neurobehavioural development in the offspring, including intelligence using the Bayley Scales of Infant Development (BSID-II, translated into Spanish³⁸) in the early years, McCarthy Scales of Children Abilities (MSCA, Spanish version) around age 4 years³⁹, Wechsler Abbreviated Scale of Intelligence (WAIS, Spanish version⁴⁰) and behaviour using the Behaviour Assessment System for Children (BASC-2, Spanish version⁴¹), executive function using the Behaviour Rating Inventory of Executive Function (BRIEF)⁴²; (4) anthropometry and body composition assessment in offspring (weight via a digital scale, height via a calibrated stadiometer, waist circumference via a non-stretchable measuring tape, skinfolds via calibrated callipers and body composition via bioimpedance analysis); (5) sexual maturation of offspring via physician-assessed Tanner staging; (6) blood pressure (left arm resting systolic and diastolic blood pressure of the mother and the child) using a standard mercury column sphygmomanometer; (7) sociodemographic information via an in-house questionnaire. [Table 3](#) summarises details of these and other research measures from each of the original cohorts, follow-up studies and pilot studies.

Patient and public involvement

During all ELEMENT research visits, a board-certified MD communicates results of tests and measurements (ie, bone lead levels, fasting glucose, lipid profile, body mass index, waist circumference) to the participants and their caregivers in order to facilitate understanding of the values in the context of normal ranges and standards. We engage in community outreach and translation via brochures and information sheets that discuss implications of our research findings in lay language. For example, in 2010, we worked with collaborators at the Instituto Nacional de Salud Pública (INSP) to develop a brochure with information on the dangers of lead exposure and ways to avoid it. This brochure was distributed to the study participants. Subsequently, we adapted this brochure for distribution to the Mexican-American community in southeast Michigan where lead exposure remains a concern. Finally, we are in the process of developing a quarterly newsletter to communicate key findings from current research in ELEMENT to the study participants. In 2014, we implemented a ‘Satisfaction Perception Questionnaire’ inquiring on participants’ perceptions and understanding of ELEMENT Project research questions, their involvement in the studies and their suggestions for improvement in study design and communication of study findings. This has helped us improve our incentives and the quality and depth of the information we give back to the participants. Additionally, as part of ELEMENT 2015 wherein dental examinations were conducted, a certified dentist on the research team discusses examination results with the participants, including explanations of the fluorosis dental images and tailored dental care advice.

FINDINGS TO DATE

ELEMENT research has generated >90 published papers as of this writing; the following captures a sample of ELEMENT’s findings. A full list of ELEMENT publications can be found here: <https://sph.umich.edu/cehc/research/publications.html>

Health consequences of perinatal exposure to lead

Perinatal outcomes

Characterising lead exposure and prevention of lead mobilisation from maternal bone have been key foci of ELEMENT.²⁶ We identified glazed ceramics as a major source of lead in this population⁴³ and discovered that maternal bone lead (eg, patella and tibia) was an endogenous source of fetal exposure due to mobilisation of stored lead into circulation^{32 34 44} during pregnancy and lactation,^{45 46} thereby increasing the risk of miscarriage,³³ lower birth weight and less weight gain at 1 month after birth,⁴⁷ smaller head circumference and shorter birth length,⁴⁸ and transfer of lead into breastmilk.⁴⁹ Calcium supplementation counteracted these adverse physiological events by reducing lead resorption,⁵⁰ blood lead

Table 3 Summary of data (denoted as ‘measure of interest (tool used)’) collected from the ELEMENT cohorts, follow-up studies and pilot studies

	Mothers	Children
Cohort 1 (delivery through 48 months postpartum)		
Biospecimen	Blood	Cord blood
	Urine	
	Breastmilk	
Questionnaire	Sociodemographic information (in-house questionnaire)	Cognition (BSID-II)
	Diet (FFQ)	IQ (WAIS)
	Home environment (HOME)	Neurodevelopment (McCarthy)
	IQ (WAIS)	
Assessments	Bone lead levels (KXRF)	Anthropometry (weight, length/height)
	Anthropometry (weight, height)	Blood pressure
	Blood pressure (manual auscultatory monitor)	
Cohort 2 (first trimester (2A) or delivery (2B) through 60 months postpartum)		
Biospecimen	Blood	Cord blood
	Urine	Blood
	Breastmilk	
Questionnaire	Sociodemographic information (in-house questionnaire)	Cognition (BSID-II)
	Diet (FFQ)	IQ (WAIS)
	Home environment (HOME)	Neurodevelopment (McCarthy)
	IQ (WAIS)	Behaviour (BASC)
Assessments	Bone lead levels (KXRF)	Diet (FFQ)
	Anthropometry (weight, height)	Anthropometry (weight, length/height)
	Blood pressure (manual auscultatory monitor)	Blood pressure (manual auscultatory monitor)
Cohort 3 (first trimester through 48 months postpartum)		
Biospecimen	Blood	Cord blood
	Urine	Blood
	Breastmilk	
Questionnaire	Sociodemographic information (in-house questionnaire)	Cognition (BSID-22)
	Diet (FFQ)	Behaviour (BASC)
	Home environment (HOME)	IQ (WAIS)
	IQ (WAIS)	Neurodevelopment (McCarthy)
Assessments	Bone lead levels (KXRF)	Diet (FFQ)
	Anthropometry (weight, height)	Anthropometry (weight, length/height)
	Blood pressure (manual auscultatory monitor)	Blood pressure (manual auscultatory monitor)
	Densitometry (BodPod)	
ELEMENT 2008 (offspring age 6–15 years)		
Biospecimen	Blood	Blood
	Urine	Urine
	Hair	Hair
Questionnaire	Sociodemographic information (in-house questionnaire)	Cognition (CANTAB)

Continued

Table 3 Continued

	Mothers	Children
	Diet (FFQ)	enrolment (BASC)
	Home environment (HOME)	IQ (WAIS)
	IQ (WAIS)	Attention (CPT-2)
		Executive function (BRIEF)
		Pre-pulse inhibition
		Diet (FFQ)
		Sexual maturation (self-reported Tanner stage)
Assessments	Anthropometry (weight, height, waist circumference, skinfold thicknesses)	Anthropometry (weight, height, waist circumference, skinfold thicknesses)
	Bone lead levels (KXRF)	Blood pressure (manual auscultatory monitor)
ELEMENT 2011 (offspring age 8–15 years)		
	Biospecimen	Blood
		Urine
	Questionnaire	Diet (FFQ)
		Sexual maturation (self-reported Tanner stage + physician evaluation)
		Physical activity (IPAQ)
Assessments		Anthropometry (weight, length, waist circumference, skinfold thicknesses)
		Blood pressure (automated oscillometric monitor)
		Sexual maturation (self-reported Tanner stage + physician evaluation)
ELEMENT 2014 (Ethnography)		
	Open-ended, long-term observations about everyday life. We collected intensive ethnographic observations about six ELEMENT families and their neighbours for 6 months each from 2014 to 2015. We continue these ongoing observations intermittently.	
ELEMENT 2015 (offspring age 13–20 years)		
	Biospecimen	Blood
		Urine
		Toenails
		Shed teeth
	Questionnaire	Sociodemographic information (in-house questionnaire)
		Attention (CPT-3)
		Behaviour (BASC-2)
		Neurobehavioural development (NIH Toolbox)
		Diet (FFQ)
		Physical activity (IPAQ)
		Sleep quality and quantity (in-house questionnaire)
Measurements		Anthropometry (weight, length, waist circumference, skinfold thicknesses)
		Blood pressure (automated oscillometric monitor)
		Body composition (BIA)
		Sexual maturation (self-reported Tanner stage + physician evaluation)

Continued

Table 3 Continued

Mothers		Children
		Dental imaging (fluorescent imaging)
		Accelerometry (Actigraph GTx3)
ELEMENT 2016 (offspring 17–20 years)		
Biospecimen		Blood
Questionnaire		Diet (FFQ)
		Physical activity (IPAQ)
		Sociodemographic information (in-house questionnaire)
Measurements		Anthropometry (weight, length, waist circumference)
		Blood pressure (automated oscillometric monitor)
		Liver fat content (MRI)
ELEMENT 2017 (mothers age 31–63 years)		
Biospecimen	Blood	
	Urine	
Questionnaire	Sociodemographic information (in-house questionnaire)	
	Diet (FFQ)	
	Health history (in-house questionnaire)	
	Physical activity (IPAQ)	
Measurements	Anthropometry (weight, height, waist circumference)	
	Body composition (iDXA, BIA)	
	Bone density (iDXA)	

BASC, Behavior Assessment System for Children; BIA, Bioelectric Impedance Assessment of Body Composition; BRIEF, Behavior Rating Inventory of Executive Function; BSID-II, Bayley Scales of Infant Development 2; CANTAB, Cambridge Neuropsychological Test Automated Battery; CPT, Conner's Continuous Performance Test; ELEMENT, Early Life Exposure in Mexico to ENvironmental Toxicants; FFQ, Food Frequency Questionnaire; HOME, Home Observation for the Measurement of the Environment; IPAQ, Integrated Physical Activity Questionnaire; KXRF, k-X-ray fluorescence; WAIS, Wechsler Abbreviated Scale of Intelligence; iDXA, Intelligent Dual X-ray Absorptiometry.

levels during pregnancy and lactation,^{18 51} and breastmilk lead concentrations.³¹

Offspring neurobehavioural development

Our studies confirmed the adverse effects of prenatal lead exposure on offspring cognitive development,⁵² with detectable adverse effects of lead exposure on cognition^{53–55} and behavioural disorders,⁴² even at low levels of exposure (eg, blood lead concentrations of <5 or 10 µg/dL). We also found that maternal self-esteem influences offspring mental and psychomotor development and may mitigate negative effects of lead exposure.⁵⁶ Genetic variants were also found to modify the relationship between lead exposure and early cognitive development.^{57 58}

Growth, maturation, blood pressure and metabolism

We found that higher cumulative prenatal lead exposure, as indicated by maternal bone lead levels, was associated with lower weight in boys but not girls during the first 5 years of life,⁵⁹ with lower body mass index (BMI) and adiposity among boys and girls undergoing adolescence⁶⁰ and with higher blood pressure in girls but not boys at 7–15 years of age.⁶¹ Additionally, while maternal lead

levels were not associated with offspring height, the child's own blood lead levels during infancy (birth to 24 months) predicted shorter stature at approximately 4 years of age,⁶² but not child adiposity at 8–6 years.⁶⁰ Prenatal lead exposure was associated with delays in pubertal onset and progression in girls only.^{60 63}

Other toxicants

Other metals

As ELEMENT evolved, its archived biorepository has been used to measure biomarkers of prenatal and postnatal exposure to an expanding range of additional toxicants. Among them have been other metals, including mercury, manganese and cadmium, as well as mixtures of metals.⁶⁴ For example, mean levels of mercury in whole blood among ELEMENT mothers during pregnancy, cord blood and offspring children were three to five times greater than values reported in population surveys from the USA and Canada,⁶⁵ with higher rates of consuming tuna and shark as suspected risk factors. Early childhood blood levels of manganese exhibited an inverted U-shaped relationship with mental development

scores.⁶⁶ Sources of and descriptive statistics surrounding dietary cadmium have also been reported in mothers and children and related to lower adiposity among girls⁶⁷ (see also, Diet).

Endocrine disrupting chemicals

ELEMENT investigators have also been examining exposures and impacts related to organics associated with plastics. So far, our studies indicate sex-specific effects of exposure to EDCs on metabolism. In boys, prenatal and concurrent exposure to certain phthalates predicted a more favourable lipid profile⁶⁸ and lower adiposity⁶⁹ during peripuberty. These unexpected results may reflect residual variability in pubertal progression. On the contrary, concurrent exposure to bisphenol A (BPA) was associated with subcutaneous adiposity in girls.⁶⁹ In studies of sexual maturation using physician-assessed Tanner staging and serum levels of sex hormones,⁷⁰ prenatal exposure to phthalates correlated with slower pubertal development in boys.^{71 72} The opposite trend was observed in girls.^{73 74} Generally, cross-sectional relationships between peripubertal exposure to these EDCs and reproductive development were less robust than perinatal exposure.^{71 74}

Pesticides

Prenatal exposure to the pesticide chlorpyrifos, as reflected by measurement of 3, 5, 6-trichloro-2-pyridinol in maternal urine samples during pregnancy as a biomarker, was found to occur at levels moderately higher than those in the USA and to also be associated with an increased risk of symptoms of attention deficit hyperactivity disorder (ADHD).⁷⁵

Fluoride

Finally, a programme of research on the potential neurotoxicity of early life exposure to fluoride was recently started, with initial findings suggesting that prenatal fluoride exposure is associated with reduced measures of intelligence⁷⁶ as well as increased symptoms of ADHD in pre-school and elementary school age children.⁷⁷ Additionally, urinary fluoride concentrations were cross-sectionally associated with later pubic hair growth and genital development in boys aged 10–17 years, but not significantly related to pubertal development in girls.⁷⁸

Diet

A key feature of ELEMENT is the repeated dietary assessments in the mothers during pregnancy, in the mothers on behalf of their children during infancy and early childhood and in the children themselves from peripuberty onward. The associated data have been used to examine dietary influences on toxicant absorption and/or distribution, such as the finding that increased levels of maternal blood and bone lead are associated with low dietary calcium.⁴⁵ A recent study reported positive correlations between intake of potatoes and urinary cadmium concentrations in children and positive correlations between dietary and urinary cadmium concentrations among

mother–child dyads who were not exposed to smoking during pregnancy.⁷⁹

Dietary factors have also been evaluated as modifiers of toxicant impacts. For example, child lead exposure was associated with lower stature, but only among individuals with zinc deficiency.⁸⁰ In our more recent work, dietary factors are also being examined as primary determinant of outcomes of interest. We found that higher cumulative intake of sugar-sweetened beverage (SSB) during the first year of life was a determinant of abdominal and overall adiposity at 8–14 years.⁸¹ During peripuberty, adherence to a prudent dietary pattern was protective against metabolic risk in boys, whereas a dietary pattern indicative of a transition towards a Westernised diet corresponded with higher adiposity among girls.⁸²

Mechanisms and mediators

Epigenetics and epigenomics

DNA methylation has surfaced as a key mechanism linking early life exposures to later health outcomes. Higher maternal patella lead corresponded with higher methylation of genes involved in growth and metabolism in girls (cord blood *IGF2* and *HSD11B2*).⁸³ Expanding our analyses to additional environmental exposures—namely, BPA and phthalates—during multiple sensitive periods (in utero, early childhood, peripuberty) revealed differential relations of each exposure with methylation of several candidate genes during peripuberty.⁸⁴ The extent to which differential methylation of these regions mediate exposure/outcome relationships⁸⁵ and the role of environmental deflection of age-related DNA methylation changes⁸⁶ are upcoming areas of interest.

Metabolomics

The ELEMENT cohort has untargeted metabolomics data assayed from fasting blood collected in ELEMENT 2011 and 2015, as well as targeted metabolomics data on amino acids and acylcarnitines in ELEMENT 2011. So far, we have identified compounds involved in lipid, amino acid and DNA pathways that were associated with higher metabolic syndrome risk in ELEMENT 2011,⁸⁷ as well as some metabolites associated with SSB intake that are also markers of elevated blood pressure during adolescence.⁸⁸

Ethnography

In the ELEMENT 2014 ('Mexican Exposures'), intensive ethnographic observation of six families in two distinct Mexico City neighbourhoods emphasised the importance of neighbourhood environments for health.⁸⁹ Our work in this area provides insights into how neoliberal economic processes (eg, the North American Free Trade Agreement) transformed the food landscape in Mexico to one composed primarily of processed foods. We have also unveiled a key role of the extended family household structure in diet-related factors, from food preparation to mealtime behaviours.⁸⁹

Statistical methods development

Given the complex data structure in the ELEMENT study, including longitudinal follow-up, multiple exposures and high-dimensional data, ELEMENT has served as fertile ground for statistical methods development and innovations in data sharing. To date, methods arising from ELEMENT data include: structural equation models^{90–93}; models to identify timing of vulnerability⁹⁴; methods for gene–environment interactions in the context of multiple correlated exposures^{95–96}; methods to identify complex interactions among multiple exposures,⁹⁷ metabolomics determinants of metabolic risk,⁸⁷ methods to characterise growth trajectories^{20–98} and derive dietary patterns¹⁹; and methods for data integration across multiple cohorts.^{99–100} Current and ongoing work involves the use of novel dimension reduction and simulation techniques to analyse accelerometry and dental imaging data.

FUTURE RESEARCH

In addition to continuing research involving the long-term follow-up of the mothers and children recruited as part of the original three cohorts, future plans include recruitment of grandchildren given that the original ‘child’ participants are now becoming parents. We recently received funds from a cohort maintenance grant to maintain and enrich resource infrastructure for existing environmental epidemiology cohorts. The overarching goals of this proposal are to investigate associations of exposure to EDCs during multiple sensitive periods of development from conception through middle-age with metabolic and reproductive health, to elucidate both within-generational and inter-generational biological pathways via epigenetic and microbiome pathways and to develop novel data management techniques and methods to improve data integration, data-sharing and cross-institute data communication to enrich the existing ELEMENT database. In parallel, we will collect data from the original mothers (ie, women in cohorts 1–3) on their cardiometabolic health as they progress through the perimenopausal transition in order to gain insight into the effects of toxicant exposure during pregnancy on long-term postpartum health.

STRENGTHS AND LIMITATIONS

Key strengths of ELEMENT include in-person collection of exposure and outcome measures using research-quality standard protocols, an extensive biorepository of specimens from gestation through adolescence, rich covariate data including physician-assessed pubertal status, newly collected high-dimensional ‘omics data and ethnographical data—all within a long-term study of early lead exposure that will soon span three generations. Together, these data make ELEMENT uniquely poised to carry out a broad range of research, from technical multi-‘omics studies to mixed-methods

analyses leveraging the ethnography data. Research projects within ELEMENT are predominantly led by students and trainees in the USA and Mexico, thereby forging fruitful international collaborations and mentorship.

This cohort also has several weaknesses. First, we have limited information on maternal perinatal conditions, with data on prepregnancy BMI in a small subset of mothers and no information on maternal glycaemia. Thus, in analyses where these characteristics may confound associations of interest, we acknowledge the potential for unmeasured confounding by maternal weight status and/or gestational glucose tolerance,¹⁰¹ and in more recent in-progress analyses, we use first trimester BMI as a proxy for prepregnancy BMI given evidence of a decent correlation between the two measures.¹⁰² Second, we have a limited remaining volume of early archived biosamples to explore new questions given the ongoing and long-term nature of ELEMENT. Third, we have limited information on the children’s biological fathers, which restricts our ability to study the contribution of paternal characteristics to developmental origins of health and disease. Finally, given that our cohort is entirely Hispanic, our findings may not be generalisable to other race/ethnicities—a weakness that we are careful to mention with each study.

COLLABORATION

Interested investigators should submit a proposal including an analysis plan to the core group of ELEMENT co-investigators, which meets bimonthly. The discussion among the co-investigator team ensures that the outside investigator has adequate knowledge of the data, understands how the proposed analysis relates to previous analyses to avoid scientific overlap between projects and optimise use of scarce resources. Once an analysis plan is approved, the data manager creates a deidentified dataset for secondary analysis. We encourage collaborations that make use of the unique structure of ELEMENT comprising long-term follow-up of a well-characterised mother–child population in a setting with relatively high toxicant exposure. Investigators interested in working with ELEMENT data can contact the ELEMENT PIs KEP at karenep@umich.edu and M-MT-R at mmtellez@insp.mx.

Author affiliations

¹Department of Epidemiology, Colorado School of Public Health, University of Colorado Denver Anschutz Medical Center, Aurora, Colorado, USA

²Department of Nutritional Sciences, University of Michigan School of Public Health, Ann Arbor, Michigan, USA

³Lifecourse Epidemiology of Adiposity and Diabetes (LEAD) Center, Colorado School of Public Health, University of Colorado Denver Anschutz Medical Center, Aurora, Colorado, USA

⁴National Council of Science and Technology, National Institute of Public Health, Mexico City, Mexico

⁵Department of Biostatistics, University of Michigan School of Public Health, Ann Arbor, Michigan, USA

⁶Department of Environmental Health Sciences, University of Michigan School of Public Health, Ann Arbor, Michigan, USA

⁷Center for Human Growth and Development, University of Michigan, Ann Arbor, MI, United States

⁸Department of Anthropology, University of Michigan, Ann Arbor, Michigan, USA

⁹Department of Cariology, Operative Dentistry and Dental Public Health, Indiana University School of Dentistry, Indianapolis, Indiana, USA

¹⁰Center for Population Health Research, National Institute of Public Health, Cuernavaca, Mexico

¹¹Department of Environmental Medicine & Public Health, Icahn School of Medicine at Mount Sinai Hospital, New York, New York, USA

¹²Division of Research in Community Interventions, Instituto Nacional de Perinatología, Mexico City, Mexico

¹³Center for Nutrition and Health Research, National Institute of Public Health, Cuernavaca, Mexico

¹⁴Department of Environmental and Occupational Health, University of Washington School of Public Health, Seattle, Washington, USA

¹⁵Dirección de Prestaciones Económicas y Sociales, Mexican Institute of Social Security, Mexico City, Mexico

Acknowledgements We are grateful for Drs Don Smith and Teresita Gonzalez de Cossio for their contributions to the ELEMENT cohorts and the American British Cowdray Medical Centre in Mexico City for providing us with research facilities.

Contributors WP and MT-0 conceived manuscript idea, wrote the initial draft of the paper and incorporated co-author comments. HH, MH-A, M-MT-R and KEP contributed funding, reagents and materials for original three cohorts and major follow-up studies (ELEMENT 2011 and 2015). LT, BS, AC, JM, DD, ER, EAM-M, HL-F, PS, AE, RW, MA, LS, DW, JG, RL, MS-G, LFB-A, AM-G, HH, MH-A, M-MT-R and KEP read various drafts of the manuscript, provided critical feedback and approved the final version for submission.

Funding This work was supported by the National Institutes of Environmental Health Sciences at the National Institutes of Health (P42ES05947, K23ES000381, P01ES022844, P30DK089503, P30ES00002, R01ES014930, R01ES07821, R24DK097153, R24ES028502, R01ES024732, R01ES021446, P30ES017885, R01ES013744, K01ES014907, K01ES016587, P42ES016454), the US Environmental Protection Agency (RD83543601); Consejo Nacional de Ciencia y Tecnología (106565, 261063, 29192-M, 37210-M, 41912-M); and Consejo de Estudios para la Restauración y Valoración Ambiental (CONSERVA), Departamento del Distrito Federal, México. National Science Foundation (NSF) 1744724 and 1430391. The funders did not play any role in the conception or writing of this paper (or any others from the ELEMENT Project).

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Bellinger D, Leviton A, Waternaux C, *et al*. Longitudinal analyses of prenatal and postnatal lead exposure and early cognitive development. *N Engl J Med* 1987;316:1037–43.
- Cummins SK, Goldman LR. Even advantaged children show cognitive deficits from low-level lead toxicity. *Pediatrics* 1992;90:995–7.
- Fergusson DM, Horwood LJ. The effects of lead levels on the growth of word recognition in middle childhood. *Int J Epidemiol* 1993;22:891–7.
- Needleman HL, Schell A, Bellinger D, *et al*. The long-term effects of exposure to low doses of lead in childhood. An 11-year follow-up report. *N Engl J Med* 1990;322:83–8.
- Silbergeld EK. Lead in bone: implications for toxicology during pregnancy and lactation. *Environ Health Perspect* 1991;91:63–70.
- Gulson BL, Mahaffey KR, Jameson CW, *et al*. Mobilization of lead from the skeleton during the postnatal period is larger than during pregnancy. *J Lab Clin Med* 1998;131:324–9.
- Smith DR, Flegal AR. Lead in the biosphere: recent trends. *Ambio* 1995;24:21–3.
- Tong S, von Schirnding YE, Prapamontol T. Environmental lead exposure: a public health problem of global dimensions. *Bull World Health Organ* 2000;78:1068–77.
- Driscoll W, Mushak P, Garfias J, *et al*. Reducing lead in gasoline. Mexico's experience. *Environ Sci Technol* 1992;26:1702–5.
- Gulson BL, Jameson CW, Mahaffey KR, *et al*. Pregnancy increases mobilization of lead from maternal skeleton. *J Lab Clin Med* 1997;130:51–62.
- Lagerkvist BJ, Ekesrydh S, Englyst V, *et al*. Increased blood lead and decreased calcium levels during pregnancy: a prospective study of Swedish women living near a smelter. *Am J Public Health* 1996;86:1247–52.
- Barker DJ, Gluckman PD, Godfrey KM, *et al*. Fetal nutrition and cardiovascular disease in adult life. *Lancet* 1993;341:938–41.
- Barker DJ, Osmond C. Infant mortality, childhood nutrition, and ischaemic heart disease in England and Wales. *Lancet* 1986;1:1077–81.
- Barker DJ, Winter PD, Osmond C, *et al*. Weight in infancy and death from ischaemic heart disease. *Lancet* 1989;2:577–80.
- Weinhold B. Lead in Mexican children: Pottery use slows reductions in blood. *Environ Health Perspect* 2004;112:A569–A.
- Schnaas L, Rothenberg SJ, Flores M-F, *et al*. Blood lead secular trend in a cohort of children in Mexico City (1987–2002). *Environ Health Perspect* 2004;112:1110–5.
- Hu H, Milder FL, Burger DE. The use of K x-ray fluorescence for measuring lead burden in epidemiological studies: high and low lead burdens and measurement uncertainty. *Environ Health Perspect* 1991;94:107–10.
- Hernandez-Avila M, Gonzalez-Cossio T, Hernandez-Avila JE, *et al*. Dietary calcium supplements to lower blood lead levels in lactating women: a randomized placebo-controlled trial. *Epidemiology* 2003;14:206–12.
- Jansen EC, Zhou L, Perng W, *et al*. Vegetables and lean proteins-based and processed meats and refined grains -based dietary patterns in early childhood are associated with pubertal timing in a sex-specific manner: a prospective study of children from Mexico City. *Nutr Res* 2018;56:41–50.
- Baek J, Zhu B, Song P. Bayesian analysis of infant's growth dynamics with in utero exposure to environmental toxicants. *Ann Appl Stat* 2019;13:297–320.
- Tang L, Zhou L, Song P. Fusion learning algorithm to combine partially heterogeneous COX models. *Comput Stat* 2019;34:395–414.
- Tang L, Song P. Fused LASSO approach in regression coefficients clustering learning parameter heterogeneity in data integration. *Journal of machine learning research : JMLR* 2016;17:1–23.
- Salud Sde, ambiental S. Limitaciones Y especificaciones sanitarias para El uso de Los compuestos de plomo 2000.
- Salud Sde, ambiental S. Limitaciones Y especificaciones sanitarias para El uso de Los compuestos de plomo 2013.
- American Academy of Pediatrics Committee on Environmental Health. Lead exposure in children: prevention, detection, and management. *Pediatrics* 2005;116:1036–46.
- U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. *Preventing lead poisoning in young children*. Atlanta: CDC, 2005.
- Centers for Disease Control and Prevention. *Interpreting and managing blood lead levels <10 ug/dL in children and reducing childhood exposures to lead*. CDC, 2007Contract No: RR08.
- Centers for Disease Control and Prevention. *Guidelines for the identification and management of lead exposure in pregnant and lactating women*. Atlanta: U.S. Department of Health and Human Services, 2010.
- Roberts EFS. Bio-ethnography: a collaborative, methodological experiment in Mexico City: Somatosphere, 2015. Available: <http://somatosphere.net/2015/02/bio-ethnography.html> [Accessed cited 19 Feb 2018].
- Roberts EFS, Sanz C. Bioethnography: a how-to guide for the twentieth century. In: Meloni M, ed. *The Palgrave Handbook of biology and society*. London: Springer, 2017.
- Ettinger AS, Téllez-Rojo MM, Amarasiriwardena C, *et al*. Influence of maternal bone lead burden and calcium intake on levels of lead in breast milk over the course of lactation. *Am J Epidemiol* 2006;163:48–56.
- Smith D, Hernandez-Avila M, Téllez-Rojo MM, *et al*. The relationship between lead in plasma and whole blood in women. *Environ Health Perspect* 2002;110:263–8.
- Lamadrid-Figueroa H, Téllez-Rojo MM, Hernández-Avila M, *et al*. Association between the plasma/whole blood lead ratio and history

- of spontaneous abortion: a nested cross-sectional study. *BMC Pregnancy Childbirth* 2007;7:22.
34. Lamadrid-Figueroa H, Téllez-Rojo MM, Hernández-Cadena L, et al. Biological markers of fetal lead exposure at each stage of pregnancy. *J Toxicol Environ Health A* 2006;69:1781–96.
 35. Ettinger AS, Téllez-Rojo MM, Amarasiwardena C, et al. Levels of lead in breast milk and their relation to maternal blood and bone lead levels at one month postpartum. *Environ Health Perspect* 2004;112:926–31.
 36. Aro AC, Todd AC, Amarasiwardena C, et al. Improvements in the calibration of 109Cd K x-ray fluorescence systems for measuring bone lead in vivo. *Phys Med Biol* 1994;39:2263–71.
 37. Hernández-Avila M, Romieu I, Parra S, et al. Validity and reproducibility of a food frequency questionnaire to assess dietary intake of women living in Mexico City. *Salud Publica Mex* 1998;40:133–40.
 38. B N. *Bayley scales of infant development manual*. 2 ed. San Antonio, TX: The Psychological Corporation, 1993.
 39. M D. *McCarthy scales of children abilities*. Madrid, Spain: TEA ediciones, 2009.
 40. Wechsler D. *WAIS-R: Wechsler adult intelligence Scale-Revised*. New York, NY: Psychological Corporation, 1981.
 41. Behavior assessment system for children. The Corsini encyclopedia of psychology:1–2.
 42. Huang S, Hu H, Sánchez BN, et al. Childhood blood lead levels and symptoms of attention deficit hyperactivity disorder (ADHD): a cross-sectional study of Mexican children. *Environ Health Perspect* 2016;124:868–74.
 43. Brown MJ, Hu H, Gonzales-Cossio T, et al. Determinants of bone and blood lead concentrations in the early postpartum period. *Occup Environ Med* 2000;57:535–41.
 44. Téllez-Rojo MM, Hernández-Avila M, Lamadrid-Figueroa H, et al. Impact of bone lead and bone resorption on plasma and whole blood lead levels during pregnancy. *Am J Epidemiol* 2004;160:668–78.
 45. Hernandez-Avila M, Gonzalez-Cossio T, Palazuelos E, et al. Dietary and environmental determinants of blood and bone lead levels in lactating postpartum women living in Mexico City. *Environ Health Perspect* 1996;104:1076–82.
 46. Téllez-Rojo MM, Hernández-Avila M, González-Cossio T, et al. Impact of breastfeeding on the mobilization of lead from bone. *Am J Epidemiol* 2002;155:420–8.
 47. González-Cossio T, Peterson KE, Sanín LH, et al. Decrease in birth weight in relation to maternal bone-lead burden. *Pediatrics* 1997;100:856–62.
 48. Sanín LH, González-Cossio T, Romieu I, et al. Effect of maternal lead burden on infant weight and weight gain at one month of age among breastfed infants. *Pediatrics* 2001;107:1016–23.
 49. Ettinger AS, Roy A, Amarasiwardena CJ, et al. Maternal blood, plasma, and breast milk lead: lactational transfer and contribution to infant exposure. *Environ Health Perspect* 2014;122:87–92.
 50. Janakiraman V, Ettinger A, Mercado-Garcia A, et al. Calcium supplements and bone resorption in pregnancy: a randomized crossover trial. *Am J Prev Med* 2003;24:260–4.
 51. Ettinger AS, Hu H, Hernandez-Avila M. Dietary calcium supplementation to lower blood lead levels in pregnancy and lactation. *J Nutr Biochem* 2007;18:172–8.
 52. Gomaa A, Hu H, Bellinger D, et al. Maternal bone lead as an independent risk factor for fetal neurotoxicity: a prospective study. *Pediatrics* 2002;110:110–8.
 53. Téllez-Rojo MM, Bellinger DC, Arroyo-Quiroz C, et al. Longitudinal associations between blood lead concentrations lower than 10 microg/dL and neurobehavioral development in environmentally exposed children in Mexico City. *Pediatrics* 2006;118:e323–30.
 54. Braun JM, Hoffman E, Schwartz J, et al. Assessing windows of susceptibility to lead-induced cognitive deficits in Mexican children. *Neurotoxicology* 2012;33:1040–7.
 55. Hu H, Téllez-Rojo MM, Bellinger D, et al. Fetal lead exposure at each stage of pregnancy as a predictor of infant mental development. *Environ Health Perspect* 2006;114:1730–5.
 56. Surkan PJ, Schnaas L, Wright RJ, et al. Exposure to lead, and child neurodevelopment. *Neurotoxicology* 2008;29:278–85.
 57. Kordas K, Ettinger AS, Bellinger DC, et al. A dopamine receptor (DRD2) but not dopamine transporter (DAT1) gene polymorphism is associated with neurocognitive development of Mexican preschool children with lead exposure. *J Pediatr* 2011;159:638–43.
 58. Pilsner JR, Hu H, Wright RO, et al. Maternal MTHFR genotype and haplotype predict deficits in early cognitive development in a lead-exposed birth cohort in Mexico City. *Am J Clin Nutr* 2010;92:226–34.
 59. Afeiche M, Peterson KE, Sánchez BN, et al. Prenatal lead exposure and weight of 0- to 5-year-old children in Mexico City. *Environ Health Perspect* 2011;119:1436–41.
 60. Liu Y, Peterson KE, Montgomery K, et al. Early lead exposure and childhood adiposity in Mexico City. *Int J Hyg Environ Health* 2019;222:965–70.
 61. Zhang A, Hu H, Sánchez BN, et al. Association between prenatal lead exposure and blood pressure in children. *Environ Health Perspect* 2012;120:445–50.
 62. Afeiche M, Peterson KE, Sánchez BN, et al. Windows of lead exposure sensitivity, attained height, and body mass index at 48 months. *J Pediatr* 2012;160:1044–9.
 63. Jansen EC, Zhou L, Song PXX, et al. Prenatal lead exposure in relation to age at menarche: results from a longitudinal study in Mexico City. *J Dev Orig Health Dis* 2018;9:467–72.
 64. Lewis RC, Meeker JD, Basu N, et al. Urinary metal concentrations among mothers and children in a Mexico City birth cohort study. *Int J Hyg Environ Health* 2018;221:609–15.
 65. Basu N, Tutino R, Zhang Z, et al. Mercury levels in pregnant women, children, and seafood from Mexico City. *Environ Res* 2014;135:63–9.
 66. Claus Henn B, Schnaas L, Ettinger AS, et al. Associations of early childhood manganese and lead coexposure with neurodevelopment. *Environ Health Perspect* 2012;120:126–31.
 67. Moynihan M, Téllez-Rojo MM, Colacino J, et al. Prenatal cadmium exposure is negatively associated with adiposity in girls not boys during adolescence. *Front Public Health* 2019;7.
 68. Perng W, Watkins DJ, Cantoral A, et al. Exposure to phthalates is associated with lipid profile in peripubertal Mexican youth. *Environ Res* 2017;154:311–7.
 69. Yang TC, Peterson KE, Meeker JD, et al. Bisphenol A and phthalates in utero and in childhood: association with child BMI z-score and adiposity. *Environ Res* 2017;156:326–33.
 70. Chavarro JE, Watkins DJ, Afeiche MC, et al. Validity of Self-Assessed sexual maturation against physician assessments and hormone levels. *J Pediatr* 2017;186:172–8.
 71. Ferguson KK, Peterson KE, Lee JM, et al. Prenatal and peripubertal phthalates and bisphenol A in relation to sex hormones and puberty in boys. *Reprod Toxicol* 2014;47:70–6.
 72. Watkins DJ, Sánchez BN, Téllez-Rojo MM, et al. Impact of phthalate and BPA exposure during in utero windows of susceptibility on reproductive hormones and sexual maturation in peripubertal males. *Environ Health* 2017;16.
 73. Watkins DJ, Sánchez BN, Téllez-Rojo MM, et al. Phthalate and bisphenol a exposure during in utero windows of susceptibility in relation to reproductive hormones and pubertal development in girls. *Environ Res* 2017;159:143–51.
 74. Watkins DJ, Téllez-Rojo MM, Ferguson KK, et al. In utero and peripubertal exposure to phthalates and BPA in relation to female sexual maturation. *Environ Res* 2014;134:233–41.
 75. Fortenberry GZ, Meeker JD, Sánchez BN, et al. Urinary 3,5,6-trichloro-2-pyridinol (TCPY) in pregnant women from Mexico City: distribution, temporal variability, and relationship with child attention and hyperactivity. *Int J Hyg Environ Health* 2014;217:405–12.
 76. Bashash M, Thomas D, Hu H, et al. Prenatal fluoride exposure and cognitive outcomes in children at 4 and 6–12 years of age in Mexico. *Environ Health Perspect* 2017;125:097017.
 77. Bashash M, Marchand M, Hu H, et al. Prenatal fluoride exposure and attention deficit hyperactivity disorder (ADHD) symptoms in children at 6–12 years of age in Mexico City. *Environ Int* 2018;121:658–66.
 78. Liu Y, Téllez-Rojo M, Hu H, et al. Fluoride exposure and pubertal development in children living in Mexico City. *Environmental Health* 2019;18.
 79. Moynihan M, Peterson KE, Cantoral A, et al. Dietary predictors of urinary cadmium among pregnant women and children. *Sci Total Environ* 2017;575:1255–62.
 80. Cantoral A, Téllez-Rojo MM, Levy TS, et al. Differential association of lead on length by zinc status in two-year old Mexican children. *Environ Health* 2015;14.
 81. Cantoral A, Téllez-Rojo MM, Ettinger AS, et al. Early introduction and cumulative consumption of sugar-sweetened beverages during the pre-school period and risk of obesity at 8–14 years of age. *Pediatr Obes* 2016;11:68–74.
 82. Perng W, Fernandez C, Peterson KE, et al. Dietary patterns exhibit sex-specific associations with adiposity and metabolic risk in a cross-sectional study in urban Mexican adolescents. *J Nutr* 2017;147:1977–85.
 83. Goodrich JM, Sánchez BN, Dolinoy DC, et al. Quality control and statistical modeling for environmental epigenetics: a study on in

- utero lead exposure and DNA methylation at birth. *Epigenetics* 2015;10:19–30.
84. Goodrich JM, Dolinoy DC, Sánchez BN, *et al.* Adolescent epigenetic profiles and environmental exposures from early life through peri-adolescence. *Environ Epigenet* 2016;2:dvw018–dvw.
 85. Bowman A, Peterson KE, Dolinoy DC, *et al.* Phthalate exposures, DNA methylation and adiposity in Mexican children through adolescence. *Front Public Health* 2019;7.
 86. Kochmanski J, Montrose L, Goodrich JM, *et al.* Environmental deflection: the impact of toxicant exposures on the aging epigenome. *Toxicol Sci* 2017;156:325–35.
 87. Perng W, Hector EC, Song PXX, *et al.* Metabolomic determinants of metabolic risk in Mexican adolescents. *Obesity* 2017;25:1594–602.
 88. Perng W, Tang L, Song PXX, *et al.* Urate and nonanoate mark the relationship between sugar-sweetened beverage intake and blood pressure in adolescent girls: a metabolomics analysis in the element cohort. *Metabolites* 2019;9:100.
 89. Roberts EFS. What gets inside: violent entanglements and toxic boundaries in Mexico City. *Cultural Anthropology* 2017;32:592–619.
 90. Sánchez BN, Budtz-Jørgensen E, Ryan LM. An estimating equations approach to fitting latent exposure models with longitudinal health outcomes. *Ann Appl Stat* 2009;3:830–56.
 91. Sánchez BN, Budtz-Jørgensen E, Ryan LM, *et al.* Structural equation models. *J Am Stat Assoc* 2005;100:1443–55.
 92. Sánchez BN, Houseman EA, Ryan LM. Residual-based diagnostics for structural equation models. *Biometrics* 2009;65:104–15.
 93. Sánchez BN, Kim S, Sammel MD. Estimators for longitudinal latent exposure models: examining measurement model assumptions. *Stat Med* 2017;36:2048–66.
 94. Sánchez BN, Hu H, Litman HJ, *et al.* Statistical methods to study timing of vulnerability with sparsely sampled data on environmental toxicants. *Environ Health Perspect* 2011;119:409–15.
 95. Sánchez BN, Kang S, Mukherjee B. A latent variable approach to study gene-environment interactions in the presence of multiple correlated exposures. *Biometrics* 2012;68:466–76.
 96. Tao Y, Sánchez BN, Mukherjee B. Latent variable models for gene-environment interactions in longitudinal studies with multiple correlated exposures. *Stat Med* 2015;34:1227–41.
 97. Ma S, Song PX-K. Varying index coefficient models. *J Am Stat Assoc* 2015;110:341–56.
 98. Perng W, Baek J, Zhou CW, *et al.* Associations of the infancy body mass index peak with anthropometry and cardiometabolic risk in Mexican adolescents. *Ann Hum Biol* 2018;1–9.
 99. Wang F, Song PX-K, Wang L. Merging multiple longitudinal studies with study-specific missing covariates: a joint estimating function approach. *Biometrics* 2015;71:929–40.
 100. Tang L, Song PXX. Fused Lasso Approach in Regression Coefficients Clustering - Learning Parameter Heterogeneity in Data Integration. *J Mach Learn Res* 2016;17.
 101. Carter M. *Associations of maternal carbohydrate intake during pregnancy and adolescent adiposity and metabolic health* [Master of Science]. Ann Arbor, MI: University of Michigan School of Public Health, 2018.
 102. Krukowski RA, West DS, DiCarlo M, *et al.* Are early first trimester weights valid proxies for preconception weight? *BMC Pregnancy Childbirth* 2016;16:357.