

Carbon Dioxide Uptake

International Edition: DOI: 10.1002/anie.201916483
German Edition: DOI: 10.1002/ange.201916483

Effective and Reversible Carbon Dioxide Insertion into Cerium Pyrazolates


Uwe Bayer, Daniel Werner, Cécilia Maichle-Mössmer, and Reiner Anwander*

Abstract: The homoleptic pyrazolate complexes $[Ce^{III}_4(Me_2pz)_{12}]$ and $[Ce^{IV}(Me_2pz)_4]_2$ quantitatively insert CO_2 to give $[Ce^{III}_4(Me_2pz \cdot CO_2)_{12}]$ and $[Ce^{IV}(Me_2pz \cdot CO_2)_4]$, respectively ($Me_2pz = 3,5$ -dimethylpyrazolato). This process is reversible for both complexes, as observed by *in situ* IR and NMR spectroscopy in solution and by TGA in the solid state. By adjusting the molar ratio, one molecule of CO_2 per $[Ce^{IV}(Me_2pz)_4]$ complex could be inserted to give trimetallic $[Ce_3(Me_2pz)_9(Me_2pz \cdot CO_2)_3(thf)]$. Both the cerous and ceric insertion products catalyze the formation of cyclic carbonates from epoxides and CO_2 under mild conditions. In the absence of epoxide, the ceric catalyst is prone to reduction by the cocatalyst tetra-*n*-butylammonium bromide (TBAB).

Introduction

Inexorably rising CO_2 levels in the earth's atmosphere—and their consequential environmental impact—have spurred much interest in combating CO_2 build-up.^[1] Capture technologies, such as carbon dioxide capture and storage (CCS) and direct air capture (DAC),^[2] and CO_2 conversion into fuels or chemical feedstocks^[3] appear promising. However, such tactics suffer from either a lack of appropriate storage and transportation of CO_2 , or overcoming the high activation barrier of CO_2 .^[4] To date, the most effective sorbents for CCS/DAC are alkali-metal/alkaline-earth metal hydroxide solutions, inorganic salts (e.g. alkali-metal carbonates),^[1,2] or high-surface supported polyamines (max. CO_2 sorption capacity ca. 3 mmol g^{-1} at 1 bar)^[5] and magnesium-based metal-organic frameworks (ca. 35 wt % or $8 \text{ mmol } CO_2 \text{ g}^{-1}$ at 1 bar).^[6]

Like alkaline-earth metals, rare-earth metals (Ln) feature a high affinity for carbon dioxide (cf. bastnaesite is the most important Ln^{III} deposit in the Earth's crust). Thus, metal-organic derivatives easily react with or insert CO_2 , as initially demonstrated by Bochkarev et al. for homoleptic silylamides



Scheme 1. Irreversible reaction of carbon dioxide with archetypal organo-rare-earth-metal complexes, with the exception of (e) as shown for $(C_5Me_5)_2Sm(SePh)(thf)$.^[22]

$Ln[N(SiMe_3)_2]_3$ ^[7,8] and alkoxides $[Ln(OnBu)_3]$ (Scheme 1 c,d).^[9,10]

Similar archetypes (including Ln^{III} aryloxides) can also be used for chemical transformations, such as the catalytic conversion of a CO_2 /epoxide mixture into cyclic carbonates^[11] or copolymers.^[10a,12,13] However, highly reactive organo-rare-earth-metal complexes such as alkyl^[14] and hydride^[10a,15] (Scheme 1 a,b) or divalent derivatives^[16] display irreversible CO_2 insertion or favor additional transformations through CO_2 post-activation (e.g. formation of CO , CO_3^{2-} , $C_2O_4^{2-}$).^[16] Recently, cerium, the most abundant rare-earth element, has gained attention for CO_2 activation.^[8,17–19] For example, while the hydrogen-bonded Ce^{IV} oxo complex $[(L_{OEt})_2Ce=O(H_2O)] \cdot MeC(O)NH_2$ ($L_{OEt}^- = [Co(\eta^5-C_5H_5)\{P(O)(OEt)_2\}_3]^-$) was shown to form the tetravalent carbonate species $[(L_{OEt})_2Ce(CO_3)]$,^[17] *ortho*-NHC-substituted aryloxide Ce^{III} complexes (NHC = N-heterocyclic carbene) insert CO_2 into the $Ce-C_{NHC}$ bond in a semireversible manner, and catalytically form propylene carbonate from propylene oxide.^[18,19]

Bulky cyclopentadienyl (Cp) derivatives (e.g. $Ln-(C_5Me_5)_3$) were shown to accommodate CO_2 insertion in a unidirectional manner, thereby forming very stable carboxylato moieties through a η^5 -to- η^1 switch in the C_5Me_5 coordination (cf. Scheme 1 a).^[20] Pyrazolates (pz), on the other hand, are dinitrogen-derived Cp counterparts, where the putative $N-CO_2$ bond may tolerate a more reversible insertion process, as seen for other CO_2 -heteroatom bonds (Scheme 1 e).^[21,22] As the tetravalent $[Ce(Me_2pz)_4]_2$ complex was recently shown to undergo reversible insertion of ketones into the $Ce-N$ bond,^[23] we extended the study toward CO_2 . Quantitative insertion of CO_2 into the $Ce-N(Me_2pz)$ bond was observed for both tetravalent and trivalent cerium Me_2pz complexes, and intriguingly the insertion process was found to be fully reversible.


[*] U. Bayer, Dr. D. Werner, Dr. C. Maichle-Mössmer, Prof. Dr. R. Anwander
Institut für Anorganische Chemie
Eberhard Karls Universität Tübingen
Auf der Morgenstelle 18, 72076 Tübingen (Germany)
E-mail: reiner.anwander@uni-tuebingen.de

Supporting information and the ORCID identification numbers for some of the authors of this article can be found under: <https://doi.org/10.1002/anie.201916483>.

© 2020 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Results and Discussion

Carbon Dioxide Insertion into a Ceric Pyrazolate: Treatment of $[\text{Ce}(\text{Me}_2\text{pz})_4]_2$ (**1**) with excess CO_2 in either toluene or thf (under 1 bar CO_2 pressure) led to a color change from dark red to orange within 5 minutes (Scheme 2). Crystallization from concentrated toluene or thf solutions at -40°C gave orange crystals of $[\text{Ce}(\text{Me}_2\text{pz}\cdot\text{CO}_2)_4]$ with either toluene (**2-toluene**, 54%) or thf (**2-thf**, 64%) within the lattice. Discounting the lattice solvent, this accounts for about 25 wt % CO_2 or 5.7 mmol CO_2 per gram.



Scheme 2. Insertion of CO_2 into the $\text{Ce}-\text{N}(\text{Me}_2\text{pz})$ bond of $[\text{Ce}(\text{Me}_2\text{pz})_4]_2$ at: a) -40°C in thf or b) ambient temperature in toluene.

The molecular structure of **2-toluene** revealed an 8-coordinate cerium(IV) center with four $\kappa^2(\text{N},\text{O})$ -coordinating $\text{Me}_2\text{pz}\cdot\text{CO}_2$ ligands (Figure 1), in contrast to the $\kappa^2(\text{O},\text{O})$ modes in carboxylates and related carbamates. The $\text{Ce}-\text{N}$ and $\text{Ce}-\text{O}$ bond lengths average 2.528 \AA and 2.255 \AA , respectively, thus matching the values found in the benzophenone-inserted product $[\text{Ce}(\text{Me}_2\text{pz})_2(\text{pdp})_2]$ ($\text{Ce}1-\text{N}1$ 2.564 \AA , $\text{Ce}1-\text{O}1$ 2.173 \AA ; $\text{pdp} = (3,5\text{-dimethylpyrazol-1-yl})\text{diphenylmethanolate}$).^[23] Other homoleptic Ce^{IV} complexes, $[\text{Ce}(\text{L})_4]$ (with L as a donor-functionalized alkoxy ligand engaged in a 5-membered chelate to cerium), also have similar $\text{Ce}-\text{O}$ bond lengths as those in **2**, thus highlighting a common chelating coordination motif.^[24,25] Seemingly, no delocalization occurs across the $\text{O}=\text{C}-\text{O}$ fragment, which exhibits average $\text{C}-\text{O}$ bond lengths of 1.207 \AA (terminal) and 1.291 \AA (bridging). Support for the localization of the $\text{C}-\text{O}$ double bond came from DRIFTS measurements of **2-toluene** and





Figure 1. Crystal structure of $[\text{Ce}(\text{Me}_2\text{pz}\cdot\text{CO}_2)_4]$ (**2-toluene**). Ellipsoids are shown at the 50% probability level. Hydrogen atoms and lattice toluene are omitted for clarity. Selected bond lengths/angles are listed in the Supporting Information.

2-thf, which showed the presence of a strong absorption band at $\tilde{\nu} = 1732$ and 1718 cm^{-1} , respectively, for the CO stretching of the $\text{C}=\text{O}$ double bond as well as a strong absorption band at $\tilde{\nu} = 1336 \text{ cm}^{-1}$ for the CO stretching of the $\text{C}-\text{O}$ single bond.

The structure was also supported by NMR spectroscopic measurements. The ^1H NMR spectrum recorded in $[\text{D}_8]$ toluene at ambient temperature revealed two distinct methyl group environments for all the pyrazolato ligands, indicative of ligand asymmetry and complete consumption of $[\text{Ce}(\text{Me}_2\text{pz})_4]_2$. The ^{13}C signal of the inserted CO_2 was detected at $\delta = 149.9 \text{ ppm}$, a region where pyrazolate $\text{N}-\text{CO}_2\text{R}$ signals are expected.^[26] ^1H NMR measurements on **2-thf** in $[\text{D}_8]$ THF at ambient temperature showed a mixture of products, which could not be assigned. Cooling the solution to -40°C under 1 bar CO_2 pressure led to a color change from red to orange and both the ^1H and ^{13}C NMR spectra recorded at -40°C showed similar signals as **2-toluene** in $[\text{D}_8]$ toluene.

Variable-temperature (VT) NMR studies of **2-toluene** and **2-thf** in $[\text{D}_8]$ toluene and $[\text{D}_8]$ THF were conducted to investigate the reversibility of CO_2 insertion (Scheme 3). In $[\text{D}_8]$ toluene, the formation of a new species at 40°C was revealed and no further liberation of CO_2 was observed even after heating above 60°C (see Figure S6 in the Supporting Information). The ^1H NMR spectrum recorded at 40°C shows two sets of signals for different Me_2pz moieties in a 1:1 ratio, which suggests the formation of putative compound $[\text{Ce}(\text{Me}_2\text{pz})_2(\text{Me}_2\text{pz}\cdot\text{CO}_2)_2]$.

Recooling the solution did not reform **2-toluene** quantitatively, likely because some of the liberated CO_2 was no longer within the reaction medium, but the addition of fresh CO_2 quantitatively reformed **2-toluene**. The $[\text{D}_8]$ THF VT NMR experiment of compound **2-thf** showed a different CO_2 -deinsertion behavior (see Figure S10). As a consequence of competitive thf coordination, displacement of CO_2 starts at 10°C and is complete at 60°C , with formation of $[\text{Ce}(\text{Me}_2\text{pz})_4(\text{thf})]$. As seen in the experiment in $[\text{D}_8]$ toluene, this reaction is fully reversible by recooling the sample and subsequently



Scheme 3. Reversible insertion of CO_2 into the $\text{Ce}-\text{N}(\text{Me}_2\text{pz})$ bond. The process is completely reversible in $[\text{D}_8]$ thf and partially reversible in $[\text{D}_8]$ toluene. The product arising from the elimination of CO_2 in $[\text{D}_8]$ toluene has not been isolated; the structure is based on ^1H NMR spectroscopic analysis at 40°C .





Figure 2. In situ IR spectroscopy of **2-thf** at 60°C in the range of $\tilde{\nu} = 1700$ to 2350 cm^{-1} . Normalized intensities are shown. Spectra were recorded every minute.

introducing CO_2 . Additionally, in situ IR measurements were performed at 60°C, which showed complete loss of inserted CO_2 and formation of free CO_2 (Figure 2 and see also Figure S58).

In the solid state, **2-toluene** is stable for several weeks at -40°C , but at ambient temperature it partially loses CO_2 over a few days or when it is exposed to vacuum, as indicated by a color change from orange to dark red. A thermogravimetric analysis (TGA), performed under a flow of Ar and heating the sample slowly from 28°C to 250°C , indicated an initial loss of mostly lattice toluene. The liberation of CO_2 and small amounts of lattice toluene was dominant between 55 and 95°C , as revealed by a step of 21.92% (theoretical proportion of CO_2 in **2-toluene** 19.98%). At 250°C , only nonvolatile parts of **2-toluene** remain, leaving a mass of 49.79% of the initial weight (theoretical value 51.61%; see Figure S59). Although the deinsertion of carbon dioxide was achieved in the solid state, bulk compound **1** did not insert any carbon dioxide when stored under 1 bar CO_2 pressure for three days. Moreover, compound **1** was hydrolyzed upon exposure to air within one hour (DRIFT spectrum, see Figure S57).

$[\text{Ce}(\text{Me}_2\text{pz})_4(\text{thf})]$ (**1-thf**) was treated with stoichiometric amounts of CO_2 to generate the putative $[\text{Ce}(\text{Me}_2\text{pz})_2(\text{Me}_2\text{pz}\cdot\text{CO}_2)_2]$ (Scheme 3). Although this species could not be isolated, it was possible to generate mono-inserted $[\text{Ce}_3(\text{Me}_2\text{pz})_9(\text{Me}_2\text{pz}\cdot\text{CO}_2)_3(\text{thf})]$ (**3**) in moderate yields of 46% (Scheme 4). The crystal structure of ceric **3** shows a ring motif with two distinct 9-coordinate and one 10-coordinate cerium atoms (Figure 3). Although all the cerium centers are coordinated by three Me_2pz ligands in an $\eta^2(N,N')$ fashion, Ce1 connects further to two oxygen atoms of neighboring $\text{Me}_2\text{pz}\cdot\text{CO}_2$ ligands as well as an additional thf molecule, 10-coordinate Ce2 is surrounded by two $\kappa^2:(N,O)$ -chelating $\text{Me}_2\text{pz}\cdot\text{CO}_2$ ligands, and Ce3 exhibits additional contacts to one $\kappa^2:(N,O)$ - $\text{Me}_2\text{pz}\cdot\text{CO}_2$ ligand and an oxygen atom of a neighboring $\text{Me}_2\text{pz}\cdot\text{CO}_2$ ligand.

Each $\text{Me}_2\text{pz}\cdot\text{CO}_2$ ligand bridges between two cerium atoms. In contrast to homoleptic **2**, all the oxygen atoms are engaged in cerium bonding, which implies delocalized O–C–O bonds (av. C–O, 1.247 \AA). The $\eta^2(N,N')$ -Ce–N(Me_2pz)



Scheme 4. Insertion of one equivalent of CO_2 into the Ce–N(Me_2pz) bond of $[\text{Ce}(\text{Me}_2\text{pz})_4(\text{thf})]$.





Figure 3. Crystal structure of $[\text{Ce}_3(\text{Me}_2\text{pz})_9(\text{Me}_2\text{pz}\cdot\text{CO}_2)_3(\text{thf})]$ (**3**). Ellipsoids are shown at the 50% probability level. Hydrogen atoms and lattice *n*-hexane are omitted for clarity. Selected bond lengths/angles are listed in the Supporting Information.

bond lengths are in the expected range.^[24,27] According to VT NMR studies carried out in $[\text{D}_8]$ toluene, the trimetallic entity **3** is retained in solution at low temperatures, with every dimethylpyrazolato ligand showing a distinct signal set in the proton NMR spectrum at -80°C (see Figure S12). The signals for the protons of the bridging $\text{Me}_2\text{pz}\cdot\text{CO}_2$ ligands are shifted upfield compared to those of the terminal $\eta^2(N,N')$ - Me_2pz ligands. As a consequence of the equilibrium between potential alternative oligomers formed in the presence of only one equivalent of CO_2 per cerium, the interpretation of the ambient-temperature NMR spectrum was difficult. This was already experienced for the insertion of benzophenone into the Ce–N(Me_2pz) bond.^[23] Upon heating to 90°C , CO_2 was liberated and $[\text{Ce}(\text{Me}_2\text{pz})_4]$ re-formed (see Figure S14). After cooling to ambient temperature, a partial reinsertion of CO_2 was observed, as was found in the VT NMR experiments on **2-toluene** and **2-thf**.

Carbon Dioxide Insertion into a Cerous Pyrazolate: To examine the role of the oxidation state of cerium (Ce^{IV} versus Ce^{III}) and, therefore, the impact of its Lewis acidity, cerous donor-free $[\text{Ce}_4(\text{Me}_2\text{pz})_{12}]$ ^[28] (**4**) was used as a precursor for CO_2 insertion (Scheme 5). Remarkably, while retaining the

Ce_4 nuclearity, all of the $Ce-N(Me_2pz)$ moieties engaged in CO_2 insertion, leading to the complex $[Ce_4(Me_2pz-CO_2)_{12}]$ (**5**; Figure 4). Compared to the six different coordination modes of the pyrazolato moieties in starting material **4**,^[28] the crystal structure of **5** revealed only three. Ce_2 , Ce_3 , and Ce_4 form a nearly equilateral triangle bridged by $\mu_3-1\kappa^2(N,O):2\kappa(O):3\kappa(O')$ - Me_2pz-CO_2 . Each of the 9-coordinate cerium centers (Ce_2 , Ce_3 , and Ce_4) is also surrounded by two terminal Me_2pz-CO_2 groups in a $\kappa^2(N,O)$ coordination mode and two oxygen atoms in a $\kappa^2(O,O')$ fashion (Figure S66). This triangle is capped by 9-coordinate Ce_1 , which is coordinated to three Me_2pz-CO_2 ligands in a $\kappa^2(N,O)$ manner (Ce_1-N_{avg} 2.710 Å, Ce_1-O_{avg} 2.395 Å) and to three bridging oxygen atoms. The $Ce-O$ bond lengths are similar to those found in $[Ce(L^R-CO_2)_3]$ ($L^R = 2-O-3,5-tBu_2-C_6H_2(1-C[N(CH)_2N(R)])$ and $R = iPr$ and Mes) reported by Arnold et al. with $Ce-O$ bond lengths of 2.466(6)–2.482(6) Å.^[18b] The $O-C-O$ bond lengths



Scheme 5. Insertion of CO_2 into the $Ce-N(Me_2pz)$ bonds of cerous $[Ce_4(Me_2pz)_{12}]$.




Figure 4. Crystal structure of $[Ce_4(Me_2pz-CO_2)_{12}]$ (**5**). Ellipsoids are shown at the 50% probability level. Hydrogen atoms and lattice toluene (ten molecules) are omitted for clarity. Cutouts of the crystal structure of **5** and a schematic view of different Me_2Pz-CO_2 binding modes, as well as selected bond lengths/angles are shown in the Supporting Information.

indicate delocalization of the bridging Me_2pz-CO_2 ligands, with two $C-O$ bond lengths in the same region (1.235(9)–1.264(9) Å) and rather localized $C-O$ single (1.26(1)–1.301(9) Å) and $C-O$ double bonds (1.21(1)–1.229(9) Å) for the capping and terminal Me_2pz-CO_2 ligands. For further comparison, the cerous carbamate $[Ce_4(O_2CNiPr_2)_{12}]$ features a lozenge arrangement of one 8-coordinate and three 7-coordinate Ce^{III} centers with $Ce-O$ bond lengths in the range 2.322(7)–2.746(7) Å (no $Ce-N$ interaction).^[29] The latter complex was obtained from the reaction of $CeCl_3(DME)$ with $HNiPr_2$ and CO_2 (Scheme 1f). In accordance with the crystal structure of cluster **5**, DRIFTS measurements show strong absorption bands for both the $C-O$ single bonds ($\tilde{\nu} = 1250$ – 1350 cm^{-1}) and $C-O$ double bonds ($\tilde{\nu} = 1600$ – 1750 cm^{-1}). 1H DOSY NMR measurements on **5** in $[D_8]$ toluene, $[D_8]$ THF, or a $[D_8]$ toluene/3,3-dimethyl-1,2-butylene oxide mixture revealed distinct diffusion coefficients (see Figures S17–S20) for the solvents employed and only one additional peak corresponding to a much larger species but correlating with every other signal in the proton NMR spectra. Calculation of the molar mass of this compound ($[D_8]$ toluene: $M_r = 1989\text{ g mol}^{-1}$; $[D_8]$ THF: $M_r = 1643\text{ g mol}^{-1}$; $[D_8]$ toluene + 3,3-dimethyl-1,2-butyleneoxide: $M_r = 2357\text{ g mol}^{-1}$) as a compact sphere-like molecule^[30] suggests it exists as a tetrametallic ($M_r = 2242\text{ g mol}^{-1}$) or a non-monometallic species in solution. Treating $[Ce(Me_2pz)_3(thf)]_2$ with CO_2 in $[D_8]$ THF gave the same NMR spectrum as that of **5-toluene**, thus indicating the formation of a multimetallic compound also in donor solvents (see Figure S16). TGA of **5-toluene** also showed an initial loss of toluene (cf. **2-toluene**), followed by a pronounced step (21.39% weight loss) in the range from 52 to 90°C, consistent with the release of CO_2 (theoretical value: 16.75%) and some lattice toluene (Figure S60). At 250°C, only the nonvolatile parts of **5-toluene** remain and a total loss of 46.82 wt% compared to the starting material fits well with the theoretical value of 45.99% for 10 molecules of toluene and 12 molecules of CO_2 eliminated from $[Ce_4(Me_2pz-CO_2)_{12}] \cdot 10\text{toluene}$ (**5-toluene**).



























































Catalytic Formation of Cyclic Carbonates from CO_2 and Oxiranes: Having established the efficiency and reversibility of CO_2 insertion into $Ce-N(Me_2pz)$ bonds, we were interested in any catalytic utilization. Accordingly, pyrazolate complexes **1** and **4** were probed as catalysts for the generation of cyclic carbonates from CO_2 and oxiranes. In the absence of CO_2 , compound **1** interacts with epoxides, as indicated by a noticeable shift in the 1H NMR spectrum upon addition of one equivalent of 3,3-dimethyl-1,2-butylene oxide (see Figure S24). This is most likely due to formation of a donor adduct, which is considered a crucial step in Lewis-acid-catalyzed cycloaddition reactions. Even though such donor adducts were shown to be isolable (e.g. $Tp^{tBu}Ca(O-2,6-iPr_2C_6H_3) \cdot (PO)$; $Tp^{tBu} = \text{tris}(3-tBu\text{-pyrazolyl})\text{borato}$, $PO = \text{propylene oxide}$ ^[31]), the putative $[Ce(Me_2pz)_4(PO)]$ (**1-PO**) could not be isolated. Tetra-*n*-butylammonium bromide (TBAB) was employed as a co-catalyst, since it was shown to promote the highest activities in such cycloaddition reactions.^[11b,32] The reaction was optimized for propylene oxide, which gave almost quantitative conversion after 24 h under mild conditions. Using 0.5 mol% **1** or 0.25 mol% **4** and

1 mol% TBAB without solvent at ambient temperature and 1 bar CO₂ pressure gave 93% conversion for the tetravalent catalyst **1** and 98% for its trivalent counterpart (Table 1, entries 1 and 3). The CO₂-insertion complexes **2** and **5** displayed similar catalytic activity (entries 2 and 4). The conversion dropped drastically on increasing the steric bulk of the substituent on the epoxides. As a result, styrene oxide, and 3,3-dimethyl-1,2-butylene oxide showed only very low conversions (entry 21) for ceric **1** and almost no conversion in the case of trivalent catalyst **4** (entry 22). Moderate conversion was observed for 1,2-*n*-hexylene oxide with the tetravalent catalyst **1** (entry 13). Conducting the catalysis at higher

temperature increased the TONs with both catalysts **1** and **4**, and resulted in nearly quantitative conversion for both systems (entries 15 and 16). Without co-catalyst TBAB, **1** showed moderate catalytic activity at 90 °C (entry 17). In almost all cases, tetravalent **1** showed higher catalytic activity than cerous **4**, which most likely results from the higher Lewis acidity of Ce^{IV} versus Ce^{III}. To further evaluate the catalytic reaction with catalyst **1** and propylene oxide, the TOFs at different stages of the catalysis were determined (see Table S1). After a short induction period, most likely corresponding to the insertion of CO₂ into **1**, the TOF reached a maximum of 11 h⁻¹ within the first 3 h. For comparison, a TOF of 155 h⁻¹ was reported by Yao and co-workers when performing the reaction under 10 bar CO₂ pressure.^[11a] Increasing the CO₂ pressure did not significantly affect the catalytic activity of compound **1** (entries 13 vs. 18 and 21 vs. 23). However, a simultaneous increase of the temperature to 90 °C and the CO₂ pressure to 10 bar led to a marked improvement in the catalytic activity, resulting in TONs of up to 300 for the sterically demanding 3,3-dimethyl-1,2-butylene oxide (entries 24 and 26). The latter conditions were also applicable for the cycloaddition of CO₂ and cyclohexene oxide, an internal epoxide (entries 27–29). Having optimized the reaction conditions, we determined the initial turnover frequencies for the different epoxides (entries 5, 12, 20, and 25). As expected, the TOFs increased as the steric of the substituents bulk decreased, ranging from 24 to 196 h⁻¹ and giving almost quantitative conversion of propylene oxide after a reaction time of one hour (entry 5). Compared to the other catalyst systems based on rare-earth metals reported by Yao and co-workers (TOFs up to 440 h⁻¹) or by Otero and co-workers (3167 h⁻¹), our system shows only moderate catalytic activity under comparable conditions.^[11a,b]

The mechanism of the cycloaddition of CO₂ and epoxides using tetraalkylammonium salts as co-catalysts has been discussed in detail.^[32,33] It is generally accepted that the epoxide is activated by coordination to a Lewis-acidic metal center followed by a nucleophilic ring-opening attack of the bromide to form a metal-alkoxy bond (Scheme 6). Subsequently, the alkoxide reacts with CO₂ and cyclizes to produce a cyclic carbonate. Hints that the mechanisms for the cycloaddition differ using ceric **1** or cerous **4** as the catalyst could be found when conducting the reactions with different amounts of catalyst loading (entries 7 and 9). This results in a change in the TONs for the tetravalent catalyst **1**, whereas the TONs remained the same for trivalent complex **4**. The occurrence of distinct reaction mechanisms is not surprising, as **1** is a monometallic complex while **4** is a tetrametallic compound in the solid state and in solution (for a more detailed possible mechanism see Scheme S1). Such a mechanism, involving multiple metal centers, was previously proposed for the bimetallic complex [Al(salen)]₂O by North and co-workers.^[32] However, any detailed information about the mechanism could not be retrieved from our catalyst system as the TOFs decreased enormously when the reactions were conducted in propylene carbonate (no significant conversion at ambient temperature after 24 h and ca. 10% conversion after 20 h at 90 °C) or any other solvent, which makes kinetic studies unfeasible.


Table 1: Catalytic formation of cyclic carbonates from epoxides and CO₂.^[a]

Entry	Catalyst	Substrate	Product	Conversion [%]	TON/Ce
1	1			93	93
2	2			93	93
3	4			98	98
4	5			96	96
5	1 ^[i]			98	196
6	1			20	20
7	1 ^[b]			4	8
8	4			13	13
9	4 ^[c]			6	12
10	4 -thf			13	13
11	4 -thf ^[c]			7	14
12	1 ^[i]			12	24
13	1			61	61
14	4			25	25
15	1 ^[d]			98	98
16	4 ^[e]			96	96
17	1 ^[f]			25	25
18	1 ^[g]			76	76
19	6			3	3
20	1 ^[i]			49	98
21	1			12	12
22	4			3	3
23	1 ^[g]			9	9
24	1 ^[h]			84	168
25	1 ^[i]			14	28
26	1 ^[i]			60	300
27	1			2	2
28	1 ^[h]			77	154
29	1 ^[i]			35	175

[a] Reaction conditions if not otherwise noted: 1 bar CO₂ pressure and 0.5 mol% catalyst (1 mol% for **2** and **6**; 0.25 mol% for **4** and **5**) and 1 mol% co-catalyst for 24 h at ambient temperature in neat epoxide.

[b] 0.25 mol% catalyst **1** and 0.5 mol% TBAB, 24 h. [c] 0.125 mol% catalyst **4** and 0.5 mol% TBAB, 24 h. [d] 0.5 mol% catalyst **1** and 1 mol% TBAB, 24 h, 90 °C. [e] 0.25 mol% catalyst **4** and 1 mol% TBAB, 24 h, 90 °C. [f] At 90 °C without TBAB as a co-catalyst. [g] 0.5 mol% catalyst **1** and 1 mol% TBAB, 24 h, 10 bar CO₂ pressure. [h] 0.25 mol% catalyst **1** and 0.5 mol% TBAB, 24 h, 90 °C, 10 bar CO₂ pressure. [i] 0.25 mol% catalyst **1** and 0.5 mol% TBAB, 1 h, 90 °C, 10 bar CO₂ pressure.

[j] 0.1 mol% catalyst **1** and 0.2 mol% TBAB, 24 h, 90 °C, 10 bar CO₂ pressure.



Scheme 6. Proposed overall mechanism for the coupling of CO₂ with epoxides.

Complex **2** underwent a reductive side reaction in the presence of TBAB to afford cerous species [Ce(Me₂pz·CO₂)₄][NBu₄] (**6**). This side reaction occurred during the catalytic studies and when combining **1** and TBAB in stoichiometric amounts in [D₈]toluene under 1 bar CO₂ pressure. Although **2** is very stable in toluene solution under these conditions (no color change was observed after several days), it underwent reduction in the presence of TBAB within hours, as evidenced by decolorization of the orange solution. Most probably bromine is formed as an oxidation product; however, no brominated product could be detected in the reaction mixture (see Figure S23). The use of epoxides as a solvent seems to stabilize the tetravalent **2**, as no paramagnetic signals were found in the ¹H NMR spectra of the catalytic reactions. The crystal structure of **6** revealed the same motif as seen in **2-toluene** and **2-thf** (see Figure S67). The 8-coordinate cerium center bears four κ²(N,O) Me₂pz·CO₂ ligands with elongated Ce–N and Ce–O bonds compared to ceric **2-toluene** and **2-thf**, as would be expected for a cerium(III) center. Cerous **6** displays poor catalytic activity compared to tetravalent **1** (Table 1, entry 19), thus underlining that it is a side product and not the active catalyst.

Conclusion

We have shown that carbon dioxide easily inserts into the Ce–N(Me₂pz) bond of both ceric [Ce(Me₂pz)₄]₂ and cerous [Ce₄(Me₂pz)₁₂] at an amount equivalent to 5.7 mmol CO₂ per gram complex and via the controlled activation of 12 molecules of CO₂ within one complex, respectively. The insertion process is reversible both in solution and in the solid state, with CO₂ desorption being complete at < 100 °C. Both trivalent and tetravalent cerium pyrazolate complexes are active catalysts for the cycloaddition of epoxides and carbon

dioxide with TBAB as a co-catalyst under mild conditions. We are currently investigating the carbon dioxide capture performance of silica-grafted variants of Ce-pyrazolates,^[34] and our findings might also stimulate research in the area of cerium-dipyrazolate-based CO₂-“breathable”/expandable MOFs.^[35,36]

Acknowledgements

We thank Prof. Doris Kunz for providing access to ReactIR 15 for in situ IR measurements, and the working group of Prof. H.-Jürgen Meyer for conducting the TGA measurements.

Conflict of interest

The authors declare no conflict of interest.

Keywords: carbon dioxide · cerium · cycloaddition · epoxides · pyrazolates

How to cite: *Angew. Chem. Int. Ed.* **2020**, *59*, 5830–5836
Angew. Chem. **2020**, *132*, 5879–5885

- [1] a) P. Falkowski et al., *Science* **2000**, *290*, 291–296; b) S. Solomon, G.-K. Plattner, R. Knutti, P. Friedlingstein, *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 1704–1709; c) R. S. Haszeldine, *Science* **2009**, *325*, 1647–1652; d) D. W. Keith, *Science* **2009**, *325*, 1654–1655; e) N. von der Assen, P. Voll, M. Peters, A. Bardow, *Chem. Soc. Rev.* **2014**, *43*, 7982–7994.
- [2] For review articles, see a) D. M. D’Alessandro, B. Smit, J. R. Long, *Angew. Chem. Int. Ed.* **2010**, *49*, 6058–6082; *Angew. Chem.* **2010**, *122*, 6194–6219; b) C.-H. Yu, C.-H. Huang, C.-S. Tan, *Aerosol Air Qual. Res.* **2012**, *12*, 745–769; c) K. Sumida, D. L. Rogow, J. A. Mason, T. M. McDonald, E. D. Bloch, Z. R. Herm, T.-H. Bae, J. R. Long, *Chem. Rev.* **2012**, *112*, 724–781; d) E. S. Sanz-Pérez, C. R. Murdock, S. A. Didas, C. W. Jones, *Chem. Rev.* **2016**, *116*, 11840–11876; e) Y. Lin, C. Kong, Q. Zhang, L. Chen, *Adv. Energy Mater.* **2017**, *7*, 1601296.
- [3] For review articles, see a) T. Sakakura, J.-C. Choi, H. Yasuda, *Chem. Rev.* **2007**, *107*, 2365–2387; b) G. Centi, S. Perathoner, *Catal. Today* **2009**, *148*, 191–205; c) M. Cokoja, C. Bruckmeier, B. Rieger, W. A. Herrmann, F. E. Kühn, *Angew. Chem. Int. Ed.* **2011**, *50*, 8510–8537; *Angew. Chem.* **2011**, *123*, 8662–8690; d) M. Aresta, A. Dibenedetto, A. Angelini, *Chem. Rev.* **2014**, *114*, 1709–1742; e) Q. Liu, L. Wu, R. Jackstell, M. Beller, *Nat. Commun.* **2015**, *6*, 5933; f) J. Artz, T. E. Müller, K. Thenert, J. Kleinekorte, R. Meys, A. Sternberg, A. Bardow, W. Leitner, *Chem. Rev.* **2018**, *118*, 434–504.
- [4] A. Paparo, J. Okuda, *Coord. Chem. Rev.* **2017**, *334*, 136–149.
- [5] a) A. Goepfert, M. Czaun, R. B. May, G. K. S. Prakash, G. A. Olah, S. R. Narayanan, *J. Am. Chem. Soc.* **2011**, *133*, 20164–20167; b) F. Inagaki, C. Matsumoto, T. Iwata, C. Mukai, *J. Am. Chem. Soc.* **2017**, *139*, 4639–4642.
- [6] a) S. R. Caskey, A. G. Wong-Foy, A. J. Matzger, *J. Am. Chem. Soc.* **2008**, *130*, 10870–10871; b) T. M. McDonald et al., *Nature* **2015**, *519*, 303–308.
- [7] Y. F. Radkov, E. A. Fedorova, S. Y. Khorshev, G. S. Kalinina, M. N. Bochkarev, G. A. Razuvaev, *Zhur. Obshchei Khimii* **1986**, *56*, 386–389.

- [8] For a recent study on Ce^{III} silylamides, see H. Yin, P. J. Carroll, E. J. Schelter, *Chem. Commun.* **2016**, 52, 9813–9816.
- [9] M. N. Bochkarev, E. A. Fedorova, Y. F. Radkov, S. Y. Khorshev, G. S. Kalinina, G. A. Razuvaev, *J. Organomet. Chem.* **1983**, 258, C29–C33.
- [10] For examples of CO₂ insertion into Ln^{III}-aryloxy bonds, see a) D. Cui, M. Nishiura, O. Tardif, Z. Hou, *Organometallics* **2008**, 27, 2428–2435; b) L. A. M. Steele, T. J. Boyle, R. A. Kemp, C. Moore, *Polyhedron* **2012**, 42, 258–264.
- [11] a) J. Qin, P. Wang, Q. Li, Y. Zhang, D. Yuan, Y. Yao, *Chem. Commun.* **2014**, 50, 10952–10955; b) J. Martínez, J. Fernández-Baeza, L. F. Sánchez-Barba, J. A. Castro-Osma, A. Lara-Sánchez, A. Otero, *ChemSusChem* **2017**, 10, 2886–2890; c) Z. Zhao, J. Qin, C. Zhang, Y. Wang, D. Yuan, Y. Yao, *Inorg. Chem.* **2017**, 56, 4568–4575.
- [12] a) D. Cui, M. Nishiura, Z. Hou, *Macromolecules* **2005**, 38, 4089–4095; b) D. V. Vitanova, F. Hampel, K. C. Hultzsich, *J. Organomet. Chem.* **2005**, 690, 5182–5197; c) B. B. Lazarov, F. Hampel, K. C. Hultzsich, *Z. Anorg. Allg. Chem.* **2007**, 633, 2367–2373; d) Z. Zhang, D. Cui, X. Liu, *J. Polym. Sci. Part A* **2008**, 46, 6810–6818; e) A. Decortes, R. M. Haak, C. Martín, M. M. Belmonte, E. Martín, J. Benet-Buchholz, A. W. Kleij, *Macromolecules* **2015**, 48, 8197–8207; f) J. Qin, B. Xu, Y. Zhang, D. Yuan, Y. Yao, *Green Chem.* **2016**, 18, 4270–4275.
- [13] For the use of organoscandium(III) complexes in the catalytic hydrosilylation of CO₂, see a) A. Berkefeld, W. E. Piers, M. Parvez, L. Castro, L. Maron, O. Eisenstein, *Chem. Sci.* **2013**, 4, 2152–2162; b) F. A. LeBlanc, W. E. Piers, M. Parvez, *Angew. Chem. Int. Ed.* **2014**, 53, 789–792; *Angew. Chem.* **2014**, 126, 808–811.
- [14] a) M. A. St. Clair, B. D. Santarsiero, *Acta Crystallogr. Sect. C* **1989**, 45, 850–852; b) H. Schumann, J. A. Meese-Marktscheffel, A. Dietrich, F. H. Görlitz, *J. Organomet. Chem.* **1992**, 430, 299–315; c) W. J. Evans, C. A. Seibel, J. W. Ziller, R. J. Doedens, *Organometallics* **1998**, 17, 2103–2112; d) F. A. LeBlanc, A. Berkefeld, W. E. Piers, M. Parvez, *Organometallics* **2012**, 31, 810–818.
- [15] a) O. T. Summerscales, C. M. Moore, B. L. Scott, M. P. Wilkerson, A. D. Sutton, *Organometallics* **2017**, 36, 4682–4685; b) D. W. Beh, W. E. Piers, I. del Rosar, L. Maron, B. S. Gelfand, C. Gendy, J.-B. Lin, *Dalton Trans.* **2018**, 47, 13680–13688.
- [16] a) W. J. Evans, C. A. Seibel, J. W. Ziller, *Inorg. Chem.* **1998**, 37, 770–776; b) W. J. Evans, J. M. Perrotti, J. C. Brady, J. W. Ziller, *J. Am. Chem. Soc.* **2003**, 125, 5204–5212; c) N. W. Davies, A. S. P. Frey, M. G. Gardinier, J. Wang, *Chem. Commun.* **2006**, 4853–4855; d) J. Andrez, J. Pécaut, P.-A. Bayle, M. Mazzanti, *Angew. Chem. Int. Ed.* **2014**, 53, 10448–10452; *Angew. Chem.* **2014**, 126, 10616–10620; e) M. Xémard, V. Goudy, A. Braun, M. Tricoire, M. Cordier, L. Ricard, L. Castro, E. Louyriac, C. E. Kefalidis, C. Clavaguéra, L. Maron, G. Nocton, *Organometallics* **2017**, 36, 4660–4668; f) A. R. Willauer, D. Toniolo, F. Fadaei-Tirani, Y. Yang, M. Laurent, M. Mazzanti, *Dalton Trans.* **2019**, 48, 6100–6110.
- [17] a) Y.-M. So, G.-C. Wang, Y. Li, H. H. Y. Sung, I. D. Williams, Z. Lin, W.-H. Leung, *Angew. Chem. Int. Ed.* **2014**, 53, 1626–1629; *Angew. Chem.* **2014**, 126, 1652–1655; b) G.-C. Wang, Y.-M. So, Y. Li, K.-L. Wong, K.-C. Au-Yeung, H. H. Y. Sung, I. D. Williams, W.-H. Leung, *Chem. Eur. J.* **2015**, 21, 16126–16135.
- [18] a) P. L. Arnold, Z. R. Turner, A. I. Germeroth, I. J. Casely, G. S. Nichol, R. Bellabarba, R. P. Tooze, *Dalton Trans.* **2013**, 42, 1333–1337; b) P. L. Arnold, R. W. F. Kerr, C. Weetman, S. R. Docherty, J. Rieb, F. L. Cruickshank, K. Wang, C. Jandl, M. W. McMullon, A. Pöthig, et al., *Chem. Sci.* **2018**, 9, 8035–8045.
- [19] For further examples of CO₂ insertion into Ln–NHC bonds, see a) P. L. Arnold, I. A. Marr, S. Zlatogorsky, R. Bellabarba, R. P. Tooze, *Dalton Trans.* **2014**, 43, 34–37; b) T. Simler, T. J. Feuerstein, R. Yadav, M. T. Gamer, P. W. Roesky, *Chem. Commun.* **2019**, 55, 222–225.
- [20] W. J. Evans, J. M. Perrotti, S. A. Kozimor, T. M. Champagne, B. I. Davis, G. W. Nycy, C. H. Fujimoto, R. D. Clark, M. A. Johnston, J. W. Ziller, *Organometallics* **2005**, 24, 3916–3931.
- [21] For an irreversible insertion of [Me₂SiO] into a Yb–Me₂pz bond, see X. Zhou, H. Ma, X. Huang, X. You, *J. Chem. Soc. Chem. Commun.* **1995**, 2483–2484.
- [22] a) For a reversible CO₂ insertion into a Sm–EPh bond (E = S, Se), see W. J. Evans, K. A. Miller, J. W. Ziller, *Inorg. Chem.* **2006**, 45, 424–429; b) a scandium(III)-bonded β-diketiminato (“nannac”)ligand framework was shown to reversibly capture CO₂: Ref. [14b].
- [23] D. Werner, G. B. Deacon, P. C. Junk, R. Anwander, *Eur. J. Inorg. Chem.* **2017**, 3419–3428.
- [24] H. C. Aspinall, J. Bacsá, A. C. Jones, J. S. Wrench, K. Black, P. R. Chalker, P. J. King, P. Marshall, M. Werner, H. O. Davies, et al., *Inorg. Chem.* **2011**, 50, 11644–11652.
- [25] J. A. Bogart, A. J. Lewis, S. A. Medling, N. A. Piro, P. J. Carroll, C. H. Booth, E. J. Schelter, *Inorg. Chem.* **2013**, 52, 11600–11607.
- [26] M. Begtrup, J. Elguero, R. Faure, P. Camps, C. Estopá, D. Ilavský, A. Fruchier, C. Marzin, J. de Mendoza, *Magn. Reson. Chem.* **1988**, 26, 134–151.
- [27] D. Werner, G. B. Deacon, P. C. Junk, R. Anwander, *Dalton Trans.* **2017**, 46, 6265–6277.
- [28] D. Werner, U. Bayer, N. E. Rad, P. C. Junk, G. B. Deacon, R. Anwander, *Dalton Trans.* **2018**, 47, 5952–5955.
- [29] U. Baisch, D. B. Dell’Amico, F. Calderazzo, L. Labella, F. Marchetti, D. Vitali, *J. Mol. Catal.* **2003**, 204–205, 259–265.
- [30] S. Bachmann, B. Gernert, D. Stalke, *Chem. Commun.* **2016**, 52, 12861–12864.
- [31] M. H. Chisholm, J. C. Gallucci, K. Phomphrai, *Inorg. Chem.* **2004**, 43, 6717–6725.
- [32] W. Clegg, R. W. Harrington, M. North, R. Pasquale, *Chem. Eur. J.* **2010**, 16, 6828–6843.
- [33] a) D. J. Darensbourg, M. W. Holtcamp, *Coord. Chem. Rev.* **1996**, 153, 155–174; b) H. Jing, T. Chang, L. Jin, M. Wu, W. Qiu, *Catal. Commun.* **2007**, 8, 1630–1634; c) J. Meléndez, M. North, R. Pasquale, *Eur. J. Inorg. Chem.* **2007**, 3323–3326; d) F. Jutz, J.-D. Grunwaldt, A. Baiker, *J. Mol. Catal. A* **2008**, 279, 94–103; e) F. Jutz, J.-D. Grunwaldt, A. Baiker, *J. Mol. Catal. A* **2009**, 297, 63–72; f) M. North, R. Pasquale, *Angew. Chem. Int. Ed.* **2009**, 48, 2946–2948; *Angew. Chem.* **2009**, 121, 2990–2992.
- [34] Y. Liang, R. Anwander, *Dalton Trans.* **2013**, 42, 12521–12545.
- [35] For Ce^{IV} carboxylate MOFs, see a) P. Ji, T. Sawano, Z. Lin, A. Urban, D. Bours, W. Lin, *J. Am. Chem. Soc.* **2016**, 138, 14860–14863; b) X.-P. Wu, L. Gagliardi, D. G. Truhlar, *J. Am. Chem. Soc.* **2018**, 140, 7904–7912.
- [36] For examples of dipyrzoyl-based MOFs, see a) H. J. Choi, M. Dinca, J. R. Long, *J. Am. Chem. Soc.* **2008**, 130, 7848–7850; b) M. Tonigold, Y. Lu, B. Bredenkötter, B. Rieger, S. Bahnmüller, J. Hitzbleck, G. Langstein, D. Volkmer, *Angew. Chem. Int. Ed.* **2009**, 48, 7546–7550; *Angew. Chem.* **2009**, 121, 7682–7687; c) N. M. Padial, E. Q. Procopio, C. Montoro, E. López, J. E. Oltra, V. Colombo, A. Maspero, N. Masciocchi, S. Galli, I. Senkowska, S. Kaskel, E. Barea, J. A. R. Navarro, *Angew. Chem. Int. Ed.* **2013**, 52, 8290–8294; *Angew. Chem.* **2013**, 125, 8448–8452; d) Z. R. Herm, B. M. Wiers, J. A. Mason, M. R. Hudson, P. Zajdel, C. M. Brown, N. Masciocchi, R. Krishna, J. R. Long, *Science* **2013**, 340, 960–964; e) J. A. Mason, J. Oktawiec, M. K. Taylor, M. R. Hudson, J. Rodriguez, J. E. Bachman, M. L. Gonzalez, A. Cervellino, A. Guagliardi, C. M. Brown, P. L. Llewellyn, N. Masciocchi, J. R. Long, *Nature* **2015**, 527, 357–363.

Manuscript received: December 23, 2019

Accepted manuscript online: January 9, 2020

Version of record online: January 30, 2020