

# Bilateral Brachial Plexus Block Using Chloroprocaine For Surgery Of Bilateral Radial Fractures [Response To Letter]

This article was published in the following Dove Press journal:  
*Local and Regional Anesthesia*

Chanchal Mangla  
Hattiyangadi Sangeetha Kamath  
Joel Yarmush

Department of Anesthesiology,  
New York Presbyterian Brooklyn  
Methodist Hospital, Brooklyn, NY, USA

## Dear editor

We would like to thank Dr Hendrickson et al for their interest in our work and writing about their opinion. We agree with the comment that mixing local anesthetics might make individual safe doses unknown, so maximum recommended doses of each local anesthetic should not be used. There are no human studies but it can be presumed to be additive based on some animal studies.<sup>1</sup> We mixed local anesthetics in our patients so that we could decrease the volume of more toxic local anesthetics (Bupivacaine) by using some less toxic ones (like chloroprocaine). We also used doses of each local anesthetic of well below the recommended toxic doses and did spacing in our blocks to avoid toxicity. Though axillary blocks along with medial brachial cutaneous and intercostobrachial block can be used,<sup>2</sup> we chose not to perform bilateral axillary blocks because it requires individual blockage of the terminal nerves which might lead to inadequate coverage, and also, performance of the block time and onset time is longer. Instead, we did infraclavicular block on one side, which greatly decreases the chances of phrenic nerve palsy. Dr Hendrickson made a very good point of using ropivacaine instead of bupivacaine due to less toxicity, but unfortunately we do not have ropivacaine available at our institution. Lastly, smaller volume of local anesthetics can be used with the use of ultrasound for a successful block. We used 30 mL volume for each block, as we wanted to ensure complete coverage of surgical anesthesia and to avoid any supplementation/deeper sedation or general anesthesia in case of an incomplete block. Also, our second block was more than two hours later, hence we avoided the overlap of peak plasma concentration of the local anesthetics from first and second blocks.

## Disclosure

The authors report no conflicts of interest in this communication.

## References

1. Yaddanapudi S. Prevention of local anesthetic systemic toxicity. *J Anaesthesiol Clin Pharm Col.* 2011;27(4):438–439. doi:10.4103/0970-9185.86566
2. Magazzeni P, Jochum D, Iohom G, Mekler G, Albuissou E, Bouaziz H. Ultrasound guided selective versus conventional block of the medial brachial cutaneous and the intercostobrachial nerves: A randomized clinical trial. *Reg Anesth Pain Med.* 2018;43:832–837. doi:10.1097/AAP.0000000000000823

Correspondence: Chanchal Mangla  
Department of Anesthesiology, New York  
Presbyterian Brooklyn Methodist Hospital,  
506 Sixth Street, Brooklyn, NY, USA  
Tel +1 718 780 3270  
Email drchanchalmangla@gmail.com

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Local and Regional Anesthesia 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Local and Regional Anesthesia editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

## Local and Regional Anesthesia

Dovepress

### Publish your work in this journal

Local and Regional Anesthesia is an international, peer-reviewed, open access journal publishing on the development, pharmacology, delivery and targeting and clinical use of local and regional anesthetics and analgesics. The journal welcomes submitted papers covering original research, basic science, clinical studies, reviews &

evaluations, guidelines, expert opinion and commentary, case reports and extended reports. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/local-and-regional-anesthesia-journal>