

Corrigendum

Corrigendum to: Diagnostic delay is common for patients with axial spondyloarthritis: results from the National Early Inflammatory Arthritis Audit

Mark D. Russell ¹, Fiona Coath², Mark Yates ¹, Katie Bechman¹, Sam Norton ¹, James B. Galloway¹, Joanna Ledingham³, Raj Sengupta⁴ and Karl Gaffney²

Rheumatology 2021;
doi:<https://doi.org/10.1093/rheumatology/keab428>

In the originally published version of this manuscript, there was an error in the y-axis of Figure 1B. It should read “Proportion of patients assessed within 3 weeks” instead of “Proportion of patients assessed within 3 months”. This error has been corrected online and in print.

¹Centre for Rheumatic Diseases, King's College London, London,

²Rheumatology Department, Norfolk and Norwich University Hospital, Norwich, ³Rheumatology Department, Portsmouth Hospitals University NHS Trust, Portsmouth and ⁴Department of Rheumatology, Royal National Hospital for Rheumatic Diseases, Bath, UK

Correspondence to: Mark Russell; Centre for Rheumatic Diseases, Weston Education Centre, King's College London, 10 Cutcombe Road, London SE5 9RJ, UK. E-mail: mark.russell@kcl.ac.uk