

POSTER PRESENTATION

Open Access

Study of changes in lipid profile and fasting blood glucose in protease inhibitor exposed HIV/AIDS patients in School of Tropical Medicine, Kolkata

Ananya Bhowmik^{1*}, Rajyasree De¹, Manotosh Mahato¹, Sujit Kumar Das¹, Subhasish Kamal Guha²

From First International Science Symposium on HIV and Infectious Diseases (HIV SCIENCE 2012)
Chennai, India. 20-22 January 2012

Background

The national second-line Anti retroviral Therapy (ART) programme was started in Kolkata in December 2008. It included a combination of Tenofovir, Lamivudine and Ritonavir-boosted Lopinavir ± Zidovudine. Dyslipidaemia and increased fasting blood sugar (FBS) often complicate protease inhibitor-containing ART. Thus a prospective study was designed to observe the above changes.

Methods

The data of 48 patients, on protease inhibitor for one year were analyzed. Body Mass Index (BMI), grip strength (GS), Triceps skin fold (TSF), 24 hour dietary recall, serum triglyceride (TG), total cholesterol (TC), HDL, LDL, VLDL and FBS were estimated for all patients at baseline, 6 months and after one year.

Results

There was a significant increase in TG, TC and VLDL levels at 1 year as compared to baseline ($p=0.013$, 0.00 and 0.00 respectively) whereas LDL significantly increased at 6 months only ($p=0.029$). HDL decreased significantly at 6 months ($p=0.019$). TSF significantly decreased both at 6 and 12 months ($p=0.00$ and 0.00 respectively). The BMI and GS showed a significant increase at both 6 months ($p=0.001$, 0.000 respectively) and 1 year ($p=0.005$ and 0.00 respectively). Four patients with normal baseline FBG and one with impaired fasting glucose progressed to overt diabetes ($FBG > 124$ mg/dl)

at 12 months. No significant change was noted in energy and protein intake of patients.

Conclusion

There is an increased incidence of dyslipidaemia and unmasking of diabetes related to protease inhibitor in this cohort. There has been an improvement in nutritional status as shown by BMI and GS.

Author details

¹Centre of Excellence in HIV, School of Tropical Medicine, Kolkata, India.

²Department of Tropical Medicine, School of Tropical Medicine, Kolkata, India.

Published: 4 May 2012

doi:10.1186/1471-2334-12-S1-P34

Cite this article as: Bhowmik et al.: Study of changes in lipid profile and fasting blood glucose in protease inhibitor exposed HIV/AIDS patients in School of Tropical Medicine, Kolkata. *BMC Infectious Diseases* 2012 12 (Suppl 1):P34.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: ananya50@gmail.com

¹Centre of Excellence in HIV, School of Tropical Medicine, Kolkata, India
Full list of author information is available at the end of the article