



Letter to the Editor

Korean J Anesthesiol 2020;73(6):566-567
<https://doi.org/10.4097/kja.20295>
pISSN 2005-6419 • eISSN 2005-7563

Received: June 5, 2020
Revised: June 11, 2020
Accepted: June 24, 2020

Corresponding author:

Emanuele Piraccini, M.D.
Department of Anesthesia, Intensive Care
Nord and Pain Management Unit, Bellaria
Hospital, via Altura 3, 40139 Bologna, Italy
Tel: +39-3396119485
Fax: +39-0543735130
Email: drpiraccini@gmail.com
ORCID: <https://orcid.org/0000-0002-4762-2312>

The efficacy of fascial plane blocks for myofascial pain syndrome: do they achieve long-term results?

Emanuele Piraccini¹, Helen Byrne²

¹Department of Anesthesia, Intensive Care Nord and Pain Management Unit, Bellaria Hospital, Bologna, Italy, ²Kingsbridge Medical Diagnostics, Kingsbridge Healthcare Group, Belfast, Northern Ireland

We carefully read the letter by Ekinici et al. [1] who replied to the detailed paper by Elsharkawy et al. [2] describing a case report on the use of rhomboid intercostal block (RIB) to manage myofascial pain syndrome (MPS).

We congratulate the authors for the clinical management of the patient and clear presentation of their results. The use of RIB for MPS is a new approach, and reports have been rising in the current literature to support the efficacy of RIB as part of the multimodal treatment [3,4].

We want to contribute to the discussion by focusing our attention not only on the benefits but also potential limitations of RIB to treat MPS.

It is crucial to obtain the correct diagnosis of MPS. MPS can be primary, which may be an overuse condition, such as lateral epicondylitis or piriformis syndrome, or secondary to other diseases or postural maladaptive changes. In secondary cases, if no therapeutic interventions are performed to treat the underlying cause, the results of fascial plane blocks as RIB are transient [5]. In the described case, Ekinici et al. [1] injected 20 ml of bupivacaine and dexamethasone which was a good idea, the authors had also performed hydrodissection of fascial planes in this way. This can provide outstanding and long-lasting results in case of fascial adhesion that is not easy to detect, but can contribute to MPS development [1,4].

The authors followed-up the patient for 4 weeks after the treatment. However, it would be interesting to have a longer follow-up, maybe 3 to 6 months, in order to establish if fascial plane blocks without any prevention strategy can ensure more consistent long-term results. In our experience, this is not as apparent, while the success of fascial plane blocks combined with physiotherapy has been reported [3-5].

Conflicts of Interest

No potential conflict of interest relevant to this article was reported.

Author Contributions

Emanuele Piraccini (Conceptualization; Writing – original draft)
Helen Byrne (Writing – review & editing)

ORCID

Emanuele Piraccini, <https://orcid.org/0000-0002-4762-2312>
Helen Byrne, <https://orcid.org/0000-0002-7190-239X>

© The Korean Society of Anesthesiologists, 2020
© This is an open-access article distributed under the terms of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

References

1. Ekinci M, Ciftci B, Alici HA, Ahiskalioglu A. Ultrasound-guided rhomboid intercostal block effectively manages myofascial pain. *Korean J Anesthesiol* 2020; 73: 564-5.
2. Elsharkawy H, Hamadnalla H, Altinpulluk EY, Gabriel RA. Rhomboid intercostal and subserratus plane block: a case series. *Korean J Anesthesiol* 2020; 73: 550-6.
3. Piraccini E, De Lorenzo E, Maitan S. Rhomboid intercostal block for myofascial pain syndrome in a patient with amyotrophic lateral sclerosis. *Minerva Anesthesiol* 2019; 85: 1367-9.
4. Piraccini E, Maitan S. Ultrasound guided rhomboid plane hydrodissection for fascial adhesion. *J Clin Anesth* 2020; 59: 13.
5. Piraccini E, Calli M, Taddei S, Byrne H, Rocchi M, Maitan S. Erector spinae plane block for myofascial pain syndrome: only a short-term relief? *Minerva Anesthesiol* 2020; 86: 888-90.