

Review Article

Rehabilitation Professional and Patient Satisfaction with Telerehabilitation of Musculoskeletal Disorders: A Systematic Review

Junaid Amin ¹, Basaruddin Ahmad ¹, Salman Amin,² Ammar Ahmed Siddiqui ^{3,4}, and Mohammad Khursheed Alam ⁵

¹School of Dental Sciences, Health Campus, University Sains Malaysia, Kubang Kerian, Malaysia

²College of Medicine & Dentistry, University of Lahore, Lahore, Pakistan

³College of Dentistry, Preventive Dental Sciences, University of Hail, Saudi Arabia

⁴Department of Community Dentistry, College of Dentistry, Bakhtawar Amin Medical and Dental College, Multan, Pakistan

⁵Department of Preventive Dentistry, College of Dentistry, Jouf University, Sakaka, Aljouf, Saudi Arabia

Correspondence should be addressed to Junaid Amin; junaid768@hotmail.com and Basaruddin Ahmad; basaruddin@usm.my

Received 27 November 2021; Revised 8 April 2022; Accepted 26 June 2022; Published 2 August 2022

Academic Editor: Rafael Scaf De Molon

Copyright © 2022 Junaid Amin et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Telerehabilitation offers an alternative healthcare delivery remotely in a patient's environment at a lower cost, better accessibility, and equivalent quality to the standard approach. Several studies had examined the effectiveness of telerehabilitation inpatients with musculoskeletal disorders, and although there is evidence that it is at least equally effective as the standard care, the patient and rehabilitation professional satisfaction with the delivery method is not conclusive. A systematic review was conducted to study the patients' and rehabilitation professionals' satisfaction with telerehabilitation for musculoskeletal disorders. A search for relevant studies on 29 April 2021 was carried out in Medline/PubMed, Scopus, and Web of Science (WOS). The search terms included "telerehabilitation," AND "satisfaction" AND "musculoskeletal disorders," "telehealth," "telemedicine," "patient experience," and "pain". Fifteen eligible studies with 12,341 patients were included in this systematic review. A report was included if it (a) assessed the satisfaction of patients or professionals or both as one of the outcomes of a telerehabilitation intervention, (b) included adults 18 years and above with musculoskeletal disorders, and (c) is an intervention study using a quantitative approach. The quality of studies was assessed using the critical appraisal checklist tool developed by Joanna Briggs Institute (JBI). Most of the studies reported that patients were satisfied with both telerehabilitation and face-to-face intervention. However, few studies reported that patients were more satisfied with telerehabilitation compared to face-to-face of intervention. Patients in one study had preferred the incorporation of telerehabilitation and face-to-face sessions. Two of three studies had reported overall satisfaction with telerehabilitation by the professionals. Overall, there is evidence that patients and rehabilitation professional are satisfied with telerehabilitation compared to face-to-face consultation.

1. Introduction

Telerehabilitation is the provision of rehabilitation services from a provider to a patient via a telecommunication system and information technology [1]. Such an alternative approach attracts the attention of the healthcare community because of several advantages. It saves time and cost related to travelling to and waiting at the healthcare centre, facili-

tates and improves access to services, and promises equal quality services to the public [2]. Telerehabilitation has been shown to be successfully performed and effective for people with medical conditions such as stroke, breast cancer patients, cardiopulmonary, and musculoskeletal disorders (MSDs) [3–6].

Patient satisfaction is one of the secondary but very important outcomes in healthcare delivery. Patients are the

main source of information to report on the quality of service provided, that is whether the standards of patient care and treatment received met the expectations. Telehealth platform changes with advancing technology and the mode of delivery can change dramatically; for example, from voice to video to multiperson conference, but regardless of the changes, a consistent patient-provider relationship must be formed [7]. As telehealth use increases, it becomes increasingly important to maintain patient satisfaction similar to the standard approach. Telehealth developers should be flexible to accommodate the patients' and healthcare professionals' perspective and needs; these should be monitored regularly. The success of a telehealth program heavily relies on patient satisfaction [7]. A variety of components underlying the satisfaction have been addressed in different studies that may include overall satisfaction and satisfaction with the application and services [8–10]. Moreover, to achieve optimal outcomes of the treatment and satisfaction of the patient, it is imperative to maintain a strong relationship between patient and professionals [11]. Therefore, it is important to take the satisfaction of both patients and the professionals for a successful and sustainable telehealth programs.

A recent systematic review reported that the evidence for the efficacy of telehealth interventions in improving musculoskeletal pain-related outcomes is comparable to the standard face-to-face interventions [12]. According to healthcare workers, online services can be a helpful addition to face-to-face therapies for chronic pain. Patients are also enthusiastic about telehealth approaches to healthcare delivery [7]. Patients who received exercises through telerehabilitation after shoulder joint replacement reported feelings of “closeness at a distance,” freedom, and increased awareness about their “body and self” [13]. Likewise, there were good levels of patient satisfaction with telehealth delivery for cognitive behavioral therapy, exercise, and pain-coping interventions [14].

A broad range of research on telemedicine, or immersive video consultations, has been conducted in various locations around the world. Telemedicine commentators often emphasize the need for more research on the protection, efficacy, and cost-effectiveness of healthcare delivery. As a result of the abundance of publications about patient satisfaction in the telemedicine literature, which are overwhelmingly optimistic, there is a tendency to believe that research in telerehabilitation is relatively of a less priority and a new area that should be focused. There is a good body of literature that examined the patient and professional satisfaction with telerehabilitation. It is imperative to take the perspectives of patients to increase the access, acceptance, and adherence with telerehabilitation in MSD management. To date, there has been no comprehensive systematic review on the satisfaction of patients and professionals with telerehabilitation for musculoskeletal disorder management. The current systematic review is aimed at determining the satisfaction of patients and professionals with telerehabilitation as compared to traditional face-to-face intervention for musculoskeletal disorders.

2. Methods

We follow the PRISMA guidelines to conduct this systematic review [15]. The PRISMA 2020 checklist can found in supplementary file 1. The study protocol was registered with PROSPERO (CRD42021252078).

2.1. Review Question. Based on PRISMA guidelines, the search questions were built on PICO format as follows: are professionals and patients satisfied with the telerehabilitation as compared to the face-to-face rehabilitation of musculoskeletal disorders? PICO format for the review questions is explained as follows: the population (patients with musculoskeletal issues)/(professionals dealing musculoskeletal issues), the intervention (telerehabilitation), comparison (face-to-face), and outcome (satisfaction).

2.2. Search Strategy. We searched the electronic databases from January 1980 to April 2021 reports in the English language. Databases searched include PubMed, Scopus, and Web of Science (WOS). The search terms were “telerehabilitation”, “satisfaction,” and “musculoskeletal disorders.” We also included common synonyms: “telehealth,” “telemedicine,” “patient experience,” and “expectations,” “pain.” Additional searches were carried out on Google Scholar and ResearchGate platforms. The search string for each variable has been provided as supplementary file 2. Moreover, the references of the included studies were also explored to find the relevant literature meeting the inclusion criteria. Two authors (JA and AAS) independently screened the abstracts based on the inclusion and exclusion criteria.

2.3. Inclusion and Exclusion Criteria. A report was included if it (a) assessed the satisfaction of patients or professionals or both as one of the outcomes of a telerehabilitation intervention, (b) included adults 18 years and above with musculoskeletal disorders, and (c) is an intervention study using a quantitative approach. A report was excluded if it is (a) not in English language and if they are (b) review articles, case reports, qualitative studies, book chapters, and articles with low quality.

3. Results

3.1. Study Selection. A total of 1091 studies were identified from Scopus ($n = 141$), Web of Science ($n = 425$), and PubMed ($n = 525$). An additional 51 reports were identified from Google Scholar, Research Gate, and reference list of the searched articles. A total of 149 duplicate studies were removed using the Reference Management Software Package (Endnote X9). A total of 728 were excluded because the studies did not meet the inclusion (satisfaction was not assessed = 291, not within the age range = 15, and not intervention study = 314) and exclusion (non-English report = 5 and not relevant report = 103) criteria. The full text of 41 studies was reviewed, and 26 studies were excluded due to missing data about satisfaction ($n = 17$), low-quality articles ($n = 5$), and review articles ($n = 4$). A total of 15 studies were included in systematic review after ensuring that those are fulfilling the




FIGURE 1: PRISMA flow diagram showing study identification and selection process.

inclusion criteria of the study. Figure 1 shows a schematic of the study identification and selection process.

3.2. Data Abstraction. Two field-based experts (JA and AAS) reviewed the full text of the report independently and used a data extraction form to record information relating to the author, year of the report, location of study, number of subjects, study design, methodology, study population, technology explored, and measures of satisfaction and findings of the study. Cohen's kappa for inter rater reliability was calculated as .74 showing substantial agreement. Disagreements were resolved by consensus and discussion or involved arbitration by the last author (MKA).

3.3. Synthesis of Results. In our review, content analysis was performed to synthesize the results. We examined articles that reported patient and professionals satisfaction. The different parameters of satisfaction were categorized into overall satisfaction, satisfaction with application, satisfaction with services, satisfaction, and patient-provider relationship. We performed a physical count of these variables to find the consistency. All parameters were displayed in number of occurrence. Moreover, the heterogeneity was also observed including sample size, study design, risk of bias, setting of studies, and outcome measures.

3.4. Quality Assessment. We utilized tools developed by the Joanna Briggs Institute (JBI) critical appraisal checklist to assess the quality of RCTs, case-control studies, cohort studies, non-RCTs, and case series [16, 17]. The JBI has well-established reliability and validity to assess the risk of bias in studies [16, 18]. Two authors independently evaluated the quality of each study. The JBI score was calculated for each study using the checklist, and score was presented as percentages. The study with a JBI score of 20-49% was considered high risk of bias and with 50-79% and 80-100% moderate and low risk of bias, respectively (Figure 2).

3.5. Article Characteristics. There were 15 eligible studies with a total of 12,341 participants. Sample sizes ranged from 3 to 10264 patients. All studies were from the developed countries including five from the United States, four from Europe, three from Canada, two from Australia, and one from Israel. There were 6 RCTs, 3 non-RCTs, 3 case-series studies, and 2 cohort studies and 1 quasiexperimental trial. Most studies ($n = 10$) used asynchronous mode through while others used synchronous videoconference mode ($n = 4$) and a combination of both ($n = 1$).

3.6. Patient Satisfaction. The majority of the studies included satisfaction of telerehabilitation as one of the secondary objectives ($n = 11$), and few studies included satisfaction in

		Risk of bias domains													Overall	JBI score	
		1	2	3	4	5	6	7	8	9	10	11	12	13			
1.	Bini SA and Mahajan J 2017 ^a	⊗	⊗	⊕	⊗	⊖	⊖	⊕	⊕	⊕	⊕	⊕	⊕	⊖	⊖		70
2.	Culliton, Sharon E., et al. 2018 ^a	⊕	⊖	⊕	⊕	⊕	⊖	⊕	⊕	⊕	⊕	⊕	⊕	⊕	⊕		84.6
3.	Doiron-Cadriu, Patrick, et al. 2020 ^a	⊕	⊕	⊕	⊖	⊖	⊕	⊕	⊕	⊕	⊕	⊕	⊕	⊕	⊕		84.6
4.	Tousignant, Michel, et al. 2011 ^a	⊗	⊗	⊕	⊕	⊕	⊖	⊕	⊕	⊗	⊕	⊕	⊕	⊖	⊖		61.53
5.	Nelson, M., et al. 2020 ^a	⊕	⊕	⊕	⊗	⊕	⊕	⊕	⊕	⊗	⊕	⊕	⊕	⊕	⊕		84.6
6.	Salazar-Fernandez, Clara Isabel, et al. 2012 ^b	⊕	⊕	⊕	⊕	⊗	⊕	⊕	⊕	⊕					⊕		88.9
7.	Katt, Brian M., et al. 2012 ^d	⊕	⊕	⊖	⊖	⊖	⊕	⊕	⊗	⊕	⊕	⊕			⊖		66.6
8.	Mayer, Naomi, et al. 2021 ^b	⊕	⊕	⊗	⊕	⊕	⊕	⊕	⊖	⊕					⊖		77.8
9.	Tsyvakh, Andriy I, and Andriy J. Hospodarsky 2018 ^c	⊕	⊕	⊕	⊕	⊕	⊕	⊖	⊕	⊖	⊕				⊕		80
10.	Moffet, Hélène, et al. 2017 ^a	⊖	⊖	⊕	⊖	⊖	⊕	⊕	⊕	⊕	⊕	⊕	⊕	⊕	⊖		69.2
11.	Pani, Danilo, et al. 2017 ^b	⊕	⊕	⊖	⊕	⊕	⊕	⊕	⊕	⊕					⊕		88.9
12.	Jansen-Kosterink, Stephanie, et al. 2015 ^c	⊕	⊕	⊕	⊕	⊕	⊕	⊖	⊕	⊕	⊕				⊕		88.9
13.	Bailey, Jeannie F., et al. 2020 ^d	⊕	⊕	⊕	⊗	⊖	⊖	⊕	⊕	⊕	⊕	⊕			⊖		72.7
14.	Peterson, Seth. 2018 ^c	⊖	⊕	⊖	⊕	⊕	⊕	⊕	⊕	⊕	⊕				⊕		80
15.	Cottrell, Michelle A., et al. 2019 ^b	⊕	⊕	⊕	⊕	⊕	⊖	⊕	⊕	⊕					⊕		88.9

- a. RCT (n = 6)
 - b. NRCT (n = 4)
 - c. CSS (n = 3)
 - d. CS (n = 2)
- ⊕ Low
 - ⊖ Moderate
 - ⊗ High

FIGURE 2: Risk of bias assessment based on Joanna Briggs Institute (JBI) critical appraisal tool.

their primary objective (n = 4). All the included studies reported the findings on patient satisfaction. Patient satisfaction was reported in both the intervention (telerehabilitation) and face-to-face groups.

The measures of satisfaction include overall satisfaction, satisfaction with application, satisfaction with services, and patient-provider relationship. A broad range of satisfaction was assessed and ranged from overall satisfaction (n = 6), satisfaction with application (n = 4), satisfaction with services (n = 4), satisfaction and patient-provider relationship (n = 11). These parameters were assessed in percentages.

The majority of the studies reported that patients were satisfied with both telerehabilitation and face-to-face [8–10, 19–22]. Three studies reported that patient satisfaction with telerehabilitation was significantly higher than that with face-to-face care [13, 21, 23]. One study reported that the patients preferred having an incorporated program that includes both telerehabilitation and face-to-face sessions [24] (Table 1).

3.7. Professional Satisfaction. There was evidence from the studies for the satisfaction of professionals with telerehabilitation. Of 15 included studies, only 3 (20%) studies assessed the satisfaction of the professionals, 2 of those had reported that there was satisfaction of professionals with telerehabilitation services [8, 22]. However, a study reported less satisfaction when dealing with new patients and favored a blended program including face-to-face visits and telerehabilitation [24] (Table 2).

3.8. Risk of Bias. Based on Joanna Briggs Institute (JBI) critical appraisal tool, the majority of the studies (n = 9) were

categorized as having a low risk of bias and 6 studies, moderate risk of bias. Details about the risk of bias are shown in Figure 2.

4. Discussion

The current systematic review found that patients in the majority of the studies are satisfied with both telerehabilitation and conventional delivery method of treatment for MSD [8–10, 19–22]. Satisfaction with telehealth approach also has been reported in systematic reviews on telermatology [30], telepsychiatry [31], and telemedicine for stroke [32].

The current review is providing promising results about satisfaction that are in line with the other clinical cases dealt with telerehabilitation. In poststroke telerehabilitation, professionals and patients reported high level of satisfaction and acceptance with remote intervention through Internet-based video conference in their home setting [3, 33]. Moreover, a study on stroke survivors reported a better relationship of patients with their providers in telegroup [34]. Patient satisfaction with telerehabilitation is imperative as it influences compliance and motivation towards intervention as patients are more engaged with their rehab professionals and confidence level is enhanced.

The high level of satisfaction with telerehabilitation among the professionals found in two studies is consistent with earlier reports [30, 32]. Three areas of satisfaction assessed, patient performance, patient-provider relationship, and quality of technology are related to the capability of professionals to adapt and accept of the new technology in their practice. However, professionals in one study are less

TABLE 1: Patient satisfaction with telerehabilitation.

Authors (years)	Country	Study design & quality of evidence ^a	Sample size	Study population	Technology explored	Variables assessed	Study demonstrate satisfaction	Findings
1 Bimi and Mahajan, 2017 [10]	USA	RCT,1	28 (intervention, n = 13; face-to-face, n = 15)	Post TKA	Asynchronous video-based mobile application	(i) Satisfaction with the experience (ii) Overall satisfaction	Yes (TR = FTF)	Patient satisfaction was high for both groups with no difference
2 Culliton et al., 2018 [19]	Canada	RCT,1	319 (intervention, n = 154; face-to-face, n = 165)	Post TKA	Online e-learning tool	(i) Patient Acceptable Symptom State (PASS) question (ii) Satisfaction with current health state	Yes (TR = FTF)	78.6% in the intervention and 78.2% in the control groups were satisfied
3 Doiron-Cadriu et al., 2020 [20]	Canada	RCT,1	22 (intervention, n = 11; face-to-face, n = 11)	Pre TKA/THA	Reacts lite application	Telecommunication applications	Yes (TR = FTF)	Patient satisfaction was excellent toward telerehabilitation without any significant association between the groups
4 Tousignant et al., 2011 [8]	Canada	RCT,1	42 (intervention, n = 22; face-to-face, n = 20)	Post TKA	Video conferencing	(i) Healthcare satisfaction questionnaire (ii) Satisfaction with the relationship with the healthcare professional (iii) Satisfaction with the services delivered, (iv) Satisfaction with the general healthcare organization	Yes (TR = FTF)	Patient satisfaction was observed in both groups (tele and comparison) without any significant association between the groups
5 Nelson et al., 2020 [21]	Australia	RCT,1	69 (intervention, n = 34; face-to-face, n = 35)	THA	eHAB application	(i) Satisfaction questionnaire of the rehabilitation program (ii) Overall satisfaction	Yes (TR > FTF)	In both groups (intervention & tele), overall satisfaction was high but the score regarding ease of attending appointment was high in the intervention group as compared to telegroup
6 Salazar-Fernandez et al., 2012 [25]	Spain	NRCT,2	992 (intervention, n = 282; face-to-face, n = 710)	Temporomandibular joint disorders	Store-and-forward telemedicine system (SFTMS)	(i) Telemedicine again (ii) Feel uncomfortable quickly (iii) Prefer to be treated (iv) Travel saving (v) Hours of lost-work saving	Yes (TR = FTF)	283 patients reported that they would like to consult by teleconsultation again There was no record of uncomfortable incidence from any patient with teleconsultation

TABLE 1: Continued.

Authors (years)	Country	Study design & quality of evidence ^a	Sample size	Study population	Technology explored	Variables assessed	Study demonstrate satisfaction	Findings
7 Katt et al., 2012 [24]	USA	PCS,2	167 (follow-up, $n = 111$; new, $n = 56$)	Upper extremity conditions	A telephone call or video	(i) Quality of time spent with the provider, preference of telehealth versus an in office encounter, or a combination of both	Yes (TR = FTF)	Both patient and physician responded that they are very satisfied with the telehealth sessions but some patients should have telehabilitation program incorporated with some face-to-face evaluation sessions as they were less satisfied while evaluating new patients
8 Mayer et al., 2021 [26]	Israel	NRCT,2	18 (intervention, $n = 9$; face-to-face, $n = 9$)	Upper limb function after fractures	Biofeedback system of elbow motion	(i) Patient satisfaction questionnaire	Yes (TR = FTF)	The telehabilitation group reported a higher level of enjoyment of the self-practice with less support from family members and no difference was found regarding satisfaction level between two groups
9 Tsvyakb, 2018 [23]	Ukraine	CSS,4	74 (intervention, $n = 48$; face-to-face, $n = 26$)	Injuries of the lower extremities	Home remote monitoring by using a smartphone	(i) Overall satisfaction	Yes (TR > FTF)	Patient satisfaction was higher for the telehabilitation as compared to traditional rehabilitation due to the reason of time saving and cost of rehabilitation
10 Moffet et al., 2017 [9]	Canada	RCT,1	182 (intervention, $n = 84$; face-to-face, $n = 98$)	Post TKA	Videoconferencing system	(i) Relationship with the professional (ii) Delivery of services (iii) Organization of services	Yes (TR = FTF)	Overall satisfaction was high in both groups without any difference in both groups
11 Pani et al., 2017 [13]	Italy	NRCT,2	40 (intervention, $n = 20$; face-to-face, $n = 20$)	Hand function impairment in rheumatic patients	Sensorized tools for hand exercises	(i) Satisfaction of the product (ii) Associated services	Yes (TR > FTF)	Most of the patients were satisfied with the services and accepted the telehabilitation system
12 Jansen-Kosterink et al., 2015 [27]	Netherlands	CSS,4	60 (intervention, $n = 41$; face-to-face, $n = 19$)	CLBP	Teleconference	(i) Rate satisfaction with service (ii) Recommend the service to another patient	Yes (TR = FTF)	About 70% telehabilitation scored 6 or higher score on a scale from 0 to 10 and 36% reported that they would like to recommend the telehabilitation services to another patient
13 Bailey et al., 2020 [28]	USA	RCS,3	Back pain ($n = 6468$), knee pain ($n = 3796$)	Low back & knee pain	Hinge health app installed	Overall satisfaction	Yes (TR = FTF)	There was an overall satisfaction among patients in Digital Care Program (DCP) with a final satisfaction score of 8.97/10

TABLE 1: Continued.

Authors (years)	Country	Study design & quality of evidence ^a	Sample size	Study population	Technology explored	Variables assessed	Study demonstrate satisfaction	Findings
14 Peterson, 2018 [29]	USA	CSS,4	3	CLBP	Mobile phone application with synchronous audio and video booster sessions	Overall satisfaction	Yes (TR = FTF)	The overall satisfaction with the telerehabilitation program was very high
15 Cottrell et al., 2019 [22]	Australia	NRCT,2	61 (intervention, n = 46; face-to-face, n = 15)	LBP & neck pain	Mobile phone application	(i) Cost (ii) Access (iii) Time (iv) Overall satisfaction	Yes (TR = FTF)	(i) The overall satisfaction between two groups was similar (ii) The satisfaction level was high for responses relating to access, cost and time saving, and overall experience in TR group (iii) The satisfaction of professionals regarding appointments was averaged 4.1 out of 5 points

TR: telerehabilitation; FTF: face-to-face; RCT: randomized clinical trial; NRCT: nonrandomized clinical trial; CCS: case-control study; CSS: case-series study; RCS: retrospective cohort study; PCS: prospective cohort study CLBP: chronic low back pain; LBP: low back pain; TKA: total knee arthroplasty; THA: total hip arthroplasty. ^aLevel of evidence. (1) Properly designed RCT or systematic review with meta-analysis; (2) well-designed controlled trial without randomization, prospective study, or comparative cohort trial; (3) case-control study or retrospective cohort study; and (4) case-series or cross-sectional study.

TABLE 2: Professional satisfaction with telerehabilitation.

Authors (years)	Country	Quality of evidence ^a	Sample size	Study population	Technology explored	Type of provider	Overall satisfaction	Findings
1 Tousignant et al., 2011 [8]	Canada	RCT,1	42 (intervention, n = 22; face-to-face, n = 20)	Post-TKA	Video conferencing	Physiotherapists	Yes	Physiotherapists were highly satisfied regarding the following: (i) Goal achievement (ii) Patient-therapist relationship (iii) Overall session satisfaction (iv) Quality and performance of the technological platform Physician responded that they are very satisfied with the telehealth sessions but reported that some patients should have telerehabilitation program incorporated with some face-to-face evaluation sessions, as they were less satisfied while evaluating new patients
2 Katt et al., 2012 [24]	USA	RCS,3	167 (follow-up, n = 111; new, n = 56)	Upper extremity conditions	A telephone call or video	Physicians	Yes	The satisfaction of professionals regarding appointments was high (averaged 4.1 out of 5 points)
3 Cottrell et al., 2019 [22]	Australia	NRCT,2	61 (intervention, n = 46; face-to-face, n = 15)	LBP & neck pain	Mobile phone application	Rehab professionals	Yes	

TR: telerehabilitation; FTF: face-to-face; RCT: randomized clinical trial; NRCT: nonrandomized clinical trial; RCS: retrospective cohort study; LBP: low back pain; TKA: total knee arthroplasty; THA: total hip arthroplasty. ^aLevel of evidence. (1) Properly designed RCT or systematic review with meta-analysis; (2) well-designed controlled trial without randomization, prospective study, or comparative cohort trial; (3) case-control study or retrospective cohort study; and (4) case-series or cross-sectional study^{*}.

atisfied with telerehabilitation when dealing with new patients but were satisfied with it in the follow-up sessions as they reported that certain conditions are hard to assess through telehealth when a radiograph is needed [24].

The above findings suggest that telehealth is an accepted service delivery method by patients and health care professionals but it is not without limitation. It may not be suitable for an initial consultation because in some cases physical examination, blood sample, and radiographic imaging are required to make a definitive diagnosis but more suitable for follow-up sessions. Telerehabilitation is more effective if a patient's record is available to the provider during a consultation session. A combination of face-to-face and telehealth consultation is perhaps a reasonable option to address the issue.

Apart from the advantages of telerehabilitation in reducing cost, time, and disparities in access and satisfying the expectation of patients and professionals [35], it also serves as a rational option for service delivery method during the current COVID-19 pandemic. The health authorities are opting the telehealth to prevent the spread of infection among communities [36, 37]. The approach reduces the number of physical visit and contact between patients and

other individuals during travel and at the healthcare centre and thus lowers the risk of infection [38]. As the health professional and patient relationship is a core element for successful treatment using telehealth, the gap has to be reduced by improving the communication skill of the provider and instilling the confidence of patients in the professionals [39]. Other issues that need consideration in improving the relationship include feasibility, acceptability, and utility [40]. These core factors indicate the willingness and capabilities of professionals to deliver the healthcare services to their patients. Accessibility, awareness, and skills relating the teletechnology among patients also influence the usage and success of telehealth delivery. The patient's expectations are also met with telerehabilitation [41]. Hence, it is imperative to enhance the efforts to use e-health technology in rehabilitation settings.

This is the first systematic review relating to satisfaction with telerehabilitation that is providing sound scientific knowledge to the body of literature and useful for policy-makers for future clinical decision-making. However, it has some limitations; thus, interpretation of the findings should be made with caution. There is wide variation in the measures of assessment and reporting of patients and

professionals' satisfaction despite the evidence being supportive of telerehabilitation. In this systematic review, we found heterogeneity in terms of study design, setting of studies, and outcome measures but there is consistency in the satisfaction and effectiveness with telerehabilitation services. Meta-analysis was not performed due to pooling of the results of different outcome measures which could lead to heterogeneity and bias meta-analysis [42]. The evidence thus far is from the developed countries, and its performance in the developing and underdeveloped countries has yet been reported; thus, further research on telerehabilitation in these countries is recommended.

5. Conclusion

In our systematic review, we contributed a substantial amount of literature about satisfaction with telerehabilitation that is relatively a new area to be explored. The findings of this review point out that telerehabilitation has similar level of satisfaction that is comparable with a face-to-face consultation among patient and professionals alike.

Data Availability

All the data related to the manuscript are included within the article.

Conflicts of Interest

The authors declare no conflict of interest.

Supplementary Materials

[1] PRISMA 2020 checklist (supplementary file 1). [2] Search strategy (supplementary file 2). (*Supplementary Materials*)

References

- [1] M. McCue, A. Fairman, and M. Pramuka, "Enhancing quality of life through telerehabilitation," *Physical Medicine and Rehabilitation Clinics*, vol. 21, no. 1, pp. 195–205, 2010.
- [2] J. Amin, A. A. Siddiqui, S. Al-Oraibi et al., "The potential and practice of telemedicine to empower patient-centered healthcare in Saudi Arabia," *International Medical Journal*, vol. 27, no. 2, pp. 151–154, 2020.
- [3] F. S. Sarfo, U. Ulasavets, O. K. Opare-Sem, and B. Ovbiagele, "Tele-rehabilitation after stroke: an updated systematic review of the literature," *Journal of Stroke and Cerebrovascular Diseases*, vol. 27, no. 9, pp. 2306–2318, 2018.
- [4] L. F. de Rezende, V. E. Francisco, and R. L. Franco, "Telerehabilitation for patients with breast cancer through the COVID-19 pandemic," *Breast cancer research and treatment*, vol. 185, no. 1, pp. 257–259, 2021.
- [5] R. Hwang, J. Bruning, N. Morris, A. Mandrusiak, and T. Russell, "A systematic review of the effects of telerehabilitation in patients with cardiopulmonary diseases," *Journal of cardiopulmonary rehabilitation and prevention*, vol. 35, no. 6, pp. 380–389, 2015.
- [6] M. A. Cottrell, O. A. Galea, S. P. O'Leary, A. J. Hill, and T. G. Russell, "Real-time telerehabilitation for the treatment of musculoskeletal conditions is effective and comparable to standard practice: a systematic review and meta-analysis," *Clinical rehabilitation*, vol. 31, no. 5, pp. 625–638, 2017.
- [7] C. S. Kruse, N. Krowski, B. Rodriguez, L. Tran, J. Vela, and M. Brooks, "Telehealth and patient satisfaction: a systematic review and narrative analysis," *BMJ Open*, vol. 7, no. 8, article e016242, 2017.
- [8] M. Tousignant, P. Boissy, H. Moffet et al., "Patients' satisfaction of healthcare services and perception with in-home telerehabilitation and physiotherapists' satisfaction toward technology for post-knee arthroplasty: an embedded study in a randomized trial," *Telemedicine and e-Health*, vol. 17, no. 5, pp. 376–382, 2011.
- [9] H. Moffet, M. Tousignant, S. Nadeau et al., "Patient satisfaction with in-home telerehabilitation after total knee arthroplasty: results from a randomized controlled trial," *Telemedicine and e-Health*, vol. 23, no. 2, pp. 80–87, 2017.
- [10] S. Bini and J. Mahajan, "Clinical outcomes of remote asynchronous telerehabilitation are equivalent to traditional therapy following total knee arthroplasty: a randomized control study," *Journal of Telemedicine and Telecare*, vol. 23, no. 2, pp. 239–247, 2017.
- [11] N. Toh, J. Pawlovich, and S. Grzybowski, "Telehealth and patient-doctor relationships in rural and remote communities," *Canadian Family Physician*, vol. 62, no. 12, pp. 961–963, 2016.
- [12] K. M. O'Brien, R. K. Hodder, J. Wiggers et al., "Effectiveness of telephone-based interventions for managing osteoarthritis and spinal pain: a systematic review and meta-analysis," *PeerJ*, vol. 6, article e5846, 2018.
- [13] D. Pani, M. Piga, G. Barabino et al., "Home tele-rehabilitation for rheumatic patients: impact and satisfaction of care analysis," *Journal of Telemedicine and Telecare*, vol. 23, no. 2, pp. 292–300, 2017.
- [14] L. G. Fernandes, H. Devan, S. J. Kamper, C. M. Williams, and B. T. Saragiotto, "Enablers and barriers of people with chronic musculoskeletal pain for engaging in telehealth interventions: protocol for a qualitative systematic review and meta-synthesis," *Systematic Reviews*, vol. 9, no. 1, pp. 1–7, 2020.
- [15] A. Liberati, D. G. Altman, J. Tetzlaff et al., "The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration," *Journal of Clinical Epidemiology*, vol. 62, no. 10, pp. e1–e34, 2009.
- [16] K. Porritt, J. Gomersall, and C. Lockwood, "JBI's systematic reviews: study selection and critical appraisal," *AJN The American Journal of Nursing*, vol. 114, no. 6, pp. 47–52, 2014.
- [17] F. Alshammari, A. A. Siddiqui, A. S. Al-Enizy et al., "Prevalence of dental fluorosis in Saudi Arabia: a meta-analysis," *Pesquisa Brasileira em Odontopediatria e Clínica Integrada*, vol. 20, 2020.
- [18] S. Moola, Z. Munn, K. Sears et al., "Conducting systematic reviews of association (etiology): the Joanna Briggs Institute's approach," *JBI Evidence Implementation*, vol. 13, no. 3, pp. 163–169, 2015.
- [19] S. E. Culliton, D. M. Bryant, S. J. MacDonald, K. M. Hibbert, and B. M. Chesworth, "Effect of an e-learning tool on expectations and satisfaction following total knee arthroplasty: a randomized controlled trial," *The Journal of Arthroplasty*, vol. 33, no. 7, pp. 2153–2158, 2018.

- [20] P. Doiron-Cadrin, D. Kairy, P.-A. Vendittoli, V. Lowry, S. Poitras, and F. Desmeules, "Feasibility and preliminary effects of a tele-prehabilitation program and an in-person prehabilitation program compared to usual care for total hip or knee arthroplasty candidates: a pilot randomized controlled trial," *Disability and Rehabilitation*, vol. 42, no. 7, pp. 989–998, 2020.
- [21] M. Nelson, M. Bourke, K. Crossley, and T. Russell, "Telerehabilitation is non-inferior to usual care following total hip replacement – a randomized controlled non-inferiority trial," *Physiotherapy*, vol. 107, pp. 19–27, 2020.
- [22] M. A. Cottrell, S. P. O'Leary, M. Raymer, A. J. Hill, T. Comans, and T. G. Russell, "Does telerehabilitation result in inferior clinical outcomes compared with in-person care for the management of chronic musculoskeletal spinal conditions in the tertiary hospital setting? A non-randomised pilot clinical trial," *Journal of Telemedicine and Telecare*, vol. 27, no. 7, pp. 444–452, 2021.
- [23] A. Tsyakh, "Telerehabilitation of patients with injuries of the lower extremities," *Annals of Physical and Rehabilitation Medicine*, vol. 61, pp. e515–e516, 2018.
- [24] B. M. Katt, C. Imbergamo, D. Fletcher et al., "Telehealth for upper extremity conditions: perceptions of the patient and provider," *JAAOS Global Research & Reviews*, vol. 4, no. 9, article e20.00127, 2020.
- [25] C. I. Salazar-Fernandez, J. Herce, A. Garcia-Palma, J. Delgado, J. F. Martín, and T. Soto, "Telemedicine as an effective tool for the management of temporomandibular joint disorders," *Journal of Oral and Maxillofacial Surgery*, vol. 70, no. 2, pp. 295–301, 2012.
- [26] N. Mayer, S. Portnoy, R. Palti, and Y. Levanon, "The efficacy of tele-rehabilitation program for improving upper limb function among adults following elbow fractures: a pilot study," *Applied Sciences*, vol. 11, no. 4, p. 1708, 2021.
- [27] S. Jansen-Kosterink, R. Huis, D. Wever, H. Hermens, and M. Vollenbroek-Hutten, "Introducing remote physical rehabilitation for patients with chronic disorders by means of telemedicine," *Health and Technology*, vol. 5, no. 2, pp. 83–90, 2015.
- [28] J. F. Bailey, V. Agarwal, P. Zheng et al., "Digital care for chronic musculoskeletal pain: 10,000 participant longitudinal cohort study," *Journal of Medical Internet Research*, vol. 22, no. 5, article e18250, 2020.
- [29] S. Peterson, "Telerehabilitation booster sessions and remote patient monitoring in the management of chronic low back pain: a case series," *Physiotherapy Theory and Practice*, vol. 34, no. 5, pp. 393–402, 2018.
- [30] J. S. Mounessa, S. Chapman, T. Braunberger et al., "A systematic review of satisfaction with teledermatology," *Journal of Telemedicine And Telecare*, vol. 24, no. 4, pp. 263–270, 2018.
- [31] M. Langarizadeh, M. S. Tabatabaei, K. Tavakol, M. Naghipour, A. Rostami, and F. Moghbeli, "Telemental health care, an effective alternative to conventional mental care: a systematic review," *Acta Informatica Medica*, vol. 25, no. 4, pp. 240–246, 2017.
- [32] T. Johansson and C. Wild, "Telemedicine in acute stroke management: systematic review," *International journal of technology assessment in health care*, vol. 26, no. 2, pp. 149–155, 2010.
- [33] K.-H. Lin, C.-H. Chen, Y.-Y. Chen et al., "Bidirectional and multi-user telerehabilitation system: clinical effect on balance, functional activity, and satisfaction in patients with chronic stroke living in long-term care facilities," *Sensors*, vol. 14, no. 7, pp. 12451–12466, 2014.
- [34] L. Piron, A. Turolla, P. Tonin, F. Piccione, L. Lain, and M. Dam, "Satisfaction with care in post-stroke patients undergoing a telerehabilitation programme at home," *Journal of Telemedicine and Telecare*, vol. 14, no. 5, pp. 257–260, 2008.
- [35] J. Kvedar, M. J. Coye, and W. Everett, "Connected health: a review of technologies and strategies to improve patient care with telemedicine and telehealth," *Health Affairs*, vol. 33, no. 2, pp. 194–199, 2014.
- [36] A. A. Siddiqui, F. Alshammary, J. Amin et al., "Knowledge and practice regarding prevention of COVID-19 among the Saudi Arabian population," *Work*, vol. 66, no. 4, pp. 767–775, 2020.
- [37] F. Alshammary, A. A. Siddiqui, J. Amin et al., "Prevention knowledge and its practice towards COVID-19 among general population of Saudi Arabia: a gender-based perspective," *Current pharmaceutical design*, vol. 27, no. 13, pp. 1642–1648, 2021.
- [38] J. R. Falvey, C. Krafft, and D. Kornetti, "The essential role of home- and community-based physical therapists during the COVID-19 pandemic," *Physical therapy*, vol. 100, no. 7, pp. 1058–1061, 2020.
- [39] W. Ngwa, I. Olver, and K. M. Schmeler, "The use of health-related technology to reduce the gap between developed and undeveloped regions around the globe," *American Society of Clinical Oncology Educational Book*, vol. 40, no. 40, pp. 227–236, 2020.
- [40] T. H. Broens, R. M. Huisin't Veld, M. M. Vollenbroek-Hutten, H. J. Hermens, A. T. van Halteren, and L. J. Nieuwenhuis, "Determinants of successful telemedicine implementations: a literature study," *Journal of Telemedicine and Telecare*, vol. 13, no. 6, pp. 303–309, 2007.
- [41] S. Peterson, J. Young, V. King, and J. Meadows, "Patient expectations for synchronous telerehabilitation visits: a survey study of telerehabilitation-naïve patients," *Telemedicine and e-Health*, vol. 28, no. 3, pp. 422–432, 2022.
- [42] M. A. Puhan, I. Soesilo, G. H. Guyatt, and H. J. Schünemann, "Combining scores from different patient reported outcome measures in meta-analyses: when is it justified?," *Health and Quality of Life Outcomes*, vol. 4, no. 1, pp. 1–8, 2006.