




SUPPLEMENT ARTICLE

State-of-the-art of measures of the obesogenic environment for children

Kun Mei^{1,2} | Hong Huang^{1,2} | Fang Xia³ | Andy Hong^{4,5} | Xiang Chen⁶ |
 Chi Zhang² | Ge Qiu⁵ | Gang Chen² | Zhenfeng Wang^{1,2} | Chongjian Wang⁷  |
 Bo Yang^{8,9} | Qian Xiao^{10,5}  | Peng Jia^{11,1,12,5} 

¹Health Assessment Center, Wenzhou Medical University, Wenzhou, China

²Zhejiang Provincial Key Laboratory of Watershed Science and Health, School of Public Health and Management, Wenzhou Medical University, Wenzhou, China

³School of Life Science, Shaoxing University, Shaoxing, China

⁴The George Institute for Global Health, University of Oxford, Oxford, UK

⁵International Institute of Spatial Lifecourse Epidemiology (ISLE), Hong Kong, China

⁶Department of Geography, University of Connecticut, Storrs, Connecticut, USA

⁷Department of Epidemiology and Biostatistics, College of Public Health, Zhengzhou University, Zhengzhou, China

⁸Institute of Lipids Medicine, Wenzhou Medical University, Wenzhou, China

⁹School of Public Health, Wenzhou Medical University, Wenzhou, China

¹⁰Department of Epidemiology, Human Genetics, and Environmental Sciences, The University of Texas Health Science Center at Houston, Houston, Texas, USA

¹¹Department of Land Surveying and Geo-Informatics, The Hong Kong Polytechnic University, Hong Kong, China

¹²Faculty of Geo-information Science and Earth Observation, University of Twente, Enschede, The Netherlands

Correspondence

Peng Jia, PhD, Director, International Institute of Spatial Lifecourse Epidemiology (ISLE); Department of Land Surveying and Geo-Informatics, The Hong Kong Polytechnic University, Hong Kong, China.
 Email: jiapengff@hotmail.com

[Correction added on 3 February 2021, after first online publication: Peng Jia's correspondence details have been updated. Also, affiliations 11 and 12 were interchanged.]

Qian Xiao, PhD, Department of Epidemiology, Human Genetics, and Environmental Sciences, The University of Texas Health Science Center at Houston, Houston, TX, USA.
 Email: qian.xiao@uth.tmc.edu

Funding information

State Key Laboratory of Urban and Regional Ecology of China, Grant/Award Number: SKLURE2018-2-5

Summary

Various measures of the obesogenic environment have been proposed and used in childhood obesity research. The variety of measures poses methodological challenges to designing new research because methodological characteristics integral to developing the measures vary across studies. A systematic review has been conducted to examine the associations between different levels of obesogenic environmental measures (objective or perceived) and childhood obesity. The review includes all articles published in the Cochrane Library, PubMed, Web of Science and Scopus by 31 December 2018. A total of 339 associations in 101 studies have been identified from 18 countries, of which 78 are cross-sectional. Overall, null associations are predominant. Among studies with non-null associations, negative relationships between healthy food outlets in residential neighbourhoods and childhood obesity is found in seven studies; positive associations between unhealthy food outlets and childhood obesity are found in eight studies, whereas negative associations are found in three studies. Measures of recreational or physical activity facilities around the participants'

Kun Mei and Hong Huang contributed equally to this study.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. Obesity Reviews published by John Wiley & Sons Ltd on behalf of World Obesity Federation

home are also negatively correlated to childhood obesity in nine out of 15 studies. Results differ by the types of measurement, environmental indicators and geographic units used to characterize obesogenic environments in residential and school neighbourhoods. To improve the study quality and compare reported findings, a reporting standard for spatial epidemiological research should be adopted.

KEYWORDS

built environment, food environment, obesity, obesogenic environment

1 | INTRODUCTION

Obesity is a leading cause of morbidity and premature mortality worldwide.¹ It has become a severe public health concern among all populations, especially children.² According to the World Health Organization (WHO), over 41 million children under the age of 5 and over 340 million children and adolescents aged 5–19 had overweight or obesity as of 2016.³ Obesity has nearly tripled worldwide since 1951. The increasing obesity rate has particularly affected upper-middle-income countries with high rates of urbanization.³ The Centers for Disease Control and Prevention (CDC) has reported that in the United States, one out of 6 children and adolescents are suffering from obesity.⁴ Childhood obesity often accompanies and leads to more serious chronic health problems, such as high blood pressure, high cholesterol, type II diabetes, asthma, sleep apnoea, fatty liver disease, gallstones, gastro-oesophageal reflux, joint problems and musculoskeletal discomfort.^{5–11} Childhood obesity is also related to contingencies in mental health, such as anxiety, depression, low self-esteem, poor quality of life and may, as a result, induce social issues, such as bullying and stigma.^{12–14} Children with overweight or obesity have increased risks of developing obesity-related comorbidities, including heart disease and cancer.¹⁵

The obesogenic environment is defined as the 'sum of the influences that the surroundings, opportunities or conditions of life have on promoting obesity in individuals and populations'.^{16,17} The obesogenic environment at the neighbourhood scale may interact with personal characteristics to influence individual's weight status. Modifiable environmental factors manifest as an indirect effect on individual's diet behaviour and physical activity. First, dietary behaviours can be shaped by the community nutrition environment (generally known as the community food environment), defined as types, locations and temporality of food outlets (e.g., supermarkets, convenience stores or fast-food restaurant) in the community.^{18,19} A quality community nutrition environment characterized by affordable and accessible food sources in the near proximity of the residential place is necessary for children and adolescents to procure nutritious food items and practice healthy diet behaviour.²⁰ Second, the proximity to a recreational or physical activity facility, such as park, playground or gym, will increase the likelihood of physical activity engagement and will decrease rates of sedentary activity, eventually mitigating risks of obesity. For example, in neighbourhoods with relatively good

walkability (e.g., more sidewalks), people are more likely to engage in physical activity such as walking and cycling, while significantly reducing time spent on sedentary activity, such as watching TV, driving and sitting.²¹ Third, there are contextual factors in the obesogenic environment that shape both diet behaviour and physical activity.²² These contextual factors include the affordability of healthy food options, peer and social supports, marketing and promotion and planning policies on the sustainability of the community design.⁴ In this review, we mainly focus on the physical aspect of the obesogenic environment and will not include these contextual factors.

Previous reviews have examined the associations between obesity and various measures of the obesogenic environment. Some studies argue that evaluations by these measures differ by age group and vary across countries. A recent review found that associations between the community food environment and obesity were less likely to be significant among children than adults in the United States and Canada.²³ Another review conducted an extended scope of work in four European and Oceanian countries (i.e., the United Kingdom, Ireland, Australia and New Zealand) and compared the findings with the North America.²⁴ Even among children, associations between the community-based obesogenic variables and obesity differed by gender, age and socio-economic status.²⁵ In addition to these regional comparisons, the association may also vary by the definition of the community or neighbourhood. Neighbourhood is loosely defined as a physical extent where individuals engage in communal activities with local residents.²⁶ This definition focused on a physical space has been further extended to the perceived neighbourhood or the geographic extent conceptualized by people as their communal space. It has been found that individuals tend to perceive their living neighbourhood as being smaller than the administrative unit (e.g., census tract and postal zone) where they reside. This means that the actual scale where the contextual factors affect individuals' health status could be very different from those derived from the administrative unit.²⁷ There have been no consensus in obesity studies about the most appropriate scales and measures where obesogenic environmental factors should be employed. For example, it was noted that the majority of food environment studies were employed at the community or neighbourhood scale in terms of schools, work sites and households²³; measures of the food environment included the availability, variety, accessibility and density of food outlets. In addition, a systematic review on green space and obesity reported that two most

common measures of the physical access were distance (Euclidean or network) to near green spaces and the count of green spaces in the vicinity of the residential place.²⁸ Despite the accumulation of research using various environmental measures, there is still lack of consensus on how to define the obesogenic environment for children.^{28,29}

This review contributes to the literature in two major aspects. First, we have systematically reviewed a full scope of literature using both objective and perceived measures of the obesogenic environment applied to childhood obesity research. Second, this review has summarized the different levels of associations between these measures and childhood obesity. This study will inform researchers about the availability, consistency and significance of these environmental measures. Furthermore, this review will shed important insights into childhood obesity research that employs a multiscale framework for intraregional and interregional comparisons.¹⁸

2 | METHODS

A systematic review was conducted in accordance with the Preferred Reporting Items for Systematic Reviews (PRISM).

2.1 | Study selection criteria

Our study inclusion criteria were as follows: (1) the study included at least one measure of the obesogenic environment, (2) the study outcome was obesity (including overweight) instead of other health outcomes, (3) the study was focused on the association with obesity rather than the obesogenic environment (e.g., food environments) *per se* or obesity-related behaviours (e.g., diet behaviour and physical activity) *per se*, (4) the study was focused on the obesity of children aged younger than 18 years and (5) the study was an original research article published in English.

2.2 | Search strategy

A keyword search was performed in four electronic bibliographic databases: Cochrane Library, PubMed, Web of Science and Scopus. The search strategy included all possible combinations of keywords, including the obesogenic environment (mainly built environment and food environment), children and adolescents and weight-related outcomes (Appendix A). To increase the coverage of the literature, we manually searched the reference lists in a snowball approach and cited relevant articles with an end search date of 31 December 2018.

Titles and abstracts of the articles identified through the keyword search were screened against the study selection criteria. The full text of potentially relevant articles was retrieved for scrutiny and integration. Two reviewers independently conducted the title and abstract screening and identified potentially relevant articles for the full-text review. Discrepancies were screened by a third reviewer. The three

reviewers jointly determined the list of articles for the full-text review through several rounds of discussion. Two reviewers then independently reviewed the full texts of all articles in the list and determined the final pool of articles included in the review.

2.3 | Data extraction

For each selected study, we adopted a standardized data extraction process to collect methodological and outcome variables, including authors, year of publication, study area, country, study year, sample size, age range/age at baseline, sample characteristics (including follow-up years), number of repeated measures, attrition rate (if applicable), statistical model, measures of the obesogenic environment (objective or perceived; residential neighbourhood or school), measures of body-weight status and key findings on the association between obesogenic environments and weight-related outcomes. Two reviewers independently extracted data from each study included in the review, and discrepancies were resolved by the third reviewer.

3 | RESULTS

3.1 | Study selection

Figure 1 shows the study selection flow chart. We identified a total of 4629 articles through the keyword search process. The search underwent title and abstract screening, by which 1697 articles were excluded. The full texts of the remaining 106 articles were reviewed against the study selection criteria. Of these full-text articles, five articles were excluded. The remaining 101 studies that examined the relationship between the obesogenic environment and weight-related outcomes were included in this review.

3.2 | Study characteristics

The main characteristics of the 101 included articles were presented in Table 1. All studies were published after 2004. The age of participants ranged from 2 to 18, with 76 cross-sectional studies, 23 longitudinal studies, and two repeated cross-sectional studies. These 101 studies covered 18 countries: 65 studies were conducted in North America, with 53 studies from the United States and 12 studies from Canada; 11 studies were from the United Kingdom; 16 were from Australia, Germany and China, with four studies from each country; two were from Brazil; and the rest were from France, Ireland, Lithuania, Malaysia, Mexico, Netherland, Portugal, South Korea, Spain, Sweden and Ukraine, with one study per country.

The geographic scales of these studies varied from country to county, while the number of participants ranged from 78 to 3 003 288. These studies were conducted at different geographic scales, including nationwide ($n = 11$), provincial ($n = 17$), multistate




FIGURE 1 Study exclusion and inclusion flowchart

($n = 1$), multicity ($n = 3$), single city ($n = 13$), multicounty ($n = 1$) and single county ($n = 7$). Most of the studies accounted for multilevel data and applied multivariable regression models for data analysis ($n = 86$, 85%), including linear regression model ($n = 27$, 27%) and logistic regression model ($n = 38$, 38%). Other methods, such as the correlation analysis ($n = 2$) and the multilevel growth curve model ($n = 3$), were also employed. Study outcomes included the absolute value of the body mass index (BMI), BMI percentile or z score, rate of obesity or overweight and change in BMI or weight.

3.3 | Diversity of measurements

The most common types of the obesogenic environment under examination were residential neighbourhoods ($n = 96$) (Table S1) and school neighbourhoods ($n = 23$) (Table S2). The investigation approaches included objective measures by Geographic Information Systems (GIS) tools ($n = 85$) or neighbourhood perceptions self-reported by the participants, their parents or the school directors ($n = 17$). Both the objective measures and the perceived measures included four environmental indicators, including availability (e.g., presence or not), count (e.g., total number), density (e.g., count/population, count/area) and proximity (e.g., straight-line/network distance). Among the 101 studies examining these indicators, count was the most common measure ($n = 72$), followed by availability ($n = 36$). More complex spatial measures such as the kernel density that weighs outlets near

participants' school ($n = 6$) or moderates the distance to the nearest retail outlet ($n = 2$) were less likely to be employed.

Studies also differed by the geographic units used to assess exposure to the obesogenic environment in residential neighbourhoods or school neighbourhoods. For instance, 22 studies measured the exposure to supermarkets in 20 different ways, and 26 studies assessed the exposure to fast-food restaurants in 17 different ways. Sixteen studies used administrative units, including census tracts ($n = 12$), postal zones ($n = 2$) and predefined grids ($n = 4$; i.e., Middle Super Output Area,²⁹ Street Segments,⁵³ Small Area Market Statistics¹⁶ and Lower Super Output Area¹¹⁵). Residential or school addresses were also used for assessing environmental exposure, buffered by a radius (and was measured either along the road network or by a set distance) (Tables S3 and S4). Buffers ranged in sizes from 0.4 to 6 km. A 1.6-km road-network buffer was the most commonly used criterion ($n = 13$), followed by a 1-km buffer ($n = 11$). Many studies performed sensitivity analyses with buffers of multiple sizes.

3.4 | Association between food environment and obesity

Sixty-five studies examined weight-related outcomes in relation to food environment measures in residential neighbourhoods ($n = 164$) (Table S1) or school neighbourhoods ($n = 72$) (Table S2). Although a high percentage ($n = 146$, 62%) of these associations were null, there

TABLE 1 Basic characteristics of the included studies

First author (year)	Study area [scale] ^a	Study design ^b	Sample size	Age at baseline (years) ^c	Sample characteristics	Statistical models	Outcome variables
Baek (2016) ³⁰	California, USA [S]	C	601 847	10–15 in 2009	FitnessGram test	Distributed lag model	BMI z score
Barrera (2016) ³¹	Cuernavaca and Guadalajara, Mexico [C2]	C	725	9–11 in 2012–2013	Elementary school children	Multiple linear regression	BMI z score
Bell (2008) ³²	Indianapolis, USA [C]	L	3831	3–16 in 1996–2002	Cohort in primary care clinic network, followed up for 2 years with two repeated measurements	Multiple linear regression	BMI z score
Berge (2014) ³³	Minneapolis/St. Paul, USA [C]	C	2682	14–16 in 2009	Eating and Activity in Teens (EAT) survey	Multiple linear regression	BMI z score
Carroll-Scott (2013) ³⁴	New Haven, USA [C]	C	1048	10–11 in 2009	Community interventions for health chronic disease prevention study	Linear regression	BMI
Carter (2012) ³⁵	Quebec, Canada [S]	L	2120	4–10 in 1997–1998	Quebec Longitudinal Study of Child Development cohort, followed up for 7 years with five repeated measurements and attrition rate of 26.1%	Linear regression	BMI z score
Casey (2012) ³⁶	Bas-Rhin, France [S]	C	3327	11–13 in 2001	France middle school students	Mixed logistic regression	Weight, BMI
Cetateanu (2014) ²⁹	UK [N]	C	3 003 288	4–5 and 10–11 in 2007–2010	National Child Measurement Program (NCMP) dataset	Stepwise linear regression	Overweight/obesity
Chaparro (2014) ³⁷	Los Angeles, USA [CT]	L	32 172	2–5 in 2005–2008	Women, Infants and Children (WIC) study, followed up for 4 years with three repeated measurements	Linear regression, multilevel linear growth model	WHZ
Cheah (2012) ³⁸	Kuching, Malaysia [C]	C	316	14–16	Secondary schools students	Univariate data analysis	BMI
Chen (2016) ³⁹	USA [N]	L	7090	11 in 2004–2007	Early Childhood Longitudinal Study-Kindergarten (ECLS-K) cohort, followed up for 4 years with two repeated measurements	Fixed-effect regression	BMI, obesity
Chiang (2017) ⁴⁰	Taiwan, China [S]	C	1458	11–16 in 2010	Nutrition and Health Survey in Taiwan	Multiple linear regression	Height z score, weight z score, BMI score, WC z score, WC/height ratio, WC/hip ratio, TSF z score, MAMC z score
Correa (2018) ⁴¹	Florianópolis, Brazil [C]	C	2195	7–14 in 2012–2013	Public and private school children	Logistic regression	BMI z score, overweight/obesity
Crawford (2010) ⁴²	Melbourne, Australia [C]	L	926	10–12 in 2001	Children's Leisure Activities Study (CLAN), followed up for 5 years with three repeated	Generalized estimating equation	BMI z score, MVPA

(Continues)

TABLE 1 (Continued)

First author (year)	Study area [scale] ^a	Study design ^b	Sample size	Age at baseline (years) ^c	Sample characteristics	Statistical models	Outcome variables
Datar (2015) ⁴³	Ft. Lewis, Ft. Carson, Ft. Drum, Ft. Bragg, Ft. Benning, Ft. Bliss, Ft. Campbell, Ft. Hood, Ft. Polk, Ft. Stewart, Ft. Sill, Ft. Riley, USA [C12]	C	903	12–13 in 2013	measurements and attrition rate of 66% Military Teenagers Environment, Exercise, and Nutrition Study	Multivariate regression	PA, BMI
Davis (2009) ⁴⁴	California, USA [S]	C	529 367	≤19 in 2002–2005	California Healthy Kids Survey	Ordinary least squares regression, logistic regression	Overweight, obesity, BMI
Duncan (2012) ⁴⁵	Boston, USA [C]	C	1034	15–18 in 2007–2008	Boston Youth Survey	Spatial regression, ordinary least squares regression	BMI
Duncan (2012) ⁴⁶	Coventry, UK [C]	C	405	14–15	Pupils	Pearson's product moment correlations	PA, BMI
Duncan (2015) ⁴⁷	Massachusetts, USA [S]	L	49 770	4–12 in 2011–2012	Pediatric practices of Harvard Vanguard Medical Associates, followed up for 1.5 years with two repeated measurements	Multivariable model	BMI z score
Dwiczaksono (2017) ⁴⁸	New York, USA [S]	C	680	In 2010–2012	Student Weight Status Category Reporting System dataset	Ordinary least squares regression, geographically weighted regression	Obesity rate
Edwards (2010) ⁴⁹	Leeds, UK [C]	C	33 594	3–13 in 2004–2005	Leeds primary care trusts record and trends study in Leeds	Geographically weighted regression	BMI
Epstein (2012) ⁵⁰	Erie, USA [CT]	L	191	8–12 in 1997–2005	Four randomized, controlled outcome studies, followed up for 2 years with two repeated measurements	Hierarchical mixed model analyses of covariance	BMI, BMI z score
Fiechtner (2016) ⁵¹	Massachusetts, USA [S]	L	498	6–12 in 2011–2013	Study of Technology to Accelerate Research trail, followed up for 3 years with two repeated measurements and attrition rate of 9%	Generalized linear mixed effects regression	BMI z score
Friedman (2009) ⁵²	Kyiv, Dniprodzerzhynsk and Mariupol, Ukraine [C3]	C	883	3 in 1993–1996	European Longitudinal Study of Pregnancy and Childhood (ELSPAC) cohort	Multivariable logistic regression	Overweight, obesity
Ghenadenik (2018) ⁵³	Quebec, Canada [S]	L	506	8–10 in 2005–2008	Quebec Adipose and Lifestyle Investigation in Youth cohort, followed up for 2 years with two repeated measurements and attrition rate of 19.3%	Multivariable linear regression	BMI z score, WHR

TABLE 1 (Continued)

First author (year)	Study area [scale] ^a	Study design ^b	Sample size	Age at baseline (years) ^c	Sample characteristics	Statistical models	Outcome variables
Gilliland (2012) ⁵⁴	London, UK [C]	C	1048	10–14	28 elementary school	Multilevel structural equation	BMI z score
Gordon-Larsen (2006) ⁵⁵	USA [N]	C	20 745	Grades 7–12 in 1994–1995	Add Health wave I	Logistic regression	Overweight
Gose (2013) ⁵⁶	Kiel, Germany [C]	L	485	6 in 2006–2012	Kiel Obesity Prevention Study (KOPS), followed up for 4 years with two repeated measurements and attrition rate of 72.6%	Generalized estimating equation	BMI standard deviation score
Grafova (2008) ⁵⁷	USA [N]	C	2482	5–18 in 2002–2003	Child Development Supplement survey	Logistic regression	BMI
Green (2018) ⁵⁸	Leeds, UK [C]	L	746	11–12 in 2005–2010	Rugby League and Athletics Development Scheme (RADS), followed up for 5 years with three repeated measurements	Multilevel linear regression	Overweight, obesity
Fiechter (2013) ⁵⁹	Massachusetts, USA [S]	C	438	2–7 in 2006–2009	High Five for Kids (HFK) study	Multivariable linear regression	BMI
Griffiths (2014) ⁶⁰	Leeds, UK [C]	C	13 291	11 in 2005–2007	RADS	Multiple linear and logistic regression	BMI
Guedes (2011) ⁶¹	Minas Gerais, Brazil [S]	C	5100	6–18 in 2007	School children	Binary logistic regression	BMI
Hamano (2017) ¹⁶	Sweden [N]	C	944 487	0–14 in 2005–2010	Swedish nationwide population and health care dataset	Multilevel logistic regression	Obesity
Harris (2011) ⁶²	Maine, USA [S]	C	552	Grades 9–12	Students at 11 Maine high schools	Logistic regression	BMI
Harrison (2011) ⁶³	Norfolk, UK [CT]	C	1724	9–10 in 2007	Sport, Physical Activity and Eating Behaviour: Environmental Determinants in Young People (SPEEDY) study	Multilevel and multivariable hierarchical regression	FMI
Howard (2011) ⁶⁴	California, USA [S]	C	879	Grade 9 in 2007	FitnessGram test	Linear regression	BMI
Hoyt (2014) ⁶⁵	California, USA [S]	L	174	8–10 in 2007–2012	Cohort Study of Young Girls' Nutrition, Environment, and Transitions (CYGNET), followed up for 4 years with at least two repeated measurements and attrition rate of 19.1%	Logistic regression	BMI, obesity
Morgan Hughey (2017) ⁶⁶	USA [CT]	L	13 469	3–5 in 2013	Children in county school district	Multilevel linear regression	BMI
Jennings (2011) ⁶⁷	Norfolk, UK [CT]	C	1669	9–10 in 2007	SPEEDY study	Poisson regression	BMI, weight, BMI z score, WC, % of body fat

(Continues)

TABLE 1 (Continued)

First author (year)	Study area [scale] ^a	Study design ^b	Sample size	Age at baseline (years) ^c	Sample characteristics	Statistical models	Outcome variables
Jerrett (2010) ⁶⁸	California, USA [S]	L	3318	9–10 in 1993 and 1996	Children's Health Study (CHS) cohort, followed up for 8 years with two repeated measurements and attrition rate of 12.9%	Multilevel growth curve model	BMI
Jerrett (2014) ⁶⁹	California, USA [S]	L	4550	5–7 in 2002–2003	A cohort of children attending kindergarten and first grade, followed up for 4 years with four repeated measurements and attrition rate of 6.4%	Multilevel linear regression	BMI
Koleilat (2012) ⁷⁰	Los Angeles, USA [CT]	C	266	3–4 in 2008	WIC study	Simple linear regression	Weight
Lakes (2016) ⁷¹	Berlin, Germany [C]	C	28 159	5–6 in 2012	Berlin children survey	Multivariate regression	% of overweight/obesity
Lange (2011) ⁷²	Kiel, Germany [C]	C	3440	13–15 in 2004–2008	KOPS	Logistic regression	BMI
Larsen (2014) ⁷³	Toronto, Canada [C]	C	943	2–20 in 2010–2011	BEAT	Logistic regression	BMI
Laska (2010) ⁷⁴	Minneapolis/St. Paul, USA [C]	C	349	10–17 in 2006–2007	Identifying Determinants of Eating and Activity Study	Multilevel regression	BMI
Leatherdale (2011) ⁷⁵	Ontario, Canada [S]	C	2449	10–13 in 2007–2008	Play-Ontario (PLAY-ON) study	Multilevel logistic regression	BMI
Leatherdale (2013) ⁷⁶	Ontario, Canada [S]	C	2331	6–9 in 2007–2008	PLAY-ON study	Multilevel logistic regression	Overweight, obesity
Leung (2011) ⁷⁷	California, USA [S]	L	444	6–7 in 2005–2008	CYGNET cohort, followed up for 3 years with two repeated measurements and attrition rate of 20.5%	Generalized linear and logistic regression	BMI z score
Li (2015) ⁷⁸	A rural BBR, USA [CT]	C	613	4–13 in 2013	School children	Multilevel models	BMI percentile
Lovasi (2013) ⁷⁹	New York, USA [C]	C	11 562	3–5 in 2004	Preschool programme	Linear and Poisson regression	BMI z score, obesity
Miller (2011) ⁸⁰	USA [N]	L	11 400	6–12 in 1998–2004	ECLS-K cohort, followed up for 7 years with two repeated measurements	Three-level growth curve model	BMI
Miller (2014) ⁸¹	Perth, Australia [C]	C	1850	5–15 in 2005–2010	Western Australian Health and Wellbeing Surveillance System database	Multivariate logistic regression	BMI
Minaker (2011) ⁸²	Alberta, Canada [S]	C	4936	11–17 in 2005	Web-Survey of Physical Activity and Nutrition study	Multinomial logistic and ordinal regressions	BMI
Molina-García (2017) ⁸³	Valencia, Spain [C]	C	325	14–18 in 2013–2015	International Physical Activity and the Environment Network adolescent study	Mixed regression	BMI, % of body fat
Nelson (2009) ⁸⁴	Ireland [N]	C	4587	15–17 in 2003–2005	Take PART study	Logistic regression	Overweight, obesity

TABLE 1 (Continued)

First author (year)	Study area [scale] ^a	Study design ^b	Sample size	Age at baseline (years) ^c	Sample characteristics	Statistical models	Outcome variables
Nesbit (2014) ⁸⁵	USA [N]	C	39 542	11–17 in 2007	National Survey of Children's Health (NSCH)	Logistic regression	BMI, obesity
Ness (2012) ⁸⁶	USA [N]	C	5342	10–19 in 2007	NSCH	Pooled and race-stratified logistic regression	BMI
Nogueira (2013) ⁸⁷	Coimbra, Portugal [CT]	C	1885	3–10 in 2009	Private and public school children	Logistic regression	BMI
Norman (2006) ⁸⁸	San Diego, USA [CT]	C	799	11–15	Health promotion intervention trial	Multiple linear regression	BMI
Ohri-Vachaspati (2013) ⁸⁹	Camden, New Brunswick, Newark and Trenton, USA [C4]	C	702	3–18 in 2009–2010	Random-digit-dial survey	Logistic regression	Overweight, obesity
Oreskovic (2009) ⁹⁰	Massachusetts, USA [S]	C	6680	2–18 in 2006	Partners HealthCare	Clustered logistic regression	Overweight/obesity
Oreskovic (2009) ⁹¹	Massachusetts, USA [S]	C	21 008	2–18 in 2006	Partners HealthCare	Multilevel logistic regression	Overweight/obesity
Park (2013) ⁹²	Seoul, South Korea [C]	C	1342	10–13 in 2011	Elementary and middle school children	Generalized estimating equation	BMI, weight status
Pearce (2017) ⁹³	South Gloucestershire, UK [S]	L	1577	7 in 2006–2012	NCMP dataset, followed up for 6 years with two repeated measurements	Multiple logistic regression	BMI, WC
Petraviciene (2018) ⁹⁴	Kaunas, Lithuania [C]	C	1498	4–6 in 2012–2013	Positive Health Effects of the Natural Outdoor Environment in Typical Populations in Different Regions in Europe project	Logistic regression	BMI z score
Pitts (2013) ⁹⁵	Greene and Pitt, USA [CT2]	C	296	11–13 in 2008–2010	Middle school children	Linear regression	BMI percentile
Poole (2017) ⁹⁶	Southampton, UK [C]	C	1748	4–5 in 2012–2013	NCMP dataset	Multilevel logistic regression	BMI percentile
Potestio (2009) ⁹⁷	Calgary, Canada [C]	C	6772	5 in 2005–2006	Public health clinics for preschool vaccinations	Two-level, random-intercept logistic regression	BMI
Rossen (2013) ⁹⁸	Baltimore, USA [C]	L	319	8–10 in 2007	Multiple Opportunities to Reach Excellence project cohort, followed up for 1 year with two repeated measurements and attrition rate of 26%	Multilevel model	BMI change, WC change
Gorski Findling (2018) ⁹⁹	USA [N]	C	3748	2–18 in 2012–2013	Food Acquisition and Purchase Survey	Logistic regression	Overweight, obesity
Sánchez (2012) ¹⁰⁰	California, USA [S]	C	926 018	2007	FitnessGram test	Log-binomial regression	BMI
Schmidt (2015) ¹⁰¹	Netherlands [N]	L	1887	4–5 in 2000–2002			BMI z score

(Continues)

TABLE 1 (Continued)

First author (year)	Study area [scale] ^a	Study design ^b	Sample size	Age at baseline (years) ^c	Sample characteristics	Statistical models	Outcome variables
Schüle (2016) ¹⁰²	Munich, Germany [C]	C	3499	5–7 in 2004–2007	Gesundheits-Monitoring-Einheiten survey	Hierarchical logistic regression	BMI, overweight, obesity
Seliske (2009) ¹⁰³	Canada [N]	C	9672	Grades 6–10 in 2005–2006	Health Behaviour in School-Aged Children survey	Multilevel regression	BMI
Seliske (2012) ¹⁰⁴	Canada [N]	C	7017	12–19 in 2007–2008	Canadian Community Health Survey	Multilevel logistic regressions	MVPA, BMI
Singh (2010) ¹⁰⁵	USA [N]	C	44 101	10–17 in 2007–2008	NSCH	Logistic regression	BMI
Slater (2013) ¹⁰⁶	USA [N]	C	11 041	Grades 8, 10 and 12 in 2010	Monitoring the Future (MTF) survey	Multivariable logistic regression	Overweight, obesity
Spence (2008) ¹⁰⁷	Edmonton, Canada [C]	C	501	4–6 in 2004	Preschool immunization	Logistic regression	BMI
Tang (2014) ¹⁰⁸	Camden, New Brunswick, Newark and Trenton, USA [C4]	C	12 954	10–17 in 2008–2009	New Jersey Childhood Obesity study	Random-effects model	BMI z score, overweight, obesity
Taylor (2014) ¹⁰⁹	13 block groups in Southeastern USA [C]	C	911	5–15	Environmental audits and a cross-sectional prevalence study of cardiovascular risk factors	Correlation analysis	Obesity, overweight, WC, WHR
Timperio (2010) ¹¹⁰	Melbourne, Australia [C]	L	409	5–6 and 10–12 in 2001–2004	CLAN; followed up for 3 years with two repeated measurements and attrition rate of 30.7%	Univariate and multivariable linear regression	BMI z score, BMI
Torres (2014) ¹¹¹	San Juan, USA [C]	C	114	12 in 2012–2013	Public school children	Spearman's correlation	BMI percentile
Veugelaers (2008) ¹¹²	Nova Scotia, Canada [S]	C	5471	10–11 in 2003	Children's Lifestyle and School-Performance Study	Multilevel linear regression	Overweight, obesity
Wall (2012) ¹¹³	Minneapolis/St. Paul, USA [C]	C	2682	12–16 in 2009–2010	EAT survey	Multiple linear regression	BMI z score
Wasserman (2014) ¹¹⁴	Kansas, USA [C]	C	12 118	4–12 in 2008–2009	School children	Hierarchical linear	BMI percentile
Williams (2015) ¹¹⁵	UK [N]	C	16 956	4–6 and 10–11 in 2010–2011	NCMP dataset	Multilevel	BMI
Wolch (2011) ¹¹⁶	California, USA [S]	L	3173	9–10 in 1993–1996	CHS cohort, followed up for 8 years with eight repeated measurements	Multilevel growth curve model	BMI change
Xu (2010) ¹¹⁷	Nanjing, China [C]	C	2375	14 in 2004	Nanjing High School Students' Health Survey	Mixed-effect logistic regression	BMI
Yang (2018) ¹¹⁸		C	41 283		Children in SCS	Multilevel logistic regression	BMI

TABLE 1 (Continued)

First author (year)	Study area [scale] ^a	Study design ^b	Sample size	Age at baseline (years) ^c	Sample characteristics	Statistical models	Outcome variables
	Shelby Count, Memphis, USA [CT]			Grades pre-K, K, 2, 4, 6, 8 and 9 in 2014–2015			
Zhang (2016) ¹¹⁹	China [N]	C	348	8–12 in 2009–2011	China Health and Nutrition Survey	Generalized estimating equation	BMI
Sallis (2018) ²¹	Maryland and King County, Washington regions, USA [S2]	C	928	12–16 in 2009–2011	Teen Environment and Neighborhood study	Mixed model linear and logistic regression	BMI percentile
Li (2014) ¹²⁰	Guangzhou and Hechi, China [C2]	C	497	8–10 in 2009–2010	Schools for routine (every 5 years) student health monitoring by local health bureau	Multiple logistic regression and linear regression	Overweight/obesity
Kepper (2016) ¹²¹	Louisiana, USA [S]	C	78	2–5	A randomized controlled trial	Multiple regression analysis	BMI z score
Crawford (2015) ¹²²	Victoria, Australia [S]	L	200	5–12 in 2007–2011	A survey on weight children in socio-economically disadvantaged neighbourhoods, followed up for 3 years with two repeated measurements and attrition rate of 41.3%	Linear and logistic regression	BMI z score, unhealthy weight gain
Powell (2007) ¹²³	USA [N]	C	73 079	13–15 in 1997–2003	MTF survey	Reduced form models	BMI, overweight
Burdette (2004) ¹²⁴	Cincinnati, USA [C]	C	7020	3–5 in 1998–2001	WIC study	Logistic regression	BMI percentile
Sturm (2005) ¹²⁵	USA [N]	L	6918	Grades K, 1 and 3 in 1998–1999	ECLS-K cohort, followed up for 4 years with two repeated measurements	Least squares and quantile regression	BMI change
Potwarka (2008) ¹²⁶	Mid-sized city in Ontario, Canada [C]	C	108	2–17 in 2006	Randomly selected	Logistic regression	Healthy weight
Galvez (2009) ¹²⁷	New York, USA [C]	C	323	6–8 in 2004	Mount Sinai Pediatrics Practice, East Harlem community health centres, community-based organizations and East Harlem schools children	Logistic regression	BMI in top tertile

Abbreviations: BMI, body mass index; FMI, fat mass index; MAMC, mid-arm muscle circumference; PA, physical activity; TSF, triceps skinfold thickness; WC, waist circumference; WHR, waist-height ratio; WHZ, weight-for-height z score.

^a[N], national; [S], state (United States) or equivalent unit (e.g., province in China); [Sn], n states or equivalent units; [CT], county or equivalent unit; [CTn], n counties or equivalent units; [C], city; [Cn], n cities.

^bC, cross-sectional study; L, longitudinal study.

^cAge in baseline year for longitudinal study and age in survey year for cross-sectional study.

were some notable findings. For example, most of the findings (seven out of nine associations in five studies for residential neighbourhoods) on healthy food outlets (e.g., supermarkets) and obesity suggested a negative association between the two, and the association was more apparent for availability, count and density measures than for distance measures. Similarly, the availability of^{39,52,108,123} and the proximity to^{51,59,73} supermarkets were inversely related to obesity. In contrast, the availability of unhealthy food outlets (e.g., convenience stores and fast-food restaurants) was positively associated with obesity in several studies (eight out of 20 associations in 15 studies for residential neighbourhood; three out of 13 associations in 11 studies for school neighbourhoods). For associations between convenience stores and obesity, seven out of 23 associations for residential neighbourhoods and six out of 11 associations for school neighbourhoods were positive. Results for fast-food restaurants were equivocal: although positive associations between fast-food availability and obesity outnumbered negative ones (seven positive vs. three negative), the majority of the associations ($n = 23$, 70%) were null. Evidence for associations with grocery stores (five positive, two negative and 15 null) and full-service restaurants (one negative, one positive and 8 null) was relatively weak.

3.5 | Association between built environment and obesity

Overall, 35 studies examined 85 associations between built environmental measures and weight-related outcomes in residential neighbourhoods (Table S1) and 18 associations in schools (Table S2). Regardless of the type of measurement, null associations were predominant. For studies examining all recreational or physical activity facilities around the participants' residential place, negative associations with obesity were reported ($n = 9$, 60%). Similar patterns emerged with built environment measures calculated for gyms and fitness centres in or around schools (three negative out of four studies). However, the results for parks were mixed. Both positive correlations and negative correlations between the availability of parks (including green spaces and playgrounds) and obesity were identified (three positive vs. six negative for residential neighbourhoods; two positive vs. two negative for school neighbourhoods). Some studies reported that travel-related built environment measures, such as dense traffic roads,^{56,63,68,69,102,110} intersections,⁴⁸ transit stations^{45,49} and traffic signs,¹¹³ had a positive correlation with obesity, whereas others found the correlations to be negative for dense traffic roads^{47,63,91} and intersections.^{47,107,110}

3.6 | Impact of geographic units on associations

The spatial delineation of geographic units affected the results to some extent. In residential neighbourhoods, there were negative associations with healthy food outlets with measures in all buffer sizes for residential neighbourhood (Table S3). On the other hand, the positive

association was dominant between the availability of unhealthy food outlets and obesity within most of geographic units ($n = 8$, 40%) especially administrative unit ($n = 4$, 80%); however, unhealthy food outlets yielded negative associations in 0.8- and 3-km road-network buffers. Some studies also identified mixed results using different geographic units, such as number of grocery store in 0.4-km straight-line buffer¹⁰⁸ and 0.4-km road-network buffer,⁷⁷ and others had even yielded opposite results using same geographic units, such as number of supermarket in postal zone.^{39,70}

To investigate the influence of geographic units on associations, 15 studies used more than one geographic unit, and they reported that the correlation between food outlet and obesity tended to be more significant when analyses were performed using smaller buffer sizes.^{54,81}

4 | DISCUSSION

This systematic review identified 101 studies that examined the associations between obesogenic environmental factors and childhood obesity. Several important findings were identified. First, there was a high degree of heterogeneity in quantifying the obesogenic environment for children. Notably, an obesogenic environment was commonly measured as either objective measures, perceived measures or both. Among the studies that employed both objective and perceived measures, the perceived measures were more likely to yield statistical significance than the objective measures. However, the effect sizes of the perceived measures were relatively small, providing only weak evidence to support a relationship between environmental factors and obesity in children.¹²⁸

Second, the majority of the studies that examined food environment and childhood obesity reported more consistent associations. Among these studies, the most commonly used objective measures were count and availability, and the results varied by the type of food outlet. Fast-food outlets and convenience stores showed more positive associations with childhood obesity. This finding resonates with the widespread concern that the frequent patronization of fast-food outlets and convenience stores has health-damaging effects.¹²⁹ This statistical linkage calls for more rigorous studies to establish the causal pathway to childhood obesity. Likewise, the proximity to supermarkets and farmers' markets showed negative associations with childhood obesity,^{39,50,60,84,88,128} and this effect could be attributed to the higher likelihood of fruit and/or vegetable intake when healthy food access is adequate. However, several studies investigating the effect of supermarkets on obesity did not reveal a significant association,^{51,59,73} implying that the association between supermarket access and obesity could be influenced by other contextual factors, such as shopping preferences, available modes of transportation and the presence of alternative food outlets.

Third, other factors of the built environment in shaping childhood obesity were rather inconclusive. Several studies recognized physical activity as an important factor in linking the obesogenic environment and childhood obesity, highlighting the health-promoting role of

recreational or physical activity facilities.^{84,88,130} Several other studies also examine the differences in transport-related environments (e.g., sidewalk, intersection and traffic) in explaining the disparity in children's physical activity and obesity.^{*} However, mixed results in terms of travel-related environmental factors were found in the literature.[†] A recent systematic review indicated that school transport interventions, such as the 'Safe Routes to School' programme in the United States, could be effective in increasing children's physical activity; however, overall quality of evidence was weak, largely due to inconsistencies across study design and short study periods.¹³² Isolating the influence of the travel-related environment on children's physical activity and obesity would be difficult because of possible interactions with other psychometric factors, such as safety perception.¹³¹ Also, some studies may be subject to residential self-selection bias¹³³ or selective daily mobility bias,¹³⁴ wherein preference or knowledge of healthy lifestyle could influence subjects' residential choice and travel patterns. The extent to which these biases also present in identifying modifiable risk factors in the built environment associated with childhood obesity remains relatively unknown, calling for further work.¹³⁵

Lastly, a large number of studies reported null associations between the obesogenic environment and childhood obesity, possibly due to the confounding effect on the individual level. Associations between environmental factors and childhood obesity could be modified by individual characteristics, such as gender, race, age, education attainment, family income and marital status. The same environment may have markedly different effects on different population groups. For example, the density of farmers' markets around the residential place was negatively associated with obesity among elementary students; the association, however, was not significant among middle/high school students.⁴⁸ For the two groups of students in the same study, the associations with the density of fast-food restaurants were the opposite. In another study, the environmental effects of supermarkets on obesity were different by gender group—girls were more likely to be affected by supermarket access than boys.³⁹ This gender difference, although being subtler in children than in adults, could be explained by the different levels of exposure and vulnerability to the obesogenic environment between genders. It originates from the physiological difference between genders in terms of body composition, hormone biology, patterns of weight gain, levels of resting energy expenditure and energy requirements, ability to engage in physical activity, levels of self-regulation in early childhood, and the susceptibility to social norms, cultures and ethnic backgrounds.¹³⁶ Likewise, socio-economic inequities in early childhood development allow children to have different opportunities of physical activity and diet quality, eventually leading to different levels of weight gain.^{137,138} Moreover, low-income families tend to be less vigilant about children's weight gain and therefore are less likely to seek appropriate interventions.^{139,140} As such, individual characteristics, notably gender

difference and socio-economic positioning, may strengthen or weaken environmental factors that contribute to childhood obesity.

This study has several limitations. First, the majority of the studies included in the review are cross-sectional. Although cross-sectional evidence is useful to test research hypotheses, further investigations using a longitudinal design will help to establish a more robust evidence base. Although prospective cohort studies are preferable, they are subject to high costs and the difficulty in capturing critical exposure over a prolonged time period or even the life course. One approach to overcome the limitation is to conduct retrospective studies linking existing administrative health records with historical geospatial data available on a global scale.¹⁴¹ Second, most of the studies in the review are focused on developed countries and do not reflect the reality of the growing obesity epidemic facing underdeveloped and developing countries.¹⁴² Especially in developing countries, rapid urbanization coupled with changing dietary patterns will likely exacerbate childhood obesity.¹⁴³ Failure to account for the obesogenic environment in underdeveloped and developing countries will lead to the omission of health risk factors posed for regions in need of obesity prevention and health intervention. Third, questionnaire-based survey methods as reviewed in this paper may have led to unreliable measurements, especially for the perceived measures. This is a common issue in survey research targeting children, as children's perception of the obesogenic environment tends to be inadvertently misrepresented in both the recruitment procedure and the survey question design.¹⁴⁴ It is thus recommended that future studies employ new technologies in a hybrid approach to offset the subjectivity in the research design.¹⁴⁵⁻¹⁴⁷ Also, active engagement of and the coproduction with children in the generation of knowledge can help minimize potential measurement biases.¹⁴⁸ Finally, the reporting quality of and comparability among future studies should be improved. The Spatial Lifecourse Epidemiology Reporting Standards (ISLE-ReSt) statement should be adopted by scientific journals in public health, geography and other relevant disciplines to increase reporting quality of such environmental health research.^{149,150}

5 | CONCLUSIONS

This systematic review reveals more significant associations of food rather than built environmental factors with weight status among children and adolescents. Heterogeneous measures in obesogenic environments for children and differences in controlling for confounding effects among studies may partly accounted for those null and inconclusive associations between some factors and weight status. This study comprehensively summarizes all existing evidence in this field and would serve as an important reference to multiple stakeholders, from new scholars in multiple relevant fields to policy makers.

ACKNOWLEDGEMENTS

We thank the International Institute of Spatial Lifecourse Epidemiology (ISLE) and the State Key Laboratory of Urban and Regional Ecology of China (SKLURE2018-2-5) for research support.

*References 48, 66, 68, 69, 104, 106, 110, 131-133.

†References 39, 45, 47, 48, 63, 66, 68, 69, 81, 91, 102, 107, 110, 113, 134.

CONFLICT OF INTEREST

No conflict of interest was declared.

ORCID

Chongjian Wang  <https://orcid.org/0000-0001-5091-6621>

Qian Xiao  <https://orcid.org/0000-0002-8388-1178>

Peng Jia  <https://orcid.org/0000-0003-0110-3637>

REFERENCES

- Zhang X, Zhang M, Zhao Z, et al. Geographic variation in prevalence of adult obesity in China: results from the 2013–2014 national chronic disease and risk factor surveillance. *Ann Intern Med.* 2020; 172(4):291–293.
- Jia P, Ma S, Qi X, Wang Y. Spatial and temporal changes in prevalence of obesity among Chinese children and adolescents, 1985–2005. *Prev Chronic Dis.* 2019;16.
- WHO. Fact sheets: obesity and overweight. <http://www.who.int/mediacentre/factsheets/fs311/en/>.
- CDC U. Childhood overweight and obesity. <http://www.cdc.gov/obesity/childhood/index.htm>.
- Cote AT, Harris KC, Panagiotopoulos C, Sandor GGS, Devlin AM. Childhood obesity and cardiovascular dysfunction. *J Am Coll Cardiol.* 62(15):1309–1319.
- Lloyd LJ, Langley-Evans SF, McMullen S, McMullen S. Childhood obesity and risk of the adult metabolic syndrome: a systematic review. *Int J Obes (Lond).* 36:1–11.
- Bacha F, Gidding SS. Cardiac abnormalities in youth with obesity and type 2 diabetes. *Curr Diab Rep.* 16(7):62.
- Mohan S, Tapp H, McWilliams A, Dulin M. Obesity and asthma: pathophysiology and implications for diagnosis and management in primary care. *Exp Biol Med (Maywood).* 239(11):1531–1540.
- Narang I, Mathew JL. Childhood obesity and obstructive sleep apnea. *J Nutr Metab.* 2012;2012:134202.
- Pollock NK. Childhood obesity, bone development, and cardiometabolic risk factors. *Mol Cell Endocrinol.* 2015;410:52–63.
- Africa JA, Newton KP, Schwimmer JB. Lifestyle interventions including nutrition, exercise, and supplements for nonalcoholic fatty liver disease in children. *Dig Dis Sci.* 2016;61(5):1375–1386.
- Morrison KM, Shin S, Tarnopolsky M, Taylor VH. Association of depression & health related quality of life with body composition in children and youth with obesity. *J Affect Disord.* 2015;172:18–23.
- Halfon N, Larson K, Slusser W. Associations between obesity and comorbid mental health, developmental, and physical health conditions in a nationally representative sample of US children aged 10 to 17. *Acad Pediatr.* 2013;13(1):6–13.
- Beck AR. Psychosocial aspects of obesity. *NASN Sch Nurse.* 2016; 31(1):23–27.
- Jensen MD, Ryan DH, Apovian CM, et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *J Am Coll Cardiol.* 2014;63(25):2985–3023.
- Jia P, Xue H, Cheng X, Wang Y, Wang Y. Association of neighborhood built environments with childhood obesity: Evidence from a 9-year longitudinal, nationally representative survey in the US. *Environ Int.* 2019;128: 158–164.
- Zhang X, Zhang M, Zhao Z, et al. Obesogenic environmental factors of adult obesity in China: a nationally representative cross-sectional study. *Environ Res Lett.* 2020;15(4):044009.
- Jia P, Xue H, Cheng X, Wang Y. Effects of school neighborhood food environments on childhood obesity at multiple scales: a longitudinal kindergarten cohort study in the USA. *Bmc Med.* 2019;17: 99.
- Wang Y, Jia P, Cheng X, Xue H. Improvement in food environments may help prevent childhood obesity: evidence from a 9-year cohort study. *Pediatr Obes.* 2019;14(10):e12536.
- Ford PB, Dziewaltowski DA. Disparities in obesity prevalence due to variation in the retail food environment: three testable hypotheses. *Nutr Rev.* 2008;66(4):216–228.
- Sallis JF, Conway TL, Cain KL, et al. Neighborhood built environment and socioeconomic status in relation to physical activity, sedentary behavior, and weight status of adolescents. *Prev Med.* 2018;110: 47–54.
- Chen X, Kwan MP. Contextual uncertainties, human mobility, and perceived food environment: the uncertain geographic context problem in food access research. *Am J Public Health.* 2015;105(9): 1734–1737.
- Cobb LK, Appel LJ, Franco M, Jones-Smith JC, Nur A, Anderson CAM. The relationship of the local food environment with obesity: a systematic review of methods, study quality, and results. *Obesity.* 2015;23(7):1331–1344.
- Pacheco AF, Balam GC, Archibald D, Grant E, Skafida V. Exploring the relationship between local food environments and obesity in UK, Ireland, Australia and New Zealand: a systematic review protocol. *BMJ Open.* 2018;8(2):e018701.
- Dunton GF, Kaplan J, Wolch J, Jerrett M, Reynolds KD. Physical environmental correlates of childhood obesity: a systematic review. *Obes Rev.* 2009;10(4):393–402.
- Carter MA, Dubois L. Neighbourhoods and child adiposity: a critical appraisal of the literature. *Health Place.* 2010;16(3):616–628.
- Jia P, Cheng X, Xue H, Wang Y. Applications of geographic information systems (GIS) data and methods in obesity-related research. *Obes Rev.* 2017;18(4):400–411.
- Lachowycz K, Jones AP. Greenspace and obesity: a systematic review of the evidence. *Obes Rev.* 2011;12(5):e183–e189.
- Cetateanu A, Jones A. Understanding the relationship between food environments, deprivation and childhood overweight and obesity: evidence from a cross sectional England-wide study. *Health Place.* 2014;27:68–76.
- Baek J, Sánchez BN, Berrocal VJ, Sanchez-Vaznaugh EV. Distributed lag models: examining associations between the built environment and health. *Epidemiology.* 2016;27(1):116–124.
- Barrera LH, Rothenberg SJ, Barquera S, Cifuentes E. The toxic food environment around elementary schools and childhood obesity in Mexican cities. *Am J Prev Med.* 2016;51(2):264–270.
- Bell JF, Wilson JS, Liu GC. Neighborhood greenness and 2-year changes in body mass index of children and youth. *Am J Prev Med.* 2008;35(6):547–553.
- Berge JM, Wall M, Larson N, Forsyth A, Bauer KW, Neumark-Sztainer D. Youth dietary intake and weight status: healthful neighborhood food environments enhance the protective role of supportive family home environments. *Health Place.* 2014;26:69–77.
- Carroll-Scott A, Gilstad-Hayden K, Rosenthal L, et al. Disentangling neighborhood contextual associations with child body mass index, diet, and physical activity: the role of built, socioeconomic, and social environments. *Soc Sci Med.* 2013;95:106–114.
- Carter MA, Dubois L, Tremblay MS, Taljaard M, Jones BL. Trajectories of childhood weight gain: the relative importance of local environment versus individual social and early life factors. *PLoS ONE.* 2012;7(10):e47065.
- Casey R, Chaix B, Weber C, et al. Spatial accessibility to physical activity facilities and to food outlets and overweight in French youth. *Int J Obes (Lond).* 2012;36(7):914–919.
- Chaparro MP, Whaley SE, Crespi CM, et al. Influences of the neighbourhood food environment on adiposity of low-income preschool-aged children in Los Angeles County: a longitudinal study. *J Epidemiol Commun H.* 2014;68(11):1027–1033.

38. Cheah WL, Chang CT, Saimon R. Environment factors associated with adolescents' body mass index, physical activity and physical fitness in Kuching south city, Sarawak: a cross-sectional study. *Int J Adolesc Med Health*. 2012;24(4):331-337.
39. Chen HJ, Wang Y. Changes in the neighborhood food store environment and children's body mass index at peripuberty in the United States. *J Adolesc Health*. 2016;58(1):111-118.
40. Chiang PH, Huang LY, Lee MS, Tsou HC, Wahlqvist ML. Fitness and food environments around junior high schools in Taiwan and their association with body composition: gender differences for recreational, reading, food and beverage exposures. *PLoS ONE*. 2017;12(8):e0182517.
41. Correa EN, Rossi CE, das Neves J, Silva DAS, de Vasconcelos FAG. Utilization and environmental availability of food outlets and overweight/obesity among schoolchildren in a city in the south of Brazil. *J Public Health-Uk*. 2018;40(1):106-113.
42. Crawford D, Cleland V, Timperio A, et al. The longitudinal influence of home and neighbourhood environments on children's body mass index and physical activity over 5 years: the CLAN study. *Int J Obes (Lond)*. 2010;34(7):1177-1187.
43. Datar A, Nicosia N, Wong E, Shier V. Neighborhood environment and children's physical activity and body mass index: evidence from military personnel installation assignments. *Child Obes*. 2015;11(2):130-138.
44. Davis B, Carpenter C. Proximity of fast-food restaurants to schools and adolescent obesity. *Am J Public Health*. 2009;99(3):505-510.
45. Duncan DT, Castro MC, Gortmaker SL, Aldstadt J, Melly SJ, Bennett GG. Racial differences in the built environment-body mass index relationship? A geospatial analysis of adolescents in urban neighborhoods. *Int J Health Geogr*. 2012;11(1):11.
46. Duncan MJ, Birch S, Woodfield L, Al-Nakeeb Y. Perceptions of the built environment in relation to physical activity and weight status in british adolescents from central England. *ISRN Obesity*. 2012;2012:903846.
47. Duncan DT, Sharifi M, Melly SJ, et al. Characteristics of walkable built environments and BMI z-scores in children: evidence from a large electronic health record database. *Environ Health Perspect*. 2015;122:1359-1365.
48. Dwicaksono A, Brissette I, Birkhead GS, Bozlak CT, Martin EG. Evaluating the contribution of the built environment on obesity among New York State students. *Health Educ Behav*. 2017;45(4):480-491.
49. Edwards KL, Clarke GP, Ransley JK, Cade J. The neighbourhood matters: studying exposures relevant to childhood obesity and the policy implications in Leeds, UK. *J Epidemiol Commun H*. 2010;64(3):194-201.
50. Epstein LH, Raja S, Daniel TO, et al. The built environment moderates effects of family-based childhood obesity treatment over 2 years. *Ann Behav Med*. 2012;44(2):248-258.
51. Fiechtner L, Kleinman K, Melly SJ, et al. Effects of proximity to supermarkets on a randomized trial studying interventions for obesity. *Am J Public Health*. 2016;106(3):557-562.
52. Friedman LS, Lukyanova EM, Serdiuk A, et al. Social-environmental factors associated with elevated body mass index in a Ukrainian cohort of children. *Int J Pediatr Obes*. 2009;4(2):81-90.
53. Ghenadenik AE, Kakinami L, Van Hulst A, Henderson M, Barnett TA. Neighbourhoods and obesity: a prospective study of characteristics of the built environment and their association with adiposity outcomes in children in Montreal, Canada. *Prev Med*. 2018;111:35-40.
54. Gilliland JA, Rangel CY, Healy MA, et al. Linking childhood obesity to the built environment: a multi-level analysis of home and school neighbourhood factors associated with body mass index. *Can J Public Health*. 2012;103(S3):S15-S21.
55. Gordon-Larsen P, Nelson MC, Page P, Popkin BM. Inequality in the built environment underlies key health disparities in physical activity and obesity. *Pediatrics*. 2006;117(2):417-424.
56. Gose M, Plachta-Danielzik S, Willii B, Johannsen M, Landsberg B, Müller MJ. Longitudinal influences of neighbourhood built and social environment on children's weight status. *Int J Env Res Pub He*. 2013;10(10):5083-5096.
57. Grafova IB. Overweight children: assessing the contribution of the built environment. *Prev Med*. 2008;47(3):304-308.
58. Green MA, Radley D, Lomax N, Morris MA, Griffiths C. Is adolescent body mass index and waist circumference associated with the food environments surrounding schools and homes? A longitudinal analysis. *BMC Public Health*. 2018;18(1):482.
59. Fiechtner L, Block J, Duncan DT, et al. Proximity to supermarkets associated with higher body mass index among overweight and obese preschool-age children. *Prev Med*. 2013;56(3-4):218-221.
60. Griffiths C, Frearson A, Taylor A, Radley D, Cooke C. A cross sectional study investigating the association between exposure to food outlets and childhood obesity in Leeds, UK. *Int J Behav Nutr Phys*. 2014;11(1):138.
61. Guedes DP, Rocha GD, Silva AJRM, Carvalhal IM, Coelho EM. Effects of social and environmental determinants on overweight and obesity among Brazilian schoolchildren from a developing region. *Rev Panam Salud Publica Pan Am J Public Health*. 2011;30:295-302.
62. Harris DE, Blum JW, Bampton M, et al. Location of food stores near schools does not predict the weight status of Maine high school students. *J Nutr Educ Behav*. 2011;43(4):274-278.
63. Harrison F, Jones AP, van Sluijs EMF, Cassidy A, Bentham G, Griffin SJ. Environmental correlates of adiposity in 9-10 year old children: considering home and school neighbourhoods and routes to school. *Soc Sci Med*. 2011;72(9):1411-1419.
64. Howard PH, Fitzpatrick M, Fulfroost B. Proximity of food retailers to schools and rates of overweight ninth grade students: an ecological study in California. *BMC Public Health*. 2011;11(1):68.
65. Hoyt LT, Kushi LH, Leung CW, et al. Neighborhood influences on girls' obesity risk across the transition to adolescence. *Pediatrics*. 2014;134(5):942-949.
66. Morgan Hughey S, Kaczynski AT, Child S, Moore JB, Porter D, Hibbert J. Green and lean: is neighborhood park and playground availability associated with youth obesity? Variations by gender, socioeconomic status, and race/ethnicity. *Prev Med*. 2017;95:S101-S108.
67. Jennings A, Welch A, Jones AP, et al. Local food outlets, weight status, and dietary intake: associations in children aged 9-10 years. *Am J Prev Med*. 2011;40(4):405-410.
68. Jerrett M, McConnell R, Chang CCR, et al. Automobile traffic around the home and attained body mass index: a longitudinal cohort study of children aged 10-18 years. *Prev Med*. 2010;50:S50-S58.
69. Jerrett M, McConnell R, Wolch J, et al. Traffic-related air pollution and obesity formation in children: a longitudinal, multilevel analysis. *Environ Health-Glob*. 2014;13:49.
70. Koleilat M, Whaley SE, Afifi AA, Estrada L, Harrison GG. Understanding the relationship between the retail food environment index and early childhood obesity among WIC participants in Los Angeles County using GeoDa. *Online J Public Health Inform*. 2012;4(1):e5.
71. Lakes T, Burkart K. Childhood overweight in Berlin: intra-urban differences and underlying influencing factors. *Int J Health Geogr*. 2016;15(1):12.
72. Lange D, Wahrendorf M, Siegrist J, Plachta-Danielzik S, Landsberg B, Müller MJ. Associations between neighbourhood characteristics, body mass index and health-related behaviours of adolescents in the Kiel Obesity Prevention Study: a multilevel analysis. *Eur J Clin Nutr*. 2011;65(6):711-719.
73. Larsen K, Cook B, Stone MR, Faulkner GEJ. Food access and children's BMI in Toronto, Ontario: assessing how the food environment relates to overweight and obesity. *Int J Public Health*. 2014;60:69-77.

74. Laska MN, Hearst MO, Forsyth A, Pasch KE, Lytle L. Neighbourhood food environments: are they associated with adolescent dietary intake, food purchases and weight status? *Public Health Nutr.* 2010;13(11):1757-1763.
75. Leatherdale ST, Poulou T, Church D, Hobin E. The association between overweight and opportunity structures in the built environment: a multi-level analysis among elementary school youth in the PLAY-ON study. *Int J Public Health.* 2011;56(3):237-246.
76. Leatherdale ST. A cross-sectional examination of school characteristics associated with overweight and obesity among Grade 1 to 4 students. *BMC Public Health.* 2013;13(1):982.
77. Leung CW, Laraia BA, Kelly M, et al. The influence of neighborhood food stores on change in young girls' body mass index. *Am J Prev Med.* 2011;41(1):43-51.
78. Li Y, Robinson LE, Carter WM, Gupta R. Childhood obesity and community food environments in Alabama's Black Belt region. *Child Care Health Dev.* 2015;41:668-676.
79. Lovasi GS, Schwartz-Soicher O, Quinn JW, et al. Neighborhood safety and green space as predictors of obesity children from low-income families in New York City among preschool. *Prev Med.* 2013;57(3):189-193.
80. Miller DP. Associations between the home and school environments and child body mass index. *Soc Sci Med.* 2011;72(5):677-684.
81. Miller LJ, Joyce S, Carter S, Yun G. Associations between childhood obesity and the availability of food outlets in the local environment: a retrospective cross-sectional study. *Am J Health Promot.* 2014;28(6):e137-e145.
82. Minaker LM, Storey KE, Raine KD, et al. Associations between the perceived presence of vending machines and food and beverage logos in schools and adolescents-diet and weight status. *Public Health Nutr.* 2011;14(8):1350-1356.
83. Molina-García J, Queralt A, Adams MA, Conway TL, Sallis JF. Neighborhood built environment and socio-economic status in relation to multiple health outcomes in adolescents. *Prev Med.* 2017;105:88-94.
84. Nelson NM, Woods CB. Obesogenic environments: are neighbourhood environments that limit physical activity obesogenic? *Health Place.* 2009;15(4):917-924.
85. Nesbit KC, Kolobe TA, Arnold SH, Sisson SB, Anderson MP. Proximal and distal environmental correlates of adolescent obesity. *J Phys Act Health.* 2014;11(6):1179-1186.
86. Ness M, Barradas DT, Irving J, Manning SE. Correlates of overweight and obesity among American Indian/Alaska Native and non-Hispanic White children and adolescents: National Survey of Children's Health, 2007. *Matern Child Health J.* 2012;16(Suppl 2):268-277.
87. Nogueira H, Ferrão M, Gama A, Mourão I, Rosado Marques V, Padez C. Perceptions of neighborhood environments and childhood obesity: evidence of harmful gender inequities among Portuguese children. *Health Place.* 2013;19:69-73.
88. Norman GJ, Nutter SK, Ryan S, Sallis JF, Calfas KJ, Patrick K. Community design and access to recreational facilities as correlates of adolescent physical activity and body-mass index. *J Phys Act Health.* 2006;3(s1):S118-S128.
89. Ohri-Vachaspati P, Lloyd K, DeLia D, Tulloch D, Yedidia MJ. A closer examination of the relationship between children's weight status and the food and physical activity environment. *Prev Med.* 2013;57(3):162-167.
90. Oreskovic NM, Kuhlthau KA, Romm D, Perrin JM. Built environment and weight disparities among children in high- and low-income towns. *Acad Pediatr.* 2009;9(5):315-321.
91. Oreskovic NM, Winickoff JP, Kuhlthau KA, Romm D, Perrin JM. Obesity and the built environment among Massachusetts children. *Clin Pediatr.* 2009;48(9):904-912.
92. Park S, Choi BY, Wang Y, Colantuoni E, Gittelsohn J. School and neighborhood nutrition environment and their association with students' nutrition behaviors and weight status in Seoul, South Korea. *J Adolesc Health.* 2013;53(5):655-662.
93. Pearce M, Bray I, Horswell M. Weight gain in mid-childhood and its relationship with the fast food environment. *J Public Health (Oxf).* 2017;40(2):237-244.
94. Petraviciene I, Grazuleviciene R, Andrusaityte S, Dedele A, Nieuwenhuijsen MJ. Impact of the social and natural environment on preschool-age children weight. *Int J Env Res Pub He.* 2018;15(3):449.
95. Pitts SB, Carr LJ, Brinkley J, Byrd JL 3rd, Crawford T, Moore JB. Associations between neighborhood amenity density and health indicators among rural and urban youth. *Am J Health Promot: AJHP.* 2013;28(1):e40-e43.
96. Poole R, Moon G. What is the association between healthy weight in 4-5-year-old children and spatial access to purposefully constructed play areas? *Health Place.* 2017;46:101-106.
97. Potestio ML, Patel AB, Powell CD, McNeil DA, Jacobson RD, McLaren L. Is there an association between spatial access to parks/green space and childhood overweight/obesity in Calgary, Canada? *Int J Behav Nutr Phy.* 2009;6(1):77.
98. Rossen LM, Curriero FC, Cooley-Strickland M, Pollack KM. Food availability en route to school and anthropometric change in urban children. *Urban Health.* 2013;90(4):653-666.
99. Gorski Findling MT, Wolfson JA, Rimm EB, Bleich SN. Differences in the neighborhood retail food environment and obesity among US children and adolescents by SNAP participation. *Obesity (Silver Spring, Md).* 2018;26:1063-1071.
100. Sánchez BN, Sanchez-Vaznaugh EV, Uscilka A, Baek J, Zhang L. Differential associations between the food environment near schools and childhood overweight across race/ethnicity, gender, and grade. *Am J Epidemiol.* 2012;175(12):1284-1293.
101. Schmidt SC, Sleddens EFC, de Vries SI, Gubbels J, Thijs C. Longitudinal association of neighborhood variables with body mass index in Dutch school-age children: the KOALA birth cohort study. *Soc Sci Med.* 2015;135:99-108.
102. Schüle SA, Fromme H, Bolte G. Built and socioeconomic neighbourhood environments and overweight in preschool aged children. A multilevel study to disentangle individual and contextual relationships. *Environ Res.* 2016;150:328-336.
103. Seliske LM, Pickett W, Boyce WF, Janssen I. Association between the food retail environment surrounding schools and overweight in Canadian youth. *Public Health Nutr.* 2009;12(9):1384-1391.
104. Seliske L, Pickett W, Janssen I. Urban sprawl and its relationship with active transportation, physical activity and obesity in Canadian youth. *Health Rep.* 2012;23:1-10.
105. Singh GK, Siahpush M, Kogan MD. Neighborhood socioeconomic conditions, built environments, and childhood obesity. *Health Aff.* 2010;29(3):503-512.
106. Slater SJ, Nicholson L, Chriqui J, Barker DC, Chaloupka FJ, Johnston LD. Walkable communities and adolescent weight. *Am J Prev Med.* 2013;44(2):164-168.
107. Spence JC, Cutumisu N, Edwards J, Evans J. Influence of neighbourhood design and access to facilities on overweight among preschool children. *Int J Pediatr Obes.* 2008;3(2):109-116.
108. Tang X, Ohri-Vachaspati P, Abbott JK, et al. Associations between food environment around schools and professionally measured weight status for middle and high school students. *Child Obes.* 2014;10(6):511-517.
109. Taylor WC, Upchurch SL, Brosnan CA, et al. Features of the built environment related to physical activity friendliness and children's obesity and other risk factors. *Public Health Nurs.* 2014;31(6):545-555.
110. Timperio A, Jeffery RW, Crawford D, Roberts R, Giles-Corti B, Ball K. Neighbourhood physical activity environments and adiposity

- in children and mothers: a three-year longitudinal study. *Int J Behav Nutr Phys*. 2010;7(1):18.
111. Torres R, Serrano M, Perez CM, Palacios C. Physical environment, diet quality, and body weight in a group of 12-year-old children from four public schools in Puerto Rico. *P R Health Sci J*. 2014;33(1):14-21.
 112. Veugelers P, Sithole F, Zhang S, Muhajarine N. Neighborhood characteristics in relation to diet, physical activity and overweight of Canadian children. *Int J Pediatr Obes*. 2008;3(3):152-159.
 113. Wall MM, Larson NI, Forsyth A, et al. Patterns of obesogenic neighborhood features and adolescent weight: a comparison of statistical approaches. *Am J Prev Med*. 2012;42(5):e65-e75.
 114. Wasserman JA, Suminski R, Xi J, Mayfield C, Glaros A, Magie R. A multi-level analysis showing associations between school neighborhood and child body mass index. *Int J Obes (Lond)*. 2014;38(7):912-918.
 115. Williams J, Scarborough P, Townsend N, et al. Associations between food outlets around schools and BMI among primary students in England: a cross-classified multi-level analysis. *PLoS ONE*. 2015;10(7):e0132930.
 116. Wolch J, Jerrett M, Reynolds K, et al. Childhood obesity and proximity to urban parks and recreational resources: a longitudinal cohort study. *Health Place*. 2011;17(1):207-214.
 117. Xu F, Li J, Liang Y, et al. Residential density and adolescent overweight in a rapidly urbanising region of mainland China. *J Epidemiol Commun H*. 2010;64:1017-1021.
 118. Yang Y, Jiang Y, Xu Y, Mzayek F, Levy M. A cross-sectional study of the influence of neighborhood environment on childhood overweight and obesity: variation by age, gender, and environment characteristics. *Prev Med*. 2018;108:23-28.
 119. Zhang J, Xue H, Cheng X, et al. Influence of proximities to food establishments on body mass index among children in China. *Asia Pac J Clin Nutr*. 2016;25(1):134-141.
 120. Li B, Adab P, Cheng KK. Family and neighborhood correlates of overweight and obesogenic behaviors among Chinese children. *Int J Behav Med*. 2014;21(4):700-709.
 121. Kepper M, Tseng TS, Volaufova J, Scribner R, Nuss H, Sothorn M. Pre-school obesity is inversely associated with vegetable intake, grocery stores and outdoor play. *Pediatr Obes*. 2016;11(5):e6-e8.
 122. Crawford D, Ball K, Cleland V, et al. Maternal efficacy and sedentary behavior rules predict child obesity resilience. *BMC Obesity*. 2015;2(1):26.
 123. Powell LM, Auld MC, Chaloupka FJ, O'Malley PM, Johnston LD. Associations between access to food stores and adolescent body mass index. *Am J Prev Med*. 2007;33(4):S301-S307.
 124. Burdette HL, Whitaker RC. Neighborhood playgrounds, fast food restaurants, and crime: relationships to overweight in low-income preschool children. *Prev Med*. 2004;38(1):57-63.
 125. Sturm R, Datar A. Body mass index in elementary school children, metropolitan area food prices and food outlet density. *Public Health*. 2005;119(12):1059-1068.
 126. Potwarka LR, Kaczynski AT, Flack AL. Places to play: association of park space and facilities with healthy weight status among children. *J Community Health*. 2008;33(5):344-350.
 127. Galvez MP, Hong L, Choi E, Liao L, Godbold J, Brenner B. Childhood obesity and neighborhood food-store availability in an inner-city community. *Acad Pediatr*. 2009;9(5):339-343.
 128. Jia P, Lakerveld J, Wu J, et al. Top 10 research priorities in spatial lifecourse epidemiology. *Environ Health Perspect*. 2019;127(7):074501.
 129. Pereira MA, Kartashov AI, Ebbeling CB, et al. Fast-food habits, weight gain, and insulin resistance (the CARDIA study): 15-year prospective analysis. *Lancet*. 2005;365(9453):36-42.
 130. Ding D, Sallis JF, Kerr J, Lee S, Rosenberg DE. Neighborhood environment and physical activity among youth: a review. *Am J Prev Med*. 2011;41(4):442-455.
 131. McMillan TE. The relative influence of urban form on a child's travel mode to school. *Transp Res Part a Policy Pract*. 2007;41(1):69-79.
 132. Larouche R, Mammen G, Rowe DA, Faulkner G. Effectiveness of active school transport interventions: a systematic review and update. *BMC Public Health*. 2018;18(1):206.
 133. Boone-Heinonen J, Gordon-Larsen P, Guilkey DK, Jacobs DR, Popkin BM. Environment and physical activity dynamics: the role of residential self-selection. *Psychol Sport Exerc*. 2011;12(1):54-60.
 134. Chaix B, Méline J, Duncan S, et al. GPS tracking in neighborhood and health studies: a step forward for environmental exposure assessment, a step backward for causal inference? *Health Place*. 2013;21:46-51.
 135. Burgoine T, Jones AP, Namenek Brouwer RJ, Benjamin Neelon SE. Associations between BMI and home, school and route environmental exposures estimated using GPS and GIS: do we see evidence of selective daily mobility bias in children? *Int J Health Geogr*. 2015;14(1):8.
 136. Sweeting HN. Gendered dimensions of obesity in childhood and adolescence. *Nutr J*. 2008;7(1):1.
 137. Gebremariam MK, Altenburg TM, Lakerveld J, et al. Associations between socioeconomic position and correlates of sedentary behaviour among youth: a systematic review. *Obes Rev*. 2015;16(11):988-1000.
 138. Watts AW, Mason SM, Loth K, Larson N, Neumark-Sztainer D. Socioeconomic differences in overweight and weight-related behaviors across adolescence and young adulthood: 10-year longitudinal findings from Project EAT. *Prev Med*. 2016;87:194-199.
 139. Hansen AR, Duncan DT, Tarasenko YN, Yan F, Zhang J. Generational shift in parental perceptions of overweight among school-aged children. *Pediatrics*. 2014;134(3):481-488.
 140. Beauchamp A, Backholer K, Magliano D, Peeters A. The effect of obesity prevention interventions according to socioeconomic position: a systematic review. *Obes Rev*. 2014;15(7):541-554.
 141. Jia P, Stein A, James P, et al. Earth observation: investigating non-communicable diseases from space. *Annu Rev Public Health*. 2019;40(1):85-104.
 142. Popkin BM, Slining MM. New dynamics in global obesity facing low- and middle-income countries. *Obes Rev*. 2013;14:11-20.
 143. Poskitt EME. Childhood obesity in low- and middle-income countries. *Paediatr Int Child H*. 2014;34(4):239-249.
 144. Einarsdóttir J. Research with children: methodological and ethical challenges. *Eur Early Child Educ Res J*. 2007;15(2):197-211.
 145. Dunton GF, Intille SS, Wolch J, Pentz MA. Children's perceptions of physical activity environments captured through ecological momentary assessment: a validation study. *Prev Med*. 2012;55(2):119-121.
 146. Jia P. Integrating kindergartener-specific questionnaires with citizen science to improve child health. *Frontiers in public health*. 2018;6:236.
 147. Jia P, Xue H, Yin L, Stein A, Wang M, Wang Y. Spatial technologies in obesity research: current applications and future promise. *Trends Endocrinol Metab*. 2019;30(3):211-223.
 148. Mirra N, Garcia A, Morrell E. *Doing Youth Participatory Action Research: Transforming Inquiry with Researchers, Educators, and Students*. UK: Routledge; 2015.
 149. Jia P, Yu C, Remais JV, et al. Spatial lifecourse epidemiology reporting standards (ISLE-ReSt) statement. *Health Place*. 2020;61:102243.
 150. Jia P. Spatial lifecourse epidemiology. *Lancet Planet Health*. 2019;3(2):e57-e59.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Mei K, Huang H, Xia F, et al. State-of-the-art of measures of the obesogenic environment for children. *Obesity Reviews*. 2021;22(S1):e13093. <https://doi.org/10.1111/obr.13093>

APPENDIX A.: SEARCH STRATEGY

The search strategy includes all possible combinations of keywords in the title/abstract from the following three groups:

1. 'built environment*', 'food environment*', 'obesogenic environment*', 'built environment factor*', 'food environment factor*', 'obesogenic environment factor*', 'built environmental factor*', 'food environmental factor*', 'obesogenic environmental factor*', 'built environment variable*', 'food environment variable*', 'obesogenic environment variable*', 'built environmental variable*', 'food environmental variable*', 'obesogenic environmental indicator*', 'built environment indicator*', 'food environment indicator*';
2. 'child*', 'juvenile*', 'pubescent*', 'pubert*', 'adolescen*', 'youth*', 'teen*', 'kid*', 'young*', 'youngster*', 'minor*', 'student*', 'pupil*', 'pediatric*', 'preschooler*', 'pre-schooler*', 'schoolchild*', 'school-child*', 'school child*', 'schoolage*', 'school-age*', 'school age*';
3. 'energy balance', 'calorie*', 'body mass index', 'BMI', 'weight', 'weight status', 'weight-related health', 'overweight', 'obese', 'obesity', 'adiposity', 'abdominal overweight', 'abdominal obesity', 'central overweight', 'central obesity', 'central adiposity', 'waist circumference', 'waist to hip', 'waist-to-hip', 'waist to height', 'waist-to-height', 'waist to stature', 'waist-to-stature', 'fatness', 'body fat', 'excess fat', 'excess weight', 'overnutrition', 'over-nutrition', 'over nutrition'.