

Original Research

Cite this article: Rekhter N and Ermasova N. Effect of the COVID-19 on perceptions of health, anticipated need for health services, and cost of health care. *Disaster Med Public Health Prep.* doi: <https://doi.org/10.1017/dmp.2021.174>.


Keywords:

perceived stress; COVID-19 pandemic; health concerns; healthcare services; healthcare access; healthcare cost

Corresponding author:

Natalia Rekhter,
Email: nrekhter@yahoo.com.

Effect of the COVID-19 on Perceptions of Health, Anticipated Need for Health Services, and Cost of Health Care

Natalia Rekhter PhD¹  and Natalia Ermasova PhD²

¹Health Administration Department, College of Health and Human Services, Governors State University, University Park, Illinois USA and ²Public Policy and Administration Unit, College of Arts and Sciences, Governors State University, University Park, Illinois USA

Abstract

Objective: This article investigates how perceived vulnerability to the coronavirus disease 2019 (COVID-19) pandemic at its early stages is associated with people's perception of their health, the need for health-care services, and expenses related to addressing the COVID-19 impact on their health.

Methods: The results are based on the analysis of surveys that were distributed among members of 26 random Facebook groups in April-May, 2020. Perceived COVID-19 pandemic related stress and health concerns were examined by using the analysis of variance (ANOVA) test.

Results: Among 315 respondents, 64% have experienced COVID-19 related stress and identified anxiety, headache, insomnia, and weight gain as their primary health concerns. The ANOVA test revealed that females are more impacted by the COVID-19 stress than males. Around 40% of respondents believed that the COVID-19 would lead to an increase in the cost of health services, and 20% of respondents anticipated that the COVID-19 pandemic would increase their need for health services.

Conclusions: Learning about how people perceive the COVID-19 pandemic impact on their health, particularly in the pandemic's early stages can allow health professionals to develop targeted interventions that can influence pandemic preventative behaviors among different population groups. This study can help understand use patterns and mitigate financial barriers that could interfere with patients' care-seeking behavior.

On March 11, 2020, the World Health Organization (WHO) declared a pandemic of the highly transmissible severe acute respiratory coronavirus 2 (SARS-CoV-2) that triggered an alarming global health crisis¹⁻⁴ In many countries, governments have set severe restrictions on daily life, mandated social distancing and health protection policies, and locked down nonessential businesses.⁵⁻⁸ Wiedemann and Dorl (2020) analyzed what and how the scientific community communicates about coronavirus disease 2019 (COVID-19) risks to the general public, media, and politics.

The regulations and managing efforts differed from country to country—these included lockdown, isolation, quarantine, travel ban, and/or closing of the nonessential business. The scale, pace, and depth of government actions varied at the federal, regional, and local levels. Despite their efforts, every country in the world has been experiencing the harsh impact of the pandemic; some countries have been hit harder than others. The perception of stress may influence the pathogenesis physical diseases by causing anxiety, headache, insomnia, and depression, which then exert direct effects on physiological processes, behavioral patterns, and overall health.⁹⁻¹¹ The perception of an individual's health and accompanying health-related expenses can affect care-seeking behavior. Therefore, learning about perceptions associated with COVID-19 can provide ground for early detection and treatment of various health conditions and for the development of targeted interventions to improve the preventative behavioral patterns and health outcomes of different population groups. Understanding the anticipated need for health services and the expected cost would help to assess the prospective use patterns and mitigate financial barriers that could interfere with care-seeking behavior.

High morbidity and mortality rates related to COVID-19, which WHO estimates to be close to 4%,² its highly contagious nature, and its broad and quick spread caused confusion, fear, and anxiety among the general public, including the United States. These perceptions are associated with COVID-19, and any perceptions related to any risk can influence people's behavior.¹² Betsch et al.¹³ wrote about the importance of behavioral insights for the COVID-19 pandemic. It includes knowledge of how people perceive the stress associated with the risk of COVID-19 and how these perceptions impact their health, protective behavior, and health-related expenses. Risk perceptions are among the core features of protection-motivation theory¹⁴ and, as such, can help to determine the public's demands for the healthcare services

during and after the pandemics, as well as health-protective behavior. However, the research that provides and analyses evidence about public perceptions of stress and health as it related to pandemics is somewhat limited. The majority of this research evidence has come from studies during the H1N1 swine flu pandemic that took place in 2009^{15,16} and the Ebola outbreak.¹⁷ According to Wise et al.,¹⁸ while some of these studies emphasized the role of risk perception, they often took place in anticipation of an outbreak or long after its emergence. A few studies that have surveyed individuals during the early stages of a pandemic have suggested that perceived personal risk of infection and how this risk affects people's health can lead to engagement in protective behavior during and after the pandemic¹⁸ as well to changes in care-seeking behavior among patients.

This observation above determined the 4 research questions of this study: (1) What was the perceived stress during the early stages of the COVID-19 pandemic?; (2) What was the perception of the impact of stress on the health status of respondents during the COVID-19 pandemic?; (3) What was the respondents' perception on the changes in their health status?; and (4) What was the respondents' perception on the cost of their healthcare services as it relates to COVID-19?

This study provides insights for medical professionals, healthcare managers, and other stakeholders into the healthcare needs of different population groups, their perceived financial impact on healthcare costs associated with COVID-19, particularly during this pandemic's first months (April-May, 2020). It helps evaluate the perceived stress and healthcare issues each group may exhibit, the perceived access to healthcare services, as well as the kind of attention each of these groups may require.

The study proceeds as follows: the first section provides methods of analysis. The next section focuses on the data and results, followed by a discussion of key research findings. The final section concludes the study and provides suggestions for future research.

Methods

Data and Sample Characteristics

Our survey was created based on the modification of the survey named Stress Management for Law Enforcement Officers Questionnaire (SMLEOQ).¹⁹ The study was reviewed and approved as an exempt study by the Institutional Review Board (IRB Approval by the Governors State University: IRB-FY2019-40). The survey was distributed through social network sites (SNS) among the general population in different social network groups. Informed consent, the procedure of maintaining confidentiality, detailed instructions on how to complete the questionnaire, the title of the study, a brief description including study purpose, inclusion criteria, and the survey link were included. To achieve broader representation, address biases, and to follow the paper topic of the general population SNS engagement, researchers worked with 41 Facebook groups with diverse and not specifically affiliated membership, such as, for instance, "Costcomans", "IsoIsolacia", Social Media sites of different cities, and similar. Facebook groups were open to people across the United States. Each groups' language was English. Researchers approached group administrators with the request to post the survey, and 26 groups' administrators approved this request. All participants read an online consent form and consented by checking "I read consent and agree to participate." The data were collected in the time-frame from April to May, 2020. These posts

Table 1. Demographic variables (N = 315)

Variables	Frequency	Percent	
Age (y)	18-29	71	22.53%
	30-39	78	24.76%
	40-49	72	22.85%
	50-55	40	12.70%
	56-60	29	9.20%
	61 and older	25	7.91%
	Total	315	100.0
Gender	Female	85	27%
	Male	230	73%
	Total	315	100.00%
Marital status	Single, never married	90	28.50%
	Married or domestic partnership	179	56.80%
	Widowed	5	1.58%
	Separated	9	2.85%
	Divorced	32	10.17%
	Total	315	100.00%
Ethnicity	White	238	75.55%
	Black or African American	50	15.87%
	Asian	14	4.44%
	Hispanic or Latino	5	1.58%
	Others	8	2.53%
	Total	315	100.00%
Education	High school diploma	31	10.10%
	Associate degree	23	7.49%
	Bachelor's degree	101	32.00%
	Master's degree	84	26.66%
	Doctoral degree	62	20.63%
	M.D. or D.O.	6	1.9%
	Other professional certificates	8	1.26%
	Total	315	100.00%

generated 336 responses, of which there were 315 completed questionnaires that qualified for the analysis. For questionnaires completed through the Web link, the data were automatically saved and converted into a database in an excel format file for analysis. The information about sample characteristics in terms of gender, age, ethnicity, educational level, and marital status are illustrated in Table 1.

As a part of the survey, respondents provided personal information regarding their age, gender, marital status, working experience, ethnicity, management experience, and current educational level. Among the 315 respondents who completed the survey, 85 (27%) were males and 230 (73%) were females (Table 1). Education was coded in 7 categories. One-third of participants had Bachelors' degree. One-fourth of respondents had Masters' degree. There were 31 respondents with a high school diploma (10.1%), 23 respondents with an Associate degree (7.49%), 6 respondents with M.D. or D.O. degrees (1.9%), and 4 respondents with certificates (1.26%).

Over half of the respondents were married. Individuals who were single/never married made up 1/3 of the sample, followed by divorced, separated, and widowed. Most participants in this study were 30-39 y old (1/4 of respondents), which was followed by the age group 40-49 and 18-29 age group, age group 50-55, and age group 56-60. The age group of 61+ had 25 participants.

Table 2. Perceived stress level during COVID-19 pandemic

Level of stress	Frequency	Percent
Caused no stress	21	6.67%
Little stress	92	29.21%
Mid-level stress	121	38.41%
High stress	81	25.71%
	315	100.00%

Measures and Analysis Strategy

Respondents were asked to indicate the perceived stress during the COVID-19 crisis as well as to identify the perceived impact of stress on their health during the COVID-19 pandemic, and the following need for healthcare services and the cost of healthcare services.

In this survey, the 4-item Perceived Stress Scale was used to assess the degree to which situations in life are perceived as stressful.²⁰ The study was constructed based on the framework created by McCreary and Thompson.²¹ The study by McCreary and Thompson²¹ was tested in studies by Ceka and Ermasova²⁰ and Ermasova et al.¹⁹ For the purpose of this study, we used the 4-item Perceived Stress Scale, that was created by McCreary and Thompson,²¹ to measure psychological stress perceptions of respondents during the COVID-19 pandemic. Analysis of stressors and stress relieving tactics during the COVID-19 outbreak questionnaire had the following Questions: “Has the COVID-19 made an impact on your stress level?” Please rate them on a scale of 1 to 4, where 1 - “caused no stress,” 2 - “a little stress,” 3 - “mid-level stress,” 4 - “high stress.” Higher scores are indicating a higher level of perceived stress. A total of 8 items (eg, “Has COVID-19 increased your level of stress, as it relates to social media?”; ... “Has COVID 19 increased your level of stress, as it relates to nonsocial media news channels?”) were measured on a 4-point scale, ranging from 1 (“caused no stress”) to 4 (“high stress”).

To test our hypotheses, first, we analyzed whether gender, ethnicity, and age had an effect on the perceived stressors, using an analysis of variance (ANOVA) test to compare groups. Second, we provided descriptive statistics on the perceived impact of stress on health, perceptions of the need for medical services, and perception of changes in the cost of health-care services, as the result of the COVID-19 pandemic.

Results

Perceptions of Overall Stress During the COVID-19 Crisis

The higher mean demonstrated that the respondents viewed COVID-19 as impacting their stress level by increasing it. More than half of the participants (64.12%) perceived stress associated with COVID-19 as “moderate” (38.41%) or “high” (25.71%) (Table 2).

Gender and Stress During COVID-19

The ANOVA test revealed that the average mean values of the male respondents in ranking the impact of stressors during COVID-19 are lower than the mean values of females, suggesting that in average female respondents (mean = 2.95) are impacted by the stressors at a higher degree than male respondents (mean = 2.44) (Table 3).

Table 3. ANOVA comparison for perceived stress by gender

Variables	Mean total	SD	Mean female	Mean male	F	Sig.
Gender	2.81	0.88	2.95	2.44	22.06	0.00****

Notes: Bartlett's test for equal variances: $\chi^2(1) = 0.85$; $\text{Prob} > \chi^2 = 0.356$. Significant at **** $P < 0.001$; *** $P < 0.01$; ** $P < 0.05$; * $P < 0.10$.

Age and Stress During COVID-19

The ANOVA test revealed no significant difference between the average mean values of the different age groups of respondents in ranking the impact of COVID-19. However, the mean of perceived stress increased with age (from mean = 1.66 for adults 18-29 y to mean = 1.8 for adults 61 and older) (Table 4).

Ethnicity and Stress During COVID-19

The ANOVA test revealed that there is no significant difference between the average mean values of the different ethnic groups of respondents in ranking the impact of stressors during COVID-19. However, the COVID-related stress perceptions were higher among Hispanic respondents than among Asian respondents (Table 5).

Perceived Impact of COVID-19 Related Stress on Health Status

The survey included a question “If COVID-19 increased your level of stress, did it also impact your health status?” Around 40% of respondents—121 respondents (38.4%)—answered that stress during COVID-19 affected their health status. The analysis of the answers to this question is presented in Table 6.

The survey had the follow-up question: “If COVID-19 impacted your health, did it manifest itself in . . .” and included a list of potential health manifestations and an option for comments. Respondents could select more than 1 answer and provide comments. Anxiety, insomnia, headaches, and weight gain were among the most often selected health concerns. The results of the survey's answers are provided in Table 7.

Among respondents, 17 people also offered comments to this question. The majority of comments discussed various health issues and what caused health issues as they relate to COVID-19. The comments are grouped by issues and self-reported causes of these issues, as they relate to COVID-19 (Table 8).

Gender and Perceived Impact of Stress on Health Status During COVID-19

The ANOVA test revealed that the average mean values of the male respondents in ranking the perceived impact of stress on health status during COVID-19 are higher than the mean values of females, suggesting that in average female respondents (mean = 1.66) had lower level of perceived impact of stress on health status during COVID-19 than male respondents (mean = 1.8) (Table 9).

Age and Perceived Impact of Stress on Health Status During COVID-19

The ANOVA test revealed no significant difference between the average mean values of the different age groups of respondents in ranking the perceived impact of stress on health status during COVID-19. However, the mean of perceived stress increased with age (from mean = 1.66 for adults 18-29 y to mean = 1.8 for adults 61 and older) (Table 10).

Table 4. ANOVA comparison for perceived overall stress by age

Variables	Mean total	SD	Mean 18-29	Mean 30-39	Mean 40-49	Mean 50-55	Mean 56-61	Mean 61 and older	F	Sig.
Age	1.7	0.6	1.66	1.65	1.73	1.74	1.75	1.8	0.39	0.85

Notes: Bartlett’s test for equal variances: $\chi^2(5) = 1.5080$; $\text{Prob} > \chi^2 = 0.912$. Significant at **** $P < 0.001$; *** $P < 0.01$; ** $P < 0.05$; * $P < 0.10$.

Table 5. ANOVA comparison for perceived stress by ethnicity

Variables	Mean total	SD	Mean White	Mean Black	Mean Asian	Mean Hispanic	Mean other	F	Sig.
Age	1.7	0.6	2.83	2.74	2.57	3.2	3.25	0.86	0.48

Notes: Bartlett’s test for equal variances: $\chi^2(4) = 1.2755$; $\text{Prob} > \chi^2 = 0.866$. Significant at **** $P < 0.001$; *** $P < 0.01$; ** $P < 0.05$; * $P < 0.10$.

Table 6. Perceived impact of stress on health status during COVID-19 pandemic

Impact of stress	Frequency	Percent
Yes	121	38.41%
No	171	54.29%
Do not know	23	7.30%
	315	100.00%

Table 7. Reported health issues (more than 1 option)

Health issues	Frequency	Percent
Anxiety	94	23.32%
Insomnia	65	16.04%
Headache	60	14.88%
Weight gain	54	13.33%
Back pain	40	9.92%
Higher blood pressure	27	6.75%
None of the above	57	14.07%
Other	6	1.4%
	403	100%

Table 8. Additional health issues of respondents that related to COVID-19

Health issues	Self-reported causes	Frequency
Anxiety, depression, TMJ, headache, escalation in symptoms related to pre-existing conditions, insomnia, fatigue, weight gain	Restricted access to healthcare services due to COVID-19.	7
	Transition to online working and learning environment	4
	Self-isolation and absence of physical interactions	1
	The need to care for immunocompromised family members, which can lead to one’s health decline	1
	Sadness and grief related to death and illness in the family	1
Weight loss	Self-isolation, hypomobility	2
Physical injuries,	Domestic violence	1

Table 9. ANOVA comparison for the perceived impact of stress on health status by gender

Variables	Mean total	SD	Mean female	Mean male	F	Sig.
Gender	1.7	0.6	1.66	1.8	3.13	0.078*

Notes: Bartlett’s test for equal variances: $\chi^2(1) = 5.45$; $\text{Prob} > \chi^2 = 0.020$. Significant at **** $P < 0.001$; *** $P < 0.01$; ** $P < 0.05$; * $P < 0.10$.

Ethnicity and Perceived Impact of Stress on Health Status During COVID-19

The ANOVA test revealed that there is no significant difference between average mean values of the different ethnic groups of respondents in ranking perceived impact of stress on health status during COVID-19 (Table 11).

Perceived Needs for Healthcare Services and Changes in Healthcare Cost as a Result of COVID-19

The respondents were asked “Do you think COVID-19 will increase your need for medical services?” Sixty-six respondents (20.9%) answered that stress during COVID-19 will impact their need for medical services. (Table 12).

The respondents were also asked “Do you think COVID-19 will increase the cost of your health-care services?” Over 40% of respondents—129 respondents (40.95%)—believed that the COVID-19 would lead to an increase in the cost of health-care services (Table 12).

Limitations

There are several limitations that may affect the generalizability of the findings in this study. One of the limitations is the limited population of respondents. Thus, the generalizations of this study’s results are limited because it is not necessarily representative of the US population. Future studies should expand the sample to include more respondents from other cities and regions in the United States. The second limitation is that more women participated in the survey than males. It can be attributed to the fact that 54.3% of the Facebook audience in the United States are females, and 45.7% are males. Females also tend to be more active and frequent in their use of Facebook: 69% of female users engage with Facebook at least once a day, compared with 54% of male users.²² Participants’ education level could also be attributed to the fact that 74% of adults with a college degree or more use Facebook, compared with 61% of those who have a high school diploma

Table 10. ANOVA comparison for the perceived impact of stress on health status by age

Variables	Mean total	SD	Mean 18-29	Mean 30-39	Mean 40-49	Mean 50-55	Mean 56-61	Mean 61 and older	F	Sig.
Age	1.7	0.6	1.66	1.65	1.73	1.74	1.75	1.8	0.39	0.86

Notes: Significant at **** $P < 0.001$; *** $P < 0.01$; ** $P < 0.05$; * $P < 0.10$.
Bartlett's test for equal variances: $\chi^2(5) = 5.21$; Prob > $\chi^2 = 0.391$.

Table 11. ANOVA comparison for perceived impact of stress on health status by ethnicity

Variables	Mean Total	SD	Mean White	Mean Black	Mean Asian	Mean Hispanic	Mean other	F	Sig.
Age	1.7	0.6	1.71	1.62	1.93	1.2	1.62	1.69	0.15

Notes: Significant at **** $P < 0.001$; *** $P < 0.01$; ** $P < 0.05$; * $P < 0.10$.
Bartlett's test for equal variances: $\chi^2(4) = 5.3560$; Prob > $\chi^2 = 0.253$.

Table 12. COVID-19 related perceived need for medical services and perception of changes in the cost of healthcare services

Impact of stress on need for medical services	Frequency	Percent
Yes	66	20.95%
No	156	49.52%
Do not know	95	30.16%
Impact of stress on health-care costs		
Yes	129	40.95%
No	107	33.97%
Do not know	79	25.08%
	315	100.00%

or less.²³ Future studies could collect data through a longitudinal or time-series studies that would provide a depiction of stress throughout the COVID-19 pandemic instead of an examination of stress in a short period of time, as done in this study.

Discussion

Based on the data analysis and finding related to the COVID-19 pandemic and stress, our study suggests that the COVID-19 pandemic leads to increased stress levels among adults of all genders, age groups, and ethnicities. This result is similar to the findings from the studies conducted in different countries among diverse population groups.^{1,9,24-27}

All population groups reported being stressed by the COVID-19 pandemic; however, females were significantly more stressed by the COVID-19 pandemic than males. These finding are consistent with the majority of research about stress as well as research about stress during the COVID-19 pandemic. Different studies proved the impact of family, personal, professional, financial problems on health concerns.²⁸⁻³⁰ Women appeared to be more susceptible to stress caused by social isolation.³¹⁻³³ However, available sex-disaggregated data for COVID-19 pandemic from Italy, China, South Korea, and the United States demonstrates that, although there is an equal number of COVID-19 cases between genders, the death rate among men is higher than among women.^{33,34} In our study, the ANOVA test revealed that females are more impacted by the COVID-19 pandemic stress than males. Additionally, it could encourage health professionals and social workers to develop gender-targeted information and behavioral interventions.

Various studies identify older people as particularly vulnerable to COVID-19. According to statistics from the CDC, 8 of 10

COVID-19 deaths reported in the United States have been in adults 65 years old or older. Adults 65 and older account for 16% of the US population, but for 80% of COVID-19 death in the United States.³⁵ These facts suggest that older people may be worse affected by the psychological distress caused by the COVID-19 pandemic, which is supported by prior research that identifies older age as the factor contributing to the stress level.^{36,37} However, our study demonstrates that, although older adults had a slightly higher stress level than other age groups, the stress level was similarly high among all age groups.

Our data also indicate that there were no significant differences in the levels of stress among different ethnic groups in first 2 months of pandemic, even though in May 2020, during the time of this study, the mortality rate among African-Americans was 73 per 100,000 compare with 36 per 100,000 among Latino persons and 22 per 100,000 among white residents.³⁸

This study's data also revealed that stress associated with the COVID-19 pandemic, manifested itself in the feelings of anxiety, depression, insomnia, headache, and similar conditions. The United States Census Bureau reported that the COVID-19 pandemic had tripled the depression rate among adults in the United States. In some cities that were harder hit by the COVID-19 pandemic, these numbers were higher.^{39,40}

Study participants also were stressed about and anticipated that the COVID-19 pandemic would impact their health in the future. This result falls into the theoretical framework of the correlation between self-reported stress and its negative effect on health status⁴¹⁻⁴³ and is congruent with the recent opinions and findings by Moreno et al.⁴⁴ and Mviena et al.⁴⁵ Respondents connected their health issues with various COVID-19 related problems and restrictions, among them inability to access medical services due to mandated restrictions on the majority of medical procedures, transitions to an online working environment, and social isolation. Many other surveys conducted in the United States and other countries described a similar cause and effect paradigm.^{46,47}

On the 1 hand, this study's participants anticipated an increase in the need for healthcare services, connecting it to postponed diagnoses, delayed care, changes in the lifestyle, and other COVID-19 related stressors. On the other hand, they also anticipated an increase in the cost of health-care services. The perceptions related to the increase in cost of healthcare services could have stemmed from different factors, such as changes in the overall economy, information about expensive hospitalizations and treatment of COVID-19 patients, particularly in the pandemic's early stages, and expectations that delays in diagnoses and in receiving medical services could result in more expensive procedures. Indeed, the American Hospital Association⁴⁸ estimated a March-June financial

impact of the COVID-19 pandemic at \$203 billion in losses. Patients are aware of these economic changes and financial demands and may conclude that these demands may increase the cost of health-care services.

These perceptions—anticipated increases in the need for healthcare services and in the cost of healthcare services—are components of the healthcare-seeking behavior model. Healthcare-seeking behavior has been defined as any action undertaken by individuals who perceive themselves to have a health problem for the purpose of finding an appropriate remedy⁴⁹ According to Oberoi et al.,⁵⁰ healthcare-seeking behavior is preceded by a decision-making process that is further governed by individuals, community norms, and expectations as well as behavior. The healthcare-seeking behavior model includes several decision-making forces that influence the individual's healthcare-seeking behavior: patients (recognition of the need, beliefs, cultural norms, age, gender, and perceived outcomes), as well as providers (behavior's influencer), and resources that enable healthcare-seeking behavior (finances, access).^{51,52}

Conclusions

Health professionals may consider focusing on addressing 2 areas of the COVID-19 pandemic stress perceptions. First, learning about people's COVID-19 perceptions about their health condition can provide ground for detecting and treating various health conditions, addressing fears, and improving access to health services. This knowledge would allow health professionals to develop targeted interventions for different population groups, including strategies that can influence COVID-19 preventative behavioral patterns. Second, studying the anticipated need for health services and the expected cost of healthcare services associated with the COVID-19 pandemic can help understand the utilization patterns, and mitigate financial barriers that could interfere with patients' care-seeking behavior.

References

1. WHO. 2020. *Novel coronavirus (2019-nCoV) situation report – 1*. World Health Organization. <https://apps.who.int/iris/handle/10665/330760>. Accessed June 12, 2021.
2. WHO. 2020. *Novel coronavirus (2019-nCoV) situation report – 5*. World Health Organization. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200125-sitrep-5-2019-ncov.pdf?sfvrsn=429b143d_8. Accessed June 12, 2021.
3. WHO. 2020. *Novel coronavirus (2019-nCoV) situation report – 11*. World Health Organization. <https://apps.who.int/iris/handle/10665/330776>. Accessed June 12, 2021.
4. WHO. 2020. *Novel coronavirus (2019-nCoV) situation report – 51*. World Health Organisation. <https://apps.who.int/iris/handle/10665/331475>. Accessed June 12, 2021.
5. Balwin R, Weder di Mauro B. *Economics in the Time of COVID-19*. London: CEPR Press; 2020.
6. McKibbin WJ, Fernando R. The global macroeconomic impacts of COVID-19: seven scenarios. Brookings. <https://www.brookings.edu/research/the-global-macroeconomic-impacts-of-covid-19-seven-scenarios/>. Published March 5, 2020. Accessed May 2, 2021.
7. Wenzel M, Stanske S, Lieberman MB. Strategic responses to crisis. *Strategic Manag J*. 2020;42(2):016-027. doi: 10.1002/smj.3161
8. Wiedemann PM, Dori W. Be alarmed. Some reflections about the COVID-19 risk communication in Germany. *J Risk Res*. 2020;23(7/8):1036-1046. doi: 10.1080/13669877.2020.1825984
9. Baloran ET. Knowledge, attitudes, anxiety, and coping strategies of students during COVID-19 pandemic. *J Loss Trauma*. 2020;25(8):635-642. doi: 10.1080/15325024.2020.1769300
10. Cohen S, Janicki-Deverts D, Miller GE. Psychological stress and disease. *JAMA*. 2007;298(14):1685-1687. doi: 10.1001/jama.298.14.1685
11. Cohen S, Janicki-Deverts D. Who's stressed? Distributions of psychological stress in the United States in probability samples from 1983, 2006, and 2009. *J Appl Soc Psychol*. 2012;42(6):1320-1334. doi: 10.1111/j.1559-1816.2012.00900.x
12. Vaughan E. Risk perceptions and self-protective behavior. *Health Psychol*. 1993;12:74-85.
13. Betsch C, Wieler LH, Habersaat K. Monitoring behavioural insights related to COVID-19. *Lancet*. 2020;395(10232):1255-1256. doi: 10.1016/S0140-6736(20)30729-7
14. Floyd DL, Prentice-Dunn S, Rogers RW. A meta-analysis of research on protection motivation theory. *J Appl Soc Psychol*. 2000;30(2):407-429. doi: 10.1111/j.1559-1816.2000.tb02323.x
15. Rudisill C. How do we handle new health risks? Risk perception, optimism, and behaviors regarding the H1N1 virus. *J Risk Res*. 2013;16(8):959-980. doi: 10.1080/13669877.2012.761271
16. Prati G, Pietrantonio L, Zani B. A social-cognitive model of pandemic influenza H1N1 risk perception and recommended behaviors in Italy. *Risk Anal*. 2011;31(4):645-656. doi: 10.1111/j.1539-6924.2010.01529.x
17. Yang JZ, Chu H. Who is afraid of the Ebola outbreak? The influence of discrete emotions on risk perception. *J Risk Res*. 2018;21(7):834-853. doi: 10.1080/13669877.2016.1247378
18. Wise T, Zbozinek TD, Michelini G, et al. Changes in risk perception and protective behavior during the first week of the COVID-19 pandemic in the United States. *PsyArXiv*. 2020. doi: 10.31234/osf.io/dz428
19. Ermasova N, Cross AD, Ermasova E. Perceived stress and coping among law enforcement officers: an empirical analysis of patrol versus non-patrol officers in Illinois, USA. *J Police Crim Psychol*. 2020;35(1):48-63. doi: 10.1007/s11896-019-09356-z
20. Ceka E, Ermasova N. Use of employee assistance programs to manage stress in policing: the effects of perceived adequacy of assistance and opportunity for promotion. *Policing (Bradford)*. 2020;44(2):291-313.
21. McCreary DR, Thompson MM. Development of two reliable and valid measures of stressors in policing: the operational and organizational police stress questionnaires. *Int J Stress Manage*. 2006;13:494-518.
22. Tankovska H. Facebook user share in the United States 2021, by gender. Statista. <https://www.statista.com/statistics/266879/facebook-users-in-the-us-by-gender/>. Published February 1, 2021. Accessed May 2, 2021.
23. Gramlich J. 10 facts about Americans and Facebook. Pew Research Center. <https://www.pewresearch.org/fact-tank/2019/05/16/facts-about-americans-and-facebook/>. Published July 31, 2020. Accessed May 2, 2021.
24. Burke RM, Midgley CM, Dratch A, et al. Active monitoring of persons exposed to patients with confirmed COVID-19 - United States, January-February 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(9):245-246. doi: 10.15585/mmwr.mm6909e1
25. Lee J. Mental health effect of school closures during COVID-19. *Lancet Child Adolesc Health*. 2020;4(6):421. doi: 10.1016/S2352-4642(20)30109-7
26. Roy D, Tripathy S, Kar SK, et al. Study of knowledge, attitude, anxiety & perceived mental healthcare need in Indian population during COVID-19 pandemic. *Asian J Psychiatr*. 2020;51:102083. doi: 10.1016/j.ajp.2020.102083
27. Wang C, Pan R, Wan X, et al. A longitudinal study on the mental health of general population during the COVID-19 epidemic in China. *Brain Behav Immun*. 2020;87:40-48. doi: 10.1016/j.bbi.2020.04.028
28. American College Health Association. *American College Health Association National College Health Assessment II: Reference Group Executive Summary Fall 2017*. Hanover: American College Health Association; 2018.
29. Bangasser DA, Eck SR, Telson AM, et al. Sex differences in stress regulation of arousal and cognition. *Physiol Behav*. 2018;187:42-50. doi: 10.1016/j.physbeh.2017.09.025
30. Liu N, Zhang F, Wei C, et al. Prevalence and predictors of PTSS during COVID-19 outbreak in China hardest-hit areas: gender differences matter. *Psychiatry Res*. 2020;287:112921. doi: 10.1016/j.psychres.2020.112921

31. **Senst L, Baimoukhametova D, Sterley T-L, et al.** Sexually dimorphic neuronal responses to social isolation. *eLife*. 2016;5:e18726. doi: [10.7554/eLife.18726](https://doi.org/10.7554/eLife.18726).
32. **van Beusekom I, Bakhshi-Raiez F, van der Schaaf M, et al.** The influence of clinical variables on the risk of developing chronic conditions in ICU survivors. *J Crit Care*. 2020 Feb;55:134-139. doi: [10.1016/j.jcrc.2019.10.014](https://doi.org/10.1016/j.jcrc.2019.10.014)
33. **Spagnolo PA, Manson JE, Joffe H.** Sex and gender differences in health: what the COVID-19 pandemic can teach us. *Ann Intern Med*. 2020; 173(5):385-386. doi: [10.7326/M20-1941](https://doi.org/10.7326/M20-1941)
34. **Sharma G, Volgman AS, Michos ED.** Sex differences in mortality From COVID-19 pandemic: are men vulnerable and women protected? *JACC Case Rep*. 2020;2(9):1407-1410. doi: [10.1016/j.jaccas.2020.04.027](https://doi.org/10.1016/j.jaccas.2020.04.027)
35. **Freed M, Cubanski J, Neuman T, et al.** What share of people who have died of COVID-19 are 65 and older – and how does it vary by state? Kaiser Family Foundation. <https://www.kff.org/coronavirus-COVID-19/issue-brief/what-share-of-people-who-have-died-of-COVID-19-are-65-and-older-and-how-does-it-vary-by-stat>. Published July 24, 2020. Accessed May 2, 2021.
36. **Martin M, Grünendahl M, Martin P.** Age differences in stress, social resources, and well-being in middle and older age. *J Gerontol B Psychol Sci Soc Sci*. 2001;56(4):P214-P222. doi: [10.1093/geronb/56.4.P214](https://doi.org/10.1093/geronb/56.4.P214)
37. **Wilson RS, Begeny CT, Boyle PA, et al.** Vulnerability to stress, anxiety, and development of dementia in old age. *Am J Geriatr Psychiatry*. 2011; 19(4):327-334. doi: [10.1097/JGP.0b013e31820119da](https://doi.org/10.1097/JGP.0b013e31820119da)
38. **Webb Hooper M, Nápoles AM, Pérez-Stable EJ.** COVID-19 and racial/ethnic disparities. *JAMA*. 2020;323(24):2466-2467. doi: [10.1001/jama.2020.8598](https://doi.org/10.1001/jama.2020.8598)
39. **Boyras G, Legros DN.** Coronavirus disease (COVID-19) and traumatic stress: probable risk factors and correlates of posttraumatic stress disorder. *J Loss Trauma*. 2020;25(6/7):503-522. doi: [10.1080/15325024.2020.1763556](https://doi.org/10.1080/15325024.2020.1763556)
40. **Twenge JM, Joiner TE.** US Census bureau-assessed prevalence of anxiety and depressive symptoms in 2019 and during the 2020 covid-19 pandemic. *Depress Anxiety*. 2020;37(10):954-956. doi: [10.1002/da.23077](https://doi.org/10.1002/da.23077)
41. **Watson D.** Intraindividual and interindividual analyses of positive and negative affect: their relation to health complaints, perceived stress, and daily activities. *J Pers Soc Psychol*. 1988;54(6):1020-1030. doi: [10.1037/0022-3514.54.6.1020](https://doi.org/10.1037/0022-3514.54.6.1020)
42. **Watson D, Pennebaker JW.** Health complaints, stress, and distress: exploring the central role of negative affectivity. *Psychol Rev*. 1989;96(2): 234-254. doi: [10.1037/0033-295x.96.2.234](https://doi.org/10.1037/0033-295x.96.2.234)
43. **Selye H.** *Stress in Health and Disease*. Boston: The Butterworth Group; 2013.
44. **Moreno C, Wykes T, Galderisi S, et al.** How mental health care should change as a consequence of the COVID-19 pandemic. *Lancet Psychiatry*. 2020;7(9):813-824. doi: [10.1016/S2215-0366\(20\)30307-2](https://doi.org/10.1016/S2215-0366(20)30307-2)
45. **Mviena JLM, Fanne M, Gondo R, et al.** How mental health care is changing in Cameroon because of the COVID-19 pandemic. *Lancet Psychiatry*. 2020;7(10):e62-e63. doi: [10.1016/S2215-0366\(20\)30390-4](https://doi.org/10.1016/S2215-0366(20)30390-4)
46. **Li J, Yang Z, Qiu H, et al.** Anxiety and depression among general population in China at the peak of the COVID-19 epidemic. *World Psychiatry*. 2020;19(2):249-250. doi: [10.1002/wps.20758](https://doi.org/10.1002/wps.20758)
47. **Fullana MA, Hidalgo-Mazzei D, Vieta E, et al.** Coping behaviors associated with decreased anxiety and depressive symptoms during the COVID-19 pandemic and lockdown. *J Affect Disord*. 2020;275:80-81. doi: [10.1016/j.jad.2020.06.027](https://doi.org/10.1016/j.jad.2020.06.027)
48. **American Health Association.** New AHA report finds financial impact of COVID-19 on hospital; health systems to be over \$200 billion through June: AHA. American Hospital Association. <https://www.aha.org/press-releases/2020-05-05-new-aha-report-finds-financial-impact-covid-19-hospitals-health-systems>. Published May 5, 2020. Accessed May 2, 2021.
49. **Ihaji E, Gerald E, Ogwuche CH.** Educational level, sex and church affiliation on health seeking behaviour among parishioners in Makurdi metropolis of Benue state. *J Educ Policy Entrepren Res*. 2014;1:311-316
50. **Oberoi S, Chaudhary N, Patnaik S, et al.** Understanding health seeking behavior. *J Family Med Prim Care*. 2016;5(2):463-464. doi: [10.4103/2249-4863.192376](https://doi.org/10.4103/2249-4863.192376)
51. **Andersen RM.** Revisiting the behavioral model and access to medical care: does it matter? *J Health Soc Behav*. 1995;36(1):1-10. doi: [10.2307/2137284](https://doi.org/10.2307/2137284)
52. **Lauver D.** A theory of care-seeking behavior. *Image J Nurs Sch*. 1992; 24(4):281-287. doi: [10.1111/j.1547-5069.1992.tb00734.x](https://doi.org/10.1111/j.1547-5069.1992.tb00734.x)
53. **Lee TMC, Chi I, Chung LWM, et al.** Ageing and psychological response during the post-SARS period. *Aging Ment Health*. 2006;10(3):303-311. doi: [10.1080/13607860600638545](https://doi.org/10.1080/13607860600638545)