



POSTER PRESENTATION

Open Access

Molecular genetics, correlation between genotype and phenotype of 65 Vietnamese patients with congenital hyperinsulinism

Dang Anh Duong^{1*}, Vu Chi Dung¹, Nguyen Phu Dat¹, Bui Phuong Thao¹, Can Thi Bich Ngoc¹,
Nguyen Ngoc Khanh¹, Tran Minh Dien¹, Nguyen Thanh Liem¹, Sarah Flanagan², Sian Ellard²

From 8th APPEs Biennial Scientific Meeting
Darwin, Australia. 29 October – 1 November 2014

Hyperinsulinemic hypoglycemia (HH) is a consequence of unregulated insulin secretion by pancreatic β -cells. Congenital HH is caused by mutations in genes involved in regulation of insulin secretion (ABCC8, KCNJ11, GLUD1, CGK, HADH, SLC16A1, HNF4A and UCP2). Severe forms of congenital HH are caused by inactivating mutations in ABCC8 and KCNJ11, which encode the two components of the pancreatic β -cell ATP-sensitive potassium channel. Our aim is to identify mutations in the ABCC8 and KCNJ11, HNF4A and GLUD genes, and to describe genotype and phenotype correlations of Vietnamese children with congenital hyperinsulinism. A prospective study was conducted on 65 cases with congenital hyperinsulinism diagnosed and treated at the National Hospital of Pediatrics from January 2007 to April 2014. Patients were selected by using inclusion criteria of Hussain K (2008). Mutations were identified in 32 cases (49.2%) including mutations of ABCC8 gene (28 cases; 43.1%), KCNJ11 (3 cases; 4.6%), HNF4A (1 case; 1.5%). 100% of cases with homozygous/compound heterozygous recessive mutations or one paternal dominant mutation of ABCC8 gene did not respond to diazoxide treatment and required 95% pancreatectomy. Molecular analysis using pancreas tissue after surgery from cases with one mutation of ABCC8 gene inherited from father confirmed focal lesion type. Other cases without identified mutations usually responded to diazoxide. In conclusions, children with congenital hyperinsulinism should be performed mutation analysis which helps in making diagnosis and treatment decision. Families of children with congenital hyperinsulinism

should be given genetic counseling. Prenatal diagnosis should be performed as well as follow - up and treatment should be given to children with congenital hyperinsulinism immediately after birth.

Authors' details

¹National Hospital of Pediatrics, Hanoi, Vietnam. ²Department of Molecular Genetics, Royal Devon & Exeter NHS Hospital. Peninsula Medical School, UK.

Published: 28 April 2015

doi:10.1186/1687-9856-2015-S1-P125

Cite this article as: Duong et al.: Molecular genetics, correlation between genotype and phenotype of 65 Vietnamese patients with congenital hyperinsulinism. *International Journal of Pediatric Endocrinology* 2015 2015(Suppl 1):P125.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹National Hospital of Pediatrics, Hanoi, Vietnam
Full list of author information is available at the end of the article