CORRECTION

Open Access



Correction: A novel tool for predicting the risk of central lymph node metastasis in patients with papillary thyroid microcarcinoma: a retrospective cohort study

Qian-wen Luo¹, Shan Gao², Xiao Lv³, Si-jia Li², Bo-fang Wang⁴, Qing-qing Han², Yun-peng Wang⁴, Quan-lin Guan^{5*†} and Tao Gong^{2*}

Correction: BMC Cancer 22, 1-12 (2022) https://doi.org/10.1186/s12885-022-09655-5

Following publication of the original article [1], the authors identified an error in the affiliations.

Professor Quan-lin Guan is affiliated to Lanzhou University First Hospital, and Professor Tao Gong is affiliated to HeBei General Hospital. The following is the correct format:

Qian-wen Luo1; Shan Gao2; Xiao Lv3; Si-jia Li2; Bofang Wang4; Qing-qing Han2; Yun-peng Wang4; Quanlin Guan 5*; Tao Gong2*

1 The First Clinical Academy of Lanzhou University, Lanzhou , Gansu 730000, China

2 Department of Gland Surgery, HeBei General Hospital, Shijiazhuang, HeBei 050051, China

3 Department of Cardiovascular Medicine, People's Hospital of Dongxihu District, Wuhan, Hubei 430040, China

The original article can be found online at https://doi.org/10.1186/s12885-022-09655-5.

[†]Quan-lin Guan is the first corresponding author.

*Correspondence: guanquanlin@163.com; gongtao0520@163.com

² Department of Gland Surgery, HeBei General Hospital, Shijiazhuang 050051, HeBei, China

⁵ Department of surgical oncology, The Lanzhou University First Hospital, Lanzhou 730030, Gansu, China

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/A.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/licenses/by/A.0/. The Creative Commons Public Domain Dedicated in a credit line to the data.

4 The Second Clinical Academy of Lanzhou University, Lanzhou 730000, Gansu, China

5 Department of surgical oncology, The Lanzhou University First Hospital, Lanzhou, Gansu 730030, China

Author details

¹The First Clinical Academy of Lanzhou University, Lanzhou 730000, Gansu, China. ²Department of Gland Surgery, HeBei General Hospital, Shijiazhuang 050051, HeBei, China. ³Department of Cardiovascular Medicine, People's Hospital of Dongxihu District, Wuhan 430040, Hubei, China. ⁴The Second Clinical Academy of Lanzhou University, Lanzhou 730000, Gansu, China. ⁵Department of surgical oncology, The Lanzhou University First Hospital, Lanzhou 730030, Gansu, China.

Published online: 24 June 2022

Reference

 Luo Q, Gao S, Lv X, et al. A novel tool for predicting the risk of central lymph node metastasis in patients with papillary thyroid microcarcinoma: a retrospective cohort study. BMC Cancer. 2022;22:606. https://doi. org/10.1186/s12885-022-09655-5.